

This is the accepted manuscript made available via CHORUS. The article has been published as:

Evidence of a New Narrow Resonance Decaying to $\chi_{c1}\gamma$ in $B \rightarrow \chi_{c1}\gamma K$

V. Bhardwaj *et al.* (Belle Collaboration)

Phys. Rev. Lett. **111**, 032001 — Published 16 July 2013

DOI: [10.1103/PhysRevLett.111.032001](https://doi.org/10.1103/PhysRevLett.111.032001)

Evidence for a new narrow resonance decaying to $\chi_{c1}\gamma$ in $B \rightarrow \chi_{c1}\gamma K$

V. Bhardwaj,³⁸ K. Miyabayashi,³⁸ I. Adachi,¹³ H. Aihara,⁶¹ D. M. Asner,⁴⁶ V. Aulchenko,³ T. Aushev,²¹ T. Aziz,⁵⁶ A. M. Bakich,⁵⁵ A. Bala,⁴⁷ B. Bhuyan,¹⁵ M. Bischofberger,³⁸ A. Bondar,³ G. Bonvicini,⁶⁶ A. Bozek,⁴² M. Bračko,^{31,22} J. Brodzicka,⁴² T. E. Browder,¹² V. Chekelian,³² A. Chen,³⁹ B. G. Cheon,¹¹ K. Chilikin,²¹ R. Chistov,²¹ K. Cho,²⁵ V. Chobanova,³² S.-K. Choi,¹⁰ Y. Choi,⁵⁴ D. Cinabro,⁶⁶ J. Dalseno,^{32,57} M. Danilov,^{21,34} Z. Doležal,⁴ Z. Drásal,⁴ A. Drutskoy,^{21,34} D. Dutta,¹⁵ K. Dutta,¹⁵ S. Eidelman,³ D. Epifanov,³ H. Farhat,⁶⁶ J. E. Fast,⁴⁶ T. Ferber,⁶ A. Frey,⁹ V. Gaur,⁵⁶ N. Gabyshev,³ S. Ganguly,⁶⁶ R. Gillard,⁶⁶ Y. M. Goh,¹¹ B. Golob,^{29,22} J. Haba,¹³ T. Hara,¹³ H. Hayashii,³⁸ Y. Horii,³⁷ Y. Hoshi,⁵⁹ W.-S. Hou,⁴¹ Y. B. Hsiung,⁴¹ H. J. Hyun,²⁷ T. Iijima,^{37,36} K. Inami,³⁶ A. Ishikawa,⁶⁰ R. Itoh,¹³ T. Iwashita,³⁸ T. Julius,³³ D. H. Kah,²⁷ J. H. Kang,⁶⁸ E. Kato,⁶⁰ T. Kawasaki,⁴⁴ H. Kichimi,¹³ C. Kiesling,³² D. Y. Kim,⁵³ J. B. Kim,²⁶ J. H. Kim,²⁵ K. T. Kim,²⁶ M. J. Kim,²⁷ Y. J. Kim,²⁵ K. Kinoshita,⁵ J. Klucar,²² B. R. Ko,²⁶ P. Kodyš,⁴ S. Korpar,^{31,22} P. Krizán,^{29,22} P. Krokovny,³ R. Kumar,⁴⁹ T. Kumita,⁶³ A. Kuzmin,³ Y.-J. Kwon,⁶⁸ J. S. Lange,⁷ S.-H. Lee,²⁶ J. Li,⁵² Y. Li,⁶⁵ C. Liu,⁵¹ Z. Q. Liu,¹⁷ D. Liventsev,¹³ P. Lukin,³ D. Matvienko,³ H. Miyata,⁴⁴ R. Mizuk,^{21,34} G. B. Mohanty,⁵⁶ A. Moll,^{32,57} R. Mussa,²⁰ E. Nakano,⁴⁵ M. Nakao,¹³ Z. Natkaniec,⁴² M. Nayak,¹⁶ E. Nedelkovska,³² N. K. Nisar,⁵⁶ S. Nishida,¹³ O. Nitoh,⁶⁴ S. Ogawa,⁵⁸ S. Okuno,²³ S. L. Olsen,⁵² P. Pakhlov,^{21,34} G. Pakhlova,²¹ E. Panzenböck,^{9,38} H. Park,²⁷ H. K. Park,²⁷ T. K. Pedlar,³⁰ R. Pestotnik,²² M. Petrič,²² L. E. Pilonen,⁶⁵ M. Ritter,³² M. Röhrken,²⁴ A. Rostomyan,⁶ H. Sahoo,¹² T. Saito,⁶⁰ K. Sakai,¹³ Y. Sakai,¹³ S. Sandilya,⁵⁶ D. Santel,⁵ L. Santelj,²² T. Sanuki,⁶⁰ Y. Sato,⁶⁰ V. Savinov,⁴⁸ O. Schneider,²⁸ G. Schnell,^{1,14} C. Schwanda,¹⁸ R. Seidl,⁵⁰ D. Semmler,⁷ K. Senyo,⁶⁷ O. Seon,³⁶ M. E. Sevier,³³ M. Shapkin,¹⁹ C. P. Shen,³⁶ T.-A. Shibata,⁶² J.-G. Shiu,⁴¹ B. Shwartz,³ F. Simon,^{32,57} J. B. Singh,⁴⁷ P. Smerkol,²² Y.-S. Sohn,⁶⁸ A. Sokolov,¹⁹ E. Solovieva,²¹ M. Starič,²² M. Steder,⁶ M. Sumihama,⁸ T. Sumiyoshi,⁶³ U. Tamponi,²⁰ K. Tanida,⁵² G. Tatishvili,⁴⁶ Y. Teramoto,⁴⁵ K. Trabelsi,¹³ T. Tsuboyama,¹³ M. Uchida,⁶² S. Uehara,¹³ T. Uglov,^{21,35} Y. Unno,¹¹ P. Urquijo,² Y. Usov,³ S. E. Vahsen,¹² C. Van Hulse,¹ P. Vanhoefer,³² G. Varner,¹² K. E. Varvell,⁵⁵ A. Vinokurova,³ M. N. Wagner,⁷ C. H. Wang,⁴⁰ M.-Z. Wang,⁴¹ P. Wang,¹⁷ M. Watanabe,⁴⁴ Y. Watanabe,²³ E. Won,²⁶ B. D. Yabsley,⁵⁵ J. Yamaoka,¹² Y. Yamashita,⁴³ S. Yashchenko,⁶ Y. Yook,⁶⁸ C. Z. Yuan,¹⁷ C. C. Zhang,¹⁷ Z. P. Zhang,⁵¹ V. Zhilich,³ V. Zhulanov,³ and A. Zupanc²⁴

(The Belle Collaboration)

¹University of the Basque Country UPV/EHU, 48080 Bilbao

²University of Bonn, 53115 Bonn

³Budker Institute of Nuclear Physics SB RAS and Novosibirsk State University, Novosibirsk 630090

⁴Faculty of Mathematics and Physics, Charles University, 121 16 Prague

⁵University of Cincinnati, Cincinnati, Ohio 45221

⁶Deutsches Elektronen-Synchrotron, 22607 Hamburg

⁷Justus-Liebig-Universität Gießen, 35392 Gießen

⁸Gifu University, Gifu 501-1193

⁹II. Physikalisches Institut, Georg-August-Universität Göttingen, 37073 Göttingen

¹⁰Gyeongsang National University, Chinju 660-701

¹¹Hanyang University, Seoul 133-791

¹²University of Hawaii, Honolulu, Hawaii 96822

¹³High Energy Accelerator Research Organization (KEK), Tsukuba 305-0801

¹⁴Ikerbasque, 48011 Bilbao

¹⁵Indian Institute of Technology Guwahati, Assam 781039

¹⁶Indian Institute of Technology Madras, Chennai 600036

¹⁷Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049

¹⁸Institute of High Energy Physics, Vienna 1050

¹⁹Institute for High Energy Physics, Protvino 142281

²⁰INFN - Sezione di Torino, 10125 Torino

²¹Institute for Theoretical and Experimental Physics, Moscow 117218

²²J. Stefan Institute, 1000 Ljubljana

²³Kanagawa University, Yokohama 221-8686

²⁴Institut für Experimentelle Kernphysik, Karlsruher Institut für Technologie, 76131 Karlsruhe

²⁵Korea Institute of Science and Technology Information, Daejeon 305-806

²⁶Korea University, Seoul 136-713

²⁷Kyungpook National University, Daegu 702-701

- ²⁸École Polytechnique Fédérale de Lausanne (EPFL), Lausanne 1015
²⁹Faculty of Mathematics and Physics, University of Ljubljana, 1000 Ljubljana
³⁰Luther College, Decorah, Iowa 52101
³¹University of Maribor, 2000 Maribor
³²Max-Planck-Institut für Physik, 80805 München
³³School of Physics, University of Melbourne, Victoria 3010
³⁴Moscow Physical Engineering Institute, Moscow 115409
³⁵Moscow Institute of Physics and Technology, Moscow Region 141700
³⁶Graduate School of Science, Nagoya University, Nagoya 464-8602
³⁷Kobayashi-Maskawa Institute, Nagoya University, Nagoya 464-8602
³⁸Nara Women's University, Nara 630-8506
³⁹National Central University, Chung-li 32054
⁴⁰National United University, Miao Li 36003
⁴¹Department of Physics, National Taiwan University, Taipei 10617
⁴²H. Niewodniczanski Institute of Nuclear Physics, Krakow 31-342
⁴³Nippon Dental University, Niigata 951-8580
⁴⁴Niigata University, Niigata 950-2181
⁴⁵Osaka City University, Osaka 558-8585
⁴⁶Pacific Northwest National Laboratory, Richland, Washington 99352
⁴⁷Panjab University, Chandigarh 160014
⁴⁸University of Pittsburgh, Pittsburgh, Pennsylvania 15260
⁴⁹Punjab Agricultural University, Ludhiana 141004
⁵⁰RIKEN BNL Research Center, Upton, New York 11973
⁵¹University of Science and Technology of China, Hefei 230026
⁵²Seoul National University, Seoul 151-742
⁵³Soongsil University, Seoul 156-743
⁵⁴Sungkyunkwan University, Suwon 440-746
⁵⁵School of Physics, University of Sydney, NSW 2006
⁵⁶Tata Institute of Fundamental Research, Mumbai 400005
⁵⁷Excellence Cluster Universe, Technische Universität München, 85748 Garching
⁵⁸Toho University, Funabashi 274-8510
⁵⁹Tohoku Gakuin University, Tagajo 985-8537
⁶⁰Tohoku University, Sendai 980-8578
⁶¹Department of Physics, University of Tokyo, Tokyo 113-0033
⁶²Tokyo Institute of Technology, Tokyo 152-8550
⁶³Tokyo Metropolitan University, Tokyo 192-0397
⁶⁴Tokyo University of Agriculture and Technology, Tokyo 184-8588
⁶⁵CNP, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061
⁶⁶Wayne State University, Detroit, Michigan 48202
⁶⁷Yamagata University, Yamagata 990-8560
⁶⁸Yonsei University, Seoul 120-749

We report measurements of $B \rightarrow \chi_{c1}\gamma K$ and $\chi_{c2}\gamma K$ decays using 772×10^6 $B\bar{B}$ events collected at the $\Upsilon(4S)$ resonance with the Belle detector at the KEKB asymmetric-energy e^+e^- collider. Evidence of a new resonance in the $\chi_{c1}\gamma$ final state is found with a statistical significance of 3.8σ . This state has a mass of $3823.1 \pm 1.8(\text{stat}) \pm 0.7(\text{syst})$ MeV/ c^2 , a value that is consistent with theoretical expectations for the previously unseen 1^3D_2 $c\bar{c}$ meson. We find no other narrow resonance and set upper limits on the branching fractions of the $X(3872) \rightarrow \chi_{c1}\gamma$ and $\chi_{c2}\gamma$ decays.

PACS numbers: 13.25.Hw, 13.20.Gd, 14.40.Pq

During the last decade, a number of new charmonium ($c\bar{c}$)-like states were observed, many of which are candidates for exotic states [1]. The first of these, the $X(3872)$, has been observed by six different experiments in the same final state [2–7]. A recent update from Belle [8] and LHCb [6] results in a world average mass at 3871.68 ± 0.17 MeV/ c^2 [9] and a stringent upper bound on its width ($\Gamma < 1.2$ MeV) [8]. The proximity of its mass to the $D^{*0}\bar{D}^0$ threshold makes it a good candidate for a $D\bar{D}^*$ molecule [10]. Other alternative models have been proposed, such as a tetraquark [11] or a $c\bar{c}g$ hybrid me-

son [12].

Radiative decays can illuminate clearly the nature of hadrons. For example, the observation of $X(3872) \rightarrow J/\psi\gamma$ confirmed the C -even parity assignment for the $X(3872)$ [13, 14]. The $X(3872) \rightarrow \chi_{c1}\gamma$ and $\chi_{c2}\gamma$ decays are forbidden by C -parity conservation in electromagnetic processes. However, if the $X(3872)$ is a tetraquark or a molecular state, it may have a C -odd partner, which could decay into $\chi_{c1}\gamma$ and $\chi_{c2}\gamma$ final states [15, 16].

In the charmonium family, the observation of a D -wave $c\bar{c}$ meson and its decay modes would test phenomenolog-

ical models [17, 18]. The as-yet undiscovered 1^3D_2 $c\bar{c}$ (ψ_2) and 1^3D_3 $c\bar{c}$ (ψ_3) states are expected to have significant branching fractions to $\chi_{c1}\gamma$ and $\chi_{c2}\gamma$, respectively [19, 20]. D -wave $c\bar{c}$ states and their properties were predicted long ago but remain unconfirmed [19, 20]. The E705 experiment reported an indication of a 1^3D_2 state in $\pi^\pm N \rightarrow J/\psi \pi^+ \pi^- + \text{anything}$, with a mass of 3836 ± 13 MeV/ c^2 [21]; however, the statistical significance of this result was below the threshold for evidence.

In this letter, we report measurements of $B \rightarrow \chi_{c1}\gamma K$ and $B \rightarrow \chi_{c2}\gamma K$ decays, where the χ_{c1} and χ_{c2} decay to $J/\psi\gamma$ [22]. These results are obtained from a data sample of 772×10^6 $B\bar{B}$ events collected with the Belle detector [23] at the KEKB asymmetric-energy e^+e^- collider operating at the $\Upsilon(4S)$ resonance [24].

The J/ψ meson is reconstructed via its decays to $\ell^+\ell^-$ ($\ell = e$ or μ). To reduce the radiative tail in the e^+e^- mode, the four-momenta of all photons within 50 mrad with respect to the original direction of the e^+ or e^- tracks are included in the invariant mass calculation, hereinafter denoted as $M_{e^+e^-(\gamma)}$. The reconstructed invariant mass of the J/ψ candidates is required to satisfy 2.95 GeV/ $c^2 < M_{e^+e^-(\gamma)} < 3.13$ GeV/ c^2 or 3.03 GeV/ $c^2 < M_{\mu^+\mu^-} < 3.13$ GeV/ c^2 . For the selected J/ψ candidates, a vertex-constrained fit is applied and then a mass-constrained fit is performed in order to improve the momentum resolution. The χ_{c1} and χ_{c2} candidates are reconstructed by combining J/ψ candidates with a photon having energy (E_γ) larger than 200 MeV in the laboratory frame. Photons are reconstructed from energy depositions in the electromagnetic calorimeter (ECL), which do not match any extrapolated charged track. To reduce the background from $\pi^0 \rightarrow \gamma\gamma$, we use a likelihood function that distinguishes an isolated photon from π^0 decays using the photon pair invariant mass, photon laboratory energy and polar angle [25]. We reject both γ 's in the pair if the π^0 likelihood probability is larger than 0.7. The reconstructed invariant mass of the χ_{c1} (χ_{c2}) is required to satisfy 3.467 GeV/ $c^2 < M_{J/\psi\gamma} < 3.535$ GeV/ c^2 (3.535 GeV/ $c^2 < M_{J/\psi\gamma} < 3.611$ GeV/ c^2). A mass-constrained fit is applied to the selected χ_{c1} and χ_{c2} candidates.

Charged kaons are identified by combining information from the central drift chamber, time-of-flight scintillation counters, and the aerogel Cherenkov counter systems. The kaon identification efficiency is 89% while the probability of misidentifying a pion as a kaon is 10%. K_S^0 mesons are reconstructed by combining two oppositely charged pions with the invariant mass $M_{\pi^+\pi^-}$ lying between 482 MeV/ c^2 and 514 MeV/ c^2 . The selected candidates are required to satisfy the quality criteria described in Ref. [26].

To reconstruct B candidates, each χ_{cJ} [27] is combined with a kaon candidate and a photon having $E_\gamma > 100$ MeV (and not used in the reconstruction of χ_{cJ}). If the invariant mass of any photon pair that includes

this photon is found to be consistent with a π^0 (i.e., 117 MeV/ $c^2 < M_{\gamma\gamma} < 153$ MeV/ c^2), this photon is rejected. Among the events containing at least one χ_{cJ} candidate, 9.0% have multiple χ_{cJ} candidates. In such cases, the γ forming the χ_{cJ} candidate with mass closest to the χ_{c1} or χ_{c2} masses [9] is not used as the additional photon. This treatment suppresses reflections from the χ_{cJ} daughter photons.

The B candidate is identified by two kinematic variables: the beam-constrained mass ($M_{bc} \equiv \sqrt{E_{\text{beam}}^{*2} - p_B^{*2}}$) and the energy difference ($\Delta E \equiv E_B^* - E_{\text{beam}}^*$). Here, E_{beam}^* is the run-dependent beam energy, and E_B^* and p_B^* are the reconstructed energy and momentum, respectively, of the B meson candidates in the center-of-mass (CM) frame. Candidates within a ΔE window of $[-28, 30]$ MeV and with $M_{bc} > 5.23$ GeV/ c^2 are selected. Of these, 9.8% (6.4%) have multiple candidates in the $B^\pm \rightarrow \chi_{c1}\gamma K^\pm$ ($B^\pm \rightarrow \chi_{c2}\gamma K^\pm$) mode; we select the B candidate with ΔE closest to zero. In order to improve the resolution in $M_{\chi_{cJ}\gamma}$, we scale the energy of the γ so that ΔE is equal to zero. This corrects for incomplete energy measurement in the ECL. To suppress continuum background, events having a ratio R_2 of the second to zeroth Fox-Wolfram moments [28] above 0.5 are rejected.

The $M_{\chi_{c1}\gamma}$ and M_{bc} projections for the $B^\pm \rightarrow \chi_{c1}\gamma K^\pm$ signal candidates are shown in Fig. 1, where a $\psi' \rightarrow \chi_{c1}\gamma$ signal is evident. In addition, there is a significant narrow peak at 3823 MeV/ c^2 , denoted hereinafter as $X(3823)$. No signal of $X(3872) \rightarrow \chi_{c1}\gamma$ is seen. We extract the signal yield from a two-dimensional unbinned extended maximum likelihood (2D UML) fit to the variables $M_{\chi_{cJ}\gamma}$ and M_{bc} .

The resolution in $M_{\chi_{cJ}\gamma}$ (M_{bc}) is parameterized by a sum of two Gaussians (Gaussian and logarithmic Gaussian [29]). MC studies show that the resolutions in both $M_{\chi_{cJ}\gamma}$ and M_{bc} for a narrow resonance in the mass range 3.8 GeV/ $c^2 < M_{\chi_{cJ}\gamma} < 4.0$ GeV/ c^2 are in good agreement with those for ψ' . The parameters of the resolution functions are determined from the MC simulation that is calibrated using the $B^\pm \rightarrow \psi'(\rightarrow \chi_{c1}\gamma)K^\pm$ signal. We take into account the ψ' natural width [9] by convolving the Breit-Wigner function and the resolution function; for the $X(3823)$ and $X(3872)$, zero natural widths are assumed. The two-dimensional probability density function (PDF) is a product of the one-dimensional distributions.

For $B^\pm \rightarrow \psi'(\rightarrow \chi_{c1}\gamma)K^\pm$ decays, the mean and width of the core Gaussian are floated and the remaining parameters are fixed according to MC. To fit the $B^\pm \rightarrow X(3823)(\rightarrow \chi_{c1}\gamma)K^\pm$ signal, we float the mean of the core Gaussian but constrain the detector resolution by using the ψ' signal results after taking into account the difference estimated from the signal MC study. For M_{bc} , the parameters are fixed to those found for the ψ' , in accordance with expectations based on the

MC simulation. For $X(3872)$, we fix the mass difference and the mass resolution change with respect to ψ' using the information from PDG [9] and MC studies. To fit $B^0 \rightarrow \psi'(\rightarrow \chi_{c1}\gamma)K^0$, $B^0 \rightarrow X(3823)(\rightarrow \chi_{c1}\gamma)K^0$, $B \rightarrow \psi'(\rightarrow \chi_{c2}\gamma)K$, $B \rightarrow X(3823)(\rightarrow \chi_{c2}\gamma)K$ and $B \rightarrow X(3872)(\rightarrow \chi_{c1}\gamma)K$, we fix all the parameters obtained from the signal MC study after correcting the PDF shapes by applying MC/data calibration factors.

To study background with a real J/ψ , we use large MC simulated $B \rightarrow J/\psi X$ samples corresponding to 100 times the integrated luminosity of the data. The non- J/ψ (non- χ_{cJ}) background is studied using $M_{\ell\ell}$ ($M_{J/\psi\gamma}$) sidebands in data. In $B \rightarrow (\chi_{cJ}\gamma)K$, the background with a broad peaking structure is mostly due to the $B \rightarrow \psi'(\rightarrow \chi_{cJ}\gamma)K$, $B \rightarrow \chi_{cJ}K^*$, $B \rightarrow J/\psi K^*$ and $B \rightarrow \psi'K^*$ decay modes. $B \rightarrow \psi'(\rightarrow \chi_{cJ}\gamma)K$ produces peaks in both distributions ($M_{\chi_{cJ}\gamma}$ and M_{bc}), while the other backgrounds are flat in $M_{\chi_{cJ}\gamma}$ but peaked in M_{bc} . We determine the PDFs from the large MC sample. The fractions of the PDF components are floated in the fit, except for $B \rightarrow \psi'(\rightarrow \chi_{cJ}\gamma)K$, whose fraction is controlled by fixing its ratio to the $B \rightarrow \psi'(\rightarrow \chi_{cJ}\gamma)K$ signal yield. For the combinatorial background, a threshold function, $(M_{\chi_{cJ}\gamma})^2 \times \exp(a(M_{\chi_{cJ}\gamma} - M_{th}) + b(M_{\chi_{cJ}\gamma} - M_{th})^2 + c(M_{\chi_{cJ}\gamma} - M_{th})^3)$, where $M_{th} = 3.543 \text{ GeV}/c^2$ ($3.585 \text{ GeV}/c^2$) for $M_{\chi_{c1}\gamma}$ ($M_{\chi_{c2}\gamma}$), is used for $M_{\chi_{cJ}\gamma}$ and an ARGUS function [30] is used for M_{bc} . The value of M_{th} is estimated from a MC study; its variation, which affects the signal yield in the fits, is incorporated in the systematic errors. The ΔE data sidebands are used to verify the background PDFs. The fractions for the signal and the background components are floated in the fit.

The results of the fits are presented in Figs. 1-3 and in Table I. The significance is estimated using the value of $-2\ln(\mathcal{L}_0/\mathcal{L}_{\max})$ where \mathcal{L}_{\max} (\mathcal{L}_0) denotes the likelihood value when the yield is allowed to vary (is set to zero). In the likelihood calculation, the χ^2 statistic uses the appropriate number of degrees of freedom (two in the case of $B^\pm \rightarrow X(3823)K^\pm$ and one for the other decay modes). The systematic uncertainty, which is described below, is included in the significance calculation [32]. We find a significant ψ' signal in all considered channels. We also obtain evidence for the $X(3823)$ in the channel $B^\pm \rightarrow \chi_{c1}\gamma K^\pm$ with a statistical significance of 3.8 standard deviations (σ). The $X(3872)$ signals are insignificant. We estimate the branching fractions according to the formula $\mathcal{B} = Y/(\epsilon \mathcal{B}_s N_{B\bar{B}})$; here, Y is the yield, ϵ is the reconstruction efficiency, \mathcal{B}_s is the secondary branching fraction taken from Ref. [9] and $N_{B\bar{B}}$ is the number of $B\bar{B}$ mesons in the data sample. Equal production of neutral and charged B meson pairs in the $\Upsilon(4S)$ decay is assumed. Measured branching fractions for the ψ' are in agreement with the world average values for all the channels [9]. We set 90% confidence level (C.L.) upper limits (U.L.) on the insignificant channels using frequentist methods based on an ensemble of pseudo-experiments.




FIG. 1: 2D UML fit projection for $B^\pm \rightarrow (\chi_{c1}\gamma)K^\pm$ decays: (a) $M_{\chi_{c1}\gamma}$ distribution for $M_{bc} > 5.27 \text{ GeV}/c^2$, (b) M_{bc} distribution for $3.660 \text{ GeV}/c^2 < M_{\chi_{c1}\gamma} < 3.708 \text{ GeV}/c^2$ (ψ' region, shown by the red dotted-dashed arrows in a), (c) M_{bc} distribution for $3.805 \text{ GeV}/c^2 < M_{\chi_{c1}\gamma} < 3.845 \text{ GeV}/c^2$ ($X(3823)$ region, shown by the magenta solid arrows in a) and (d) M_{bc} distribution for $3.84 \text{ GeV}/c^2 < M_{\chi_{c1}\gamma} < 3.89 \text{ GeV}/c^2$ ($X(3872)$ region). The curves used in the fits are described in [31].




FIG. 2: 2D UML fit projection for $B^\pm \rightarrow (\chi_{c2}\gamma)K^\pm$ decays: (a) $M_{\chi_{c2}\gamma}$ distribution for $M_{bc} > 5.27 \text{ GeV}/c^2$, (b) M_{bc} distribution for the ψ' region, (c) M_{bc} distribution for the $X(3823)$ region and (d) M_{bc} distribution for the $X(3872)$ region. The curves used in the fits are described in [31].

A correction for small differences in the signal detection efficiency between MC and data has been applied for the lepton and kaon identification requirements. Uncertainties in these corrections are included in the sys-




FIG. 3: 2D UML fit projection for $B^0 \rightarrow (\chi_{c1}\gamma)K_S^0$ decays: (a) $M_{\chi_{c1}\gamma}$ distribution for $M_{bc} > 5.27$ GeV/c^2 , (b) $M_{\chi_{c2}\gamma}$ distribution for $M_{bc} > 5.27$ GeV/c^2 , (c) and (d) M_{bc} distribution for the ψ' region, (e) and (f) M_{bc} distribution for the $X(3823)$ region, and (g) and (h) M_{bc} distribution for the $X(3872)$ region. The curves used in the fits are described in [31].

tematic error. The $e^+e^- \rightarrow e^+e^-\ell^+\ell^-$ ($\ell = e$ or μ) and $D^{*+} \rightarrow D^0(K^-\pi^+)\pi^+$ samples are used to estimate the lepton identification correction and the kaon (pion) identification correction, respectively. To estimate the correction and residual systematic uncertainty for K_S^0 reconstruction, $D^{*+} \rightarrow D^0(\rightarrow K_S^0\pi^+\pi^-)\pi^+$ samples are used. The errors on the PDF shapes are obtained by varying all fixed parameters by $\pm 1\sigma$ and taking the change in the yield as the systematic uncertainty. The uncertainties due to the secondary branching fractions are also taken into account. The uncertainties of the tracking efficiency and $N_{B\bar{B}}$ are estimated to be 0.35% per track and 1.4%, respectively. The uncertainty on the photon identification is estimated to be 2.0%/photon. The sys-




FIG. 4: 2D UML fit projection of $M_{\chi_{c1}\gamma}$ distribution for the simultaneous fit of $B^\pm \rightarrow (\chi_{c1}\gamma)K^\pm$ and $B^0 \rightarrow (\chi_{c1}\gamma)K_S^0$ decays for $M_{bc} > 5.27$ GeV/c^2 . The curves used in the fits are described in [31].

TABLE I: Summary of the results. Signal yield (Y) from the fit, significance (\mathcal{S}) with systematics included, corrected efficiency (ϵ) and measured \mathcal{B} . For \mathcal{B} , the first (second) error is statistical (systematic). In the neutral B decay, the efficiency below includes the $K_S^0 \rightarrow \pi^+\pi^-$ branching fraction but does not include the factor of two for $K^0 \rightarrow K_S^0$ or K_L^0 .

Decay	Yield (Y)	$\mathcal{S}(\sigma)$	$\epsilon(\%)$	Branching fraction
$B^\pm \rightarrow \psi'(\rightarrow \chi_{c1}\gamma)K^\pm$				$\mathcal{B}(10^{-4})$
χ_{c1}	193.2 ± 19.2	14.8	8.6	$7.7 \pm 0.8 \pm 0.9$
χ_{c2}	59.1 ± 8.4	7.8	6.0	$6.3 \pm 0.9 \pm 0.6$
$B^0 \rightarrow \psi'(\rightarrow \chi_{c1}\gamma)K^0$				
χ_{c1}	50.3 ± 7.3	7.2	5.1	$6.8 \pm 1.0 \pm 0.7$
χ_{c2}	12.9 ± 4.4	2.9	3.5	$4.7 \pm 1.6 \pm 0.8$
$B^\pm \rightarrow X(3823)(\rightarrow \chi_{c1}\gamma)K^\pm$				$\mathcal{B}(10^{-6})$
χ_{c1}	33.2 ± 9.7	3.8	10.9	$9.7 \pm 2.8 \pm 1.1$
χ_{c2}	0.3 ± 3.9	0.1	8.8	< 3.6
$B^0 \rightarrow X(3823)(\rightarrow \chi_{c1}\gamma)K^0$				
χ_{c1}	3.9 ± 3.4	1.2	6.0	< 9.9
χ_{c2}	5.3 ± 2.9	2.4	5.0	< 22.8
$B^\pm \rightarrow X(3872)(\rightarrow \chi_{c1}\gamma)K^\pm$				
χ_{c1}	-0.9 ± 5.1		11.1	< 1.9
χ_{c2}	4.7 ± 4.4	1.3	9.3	< 6.7
$B^0 \rightarrow X(3872)(\rightarrow \chi_{c1}\gamma)K^0$				
χ_{c1}	4.6 ± 3.0	1.6	6.2	< 9.6
χ_{c2}	2.3 ± 2.2	1.1	5.2	< 12.2

tematic uncertainty associated with the difference of the π^0 veto between data and MC is estimated to be 1.2% from a study of the $B^\pm \rightarrow \chi_{c1}(\rightarrow J/\psi\gamma)K^\pm$ sample.

To improve the mass determination of the $X(3823)$, a simultaneous fit to $B^\pm \rightarrow (\chi_{c1}\gamma)K^\pm$ and $B^0 \rightarrow (\chi_{c1}\gamma)K_S^0$ is performed, assuming that $\mathcal{B}(B^\pm \rightarrow X(3823)K^\pm)/\mathcal{B}(B^0 \rightarrow X(3823)K^0) = \mathcal{B}(B^\pm \rightarrow \psi'K^\pm)/\mathcal{B}(B^0 \rightarrow \psi'K^0)$. The ψ' peak position and $M_{\chi_{c1}\gamma}$ resolution are common for both charged and neutral B candidates. From this fit, we estimate the significance for $X(3823)$ to be 4.0σ (including systematic uncertainties).

We determine the mass of the signal peak relative to the well-measured ψ' mass :

$$M_{X(3823)} = M_{X(3823)}^{\text{meas}} - M_{\psi'}^{\text{meas}} + M_{\psi'}^{\text{PDG}} \\ = 3823.1 \pm 1.8 \pm 0.7 \text{ MeV}.$$

Here, the first uncertainty is statistical and the second is systematic. Because of the mass-constrained fit to the χ_{c1} candidate, the systematic uncertainty of $M_{X(3823)}$ is dominated by the additional photon's energy scale. This photon energy scale uncertainty is estimated by the difference between the $\chi_{c1} \rightarrow J/\psi\gamma$ candidates' mass without any constraint and the χ_{c1} nominal mass [9], which results in 0.7 MeV as the $M_{X(3823)}$ systematic error. In order to estimate the width, we float this parameter and find no sensitivity with the available statistics: the width is 1.7 ± 5.5 MeV. Using pseudo-experiments generated with different width hypotheses for the $X(3823)$, the U.L. at 90% C.L. on its width is estimated to be 24 MeV.

The mass of the $X(3823)$ is near potential model expectations for the centroid of the 1^3D_J states: the Cornell [17] and the Buchmüller-Tye [18] potentials give 3810 MeV/ c^2 . Other models predict the mass of ψ_2 (the 1^3D_2 $c\bar{c}$ state, having $J^{PC} = 2^{--}$) to be 3815-3840 MeV/ c^2 [33–36]. The $X(3823)$ mass agrees quite well with these models. In addition, since no peak has been seen around $X(3823)$ in the $D\bar{D}$ final state [37], one expects that ψ_2 does not decay to $D\bar{D}$ [35]. The ratio $R_B = \frac{\mathcal{B}(X(3823) \rightarrow \chi_{c2}\gamma)}{\mathcal{B}(X(3823) \rightarrow \chi_{c1}\gamma)} < 0.41$ (at 90% C.L.) is consistent with the expectation ($R_B \sim 0.2$) for ψ_2 [34, 38, 39]. The limited statistics preclude an angular analysis to determine the J^{PC} of the $X(3823)$. The product of branching fractions for the $X(3823)$ is approximately two orders of magnitude lower than for the ψ' , as shown in Table I; it is consistent with the interpretation of the $X(3823)$ as $\psi_2(1^3D_2)$, whose production rate is suppressed by the factorization [40] in the two-body B meson decays.

In summary, we obtain the first evidence of a narrow state, $X(3823)$, that decays to $\chi_{c1}\gamma$ with a mass of $3823.1 \pm 1.8(\text{stat}) \pm 0.7(\text{syst})$ MeV/ c^2 and a significance of 3.8σ , including systematic uncertainties. We measure the branching fraction product $\mathcal{B}(B^\pm \rightarrow X(3823)K^\pm)\mathcal{B}(X(3823) \rightarrow \chi_{c1}\gamma) = (9.7 \pm 2.8 \pm 1.1) \times 10^{-6}$. No evidence is found for $X(3823) \rightarrow \chi_{c2}\gamma$ and we set an U.L. on its branching fraction product \mathcal{B} as well as the ratio $R_B \equiv \frac{\mathcal{B}(X(3823) \rightarrow \chi_{c2}\gamma)}{\mathcal{B}(X(3823) \rightarrow \chi_{c1}\gamma)} < 0.41$ at 90% C.L. The properties of the $X(3823)$ are consistent with those expected for the ψ_2 (1^3D_2 $c\bar{c}$) state. We also determine an U.L. on the product of branching fractions, $\mathcal{B}(B^\pm \rightarrow X(3872)K^\pm)\mathcal{B}(X(3872) \rightarrow \chi_{c1}\gamma) < 1.9 \times 10^{-6}$ at 90% C.L.; this is less than one quarter of the corresponding value in $X(3872) \rightarrow J/\psi\pi^+\pi^-$ [9]. Our results show that the production of the $X(3872)$'s C -odd partner in two-body B decays and its decay to $\chi_{c1}\gamma$ are considerably suppressed.

We thank the KEKB group for excellent operation

of the accelerator; the KEK cryogenics group for efficient solenoid operations; and the KEK computer group, the NII, and PNNL/EMSL for valuable computing and SINET4 network support. We acknowledge support from MEXT, JSPS and Nagoya's TLPSC (Japan); ARC and DIISR (Australia); FWF (Austria); NSFC (China); MSMT (Czechia); CZF, DFG, and VS (Germany); DST (India); INFN (Italy); MEST, NRF, GSDC of KISTI, and WCU (Korea); MNiSW and NCN (Poland); MES and RFAAE (Russia); ARRS (Slovenia); IKERBASQUE and UPV/EHU (Spain); SNSF (Switzerland); NSC and MOE (Taiwan); and DOE and NSF (USA). This work is partly supported by MEXT's Grant-in-Aid for Scientific Research on Innovative Areas ("Elucidation of New hadrons with a Variety of Flavors").

-
- [1] N. Brambilla *et al.*, Eur. Phys. J. C **71**, 1534 (2011).
 - [2] S.-K. Choi *et al.* (Belle Collaboration), Phys. Rev. Lett. **91**, 262001 (2003).
 - [3] D. Acosta *et al.* (CDF Collaboration), Phys. Rev. Lett. **93**, 072001 (2004).
 - [4] V.M. Abazov *et al.* (D0 Collaboration), Phys. Rev. Lett. **93**, 162002 (2004).
 - [5] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. D **71**, 071103 (2005).
 - [6] R. Aaij *et al.* (LHCb Collaboration), Eur. Phys. J. C **72**, 1972 (2012).
 - [7] V. Chiochia *et al.* (CMS Collaboration), arXiv:hep-ex/1201.6677.
 - [8] S.-K. Choi *et al.* (Belle Collaboration), Phys. Rev. D **84**, 052004 (2011).
 - [9] J. Beringer *et al.* (Particle Data Group), Phys. Rev. D **86**, 010001 (2012).
 - [10] E.S. Swanson, Phys. Lett. B **598**, 197 (2004); E.S. Swanson, Phys. Rept. **429**, 243 (2006).
 - [11] L. Maiani, F. Piccinini, A.D. Polosa and V. Riquer, Phys. Rev. D **71**, 014028 (2005).
 - [12] B.A. Li, Phys. Lett. B **605**, 306 (2005).
 - [13] B. Aubert *et al.* (BaBar Collaboration), Phys. Rev. Lett. **102**, 132001 (2009).
 - [14] V. Bhardwaj *et al.* (Belle Collaboration), Phys. Rev. Lett. **107**, 091803 (2011).
 - [15] K. Terasaki, Prog. Theor. Phys. **127**, 577 (2012).
 - [16] J. Nieves and M. PavonValderrama, Phys. Rev. D **86**, 056004 (2012).
 - [17] E. Eichten, K. Gottfried, T. Kinoshita, K.D. Lane and T.M. Yan, Phys. Rev. D **17**, 3090 (1978); E. Eichten, K. Gottfried, T. Kinoshita, K.D. Lane and T.M. Yan, Phys. Rev. D **21**, 203 (1980).
 - [18] W. Buchmüller and S.-H.H. Tye, Phys. Rev. D **24**, 132 (1981).
 - [19] E.J. Eichten, K. Lane and C. Quigg, Phys. Rev. Lett. **89**, 162002 (2002).
 - [20] P. Cho and M.B. Wise, Phys. Rev. D **51**, 3352 (1995).
 - [21] L. Antoniazzi *et al.* (E705 Collaboration), Phys. Rev. D **50**, 4258 (1994). Note that no hint of the $\psi(3836)$ is seen in $J/\psi\pi^+\pi^-$ studies focused on the $X(3872)$ [2–7].
 - [22] Charge-conjugate and neutral modes are included

throughout the paper unless stated otherwise.

- [23] A. Abashian *et al.* (Belle Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 117 (2002); also see detector section in J. Brodzicka *et al.*, Prog. Theor. Exp. Phys. (2012) 04D001.
- [24] S. Kurokawa and E. Kikutani, Nucl. Instrum. Methods Phys. Res., Sect. A **499**, 1 (2003), and other papers included in this Volume; T. Abe *et al.*, Prog. Theor. Exp. Phys. (2013) 03A001 and following articles up to 03A011.
- [25] P. Koppenburg *et al.* (Belle Collaboration), Phys. Rev. Lett. **93**, 061803 (2004).
- [26] K.-F. Chen *et al.* (Belle Collaboration), Phys. Rev. D **72**, 012004 (2005).
- [27] Hereinafter, χ_{cJ} refers to either χ_{c1} or χ_{c2} , depending on which is reconstructed.
- [28] G.C. Fox and S. Wolfram, Phys. Rev. Lett. **41**, 1581 (1978).
- [29] The logarithmic Gaussian is parameterized as:

$$f(x) = \frac{N_0}{\sqrt{2\pi}\sigma_0(\epsilon-x)} \exp - \frac{(\ln(\frac{\epsilon-x}{\epsilon-x_p}))^2}{2\sigma_0^2}, \text{ where } \epsilon = \frac{\sigma}{a} + x_p$$
and $\sigma_0 = \frac{\ln(a\sqrt{2\ln 2} + \sqrt{1+2a^2\ln 2})}{\sqrt{2\ln 2}}$. Here, N_0 is the normalization, σ is the standard deviation, x_p is the mean and a is the asymmetry.
- [30] H. Albrecht *et al.* (ARGUS Collaboration), Phys. Lett. B **241**, 278 (1990).
- [31] The curves show the signal [red large-dashed for ψ' , magenta dashed for $X(3823)$ and violet dotted for $X(3872)$] and the background component [black dotted-dashed for combinatorial, dark green two dotted-dashed for $B \rightarrow \psi'(\rightarrow \chi_{cJ}\gamma)K$ and cyan three dotted-dashed for peaking component] as well as the overall fit [blue solid]. $B \rightarrow \psi'(\rightarrow \chi_{cJ}\gamma)K$ is specific to the decay mode under study.
- [32] R.D. Cousins and V.L. Highland, Nucl. Instrum. Methods Phys. Res., Sect. A **320**, 331 (1992).
- [33] S. Godfrey and N. Isgur, Phys. Rev. D **32**, 189 (1985).
- [34] D. Ebert, R.N. Faustov and V.O. Galkin, Phys. Rev. D **67**, 014027 (2003).
- [35] E.J. Eichten, K. Lane and C. Quigg, Phys. Rev. D **69**, 094019 (2004).
- [36] M. Blank and A. Krassnigg, Phys. Rev. D **84**, 096014 (2011).
- [37] P. Pakhlov *et al.* (Belle Collaboration), Phys. Rev. Lett. **100**, 202001 (2008).
- [38] P. Ko, J. Lee and H.S. Song, Phys. Lett. B **395**, 107 (1997).
- [39] C.-F. Qiao, F. Yuan and K.-T. Chao, Phys. Rev. D **55**, 4001 (1997).
- [40] M. Suzuki, Phys. Rev. D **66**, 037503 (2002); P. Colangelo, F. De Fazio and T.N. Pham, Phys. Lett. B **542**, 71 (2002).