

This is the accepted manuscript made available via CHORUS. The article has been published as:

Third Law of Thermodynamics and The Shape of the Phase Diagram for Systems With a First-Order Quantum Phase Transition

T. R. Kirkpatrick and D. Belitz

Phys. Rev. Lett. **115**, 020402 — Published 8 July 2015

DOI: [10.1103/PhysRevLett.115.020402](https://doi.org/10.1103/PhysRevLett.115.020402)

The third law of thermodynamics and the shape of the phase diagram for systems with a first-order quantum phase transition

T. R. Kirkpatrick¹ and D. Belitz^{2,3}

¹ *Institute for Physical Science and Technology,
and Department of Physics, University of Maryland,
College Park, MD 20742, USA*

² *Department of Physics and Institute of Theoretical Science,
University of Oregon, Eugene, OR 97403, USA*

³ *Materials Science Institute,
University of Oregon, Eugene, OR 97403, USA*

(Dated: June 5, 2015)

The Third Law of Thermodynamics constrains the phase diagram of systems with a first-order quantum phase transition. For zero conjugate field, the coexistence curve has an infinite slope at $T = 0$. If a tricritical point exists at $T > 0$, then the associated tricritical wings are perpendicular to the $T = 0$ plane, but not to the zero-field plane. These results are based on the third law and basic thermodynamics only, and are completely general. As an explicit example we consider the ferromagnetic quantum phase transition in clean metals, where a first-order quantum phase transition is commonly observed.

PACS numbers:

First-order phase transitions are ubiquitous in nature, the solid-to-liquid and liquid-to-gas transitions being the most commonly observed ones. Another common example of a first-order transition is the ferromagnetic transition below the Curie temperature as a function of an external magnetic field. First-order transitions are characterized by a coexistence curve in the phase diagram along which the two phases coexist in thermodynamic equilibrium. (The coexistence curve may be the projection of a higher-dimensional coexistence manifold into a particular plane in the phase diagram.)

It has long been known that the curvature of the coexistence curve is determined by the discontinuities of certain observables across it. The Clapeyron-Clausius (CC) equation relates the slope of the coexistence curve in the pressure-temperature (p - T) plane to the discontinuities of the entropy and the volume [1]:

$$\left(\frac{dp}{dT}\right)_H = \frac{\Delta s}{\Delta v}, \quad (1)$$

where $\Delta s = s_1 - s_2$ and $\Delta v = v_1 - v_2$ with $s_{1,2}$ and $v_{1,2}$ the specific entropy and volume per particle, respectively, in the two phases. For definiteness, let 1 and 2 label the ordered and disordered phases, respectively, and for later reference we indicate that an appropriate external field H , if any, is held constant in taking the derivative.

The CC equation (1) and its analogs in different planes of the phase diagram are very general, as they rely only on basic thermodynamic arguments. In this Letter we show that for *quantum* phase transitions, when combined with the Third Law of Thermodynamics, they provide interesting constraints on the shape of the phase diagram. We will consider a pressure-driven transition at $T = 0$ that is first order, remains first order at low T , and turns




Figure 1: Schematic phase diagram showing a line of first-order transitions at low T separated from a line of second-order transitions at higher T by a tricritical point (TCP). In a nonzero conjugate field H tricritical wings emerge from the TCP. These are surfaces of first-order transitions that are bounded by lines of second-order transitions and terminate in two quantum wing-critical points (QWCP) in the $T = 0$ plane.

second order at higher T via a tricritical point (TCP). The schematic phase diagram in the space spanned by T , p , and H , where H is the field conjugate to the order parameter, is shown in Fig. 1. As we will see, the detailed shape of this phase diagram at low T is constrained by thermodynamics. Our arguments leading to this conclusion are completely general; however, as an explicit example we will discuss the quantum ferromag-

netic transition in clean metals [2]. Another example of a first-order quantum phase transition with a TCP in the phase diagram is the Ising antiferromagnet dysprosium aluminum garnet [3].

We are interested in a system with T , p , and H as independent variables. Denoting the order parameter by M , the appropriate thermodynamic potential is the generalized Gibbs free energy [4]

$$\begin{aligned}\tilde{G} &= U - TS + pV - HM \\ &= \mu N ,\end{aligned}\quad (2a)$$

whose differential is

$$d\tilde{G} = -SdT + Vdp - MdH + \mu dN . \quad (2b)$$

Here S , V , and μ are the system's entropy, volume, and chemical potential, respectively, and N is the particle number. From Eqs. (2) we obtain the Gibbs-Duhem relation

$$d\mu = d\tilde{g} = -sdT + vdp - mdH , \quad (3)$$

where \tilde{g} , s , v , and m are the generalized Gibbs free energy, entropy, volume, and order parameter per particle, respectively. On the coexistence curve the chemical potentials of the two phases must coincide. Using this condition with Eq. (3) at fixed external field leads to Eq. (1). An analogous argument yields

$$\left(\frac{dT}{dH}\right)_p = -\frac{\Delta m}{\Delta s} . \quad (4)$$

The CC equations (1,4) are completely general. When applied to a quantum phase transition, the Third Law provides the additional constraint $\Delta s(T \rightarrow 0) \rightarrow 0$. To be specific, let us assume that in either phase the entropy vanishes as $s(T \rightarrow 0) = \gamma T^n$. In particular, if the phases are Fermi liquids (see below) then $n = 1$ and γ is the specific-heat coefficient. For asymptotically low temperatures we thus have

$$\left(\frac{dT}{dp}\right)_H = \frac{1}{T^n} \frac{\Delta v}{\Delta \gamma} , \quad (5a)$$

$$\left(\frac{dT}{dH}\right)_p = \frac{-1}{T^n} \frac{\Delta m}{\Delta \gamma} . \quad (5b)$$

In addition, we obtain from the equilibrium condition in conjunction with Eq. (2b) a third CC equation,

$$\left(\frac{dH}{dp}\right)_T = \frac{\Delta v}{\Delta m} . \quad (5c)$$

These three CC equations are the basis of our discussion.

Let us start by briefly discussing the obvious coexistence region in the T - p plane, which is labeled “ordered” in Fig. 1. The first-order transition across this plane, which is driven by the external field, does not involve

any change in either entropy or specific volume. We thus have $\Delta s = \Delta v = 0$. Equations (5b, 5c) then imply $(dH/dT)_p = (dH/dp)_T = 0$. This identifies the $H = 0$ plane as the locus of the coexistence curves. $(dT/dp)_{H=0}$, Eq. (5a), is indeterminate, which is consistent with the fact that *any* curve in the $H = 0$ plane below the transition temperature is a coexistence curve. The CC equations thus correctly describe the coexistence plane, but do not provide any nontrivial information.

This changes as we consider the other coexistence surfaces, viz., the tricritical wings. Obviously, we have $\Delta m > 0$ across any first-order transition, but now Δs and Δv will not be zero. To find Δv , we turn to scaling theory. Scaling is often thought of as valid only at second-order transitions. However, Fisher and Berker [5] have shown that finite-size scaling considerations allow for the definition of a diverging length scale even at a first-order transition. Consequently, a classical first-order transition can be considered a limiting case of a second-order transition, and the homogeneity laws, exponent relations, etc., that are known from the scaling description of second-order transitions still hold. This formalism has recently been generalized to the case of quantum first-order transitions [6], and we now apply it to the problem under consideration. Let $r = (p - p^*)/p^*$ be the dimensionless distance from the transition at $T = 0$. Then the generalized Gibbs free energy obeys a homogeneity law [6]

$$\tilde{g}(r, H, T) = b^{-(d+z)} \Phi_{\tilde{g}}(r b^{d+z}, H b^{d+z}, T b^z) . \quad (6)$$

Here $b > 0$ is the length scaling parameter, $\Phi_{\tilde{g}}$ is a scaling function, and we have made use of several exponent values that characterize a first-order quantum phase transition (see Ref. 6 for details): z is the relevant dynamical exponent [7], the inverse correlation length exponent has its largest possible value $1/\nu = d + z$, and the scale dimension of the field, $[H] = \beta\delta/\nu$, reflects the fact that the order-parameter exponents are $\beta = 0$, $\delta = \infty$, such that $\beta\delta = 1$. This is a generalization of the scaling description of classic first-order transitions given by Fisher and Berker [5]. Differentiating \tilde{g} with respect to $r \propto p$, we see that the scaling part of the specific volume, $v = \partial\tilde{g}/\partial p = \partial\tilde{g}/\partial r$, has a zero scale dimension. This implies a discontinuity of the specific volume across the transition, and a corresponding δ -function contribution to the compressibility $\kappa = -(\partial v/\partial p)/v$. This is in direct analogy to the latent heat at a classical first-order transition and can be interpreted as a “latent volume” at a pressure-driven QPT, i.e., the volume changes by a finite amount upon an infinitesimal change in pressure [8]. We also note that differentiating with respect to r again we obtain $\kappa \propto |r|^{-1}$; see Ref. [5] for an interpretation of this power-law divergence at a first-order transition in terms of finite-size scaling. The compressibility is positive in thermodynamic equilibrium, and we thus have $\Delta v > 0$ across any coexistence curve contained in the wings. The only remaining question is the sign of $\Delta\gamma$. For a transi-




Figure 2: Measured phase diagram of UGe_2 , with the same notation as in Fig. 1. The main figure is adapted from Ref. 10; the squares represent data points, the lines and surfaces are guides to the eye. The inset shows the $H = 0$ plane with data from Ref. 11. Note the extremely steep drop of the Curie temperature past the tricritical point. The ferromagnetic phase consists of two phases, FM1 and FM2, separated by a line of first-order transitions at low temperatures that ends in a critical point (CP).

tion from an ordered phase to a disordered one, one would naively expect $\Delta\gamma < 0$. We will adopt this expectation for now and give a more detailed discussion below.

From Eq. (5a) we conclude that the $(\partial T/\partial p)_H < 0$, i.e., the slope of any coexistence curve at fixed H is negative, and it approaches $-\infty$ as $1/T^n$ for $T \rightarrow 0$. Similarly, Eq. (5b) yields $(\partial T/\partial H)_p > 0$, and it approaches $+\infty$ as $1/T^n$ for $T \rightarrow 0$. This means that the wings are necessarily perpendicular to the $T = 0$ plane, and in particular the coexistence curve in zero field has an infinite slope at the QPT. Finally, Eq. (5c) implies that $(\partial H/\partial p)_T$ is positive and finite, which implies that the wings are tilted in the direction of the disordered phase and are *not* perpendicular to the p -axis.

We now turn to an explicit example that illustrates all of the above considerations, namely, the quantum phase transition in clean metallic ferromagnets [2, 9]. In this case, the exponent n in Eqs. 5 is $n = 1$, and the dynamical exponent z in Eq. (6) is $z = 1$. There is a second dynamical exponent $z = 3$, but for our purposes $z = 1$ yields the dominant contribution (see Refs. [2, 6] for a detailed discussion of this point). The order parameter m is the magnetization, H is the external magnetic field, and the phase diagram is generically observed to have the topology shown in Fig. 1. The features discussed above are indeed universally observed in all cases where the tricritical wings have been mapped out in detail. As an example, we show the experimentally determined wings in UGe_2 in Fig. 2; for other examples, see Ref. 2. Note the extremely sharp drop of the Curie temperature for

pressures above the tricritical pressure that is apparent in the inset. Also of interest is the transition inside the ferromagnetic phase (from FM1 to FM2), which is first order at low temperatures. Our considerations apply to this transition as well, and the steep drop of the transition temperature is again consistent with an infinite slope of the coexistence curve at $T = 0$.

We now return to the issue of the sign of $\Delta\gamma$, or more generally Δs , across the coexistence curve. From Eq. (3) we see that $(\partial s/\partial p)_{T,H} = -(\partial v/\partial T)_{p,H} = -v\alpha_p$, with $\alpha_p = (\partial v/\partial T)_p/v$ the thermal expansion coefficient. An increase in entropy with increasing pressure thus implies $\alpha_p < 0$. Returning to Eq. (5a), and remembering that $\Delta v > 0$ since the compressibility is necessarily positive, we see that a decreasing Curie temperature with increasing pressure implies a negative thermal expansion coefficient, and vice versa. Consistent with this, the thermal expansion coefficient at low T is indeed negative in UGe_2 [12], MnSi [13], and ZrZn_2 [14], which all are low- T ferromagnets with qualitatively identical phase diagrams. It is interesting that $\alpha_p < 0$ by itself implies that the high-pressure phase must be the paramagnetic one. We also note that the volume is discontinuous if the coexistence curve is crossed at fixed p as a function of T as well as at fixed T as a function of p . This is intuitively obvious and also follows from Eqs. (3, 6). Accordingly, α_p at the first-order transition has a δ -function contribution that reflects the same “latent volume” as the corresponding δ -function contribution to the compressibility. This is consistent with the experiment by Kabeya et al [12], who observed a pronounced negative peak in α_p at the transition, which they attributed to a broadened first-order transition.

While in most quantum ferromagnets hydrostatic pressure destroys the ferromagnetic order, there are systems in which the opposite occurs. An example is YbCu_2Si_2 , which is paramagnetic at ambient pressure, but becomes ferromagnetic upon the application of hydrostatic pressure of roughly 10 GPa [15], see Fig. 3. The thermodynamic arguments presented above then predict that α_p in this material, at this pressure and at low temperature, must be positive. The thermal expansion data of Ref. 16 are consistent with this prediction, although not quite conclusive, as they focused on a higher temperature region. There also are materials where hydrostatic pressure drives the system away from ferromagnetic order, while uniaxial stress favors it; an example is UCoAl [17]. This can be understood by realizing that in many solids the thermal expansion coefficient is anisotropic to the point of being positive along some crystal axes, but negative along others [18].

We finally briefly discuss an explicit equation of state that has been used to describe the qualitative phase diagram of metallic quantum ferromagnets and leads to a schematic phase diagram as shown in Fig. 1 [19]. It is




Figure 3: Temperature-pressure phase diagram of YbCu_2Si_2 . Ferromagnetism is induced by hydrostatic pressure $p > P_c \approx 8.25$ GPa. From Ref. 15.

derived by minimizing a generalized Landau functional

$$f(m) = -Hm + rm^2 + wm^4 \ln(m^2 + T^2) + um^4 \quad (7)$$

with respect to the magnetization m . Here m , H , and T are measured in suitable microscopic units, and r , u , and w are parameters of the generalized Landau theory. The physical origin of the term with coupling constant w is due to soft fermionic excitations that couple to the magnetization; this has been discussed in detail before [2, 20, 21] and will not be repeated here. In zero field, $H = 0$, the logarithmic term leads to a first-order transition at $r = r_1 = w e^{-1-u/w}$ where the magnetization value is $m_1 = \sqrt{r_1}$, and to a tricritical point at $T_{tc} = e^{-u/2w}$. The coexistence curve can easily be obtained explicitly [19]; here we just quote the asymptotic behavior for $r \rightarrow r_1$,

$$T(r \rightarrow r_1) = \frac{1}{\sqrt{w}} (r_1 - r)^{1/2} \quad (8)$$

which yields

$$dT/dr|_{r \rightarrow r_1} = -1/2wT, \quad (9)$$

in agreement with the general conclusions drawn above from thermodynamics, see Eq. (5a) and the related discussion. In the presence of a small magnetic field, one finds for the coexistence curve in the $T = 0$ plane

$$H = m_1 \left(1 + \frac{3}{11} \frac{u}{w} \right) \delta r + O(\delta r^2), \quad (10)$$

where $\delta r = r - r_1$. This reflects the linear slope of the tricritical wings with respect to the r -axis that follows from Eq. (5c). A more involved, but elementary analysis shows that the tricritical wings are perpendicular to the $T = 0$ plane everywhere. We stress that these properties are not tied to the specific physical mechanism that

underlies the free-energy function (7); they must be true for *any* model that leads to a first-order quantum phase transition and correctly reflects thermodynamics.

We conclude with some additional discussion points. (1) The most often observed shape of the phase diagram in quantum ferromagnets, with increasing hydrostatic pressure driving the system into the disordered phase, is not what one might naively expect. In a fluid analogy, this is equivalent to what is observed in H_2O and H_2S , while in most fluids increased pressure stabilizes the ordered phase. In quantum ferromagnets the latter can also occur, see Fig. 3, but it is not common. As we have shown, this feature of the phase diagram is tied to the sign of the thermal expansion coefficient, which tends to be negative in low-temperature ferromagnets. Regardless of whether hydrostatic pressure induces or destroys ferromagnetism, Eq. (5c) implies that the tricritical wings must always extend in the direction of the paramagnetic phase. (2) All actually measured tricritical wings show all of the structural features discussed above, as they must, since the former hinge on basic thermodynamics only. However, occasionally schematic drawings of wings in the literature violate these thermodynamic requirements: They show wings that are perpendicular to the pressure axis and/or not perpendicular to the $T = 0$ plane. (3) The discontinuity of the specific-heat coefficient across the first-order transition, $0 < \Delta\gamma < \infty$, has an interesting implication for the dynamical critical exponents in the system. Since the static and dynamic specific-heat exponents $\bar{\alpha}$ and z_c are related by a hyperscaling relation $\bar{\alpha} = \nu(z_c - d)$, with ν the correlation-length exponent and d the spatial dimensionality of the system [6], a discontinuous specific-heat coefficient ($\bar{\alpha} = 0$) implies $z_c = d$. This is indeed the value of z_c within an explicit theory that describes the first-order transition, see Ref. 6.

This work was supported by the NSF under grant Nos. DMR-1401410 and DMR-1401449.

-
- [1] L. D. Landau and E. M. Lifshitz, *Statistical Physics Part 1* (Butterworth Heinemann, Oxford, 1980).
 - [2] M. Brando, D. Belitz, F. M. Grosche, and T. R. Kirkpatrick, arXiv:1502.02898.
 - [3] N. Giordano and W. P. Wolf, Phys. Rev. Lett. **39**, 342 (1977).
 - [4] This is thermodynamic potential appropriate for a situation where H is an independent variable. It is the Gibbs version of the thermodynamic potential denoted by $\tilde{\mathcal{F}}$ in Ref. 22, $\tilde{G} = \tilde{\mathcal{F}} + pV$. Note that Ref. 22 considers ferromagnets, with M the magnetic moment and H the magnetic field, but Eq. (2a) remains valid for any order-parameter-conjugate-field pair.
 - [5] M. E. Fisher and A. N. Berker, Phys. Rev. B **26**, 2507 (1982).
 - [6] T. Kirkpatrick and D. Belitz, arXiv:1503.04175 (Phys. Rev. B in press).

- [7] In general there may be more than one dynamical critical exponent, and more than one scaling function, see Ref. [6]. If this is the case, one needs to determine which yield the dominant contribution for the problem at hand.
- [8] In Ehrenfest's classification of phase transitions (P. Ehrenfest, *Verhandlungen der Koninklijke Akademie van Wetenschappen (Amsterdam)* **36**, 153 (1933)), a first-order transition was characterized by a latent heat. This is commonly interpreted as requiring only that *some* derivative of an appropriate free energy shows a discontinuity; see, e.g., G. Jaeger, *Arch. Hist. Exact Sci.* **53**, 61 (1998). This definition is trivially generalized to $T = 0$.
- [9] Our conclusions hold as long as the transition at $T = 0$ is first order, irrespective of whether the material is clean or not. However, only for weakly disordered metals is the ferromagnetic quantum phase transition theoretically predicted and experimentally observed to be generically of first order, see Ref. [2]. The evolution of the phase diagram with increasing disorder strength has been discussed in Ref. [23].
- [10] H. Kotegawa, V. Taufour, D. Aoki, G. Knebel, and J. Flouquet, *J. Phys. Soc. Japan* **80**, 083703 (2011).
- [11] V. Taufour, D. Aoki, G. Knebel, and J. Flouquet, *Phys. Rev. Lett.* **105**, 217201 (2010).
- [12] N. Kabeya, R. Iijima, E. Osaki, S. Ban, K. Imura, K. Deguchi, N. Aso, Y. Homma, Y. Shiokawa, and N. Sato, *J. Phys: Conf. Series* **200**, 032028 (2010).
- [13] A. Miyaki, A. Villaume, Y. Haga, G. Knebel, B. Salce, G. Lapertot, and J. Flouquet, *J. Phys. Soc. Japan* **78**, 044703 (2009).
- [14] S. Ogawa, *Physica B* **119**, 68 (1983).
- [15] N. Tateiwa, T. D. Matsuda, Y. Haga, and Z. Fisk, *Phys. Rev. B* **89**, 035127 (2014).
- [16] Y. Uwatoko, G. Oomi, J. Thompson, P. Canfield, and Z. Fisk, *Physica B* **186**, 593 (1993).
- [17] Y. Shimizu, B. Salce, T. Combier, D. Aoki, and J. Flouquet, *J. Phys. Soc. Jpn.* **84**, 023704 (2015).
- [18] See, e.g., Ref. 24. A detailed analysis of the consequences of anisotropy for thermodynamics requires a more thorough treatment, see, e.g., Ref. 25.
- [19] D. Belitz, T. R. Kirkpatrick, and J. Rollbühler, *Phys. Rev. Lett.* **94**, 247205 (2005).
- [20] D. Belitz, T. R. Kirkpatrick, and T. Vojta, *Phys. Rev. Lett.* **82**, 4707 (1999).
- [21] T. R. Kirkpatrick and D. Belitz, *Phys. Rev. B* **85**, 134451 (2012).
- [22] L. D. Landau and E. M. Lifshitz, *Electrodynamics of Continuous Media* (Pergamon, Oxford, 1984).
- [23] Y. Sang, D. Belitz, and T. R. Kirkpatrick, *Phys. Rev. Lett.* **113**, 207201 (2014).
- [24] B. Childs, *Rev. Mod. Phys.* **25**, 665 (1953).
- [25] H. Callen, *Thermodynamics and an introduction to thermostatistics* (Wiley (New York), 1985).