



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Long-Lived Gluinos and Stable Axinos

Stuart Raby

Phys. Rev. Lett. **115**, 231801 — Published 1 December 2015

DOI: [10.1103/PhysRevLett.115.231801](https://doi.org/10.1103/PhysRevLett.115.231801)

Glauino LOSP with Axino LSP

Stuart Raby

Department of Physics
The Ohio State University
191 W. Woodruff Ave, Columbus, OH 43210, USA

October 26, 2015

Abstract

In this letter we have presented a novel version of “long-lived” gluinos in supersymmetric models with the gluino the lightest ordinary supersymmetric particle [LOSP] and axino LSP. Within certain ranges of the axion decay constant $f_a < 1 \times 10^{10}$ GeV, the gluino mass bounds are reduced to less than 1000 GeV. The best limits can be obtained by looking for decaying R-hadrons in the detector where the gluino decays to a gluon and axino in the calorimeters. SUSY models with a gluino LOSP can occur over a significant region of parameter space in either *mirage-mediation* or general gauge-mediated SUSY breaking models. The gluino LOSP is not constrained by cosmology, but in this scenario the axion/axino may be good dark matter candidates.

1 Introduction

Supersymmetric models with non-universal gaugino masses are well-defined even in the range of parameters for which gluinos are the LSP [1–5]. The most stringent constraint in this case comes from cosmology. Although gluinos are Majorana particles and annihilate with a strong interaction rate, there are still too many of them left over in the early universe. They form stable, hadronic bound states and searches for heavy hydrogen essentially rule out this possibility [6–8]. In this paper we show that when gluinos are the Lightest Ordinary Supersymmetric Particle with an axino LSP we clearly evade these cosmological bounds. Moreover we show that collider bounds on the gluino mass can be significantly reduced for an axion decay constant in the allowed region with $f_a < 1 \times 10^{10}$ GeV.

There are presently strong limits on gluino masses coming from the first run of the LHC. In particular, we are interested in the bounds for long-lived gluinos. We consider recent CMS results. These come from searches for R-hadrons which travel through the detector [9] or for R-hadrons which stop in the detector and then decay [10]. For long-lived gluinos the bounds depend on the fraction, f , which initially hadronize as a gluino-gluon bound state [9]. For $f = 0.1$, the bounds require $m_{\tilde{g}} > 1233$ GeV or even greater, depending on the model for propagation. These bounds only require data from the central tracker. The track that the charged R-hadron leaves in the central tracker must satisfy $|\eta| < 2.1$, $p_T > 45$ GeV/c and have significant dE/dx . The track must also be isolated in both the tracker and calorimeter with $\sum p_T < 50$ GeV/c where the sum is over all tracks (except the candidate track) within a cone $\Delta R < 0.3$ radians. Thus if the R-hadron should decay in the tracker or calorimeter it will not pass these cuts. However, if the lifetime is long enough such that the gluino decays outside the window of the collision region, it will be counted in this event sample. Therefore, in order to evade these bounds, the gluino must decay fast enough so that it leaves energy in the calorimeters. Thus we need $\langle\beta\rangle c\tau_{\tilde{g}} \lesssim 2.5$ meters or $\tau_{\tilde{g}} \lesssim 8.3 (\langle\beta\rangle)^{-1} \times 10^{-9}$ s. If the gluino does live long enough to make it to the muon system, then it will be constrained by other data sets which give just as strong bounds on the gluino mass. The limits are slightly less strong for gluinos which are long-lived and stop in the detector [10]. In this case the most stringent bound is $m_{\tilde{g}} \geq 1000$ GeV for gluino life-times in the range, 10^{-6} s $< \tau_{\tilde{g}} < 10^3$ s. However, any event which satisfies the latter stopped bounds were already constrained by the former long-lived bounds.

These limits can be applied to any theory where the gluino is the NLSP which decays into a gluon and neutral LSP. It has applications to gauge-mediated SUSY breaking models where the gluino decays to a gluon and a goldstino [11] or to split SUSY models where the gluino decays to a quark-anti-quark pair and neutralino LSP [12].

In this letter we consider another scenario where the gluino is the lightest ordinary supersymmetric particle [LOSP] with an axino LSP. The strong CP problem has a natural solution in terms of the Peccei-Quinn-Weinberg-Wilczek axion [13–15]. Its modern form is in terms of an invisible axion of the KSVZ type [16, 17] or of the DFSZ type [18, 19]. In a supersymmetric theory the axion has a scalar partner, the saxion, and a fermionic partner, the axino [20, 21]. The saxion typically has mass of order the gravitino mass while the axino can be much lighter [22–25].

Supersymmetric models with non-universal gaugino masses, such as *mirage-mediation* SUSY breaking or general gauge-mediation have significant ranges of parameters where the

gluino is the LSP. For example, in a recent study of *mirage-mediation* in an SO(10) GUT [5] the gaugino mass formula at the GUT scale is given by

$$M_i = \left(1 + \frac{g_G^2 b_i \alpha}{16\pi^2} \log \left(\frac{M_{Pl}}{m_{16}} \right) \right) M_{1/2}, \quad (1)$$

where $b_i = (33/5, 1, -3)$ for $i = 1, 2, 3$, $M_{1/2}$ is the overall mass scale, and α is the ratio of the anomaly mediation to gravity mediation contributions. For α in the range, $3 \leq \alpha \leq 4$, we find that the gluino is the LSP. Also in the case of general gauge-mediation (see Ref. [3]), for messenger scales of order the GUT scale, the gravitino is heavy and we find that it is quite natural to have a gluino LSP. We note that these models have the very interesting property of precision gauge coupling unification [4, 26], i.e. the gauge couplings unify at the GUT scale to high accuracy. Thus requiring little or no threshold corrections at M_{GUT} . However, as discussed earlier, such theories are unacceptable cosmologically. So therefore it becomes advantageous to combine these models with the axion solution to the strong CP problem. Then the gluino is the LOSP with an axino LSP.

2 Gluino - Axino coupling and gluino life-time

We consider the supersymmetrized axion multiplet coupling to the SU(3) gauge sector. We have [27, 28]

$$\mathcal{L}^{eff} = -\frac{\alpha_s}{2\sqrt{2}\pi f_a} \int d^2\theta A W^{a\alpha} W_\alpha^a + h.c. \quad (2)$$

where the axion superfield,

$$A = \frac{s + ia}{\sqrt{2}} + \sqrt{2}(\theta \psi_a) + (\theta)^2 F_A, \quad (3)$$

and the gauge superfield strength is given by

$$W_\alpha^a = -i\lambda_\alpha^a + [\delta_\alpha^\beta D^a - \frac{i}{2}(\sigma^\mu \bar{\sigma}^\nu)_\alpha^\beta G_{\mu\nu}^a] \theta_\beta + (\theta)^2 \sigma_{\alpha\dot{\alpha}}^\mu D^\mu \bar{\lambda}^{\dot{\alpha}}. \quad (4)$$

We define the four component Majorana spinors for the gluino and axino fields by

$$\tilde{g} = \begin{pmatrix} -i\lambda \\ i\bar{\lambda} \end{pmatrix}, \quad \tilde{a} = \begin{pmatrix} \psi_a \\ \bar{\psi}_a \end{pmatrix}. \quad (5)$$

With this notation, we have

$$\begin{aligned} \mathcal{L}^{eff} = & \frac{\alpha_s}{8\pi f_a} [a (G^{a\mu\nu} \tilde{G}_{\mu\nu}^a + D_\mu \bar{g} \gamma^\mu \gamma_5 \tilde{g}) \\ & + s (G^{a\mu\nu} G_{\mu\nu}^a - 2D^a D^a + 2\tilde{g} \gamma^\mu D_\mu \tilde{g}) \\ & + i\tilde{a} G_{\mu\nu}^a \frac{[\gamma^\mu, \gamma^\nu]}{2} \gamma_5 \tilde{g} - 2\tilde{a} \tilde{g} D^a] \\ & + \sqrt{2} F_A (\lambda \lambda) + h.c.. \end{aligned} \quad (6)$$

We then find the gluino decay rate (neglecting the axino mass) given by

$$\Gamma_{\tilde{g} \rightarrow \tilde{a} g} = \frac{\alpha_s^2 m_{\tilde{g}}^3}{128\pi^3 f_a^2}. \quad (7)$$

Thus the gluino lifetime for particular values of f_a and gluino mass is given by

$$\tau_{\tilde{g} \rightarrow \tilde{a} g} = 3 \times 10^{-8} s \frac{(f_a/10^{10} \text{ GeV})^2}{(m_{\tilde{g}}/850 \text{ GeV})^3}. \quad (8)$$

Therefore, as an example, for values of $f_a \leq 0.7 \times 10^{10}$ GeV and an average gluino velocity, $\beta \sim \frac{1}{2}$, the gluino mass bound may reduce to $m_{\tilde{g}} \gtrsim 850$ GeV.

3 Conclusion

We have presented a novel version of gluinos LOSPs in supersymmetric models with an axino LSP. Within certain ranges of the axion decay constant $f_a < 1 \times 10^{10}$ GeV, the gluino mass bounds are reduced to less than 1000 GeV. The best limits can be obtained by looking for decaying R-hadrons in the hadronic calorimeters. Such decays will produce jets deep in the calorimeter. SUSY models with a gluino LOSP can occur over a significant region of parameter space in either *mirage-mediation* or general gauge-mediate SUSY breaking models. Since the gluino LOSP is no longer constrained by cosmology, it would now be interesting to analyze the possibility of axion/axino dark matter in this scenario.

Acknowledgments

S.R. acknowledges stimulating conversations with L.J. Hall, F. D’Eramo and significant feedback from C.S. Hill during the preparation of this letter. S.R. received partial support for this work from DOE/ DE-SC0011726. The idea was first presented while S.R. was supported by the Munich Institute for Astro- and Particle Physics (MIAPP) of the DFG cluster of excellence ”Origin and Structure of the Universe”.

References

- [1] H. Baer, K.-m. Cheung, and J. F. Gunion, “A Heavy gluino as the lightest supersymmetric particle,” *Phys. Rev.* **D59** (1999) 075002, hep-ph/9806361.
- [2] A. Mafi and S. Raby, “An Analysis of a heavy gluino LSP at CDF: The Heavy gluino window,” *Phys. Rev.* **D62** (2000) 035003, hep-ph/9912436.
- [3] S. Raby and K. Tobe, “The Phenomenology of SUSY models with a gluino LSP,” *Nucl. Phys.* **B539** (1999) 3–22, hep-ph/9807281.
- [4] S. Raby, M. Ratz, and K. Schmidt-Hoberg, “Precision gauge unification in the MSSM,” *Phys. Lett.* **B687** (2010) 342–348, 0911.4249.
- [5] A. Anandakrishnan, B. C. Bryant, and S. Raby, “LHC Phenomenology of SO(10) Models with Yukawa Unification II,” *Phys.Rev.* **D90** (2014) 015030, 1404.5628.
- [6] R. A. Muller, L. W. Alvarez, W. R. Holley, and E. J. Stephenson, “Quarks with Unit Charge: A Search for Anomalous Hydrogen,” *Science* **196** (1977) 521–523.
- [7] P. F. Smith and J. R. J. Bennett, “A SEARCH FOR HEAVY STABLE PARTICLES,” *Nucl. Phys.* **B149** (1979) 525–533.
- [8] P. F. Smith, J. R. J. Bennett, G. J. Homer, J. D. Lewin, H. E. Walford, and W. A. Smith, “A SEARCH FOR ANOMALOUS HYDROGEN IN ENRICHED D-2 O, USING A TIME-OF-FLIGHT SPECTROMETER,” *Nucl. Phys.* **B206** (1982) 333–348.
- [9] CMS Collaboration, S. Chatrchyan *et al.*, “Searches for long-lived charged particles in pp collisions at $\sqrt{s}=7$ and 8 TeV,” *JHEP* **1307** (2013) 122, 1305.0491.
- [10] CMS Collaboration, V. Khachatryan *et al.*, “Search for decays of stopped long-lived particles produced in protonproton collisions at $\sqrt{s} = 8$ TeV,” *Eur. Phys. J.* **C75** (2015), no. 4, 151, 1501.05603.
- [11] S. Dimopoulos, M. Dine, S. Raby, and S. D. Thomas, “Experimental signatures of low-energy gauge mediated supersymmetry breaking,” *Phys. Rev. Lett.* **76** (1996) 3494–3497, hep-ph/9601367.
- [12] N. Arkani-Hamed and S. Dimopoulos, “Supersymmetric unification without low energy supersymmetry and signatures for fine-tuning at the LHC,” *JHEP* **06** (2005) 073, hep-th/0405159.
- [13] R. D. Peccei and H. R. Quinn, “CP Conservation in the Presence of Instantons,” *Phys. Rev. Lett.* **38** (1977) 1440–1443.
- [14] S. Weinberg, “A New Light Boson?,” *Phys. Rev. Lett.* **40** (1978) 223–226.
- [15] F. Wilczek, “Problem of Strong p and t Invariance in the Presence of Instantons,” *Phys. Rev. Lett.* **40** (1978) 279–282.

- [16] J. E. Kim, “Weak Interaction Singlet and Strong CP Invariance,” *Phys. Rev. Lett.* **43** (1979) 103.
- [17] M. A. Shifman, A. I. Vainshtein, and V. I. Zakharov, “Can Confinement Ensure Natural CP Invariance of Strong Interactions?,” *Nucl. Phys.* **B166** (1980) 493.
- [18] M. Dine, W. Fischler, and M. Srednicki, “A Simple Solution to the Strong CP Problem with a Harmless Axion,” *Phys. Lett.* **B104** (1981) 199.
- [19] A. R. Zhitnitsky, “On Possible Suppression of the Axion Hadron Interactions. (In Russian),” *Sov. J. Nucl. Phys.* **31** (1980) 260. [*Yad. Fiz.*31,497(1980)].
- [20] H. P. Nilles and S. Raby, “Supersymmetry and the strong CP problem,” *Nucl. Phys.* **B198** (1982) 102–112.
- [21] J. E. Kim and H. P. Nilles, “The mu Problem and the Strong CP Problem,” *Phys. Lett.* **B138** (1984) 150.
- [22] P. Moxhay and K. Yamamoto, “Peccei-Quinn Symmetry Breaking by Radiative Corrections in Supergravity,” *Phys. Lett.* **B151** (1985) 363.
- [23] K. Rajagopal, M. S. Turner, and F. Wilczek, “Cosmological implications of axinos,” *Nucl. Phys.* **B358** (1991) 447–470.
- [24] E. J. Chun, J. E. Kim, and H. P. Nilles, “Axino mass,” *Phys. Lett.* **B287** (1992) 123–127, hep-ph/9205229.
- [25] J. E. Kim and M.-S. Seo, “Mixing of axino and goldstino, and axino mass,” *Nucl. Phys.* **B864** (2012) 296–316, 1204.5495.
- [26] S. Krippendorff, H. P. Nilles, M. Ratz, and M. W. Winkler, “Hidden SUSY from precision gauge unification,” *Phys.Rev.* **D88** (2013) 035022, 1306.0574.
- [27] A. Strumia, “Thermal production of axino Dark Matter,” *JHEP* **06** (2010) 036, 1003.5847.
- [28] K.-Y. Choi, L. Covi, J. E. Kim, and L. Roszkowski, “Axino Cold Dark Matter Revisited,” *JHEP* **04** (2012) 106, 1108.2282.