



This is the accepted manuscript made available via CHORUS. The article has been published as:

Universality in Higher Order Spin Noise Spectroscopy

Fuxiang Li and N. A. Sinitsyn

Phys. Rev. Lett. **116**, 026601 — Published 15 January 2016

DOI: [10.1103/PhysRevLett.116.026601](https://doi.org/10.1103/PhysRevLett.116.026601)

Universality in higher order spin noise spectroscopy

Fuxiang Li^{1,2}, and N. A. Sinitsyn^{2,*}

¹ *Center for Nonlinear Studies, Los Alamos National Laboratory, Los Alamos, NM 87545 USA and*

² *Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM 87545 USA*

Higher order time-correlators of spin fluctuations reveal considerable information about spin interactions. We argue that in a broad class of spin systems one can justify a phenomenological approach to explore such correlators. We predict that the 3rd and 4th order spin cumulants are described by a universal function that can be parametrized by a small set of parameters. We show that the fluctuation theorem constrains this function so that such correlators are fully determined by lowest nonlinear corrections to the free energy and the mean and variance of microscopic spin currents. We also provide an example of microscopic calculations for conduction electrons.

Spin noise spectroscopy (SNS) [1, 2] is a quickly evolving interdisciplinary field of research. It explores spin interactions by tracing dynamics of spontaneous spin fluctuations at or near the thermodynamic equilibrium without the need to intentionally polarize spins. The SNS has been successfully applied to semiconductors [3–5], quantum dots [6–8], hot and ultra-cold atomic gases [2, 9, 10].

So far the SNS has been focused on studies of the 2nd order spin correlator, $\langle S_z(t)S_z(0) \rangle$, of a local time-dependent spin polarization $S_z(t)$, or rather on its Fourier transform called the spin noise power spectrum:

$$C_2(\omega) \equiv \langle |S_z(\omega)|^2 \rangle, \quad (1)$$

where $S_z(\omega) = \frac{1}{\sqrt{T_m}} \int_0^{T_m} dt S_z(t) e^{i\omega t}$, and averaging is over many repeated time intervals of duration T_m . The information content of the correlator (1) is strongly restricted. Hence, one of the promising future directions to extend the SNS is to measure higher order spin correlators [11–14], the most accessible of which are the 3rd and 4th order ones:

$$C_3 \equiv \langle S_z(\omega_1)S_z(\omega_2)S_z(-\omega_1 - \omega_2) \rangle, \quad (2)$$

$$C_4 \equiv \langle |S_z(\omega_1)|^2 |S_z(\omega_2)|^2 \rangle - \langle |S_z(\omega_1)|^2 \rangle \langle |S_z(\omega_2)|^2 \rangle, \quad (3)$$

which depend on two frequencies, ω_1 and ω_2 .

Unlike the noise power (1) that describes the spectral frequency weights, the bi-spectra (2)-(3) tell how different frequencies ‘talk’ to each other. These correlators are sensitive to many-body interactions [15] and quantum effects [16], suggesting that their studies by the SNS can reveal essentially new information about correlated spin systems. As the noise of a single spin in a quantum dot [17] and the noise of only a few hundreds of spins of conduction electrons in a 2D electron gas [18] have already been studied experimentally, the goal to obtain the 3rd and 4th order spin correlators experimentally becomes achievable [16]. However, very little is known about properties of C_3 and C_4 in basic systems studied by the SNS, such as conduction electrons and atomic gases, e.g., about how (2)-(3) are influenced by the Pauli principle, scatterings, spin orbit coupling, and external magnetic field. There have been no quantum mechanically justified studies of such correlators in interacting electronic systems.

An important observation made throughout all known SNS applications is that the spin noise often shows well recognized patterns. For example, the noise power spectrum often consists of one or several peaks having the Lorentzian shape. The position and the width of such a peak determine useful parameters: the g-factor of the spin resonance and its life time [2]. This universality is not a coincidence and it is well understood: the Lorentzian shape of a peak indicates exponential relaxation in time that happens due to fast uncorrelated interactions. For example, conduction electrons that experience fast fluctuations of the spin-orbit field usually demonstrate a Lorentzian shape of the spin noise power spectrum [19]. In fact, the commonly used Bloch equation and various relaxation time approximations are justified by exactly this type of universality.

In this letter, we argue that a similar universality exists on the level of higher order spin correlators, namely, under the conditions that the noise power spectrum is Lorentzian, the 3rd and 4th spin correlators can also be parametrized by a small set of parameters with a clear physical meaning. In addition, we show that, as a consequence of the fluctuation theorem, such parameters are not independent when the system is probed at or near the thermodynamic equilibrium.

A Lorentzian peak in the noise power spectrum indicates that the dynamics of the spin fluctuation follows the Bloch-Langevin equation:

$$\dot{\mathbf{S}} = \mathbf{B} \times \mathbf{S} - \gamma \mathbf{S} + \boldsymbol{\eta}(t), \quad (4)$$

where γ is the relaxation rate, and $\boldsymbol{\eta}$ is the noise term:

$$\langle \boldsymbol{\eta} \rangle = 0, \quad \langle \eta_i(t) \eta_j(t') \rangle = 2D_{ij} \delta(t - t'); \quad i, j = x, y, z. \quad (5)$$

For simplicity, we assumed an isotropic relaxation rate, and we absorbed the g-factor in the definition of the magnetic field \mathbf{B} . Although phenomenological, Eq. (4) has been highly successful to describe diverse spin systems, including conduction electrons, nuclear spins, warm atomic gases, and hole spins of quantum dots [2, 3, 7, 9]. For small spontaneous fluctuations in a region with $N \gg 1$ spins, we can disregard the dependence of the relaxation rate γ on \mathbf{S} . The probability of a fluctuation $\mathbf{S} \neq 0$ at the thermodynamic equilibrium is determined by the

free energy $F(\mathbf{S})$ of spins in the region at known polarization \mathbf{S} :

$$P(\mathbf{S}) \sim e^{-\beta F(\mathbf{S})}, \quad F = \frac{a_0}{2} \mathbf{S}^2 + \frac{b_0}{4} \mathbf{S}^4 + \dots, \quad (6)$$

where $\beta = 1/k_B T$. Hence, D_{ij} are not always independent parameters. In order to reproduce the Gaussian part in (6), one should set $D_{ij} = D_0 \delta_{ij}$, where $D_0 = \gamma/(\beta a_0)$.

We will assume that the magnetic field is applied along the y -axis, which is transverse to the measurement z -axis. The noise power spectrum produced by Eq. (4) is then well known. It consists of two Lorentzian peaks [19]:

$$C_2(\omega) = \sum_{s=\pm} \frac{D_0}{(\omega - s\omega_L)^2 + \gamma^2}, \quad \omega_L \equiv |\mathbf{B}|. \quad (7)$$

Equation (4) with constant parameters γ and D_0 predicts zero values for all higher than 2nd order spin cumulants.

Our first observation is that it is straightforward to generalize Eq. (4) to include nonlinear effects. Correlators (2)-(3) must follow then from higher order corrections to parameters that, for an isotropic system, read

$$\gamma = \gamma(\mathbf{S}) = \gamma_0 + \gamma_2 \mathbf{S}^2 + O(\mathbf{S}^4), \quad (8)$$

$$D_{ij} = (D_0 + D_1 \mathbf{S}^2) \delta_{ij} + D_2 (S_i S_j - \mathbf{S}^2 \delta_{ij}) + O(\mathbf{S}^4) \quad (9)$$

Here D_1 corresponds to renormalization of the noise part related to dissipative spin relaxation, and the term with D_2 describes angular diffusion of a spin fluctuation without relaxation of its absolute value [20].

Microscopic calculations of the leading corrections to the nonlinear relaxation rate γ and the 2nd order noise correlator D_{ij} can be done within the approach developed in [15]. The biggest complication, however, is that the knowledge of $\gamma(\mathbf{S})$ and $D_{ij}(\mathbf{S})$ is insufficient to determine C_3 and C_4 because the noise term $\boldsymbol{\eta}$ in (4) can no longer be considered Gaussian, while a microscopic quantum theory to obtain non-Gaussian statistics of $\boldsymbol{\eta}$ is generally missing. Below we show, and this will be one of our key results, that this problem can be avoided for fluctuations at the thermodynamic equilibrium because non-Gaussian correlations of $\boldsymbol{\eta}$ are then uniquely constrained, i.e. they can be derived without additional microscopic calculations.

To address this problem, we note that, for a mesoscopic system, one can choose the time step δt such that the number of spin flips in the system is large but still much smaller than the typical size of the spin fluctuation \mathbf{S} . Let $P[\delta \mathbf{S}|\mathbf{S}]$ be the probability of observing the change of the spin polarization by the amount $\delta \mathbf{S}$ during δt given that initially the spin fluctuation size is \mathbf{S} . The law of large numbers guarantees that cumulants of $\delta \mathbf{S}$ grow linearly with δt as far as $\langle \delta \mathbf{S} \rangle \ll \mathbf{S}$ [21]. This fact can be expressed by introducing the cumulant generating function $\mathcal{H}(\boldsymbol{\chi}, \mathbf{S})$:

$$P[\delta \mathbf{S}|\mathbf{S}] = \int d\boldsymbol{\chi} e^{i\boldsymbol{\chi} \cdot \delta \mathbf{S}} e^{\delta t \mathcal{H}(\boldsymbol{\chi}, \mathbf{S})}. \quad (10)$$




FIG. 1. Fourth order cumulants in the presence of a magnetic field for different values of parameters: (a) $D_1 = 1$, $D_2 = 0$, $\omega_L = 0.0$; (b) $D_1 = 1$, $D_2 = 0$, $\omega_L = 0.1$; (c) $D_1 = 0$, $D_2 = 1$, $\omega_L = 0$; (d) $D_1 = 0$, $D_2 = 1$, $\omega_L = 0.1$. Other parameters are: $\gamma_0 = 0.1$, $a_0 = 1$, $b_0 = 0$, $\beta = 1$ and $\gamma_2 = 0$.

Next, we note that fluctuations near the thermodynamic equilibrium should satisfy the detailed balance constraints, so that probabilities of spin polarization changes by $\delta \mathbf{S}$ and $-\delta \mathbf{S}$ are related by [22, 23]

$$\frac{P[\delta \mathbf{S}|\mathbf{S}]}{\tilde{P}[-\delta \mathbf{S}|\mathbf{S} + \delta \mathbf{S}]} = e^{\beta(F(\mathbf{S}) - F(\mathbf{S} + \delta \mathbf{S}))} \approx e^{-\beta \delta \mathbf{S} \cdot \boldsymbol{\mu}}, \quad \boldsymbol{\mu} = \frac{\partial F}{\partial \mathbf{S}}, \quad (11)$$

where tilde sign in $\tilde{P}[-\delta \mathbf{S}|\mathbf{S} + \delta \mathbf{S}]$ means the probability at time-reversed values of model parameters. In combination with (10), Eq. (11) leads to the fluctuation relation (which is a special case of Eq. (3.2.51) in [24]):

$$\mathcal{H}(\boldsymbol{\chi}, -\mathbf{S}, -\mathbf{B}) = \mathcal{H}(\boldsymbol{\chi} + i\beta \boldsymbol{\mu}, \mathbf{S}, \mathbf{B}), \quad (12)$$

where \mathcal{H} is defined in (10). Next, we recall that if $\mathcal{H}(\boldsymbol{\chi}, \mathbf{S})$ is known, arbitrary spin correlator can be derived by the method of the stochastic path integral [15, 21]. The latter is the sum, Z , over all possible stochastic trajectories, discretized in time-steps δt , of random variables $\delta \mathbf{S}(t)$ and $\mathbf{S}(t)$ weighted by probabilities (10) and delta-functions $\delta(\dot{\mathbf{S}} - \delta \mathbf{S}/\delta t)$ at each elementary time interval. Integrals over $\delta \mathbf{S}(t)$ can be performed explicitly with an expense of introducing a conjugated to $\mathbf{S}(t)$ variable $\boldsymbol{\chi}(t)$. Following [15, 21], we find:

$$Z = \int D\mathbf{S}(t) \int D\boldsymbol{\chi}(t) e^{\int dt (i\boldsymbol{\chi} \cdot \dot{\mathbf{S}} + \mathcal{H}(\boldsymbol{\chi}, \mathbf{S}))}. \quad (13)$$

In order to derive an n -th order spin correlator in the $N \gg 1$ limit, it is enough to keep only terms up to the n -th power of variables $\boldsymbol{\chi}$, \mathbf{S} in $\mathcal{H}(\boldsymbol{\chi}, \mathbf{S})$. Assuming an isotropic paramagnetic system, the most general form of

$\mathcal{H}(\chi, \mathbf{S})$ up to the 4th power of variables is

$$\mathcal{H} = i\gamma(\mathbf{S})\chi \cdot \mathbf{S} + i\chi \cdot (\mathbf{S} \times \mathbf{B}) - \hat{D}(\mathbf{S}, \chi) + iD_3\chi^2(\mathbf{S} \cdot \chi) + D_4\chi^4, \quad (14)$$

where

$$\hat{D}(\mathbf{S}, \chi) = D_0\chi^2 + (D_1 - D_2)\mathbf{S}^2\chi^2 + D_2(\mathbf{S} \cdot \chi)^2. \quad (15)$$

Here constants D_0, D_1, D_2 have the same physical meaning as in (9), and $\gamma(\mathbf{S})$ is given by the two first terms in (8). Importantly, applying the symmetry (12) to (14), we find the standard fluctuation-dissipation theorem prediction $D_0 = \gamma_0/(\beta a_0)$, as well as the additional constraints relating higher order coefficients:

$$D_4 = \frac{\beta b_0 D_0 + \beta a_0 D_1 - \gamma_2}{\beta^3 a_0^3}, \quad D_3 = -2\beta a_0 D_4. \quad (16)$$

Equation (14) with constraints (16) are the central results of this article. They show that the information about up to the 4th order spin correlators is contained in a simple function \mathcal{H} that depends on a small set of parameters. Moreover, note that terms of higher than 2nd power of χ in (14) characterize higher order correlators of the noise term η in (4). The fact that corresponding coefficients D_3 and D_4 can be written in terms of coefficients at non-linear corrections to the 2nd order correlators of η and the quartic correction to the free energy in (6) considerably simplifies the goal of their microscopic calculation.

Remaining parameters depend on microscopic spin dynamics. Nevertheless, the universality that we found allows us to look at *possible patterns* of 3rd and 4th correlators. They can be calculated by switching to the frequency domain in the action of the path integral and treating 4th order terms in (14) as a small perturbation:

$$C_4(\omega_1, \omega_2) = \langle |S_z(\omega_1)|^2 |S_z(\omega_2)|^2 \mathcal{R}_4 \rangle_0 \quad (17)$$

where $\mathcal{R}_4 = \int dt \mathcal{H}_4$ and \mathcal{H}_4 is the quartic part of the Hamiltonian in Eq. (14), and $\langle \dots \rangle_0$ means that the average is taken over the quadratic action in the path integral. After applying the Wick's rule, we find a relatively complex expression that we provide in supplementary material [20]. In Fig. 1, we show examples of the obtained 4th correlator shapes at different values of independent parameters, including the magnetic field.

Analogously, we can explore the form of the 3rd order correlator. It becomes nonzero at nonequilibrium conditions, for example, when a finite spin density is induced in conduction electors by a resonant optical pumping. We will consider the limit of a weak intensity of the pumping beam so that spin generation happens in uncorrelated events that have a Poisson distribution, which contributes to the Hamiltonian in the action of Eq. (13) with a term [21]:

$$\mathcal{H}_p = k_p \left(e^{-i\chi_z} - 1 \right), \quad (18)$$

where k_p is the generation rate of spin polarization by an optical beam. Since the 3rd order correlator is zero

at $k_p = 0$, and since we are interested in the linear, in k_p , response contribution, we can disregard the effect of small renormalization of all other terms in the action of the path integral (13) on C_3 . The saddle point equations with the total Hamiltonian $\mathcal{H}_T = \mathcal{H} + \mathcal{H}_p$ read:

$$\delta\mathcal{H}_T/\delta\chi_i = 0, \text{ and } \delta\mathcal{H}_T/\delta S_i = 0, \text{ with } i = x, z.$$

They have a solution $\bar{\chi}_i = 0$ and

$$\bar{S}_z = \frac{k_p \gamma_0}{\gamma_0^2 + \omega_L^2}, \quad \bar{S}_x = \frac{k_p \omega_L}{\gamma_0^2 + \omega_L^2}. \quad (19)$$




FIG. 2. The real (a) and imaginary (b) parts of third order cumulant in the regime of a continuous spin pumping along z axis with $k_p = 0.3$. Parameters are: $D_1 = D_2 = 1$, $\gamma_0 = 0.1$, $\omega_L = 1$, $a_0 = 1$, $b_0 = 0$, $\beta = 1$ and $\gamma_2 = 0$.

By expanding the action in powers of small fluctuations $\delta\mathbf{S}$ and χ from the steady state, the 3rd order in $\delta\mathbf{S}$ and χ part of the Hamiltonian in the path integral reads:

$$\begin{aligned} \mathcal{H}_{T,3} = & -\frac{1}{2}k_p\chi_z^2\delta S_z - 2(D_1 - D_2)(\bar{\mathbf{S}}\delta\mathbf{S})\chi^2 \\ & - 2D_2(\bar{\mathbf{S}} \cdot \chi)(\chi \cdot \delta\mathbf{S}) + iD_3(\bar{\mathbf{S}} \cdot \chi)\chi^2 \\ & + i\gamma_2((\bar{\mathbf{S}} \cdot \chi)(\delta\mathbf{S})^2 + 2(\delta\mathbf{S} \cdot \bar{\mathbf{S}})(\chi \cdot \mathbf{S})). \end{aligned} \quad (20)$$

We find that corresponding correlator C_3 is generally complex valued. In Fig. 2(a) and (b) we plot, respectively, the real and imaginary parts of a typical pattern of $C_3(\omega_1, \omega_2)$.

As a demonstration of a microscopic estimate of independent parameters, we consider a model of conduction electrons, for which spin relaxation and fluctuations are caused by scatterings on weak randomly distributed static magnetic impurities. We assume that energy-momentum degrees of freedom of electrons equilibrate to the Fermi-Dirac distribution at the ambient temperature very quickly, while the spin degrees of freedom equilibrate at a much slower rate. For an open isotropic region around the studied spot, we can introduce a local instantaneous vector chemical potential $\mu(t)$ to describe the single particle density matrix of electrons:

$$\hat{\rho}(\epsilon) = \frac{1}{1 + e^{\beta(\epsilon - \mu \cdot \hat{\sigma}/2)}}, \quad (21)$$

where $\hat{\sigma}$ is the vector made of Pauli matrices acting in the spin space. The potential μ is a slow variable that changes at the spin relaxation time scale. We assume

that the observation region is much smaller than the spin diffusion length. The average spin polarization density is then given by $\mathbf{S}(t) = d_s \int d\epsilon \text{Tr}[\hat{\boldsymbol{\sigma}}\hat{\rho}]/2 = d_s \boldsymbol{\mu}(t)/2$, where d_s is the density of states per unit energy. Fast transitions of electronic spins through the observation region and quantum measurement effects will induce fast noise $\xi(t)$ in the measured Faraday rotation angle θ_F of the probe beam. However, averaging the signal over an interval δt that is smaller but comparable to spin relaxation time, one would find that $\langle \theta_F \rangle \sim S_z \delta t$, i.e. the information about S_z accumulates with time and dominates the measured correlations at spin relaxation time scale, while $\xi(t)$ contributes to the background noise and the high frequency tail of the spectrum, which we will not study here.

For a fermionic system with the density matrix (21), the free energy is a quadratic function of a parameter \mathbf{S} : $F = \int \boldsymbol{\mu} d\mathbf{S} = \mathbf{S}^2/(d_s)$. Let $\hat{\Psi} \equiv (\hat{a}_\uparrow, \hat{a}_\downarrow)$, where $\hat{a}_{\uparrow,\downarrow}$ are the annihilation operators of electrons in two degenerate eigenstates of the Hamiltonian that includes non-magnetic disorder. Weak interaction with magnetic impurities couples these states so that, in the Dirac picture, annihilation operators evolve with time:

$$\hat{\Psi}(\delta t) = (\cos \theta + i \mathbf{m} \cdot \hat{\boldsymbol{\sigma}} \sin \theta) \hat{\Psi}(0), \quad (22)$$

where θ is a random parameter of the evolution matrix, such that averaging over all scattering channels $\langle \theta^2 \rangle \sim \delta t$ and, for each scattering channel, \mathbf{m} is a randomly directed unit vector. Let $\hat{\mathbf{s}} = \frac{1}{2} \hat{\Psi}^\dagger \hat{\boldsymbol{\sigma}} \hat{\Psi}$ be the spin operator in the subspace of the two states. As we showed before, to determine parameters of the path integral action we need to know only up to 2nd order correlators of the spin change during a small time interval δt , which can be obtained by a simplified procedure, known to work only for such lowest order correlators [24]. Namely, we introduce the operator of the change of the spin: $\delta \hat{\mathbf{s}} \equiv \hat{\mathbf{s}}(\delta t) - \hat{\mathbf{s}}(0)$. Explicitly:

$$\begin{aligned} \delta \hat{\mathbf{s}} = & \frac{1}{2} \sin^2 \theta \left[\Psi^\dagger (\mathbf{m} \cdot \hat{\boldsymbol{\sigma}}) \hat{\boldsymbol{\sigma}} (\mathbf{m} \cdot \hat{\boldsymbol{\sigma}}) \Psi - \Psi^\dagger \hat{\boldsymbol{\sigma}} \Psi \right] \\ & + \frac{i}{2} \sin \theta \cos \theta \left[\Psi^\dagger \hat{\boldsymbol{\sigma}} (\mathbf{m} \cdot \hat{\boldsymbol{\sigma}}) \Psi - \Psi^\dagger (\mathbf{m} \cdot \hat{\boldsymbol{\sigma}}) \hat{\boldsymbol{\sigma}} \Psi \right]. \end{aligned}$$

Then the average change of the spin density and its variance are given by $\langle \delta \mathbf{S} \rangle = d_s \int d\epsilon \text{Tr}[\hat{\rho} \delta \hat{\mathbf{s}}]$ and $\langle \delta S_\alpha \delta S_\beta \rangle = d_s \int d\epsilon \text{Tr}[\hat{\rho} \{\delta \hat{s}_\alpha, \delta \hat{s}_\beta\}/2]$, where curly brackets denote the anticommutator. Using the density matrix (21) in the secondary quantized form, we obtain spin fluctuations due to scatterings between particular degenerate spin states. Finally, we assume that all scattering channels are independent during time δt , average over

parameters θ and \mathbf{m} , and integrate over ϵ . We find $\langle \delta \mathbf{S} \rangle / \delta t \equiv -\gamma_0 \mathbf{S} = -\frac{4}{3} \mathbf{S} \langle \theta^2 \rangle / \delta t$, and

$$\langle \delta S_\alpha \delta S_\beta \rangle = \delta_{\alpha\beta} \frac{4}{3} \langle \theta^2 \rangle \left(\frac{d_s}{\beta} + \frac{\beta}{3d_s} \mathbf{S}^2 \right).$$

Comparing with Eq. (4) and (9), we identify

$$D_0 = \gamma_0 d_s / (2\beta), \quad D_1 = \gamma_0 \beta / (6d_s), \quad D_2 = 0. \quad (23)$$

In this case, the coefficient in the free energy $a_0 = 2/d_s$. Therefore, we recover the fluctuation-dissipation relation $D_0 = \frac{\gamma_0}{a_0 \beta}$. Using Eq. (16), and considering the fact that $b_0 = 0$ and $\gamma_2 = 0$ we find:

$$D_4 = d_s \gamma_0 / (24\beta), \quad D_3 = -\gamma_0 / 6. \quad (24)$$

The resulting Hamiltonian in the path integral for this model reads:

$$\begin{aligned} \mathcal{H} = & i\gamma_0 \boldsymbol{\chi} \cdot \mathbf{S} + i\boldsymbol{\chi} \cdot (\mathbf{S} \times \mathbf{B}) - \frac{\gamma_0 d_s}{2\beta} \boldsymbol{\chi}^2 - \frac{\gamma_0 \beta}{6d_s} \mathbf{S}^2 \boldsymbol{\chi}^2 \\ & - i \frac{\gamma_0}{6} (\boldsymbol{\chi} \cdot \mathbf{S}) \boldsymbol{\chi}^2 + \frac{d_s \gamma_0}{24\beta} \boldsymbol{\chi}^4. \end{aligned} \quad (25)$$

Thus, this model corresponds to the case with $D_2 = 0$, which is illustrated in Fig. 1(a) and (b).

One additional consequence of this result is the prediction that the 4th correlator of a Fermi liquid scales linearly with temperature. For example, the D_1 -term contribution to the 4th cumulant, according to the Wick's theorem, is of the order of $C_4^{(1)} \sim D_1 \langle \mathbf{S}\boldsymbol{\chi} \rangle_0^2 \times \langle \mathbf{S}\mathbf{S} \rangle_0^2 \sim \beta \times 1 \times \frac{1}{\beta^2} = \frac{1}{\beta}$, where we used the fact that, for the second order correlators, the temperature dependence scales as $\langle \mathbf{S}\boldsymbol{\chi} \rangle_0 \sim \beta^0$ and $\langle \mathbf{S}\mathbf{S} \rangle_0 \sim \beta^{-1}$.

In conclusion, we developed a phenomenological approach that extends the Bloch-Langevin equation for spin dynamics to include the 3rd and 4th order spin correlations. This approach is justified by the law of large numbers and the higher order fluctuation relations. Our theory should be applicable practically to all spin systems, near the thermodynamics equilibrium, that exhibit Lorentzian peaks in the spin noise power spectrum. Such cases are ubiquitous. Therefore, although the microscopic theory of higher order spin correlations is at the early stage of development, our results make a valuable insight into the possible forms of such correlators and their dependence on temperature and optical spin pumping.

Acknowledgments. Authors thank A. V. Andreev for useful discussions. The work was carried out under the auspices of the National Nuclear Security Administration of the U.S. Department of Energy at Los Alamos National Laboratory under Contract No. DE-AC52-06NA25396. Authors also thank the support from LDRD program at LANL.

[1] E. Aleksandrov and V. Zapasskii, Zh. Eksp. Teor. Fiz **81**, 132 (1981).

[2] S. Crooker, D. Rickel, A. Balatsky, and D. Smith, Nature

- 431**, 49 (2004).
- [3] M. Oestreich, M. Römer, R. Haug, and D. Hagele, *Phys. Rev. Lett.* **95**, 216603 (2005).
 - [4] S. A. Crooker, L. Cheng, and D. L. Smith, *Phys. Rev. B* **79**, 035208 (2009).
 - [5] G.M.Müller, M. Römer, D. Schuh, W. Wegscheider, J. Hübner, and M. Oestreich, *Phys. Rev. Lett.* **101**, 206601 (2008).
 - [6] S. Crooker, J. Brandt, C. Sandfort, A. Greilich, D. Yakovlev, D. Reuter, A. Wieck, and M. Bayer, *Phys. Rev. Lett.* **104**, 036601 (2010).
 - [7] Y. Li, N. Sinitsyn, D. Smith, D. Reuter, A. Wieck, D. Yakovlev, M. Bayer, and S. Crooker, *Phys. Rev. Lett.* **108**, 186603 (2012).
 - [8] L. Yang, P. Glasenapp, A. Greilich, D. Reuter, A. Wieck, D. Yakovlev, M. Bayer, and S. Crooker, *Nat. Comm.* **5**, 4949 (2014).
 - [9] G. Katsoprinakis, A. Dellis, and I. Kominis, *Phys. Rev. A* **75**, 042502 (2007).
 - [10] V. S. Zapasskii, A. Greilich, S. A. Crooker, Y. Li, G. G. Kozlov, D. R. Yakovlev, D. Reuter, A. D. Wieck, and M. Bayer, *Phys. Rev. Lett.* **110**, 176601 (2013).
 - [11] S. Cundiff and S. Mukamel, *Phys. Today* July, 44 (2013).
 - [12] A. Bechtold, D. Rauch, F. Li, T. Simmet, P. Ardelt, A. Regler, K. Müller, N. A. Sinitsyn & J. J. Finley, *Nature Phys.* (2015).
 - [13] N. Ubbelohde, C. Fricke, C. Flindt, F. Hohls, and R. J. Haug, *Nat. Comm.* **3**, 612 (2012).
 - [14] A. M. Branczyk, D. B. Turner, and G. D. Scholes, *Annalen der Physik* **526**, 31 (2014).
 - [15] F. Li, A. Saxena, D. Smith, and N. A. Sinitsyn, *New J. Phys.* **15**, 113038 (2013).
 - [16] A. Bechtold, *et al.*, Preprint arXiv:1511.03684 (2015).
 - [17] R. Dahbashi, J. Hübner, F. Berski, K. Pierz, and M. Oestreich, *Phys. Rev. Lett.* **112**, 156601 (2014).
 - [18] S. V. Poltavtsev, *et al.*, *Phys. Rev. B* **89**, 081304 (2014).
 - [19] F. Li, Y. V. Pershin, V. A. Slipko, and N. A. Sinitsyn, *Phys. Rev. Lett.* **111**, 067201 (2013).
 - [20] Supplemental material.
 - [21] S. Pilgram, A. N. Jordan, E. V. Sukhorukov, and M. Büttiker, *Phys. Rev. Lett.* **90**, 206801(2003)
 - [22] L. Bertini, A. De Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim, *Rev. Mod. Phys.* **87**, 593 (2015).
 - [23] G. E. Crooks, *Phys. Rev. E* **60**, 2721 (1999).
 - [24] R. L. Stratonovich, *Nonlinear Nonequilibrium Thermodynamics I* (Springer, 1992).