

This is the accepted manuscript made available via CHORUS. The article has been published as:

Synthetic Spin-Orbit Coupling in an Optical Lattice Clock

Michael L. Wall, Andrew P. Koller, Shuming Li, Xibo Zhang, Nigel R. Cooper, Jun Ye, and

Ana Maria Rey

Phys. Rev. Lett. **116**, 035301 — Published 22 January 2016

DOI: [10.1103/PhysRevLett.116.035301](https://doi.org/10.1103/PhysRevLett.116.035301)

Synthetic spin-orbit coupling in an optical lattice clock

Michael L. Wall,^{1,2} Andrew P. Koller,^{1,2,3} Shuming Li,^{1,3} Xibo Zhang,^{1,3,*} Nigel R. Cooper,⁴ Jun Ye,^{1,3} and Ana Maria Rey^{1,2,3}

¹*JILA, NIST and University of Colorado, 440 UCB, Boulder, CO 80309, USA*

²*Center for Theory of Quantum Matter, University of Colorado, Boulder, Colorado 80309, USA*

³*Department of Physics, University of Colorado, 440 UCB, Boulder, CO 80309, USA*

⁴*T.C.M. Group, Cavendish Laboratory, J.J. Thomson Avenue, Cambridge CB3 0HE, United Kingdom*

(Dated: December 16, 2015)

We propose the use of optical lattice clocks operated with fermionic alkaline-earth-atoms to study spin-orbit coupling (SOC) in interacting many-body systems. The SOC emerges naturally during the clock interrogation when atoms are allowed to tunnel and accumulate a phase set by the ratio of the “magic” lattice wavelength to the clock transition wavelength. We demonstrate how standard protocols such as Rabi and Ramsey spectroscopy, that take advantage of the sub-Hertz resolution of state-of-the-art clock lasers, can perform momentum-resolved band tomography and determine SOC-induced s -wave collisions in nuclear spin polarized fermions. With the use of a second counter-propagating clock beam, we propose a method to engineer controlled atomic transport and study how it is modified by p - and s -wave interactions. The proposed spectroscopic probes provide clean and well-resolved signatures at current clock operating temperatures.

The recent implementation of synthetic gauge fields and spin-orbit coupling (SOC) in neutral atomic gases [1–7] is a groundbreaking step towards using these fully-controllable systems to synthesize and probe novel topological states of matter. So far optical Raman transitions have been used to couple different internal (e.g. hyperfine) states while transferring net momentum to the atoms. However, in alkali atoms Raman-induced spin-flips inevitably suffer from heating mechanisms associated with spontaneous emission. While this issue has not yet been an impediment for the investigation of non-interacting processes or mean field effects [4, 5], it could limit the ability to observe interacting many-body phenomena that manifest at longer timescales. Finding alternative, more resilient methods for generating synthetic SOC, and probing its interplay with interactions, is thus highly desirable.

To reduce heating, the use of atoms with richer internal structure such as alkaline-earth-atoms (AEAs) [3, 7] or lanthanide atoms such as Dy and Er [8] have been suggested. In addition to long-lived electronic states AEAs also offer accessible electronic state-dependent trapping potentials with applications that range from the generation of synthetic gauge fields [9] and particle number fractionalization [10, 11] to quantum information processing [12, 13]. So far most of the existing proposals require complicated experimental set-ups including laser-assisted tunneling and rectification protocols or have been targeted to study the non-interacting regime [7]. Here, we demonstrate that SOC emerges naturally in cold AEA optical lattice clocks (OLCs) [14] that use a “magic” (state independent) lattice. This is because the clock laser imprints a phase that varies significantly from one lattice site to the next as it drives an ultra-narrow optical transition. Our implementation (1) uses a direct transition to a long-lived electronic clock state with nat-




FIG. 1. (Color online) (a) A clock laser along the Z direction of wavelength λ and Rabi frequency Ω interrogates the $^1S_0(g)$ - $^3P_0(e)$ transition in fermionic alkaline-earth atoms trapped in an optical lattice with magic wavelength λ_m . The transverse confinement is provided by Gaussian curvature of the lattice beams with harmonic frequency $h\nu_{\perp}$. Many transverse modes n are populated at current operating temperatures. (b) The phase difference ϕ between adjacent sites j and $(j+1)$ induces SOC when atoms can tunnel with mode-dependent tunnel-coupling J_n , realizing a synthetic two-leg ladder with flux ϕ per plaquette.

ural lifetime $\gtrsim 10^2$ s [15], and thus heating from spontaneous emission is negligible; (2) takes advantage of the sub-Hz resolution of clock lasers [16–19]; and (3) accesses the regime where the interaction energy per particle, U , is weak compared to the characteristic trapping energies [20–25] but comparable to SOC scales determined by J , the tunneling, and Ω , the clock Rabi frequency. This regime allows us to investigate the interplay of interactions and SOC, to carry out controllable analytic calculations and make concrete predictions and to devise feasible interaction sensitive probing protocols accessible and testable in current experiments.

In the first protocol, we demonstrate that momentum-

resolved tomography of chiral band structures can be performed using Rabi spectroscopy in the parameter regime $J \sim \Omega \gg U$. In the second protocol we show that the modification of collisional properties by SOC [26, 27] is manifest in standard Ramsey spectroscopy, focusing on the parameter regime $J \gtrsim U$. In the final protocol, controlled and spatially resolved atomic transport [28–35] is induced by an additional counter-propagating clock beam that exhibits a controllable phase difference with respect to the original probe beam. Interaction effects beyond mean field are shown to modify the dynamics in the parameter regime $\Omega \gg J \sim U$. Inelastic collisions in the excited state impose limitations in the probing time [24, 25, 36, 37], however, they can be dynamically accounted for during clock interrogation as demonstrated in Refs. [20, 24]. We also show that losses can be used as a resource for state preparation and readout.

Current OLCs interrogate the $^1S_0(g)\text{--}^3P_0(e)$ transition of ensembles of thousands of nuclear spin polarized fermionic AEAs trapped in a deep 1D optical lattice that splits the gas in arrays of 2D pancakes [14] (see Fig. 1). The lattice potential uses the magic-wavelength, λ_m to generate identical trapping conditions for the two states. At current operating temperatures, $T \sim \mu\text{K}$ [20] the population of higher axial bands is negligible ($\lesssim 5\%$). On the other hand, along the transverse directions, where the confinement is provided wholly by the Gaussian curvature of the optical lattice beams, modes are thermally populated with an average number of mode quanta $\langle n \rangle \sim 50$. To generate SOC coherent tunneling between lattice sites is required. Our proposal is to superimpose a running-wave beam on the lattice potential:

$$V_{\text{ext}}(\mathbf{R}) = -\exp\left(-\frac{2R^2}{w_0^2}\right) \left[V_{\text{const}} + V \cos^2\left(\frac{2\pi Z}{\lambda_m}\right) \right]; \quad (1)$$

this increases the transverse confinement without significantly affecting the axial motion [40]. Here, w_0 is the beam waist, R the transverse radial coordinate, Z the axial coordinate, V the axial lattice corrugation, and V_{const} the running-wave induced potential. By increasing V_{const} as V is lowered, the transverse confinement frequency $\nu_{\perp} \sim \sqrt{\frac{1}{m\pi^2 w_0^2}(V_{\text{const}} + V)}$, with m the atom mass, is kept constant while the tunneling rate along the axial direction increases.

Since $V_{\text{ext}}(\mathbf{R})$ is discretely translationally invariant along the axial direction, atoms trapped in the lowest axial lattice band are governed by the Hamiltonian $\hat{H}^0 = \sum_{\mathbf{n}} \hat{H}_{\mathbf{n}}^0$ [41]

$$\hat{H}_{\mathbf{n}}^0 = \sum_{q,\alpha} E_{\alpha,\mathbf{n},q} \hat{n}_{\alpha,\mathbf{n},q} - \sum_q \left[\frac{\Omega_{\mathbf{n}}}{2} \hat{a}_{+,\mathbf{n},q+\phi}^\dagger \hat{a}_{-,\mathbf{n},q} + \text{H.c.} \right] \quad (2)$$

Throughout, $q = \tilde{q}a$ is the dimensionless product of axial quasimomentum \tilde{q} and lattice spacing $a = \lambda_m/2$, $\hat{a}_{\alpha,\mathbf{n},q}$ annihilates a fermion in the two-dimensional transverse mode \mathbf{n} , quasimomentum q , state $\alpha = \pm$ (for e and




FIG. 2. (Color online) (a) SOC band structure for $\delta = -2J_0$, $\Omega_0 = J_0$ (solid lines) and $\Omega_0^p = 0.05J_0$ (dashed lines). The axial depth is $V = 12$ recoils, $J_0/h = 42$ Hz and $\nu_{\perp} \approx 900$ Hz. Colors correspond to state character, with g (e) being more blue (red). (b) Chiral Bloch vector angle, θ_{0q} , in the xz plane extracted from Rabi spectroscopy using the protocol explained in the text [41]. The figure shows three temperatures for the parameters of (a).

g), and $\hat{n}_{\alpha,\mathbf{n},q} = \hat{a}_{\alpha,\mathbf{n},q}^\dagger \hat{a}_{\alpha,\mathbf{n},q}$. The energy $E_{\alpha,\mathbf{n},q}(q) = \alpha \frac{\delta}{2} + \bar{E}_{\mathbf{n}} - 2J_{\mathbf{n}} \cos(q)$ has contributions from the mode dependent tunneling $J_{\mathbf{n}}$, the average energy of the transverse mode \mathbf{n} , $\bar{E}_{\mathbf{n}}$, and the laser detuning δ . $\Omega_{\mathbf{n}}$ is the Rabi frequency for mode \mathbf{n} . The clock laser with wavelength λ imprints a phase that varies between adjacent lattice sites by $\phi = \pi\lambda_m/\lambda$.

If one views the two internal states as a discrete synthetic dimension [42, 43], as shown in Fig. 1, $\hat{H}_{\mathbf{n}}^0$ describes the motion of a charged particle on a two-leg ladder in a magnetic field with flux ϕ per lattice plaquette. \hat{H}^0 hence has the interpretation of many copies of those ladders – one for each transverse mode. By performing a gauge transformation $\hat{a}_{+,q+\phi,\mathbf{n}} \rightarrow \hat{a}_{+,q,\mathbf{n}}$, $\hat{H}_{\mathbf{n}}^0$ becomes diagonal in momentum space with the excited state dispersion shifted by ϕ , $q \rightarrow q + \phi$. The latter can be then conveniently written in terms of spin-1/2 operators acting on the populated modes

$$\hat{H}^0 = - \sum_{\mathbf{n},q} \mathbf{B}_{\mathbf{n}q} \cdot \hat{\mathbf{S}}_{\mathbf{n}q} \quad (3)$$

where $\mathbf{B}_{\mathbf{n}q} = (\Omega_{\mathbf{n}}, 0, \Delta E_{\mathbf{n}}(q, \phi) + \delta)$, $\Delta E_{\mathbf{n}}(q, \phi) \equiv 2J_{\mathbf{n}} [\cos(q) - \cos(q + \phi)]$, and $\hat{\mathbf{S}}_{\mathbf{n}q}^{x,y,z}$ are spin-1/2 angular momentum operators. The eigenstates are described by Bloch vectors pointing in the xz plane, with a direction specified by a single angle $\theta_{\mathbf{n}q} = \arctan\left(\frac{\Omega_{\mathbf{n}}}{\Delta E_{\mathbf{n}}(q, \phi) + \delta}\right)$ (Fig. 2(a)). The q dependence of this angle is a manifestation of chiral spin-momentum locking, which is directly connected to the topological chiral edge modes of the two-dimensional Harper-Hofstadter model [43, 44].

We first discuss the use of Rabi spectroscopy to probe the non-interacting chiral band structure of Eq. (3). In the regime $\Omega_0 \ll J_0$, there exists a finite window of δ s where the two dispersions cross at two special quasimomentum points $q_{\mathbf{n}}^*$ (Fig. 2(a)). The window, whose width

is $8J_n|\sin \frac{\phi}{2}|$, is signaled in the carrier linewidth, and thus when resolved it can be used to determine ϕ . At finite temperature, many transverse modes are populated and hence the dependence of Ω_n and J_n on \mathbf{n} could broaden the line and in general prevent an accurate determination of ϕ . However, a direct simulation of the Rabi lineshape using the potential Eq. (1) demonstrates that the features of the ideal, zero-temperature lineshape are captured even for a temperature of $3\mu\text{K}$ [41].

The ability to resolve the q_n^* resonances that appear across the entire Brillouin zone (BZ) as the detuning is varied can be used to perform momentum resolved spectroscopy and to precisely determine the chiral Bloch vector angle θ_{0q} for given values of Ω_0 and δ from Rabi oscillations. We propose the use of a three spectroscopic sequences [41]. One sequence selectively excites atoms at q_n^* from g to e and induces Rabi oscillations. Another filters the dynamics of the excited atoms from the remaining g atoms and the third is used to “correct” imperfections arising from finite temperature. As shown in Fig. 2(b), this protocol allows us to extract θ_{0q} over the BZ. This method is not restricted to OLCs, requiring only a stable probe, and complements other techniques for measuring band structures [45–47] used in degenerate Fermi gases.

To access the interplay between SOC and interactions we propose to use Ramsey spectroscopy (Fig. 3). The first pulse rotates the Bloch vector, initially pointing down, by an angle θ_1 , set by the pulse area, and the second converts the accumulated phase during the dark time τ into a $g - e$ population difference measured as Ramsey fringes. Interactions induce a density-dependent frequency shift in the fringes.

Interactions between two nuclear spin polarized atoms depend on the motional and electronic degrees of freedom [23]. When the atoms collide they experience s -wave interactions, characterized by the elastic scattering length a_{eg} , when their electronic state is antisymmetric ($|eg\rangle - |ge\rangle)/\sqrt{2}$. They can also collide via p -wave interactions, described by the corresponding p -wave elastic scattering volumes b_{gg}^3 , b_{ee}^3 , and b_{eg}^3 , in the three possible symmetric electronic configurations $|gg\rangle$, $|ee\rangle$, $(|eg\rangle + |ge\rangle)/\sqrt{2}$, respectively. In addition to elastic interactions, atoms can also exhibit inelastic collisions. In ^{87}Sr only the ee type has been observed to give rise to measurable losses [22, 41] while in $^{173,171}\text{Yb}$, both ee and eg losses have been reported [24, 25, 36].

When tunneling is suppressed the differential phase imparted by the laser is irrelevant, and as long as Ω_n is the same for all modes – a condition well-satisfied in current OLCs [20] – the collective spin of the atoms within each lattice site remains fully symmetric after the pulse and only p -wave collisions occur during the dark time. Measurements under this condition [20, 21] indeed observed a frequency shift linearly dependent on the excitation fraction of atoms, $(1 - \cos \theta_1)/2$ and fully consistent with




FIG. 3. (Color online) A clock laser pulse with pulse area θ_1 imprints a phase difference ϕ between atoms in neighboring sites. Atom tunneling, J_n , allows for s -wave interactions, $\propto U_{eg}^-$, which are signaled as a density shift in Ramsey spectroscopy after a second pulse of area θ_2 is applied.

a p -wave interacting model [23]. If instead tunneling is allowed during the dark time, atoms become sensitive to the spatially inhomogeneous spin rotation from the site-dependent laser phase, which in turn allows for s -wave collisions after a tunneling event (see Fig. 3). The s -wave collisions in SOC-coupled spin polarized fermions can lead to exotic phases of matter including topological quantum liquids [38, 39].

In the regime of weak interactions compared to tunneling, we compute the dynamics perturbatively, and find that SOC manifests itself in the density shift at short times as [41]

$$\Delta\nu = \Delta\nu_0 \left[1 + \frac{4\langle J^2 \rangle_{T_R} \tau^2 \zeta \cos \theta_1 \sin^2 \frac{\phi}{2}}{3\hbar^2(C - \chi \cos \theta_1)} \right], \quad (4)$$

where $\Delta\nu_0 = N(C - \chi \cos \theta_1)$ is the density shift in the absence of tunneling [20, 23], N the mean atom number per pancake, $\langle J^2 \rangle_{T_R}$ the thermally averaged squared tunneling rate, $\zeta = (V^{eg} - U^{eg})/2$, $\chi = (V^{ee} + V^{gg} - 2V^{ge})/2$, and $C = (V^{ee} - V^{gg})/2$, with $V^{\alpha\alpha'} = b_{\alpha\alpha'}^3 \langle P \rangle_{T_R}$ and $U^{eg} = a_{eg} \langle S \rangle_{T_R}$. Here, $\langle P \rangle_{T_R} \propto (T_R)^0$ and $\langle S \rangle_{T_R} \propto T_R^{-1}$ correspond to the thermal averages of the p -wave and s -wave mode overlap coefficients respectively [23], and T_R is the radial temperature. For the JILA ^{87}Sr clock operated at $T_R \sim (1 - 5)\mu\text{K}$ and $\theta_1 \ll \pi$ for $\tau \sim 80$ ms, $\Delta\nu_0 \sim -5\text{Hz}$. Since SOC introduces contributions from s -wave interactions, which can be one order of magnitude larger than p -wave at $T_R \sim 1\mu\text{K}$, then from Eq. 4 we expect significant modifications of the density shift. In the JILA ^{87}Sr clock the ratio of elastic to inelastic p -wave collision rates has been measured to be ~ 2 under typical conditions [20, 22, 41], and losses have been compensated during clock operation by dynamically tracking the population decay during the dark time [20, 21]. In Ramsey interrogated SOC systems with dominant s -wave elastic collisions, inelastic processes are expected to become even less relevant.

The third interrogation protocol we discuss is accomplished by generating a sliding superlattice via a pair of counter-propagating beams close to resonance with the clock transition and with a global phase difference $\Upsilon(t)$ which can be controlled in time (see Fig. 4(a)). The non-interacting Hamiltonian, written using a Wannier orbital

basis along the lattice direction, is

$$\hat{H}_L^0 = - \sum_{\mathbf{n}, j, \alpha} \left(J_{\mathbf{n}} \left[\hat{a}_{\alpha, \mathbf{n}, j}^\dagger \hat{a}_{\alpha, \mathbf{n}, j+1} + \text{H.c.} \right] - \frac{\delta}{2} \alpha \hat{n}_{\alpha, \mathbf{n}, j} \right) - \sum_{\mathbf{n}, j} \left(\Omega_{\mathbf{n}} \cos(\Upsilon(t) - j\phi) \left[\hat{a}_{+, \mathbf{n}, j}^\dagger \hat{a}_{-, \mathbf{n}, j} + \text{H.c.} \right] \right). \quad (5)$$

As $\Upsilon(t)$ is changed, the clock laser standing wave “slides” with respect to the optical lattice. This allows for the minimum realization of a topological pump when $\Upsilon(t)$ is adiabatically varied from $0 \rightarrow 2\pi$ [28, 32–35]. In the weak tunneling limit the quantized nature of particle transport can be directly linked to spatially isolated tunneling resonances [28]. We now show how those resonances can be spectroscopically measured in OLCs.

Let us first consider the case $J_{\mathbf{n}} = \delta = 0$ and set $\phi = 7\pi/6$, relevant for the ^{87}Sr system. We write $\Upsilon = (2\pi s + \Delta\Upsilon)/12$, with $0 \leq \Delta\Upsilon < 2\pi$ and s an integer. Under these conditions, the localized dressed eigenstates are spin-polarized along $\pm x$ alternating between neighboring sites except for “defects” at $j_d(r) = 6r + 3 + s$ (r an integer) when $\cos(\Upsilon - \Delta\Upsilon/12 - j_d\phi) = 0$ and the ground states at $j_d(r)$ and $j_d(r) + 1$ point along the same direction. Since tunneling preserves polarization, it is suppressed when $J_{\mathbf{n}} \ll \Omega_{\mathbf{n}}$ due to the energy offset $\sim \Omega$ between neighboring sites. The one exception is the case $\Delta\Upsilon = \pi$ where the defect pair $j_d(r)$ and $j_d(r) + 1$ is resonantly tunnel-coupled. Quantized transport occurs when $\Delta\Upsilon$ is slowly varied across the resonance. Instead of adiabatic transport we propose to spectroscopically resolve the resonance using a Ramsey-type protocol. Here, atoms initially prepared in g at $|\delta| \gg \Omega_0$ are adiabatically transferred to the ground dressed state by slowly turning off δ at $\Upsilon = (2\pi s + \Delta\Upsilon_p)/12$ with $\Delta\Upsilon_p \neq \pi$. Then Υ is quenched so resonant tunneling is allowed, $s \rightarrow s + 1$ and $\Delta\Upsilon = \pi$, and the system evolves for a time τ . Following this evolution tunneling is turned off and the phase switched back, $s + 1 \rightarrow s$ and $\Delta\Upsilon \rightarrow \Delta\Upsilon_p$. Those atoms which have tunneled at the resonant sites are now in an excited state of the local dressed basis. These excitations can be measured by adiabatically converting the dressed excitations to “bare” e excitations by adiabatically ramping on $\delta \gg \Omega_0$; this leads to a measurable excited state population $n_e(t)$. In Fig. 4(c) we show these tunneling resonances are clearly observable even at finite temperature. Moreover, since resonances are spatially well-separated (every six sites), they can be resolved with low-resolution imaging.

Interactions modify the transport dynamics in this pumping protocol. The sliding superlattice simplifies the treatment of interactions by isolating resonant site pairs $\{j_d, j_d + 1\}$. We consider the case where at most two atoms occupy the resonant sites, a condition which can be achieved by decreasing the atomic density with a large-volume dipole trap [16]. The two particle states can be classified in terms of the atoms’ spin polarization along x




FIG. 4. (a) An additional counter-propagating probing beam with a differential phase Υ generates a sliding superlattice potential, shown for $\phi = 7\pi/6$, corresponding to an ^{87}Sr OLC. For weak tunneling $J \ll \Omega$ transport is energetically suppressed except at resonant defect points (circled). (b) Two-particle interaction sectors classified by the total polarization M_x and spatial symmetry of the dressed states. An oval (figure-eight) denotes a symmetric (antisymmetric) spatial wavefunction. (c) Dynamics of a single particle at the tunneling resonance for two temperatures (red solid and blue dashed) and an off-resonant site (black dotted) for $J_0/h = 8\text{Hz}$, $\Omega_0/h = 1\text{kHz}$, $\phi = 7\pi/6$. (d) Normalized differential excitation extracted as explained in the text with the interaction parameters of Ref. [22]. The $M_x = 0$ components (solid lines) involve the s -wave sector, and so display a strong dependence of contrast on temperature, while the $M_{\pm 1}$ sectors (dashed lines) experience only weaker single-particle thermal dephasing.

in four sectors: three symmetric ones (triplets) with total x polarization $M_x = +1, -1, 0$ and a $M_x = 0$ singlet. Within each sector, the dressed states’ interaction parameters are given by [41] $V_{+1} = V_{-1} = (V^{gg} + V^{ee} + 2V^{eg})/4$, $V_0 = (V^{gg} + V^{ee})/2$ and $U_0 = U^{eg}$ (see Fig. 4(b)). Since the p -wave parameters are not $\text{SU}(2)$ symmetric, i.e. $b_{ee} \neq b_{gg} \neq b_{eg}$, the triplet sectors are coupled. However, in the weakly interacting limit $V^{\alpha\beta} \ll \Omega$, the triplets are separated by energy gaps $\sim \Omega$ and coupling between the sectors can be neglected. The singlet sector is always decoupled from the triplets.

Within each sector interactions modify the dynamics, making it sensitive to temperature and density. The modifications can be extracted by performing measurements of the atom number-normalized excitation fraction for different densities and then differentiating the high-density and low-density results [41]. This procedure removes the single-particle contribution and is particularly suitable for characterizing the role of interactions in clock experiments [20, 22]. The normalized differential excitation is shown in Fig. 4(d). For the adiabatic dressed state preparation, all four manifolds, and hence all interaction parameters $V_{\pm,0}, U_0$, contribute to the dynamics. A filtering protocol that uses the ee losses can be used to separate the dynamics of the various sectors. For example, by transferring all atoms to the e state and holding before

the adiabatic ground state preparation, the doubly occupied $M_x = \pm 1$ triplet sectors will be removed and only the $M_x = 0$ singlet and triplet remain and contribute (here the ground dressed states have one atom at j_d and $j_d + 1$). As shown in Fig. 4 the dynamics of the $V_{\pm,0}$ and U_0 sectors can be distinguished by the different scaling of the p - and s -wave interaction parameters with temperature T_R , $V_{\pm,0}(T_R) \sim \text{const.}$ and $U_0(T_R) \sim T_R^{-1}$ [23] (See Fig.4(d)). By comparing these dynamics to that without the holding time, information about the $M_x = \pm 1$ dynamics can be extracted. In general s -wave interactions, purely elastic for nuclear-spin-polarized Sr, dominate the normalized differential contrast, with p -wave contributions (including losses) relevant only at hotter temperatures.

Summary We have described approaches to implement and probe SOC and its interplay with s - and p - wave interactions in OLCs. These open the unique opportunity of exploring, for the first time, a controllable interplay of interactions, spin-orbit coupling, and non-trivial topology in regimes where a complete theoretical treatment can also be performed. While the described protocols work at the temperatures achievable in current OLCs, their sensitivity and applicability are expected to significantly improve when operating the clock in the quantum degenerate regime. Moreover, if nuclear spins are included, they can open a window for the investigation of SOC with SU(N)-symmetric collisions [48].

Acknowledgments We thank Ed Marti and James Thompson for comments on the manuscript, and Leonid Isaev and the JILA Sr clock team for discussions. This work was supported by the NSF (PIF-1211914 and PFC-1125844), AFOSR, AFOSR-MURI, NIST and ARO, the EPSRC Grant EP/K030094/1, and by the JILA Visiting Fellows Program. MLW thanks the NRC postdoctoral fellowship program for support. AK was supported by the Department of Defense through the NDSEG program.

* Present Address: International Center for Quantum Materials, School of Physics, Peking University, Beijing 100871, China

- [1] V. Galitski and I. B. Spielman, Spin-orbit coupling in quantum gases, *Nature* **494**, 49–54 (2013).
- [2] N. Cooper, Quantum Hall States of Ultracold Gases, In *Many-Body Physics with Ultracold Gases: Lecture Notes of the Les Houches Summer School: Volume 94, 28 June – 23 July 2010*, Salomon, C., Shlyapnikov, G. V., and Cugliandolo, L. F., editors. Oxford University Press, Oxford (2013).
- [3] J. Dalibard, F. Gerbier, G. Juzeliūnas, and P. Öhberg, Colloquium: Artificial gauge potentials for neutral atoms, *Rev. Mod. Phys.* **83**, 1523–1543 (2011).
- [4] N. Goldman, G. Juzeliūnas, P. Öhberg, and I. B. Spielman, Light-induced gauge fields for ultracold atoms, *Rep. Prog. Phys.* **77** 126401 (2014).
- [5] H. Zhai, Degenerate quantum gases with spin-orbit coupling: a review *Rep. Prog. Phys.* **78**, 026001 (2015).
- [6] B. K. Stuhl, H.-I. Lu, L. M. Ayccock, D. Genkina, and I. B. Spielman, Visualizing edge states with an atomic Bose gas in the quantum Hall regime, arxiv:1502.02496 (2015).
- [7] M. Mancini, G. Pagano, G. Cappellini, L. Livi, M. Rider, J. Catani, C. Sias, P. Zoller, M. Inguscio, M. Dalmonte, and L. Fallani, Observation of chiral edge states with neutral fermions in synthetic Hall ribbons, arXiv:1502.02495 (2015).
- [8] X. Cui, B. Lian, T.L. Ho, B. Lev, and H. Zhai, Synthetic gauge field with highly magnetic lanthanide atoms *Phys. Rev. A*, **88**, 011601 (2013).
- [9] F. Gerbier and J. Dalibard, *New J. Phys.* **12**, 033007 (2010).
- [10] Janne Ruostekoski, Gerald V. Dunne, and Juha Javanainen, *Phys. Rev. Lett.* **88**, 180401 (2002).
- [11] J. Ruostekoski, J. Javanainen, and G. V. Dunne, *Phys. Rev. A* **77**, 013603 (2008).
- [12] Andrew J. Daley, Martin M. Boyd, Jun Ye, and Peter Zoller, *Phys. Rev. Lett.* **101**, 170504 (2008).
- [13] A. V. Gorshkov, A. M. Rey, A. J. Daley, M. M. Boyd, J. Ye, P. Zoller, and M. D. Lukin, *Phys. Rev. Lett.* **102**, 110503 (2009).
- [14] Andrew D. Ludlow, Martin M. Boyd, Jun Ye, E. Peik, and P. O. Schmidt, Optical atomic clocks, *Rev. Mod. Phys.* **87**, 637 (2015).
- [15] M. M. Boyd, T. Zelevinsky, A. D. Ludlow, S. M. Foreman, S. Blatt, T. Ido, and J. Ye, *Science* **314**, 1430 (2006).
- [16] B. J. Bloom, *et al.*, *Nature* **506**, 71 (2014).
- [17] T. L. Nicholson, *et al.*, *Nat. Commun.* **6**, 6896 (2015).
- [18] N. Hinkley, *et al.*, *Science* **341**, 1215 (2013).
- [19] U. Ichiro *et al.*, Cryogenic optical lattice clocks, *Nat. Photon.*, **9**, 185-189 (2015).
- [20] M. J. Martin, M. Bishof, M. D. Swallows, X. Zhang, C. Benko, J. von-Stecher, A. V. Gorshkov, A. M. Rey, and Jun Ye, A quantum many-body spin system in an optical lattice clock, *Science* **341**, 632 - 636 (2013).
- [21] N. D. Lemke, J. von Stecher, J. A. Sherman, A. M. Rey, C. W. Oates, and A. D. Ludlow, p -Wave cold collisions in an optical lattice clock, *Phys. Rev. Lett.* **107**, 103902 (2011).
- [22] X. Zhang, M. Bishof, S. L. Bromley, C. V. Kraus, M. S. Safronova, P. Zoller, A. M. Rey, and J. Ye, Spectroscopic observation of SU(N)-symmetric interactions in Sr orbital magnetism, *Science* **345**, 1467 (2014).
- [23] A. M. Rey, A.V. Gorshkov, C.V. Kraus, M. J. Martin, M. Bishof, M. D. Swallows, X. Zhang, C. Benko, J. Ye, N.D. Lemke, and A.D. Ludlow, Probing many-body interactions in an optical lattice clock, *Annals Phys.* **340**, 311 - 351 (2014).
- [24] M. Bishof *et al.*, Inelastic collisions and density-dependent excitation suppression in a 87Sr optical lattice clock, *Phys. Rev. A* **84**, 052716 (2011).
- [25] A. D. Ludlow, N. D. Lemke, J. A. Sherman, C. W. Oates, G. Qumner, J. von Stecher, and A. M. Rey, Cold-collision-shift cancellation and inelastic scattering in a Yb optical lattice clock, *Phys. Rev. A* **84**, 052724 (2011).
- [26] R. A. Williams *et al.*, Synthetic Partial Waves in Ultracold Atomic Collisions, *Science* **335**, 314-317 (2012).
- [27] Z. Fu *et al.*, Production of Feshbach molecules induced

- by spin-orbit coupling in Fermi gases, *Nature Physics* **10**, 110 (2014).
- [28] N. R. Cooper and A. M. Rey, Adiabatic Control of Atomic Dressed States for Transport and Sensing, *Phys. Rev. A* **92**, 021401(R) (2015).
 - [29] Michael Lohse, Christian Schweizer, Oded Zilberberg, Monika Aidelsburger, and Immanuel Bloch, A Thouless Quantum Pump with Ultracold Bosonic Atoms in an Optical Superlattice, arXiv:1507.02225 (2015).
 - [30] Shuta Nakajima, Takafumi Tomita, Shintaro Taie, Tomohiro Ichinose, Hideki Ozawa, Lei Wang, Matthias Troyer, and Yoshiro Takahashi, Topological Thouless Pumping of Ultracold Fermions, arXiv:1507.02223 (2015).
 - [31] Hsin-I Lu *et al.* Geometrical pumping with a Bose-Einstein condensate, arXiv:1508.04480 (2015).
 - [32] D. J. Thouless, Quantization of particle transport, *Phys. Rev. B* **27**, 6083 (1983).
 - [33] Lei Wang, Matthias Troyer, and Xi Dai, Topological Charge Pumping in a One-Dimensional Optical Lattice, *Phys. Rev. Lett.* **111**, 026802 (2013).
 - [34] Feng Mei *et al.* Topological insulator and particle pumping in a one-dimensional shaken optical lattice, *Phys. Rev. A* **90**, 063638 (2014).
 - [35] Ran Wei and Erich J. Mueller, Anomalous charge pumping in a one-dimensional optical superlattice, *Phys. Rev. A* **92**, 013609 (2015).
 - [36] F. Scazza, *et al.* Observation of two-orbital spin-exchange interactions with ultracold SU(N)-symmetric fermions *Nature Phys.* **10**, 779-784 (2014).
 - [37] F. Cappelinni, *et al.* Direct observation of coherent inter-orbital spin-exchange dynamics *Phys. Rev. Lett.* **113**, 120402 (2014).
 - [38] C. Zhang, S. Tewari, R. M. Lutchyn, and S. Das Sarma, $p_x + ip_y$ Superfluid from s -wave Interactions of Fermionic Cold Atoms, *Phys. Rev. Lett.* **101**, 160401 (2008).
 - [39] B. Juliá-Díaz, T. Groß, O. Dutta, D.E. Chang, and M. Lewenstein, Engineering p -wave interactions in ultracold atoms using nanoplasmonic traps, *Nat. Comm.* **4**, 2046 (2013).
 - [40] Michael L. Wall, Kaden R. A. Hazzard, and Ana Maria Rey, Effective many-body parameters for atoms in non-separable Gaussian optical potentials, *Phys. Rev. A* **92**, 013610 (2015).
 - [41] See the Supplementary Material at [xxx] for a derivation of the density shift in Ramsey spectroscopy, the extraction of the chiral Bloch vector angle in Rabi spectroscopy, and the interaction-induced contrast decay with the sliding clock superlattice.
 - [42] A. Celi, P. Massignan, J. Ruseckas, N. Goldman, I.B. Spielman, G. Juzeliunas, and M. Lewenstein, Synthetic gauge fields in synthetic dimensions, *Phys. Rev. Lett.* **112**, 043001 (2014).
 - [43] Dario Hügél and Belén Paredes, Chiral Ladders and the Edges of Chern Insulators, *Phys. Rev. A* **89**, 023619 (2014).
 - [44] Y. Hatsugai, Topological aspects of the quantum Hall effect, *J. Phys.: Condens. Matter* **9**, 2507 (1997).
 - [45] L. Tarruell, *et al.*, Creating, moving and merging Dirac points with a Fermi gas in a tunable honeycomb lattice, *Nature* **483**, 302 (2012).
 - [46] M. Atala, *et al.*, Direct measurement of the Zak phase in topological Bloch bands, *Nat. Phys.* **9**, 795 (2013).
 - [47] L. Duca, *et al.*, An Aharonov-Bohm interferometer for determining Bloch band topology, *Science* **347**, 288 (2015).
 - [48] M. Cazalilla and A. M. Rey, Ultracold Fermi Gases with Emergent SU(N) Symmetry, *Rep. Prog. Phys.* **77**, 124401 (2014).