

This is the accepted manuscript made available via CHORUS. The article has been published as:

First Observation of the Doubly Cabibbo-Suppressed Decay of a Charmed Baryon: $\Lambda_c^+ \rightarrow p K^+ \pi^-$

S. B. Yang *et al.* (Belle Collaboration)

Phys. Rev. Lett. **117**, 011801 — Published 27 June 2016

DOI: [10.1103/PhysRevLett.117.011801](https://doi.org/10.1103/PhysRevLett.117.011801)

First Observation of the Doubly Cabibbo-Suppressed Decay of a Charmed Baryon:

$$\Lambda_c^+ \rightarrow p K^+ \pi^-$$

S. B. Yang,⁵⁷ K. Tanida,⁵⁷ B. H. Kim,⁵⁷ I. Adachi,^{15,12} H. Aihara,⁷⁰ D. M. Asner,⁵⁴ V. Aulchenko,^{3,52} T. Aushev,⁴¹ V. Babu,⁶⁴ I. Badhrees,^{63,28} A. M. Bakich,⁶² E. Barberio,³⁹ V. Bhardwaj,⁶⁰ B. Bhuyan,¹⁸ J. Biswal,²⁵ G. Bonvicini,⁷⁵ A. Bozek,⁴⁸ M. Bračko,^{37,25} T. E. Browder,¹⁴ D. Červenkov,⁴ V. Chekelian,³⁸ A. Chen,⁴⁵ B. G. Cheon,¹³ K. Chilikin,⁴⁰ R. Chistov,⁴⁰ K. Cho,²⁹ V. Chobanova,³⁸ Y. Choi,⁶¹ D. Cinabro,⁷⁵ J. Dalseno,^{38,65} M. Danilov,⁴⁰ N. Dash,¹⁷ Z. Doležal,⁴ Z. Drásal,⁴ D. Dutta,⁶⁴ S. Eidelman,^{3,52} H. Farhat,⁷⁵ J. E. Fast,⁵⁴ T. Ferber,⁷ B. G. Fulsom,⁵⁴ N. Gabyshev,^{3,52} A. Garmash,^{3,52} V. Gaur,⁶⁴ R. Gillard,⁷⁵ Y. M. Goh,¹³ P. Goldenzweig,²⁷ D. Greenwald,⁶⁶ J. Grygier,²⁷ J. Haba,^{15,12} P. Hamer,¹¹ T. Hara,^{15,12} K. Hayasaka,⁴³ H. Hayashii,⁴⁴ W.-S. Hou,⁴⁷ T. Iijima,^{43,42} K. Inami,⁴² G. Inguglia,⁷ A. Ishikawa,⁶⁸ R. Itoh,^{15,12} Y. Iwasaki,¹⁵ W. W. Jacobs,²⁰ I. Jaegle,¹⁴ H. B. Jeon,³² K. K. Joo,⁵ T. Julius,³⁹ K. H. Kang,³² E. Kato,⁶⁸ P. Katrenko,⁴¹ C. Kiesling,³⁸ D. Y. Kim,⁵⁹ H. J. Kim,³² J. B. Kim,³⁰ K. T. Kim,³⁰ M. J. Kim,³² S. H. Kim,¹³ S. K. Kim,⁵⁷ Y. J. Kim,²⁹ K. Kinoshita,⁶ N. Kobayashi,⁷¹ P. Kodyš,⁴ S. Korpar,^{37,25} P. Križan,^{34,25} P. Krokovny,^{3,52} T. Kuhr,³⁵ A. Kuzmin,^{3,52} Y.-J. Kwon,⁷⁷ J. S. Lange,⁹ I. S. Lee,¹³ C. H. Li,³⁹ H. Li,²⁰ L. Li,⁵⁶ Y. Li,⁷⁴ L. Li Gioi,³⁸ J. Libby,¹⁹ D. Liventsev,^{74,15} M. Lubej,²⁵ M. Masuda,⁶⁹ D. Matvienko,^{3,52} K. Miyabayashi,⁴⁴ H. Miyata,⁵⁰ R. Mizuk,^{40,41} G. B. Mohanty,⁶⁴ A. Moll,^{38,65} H. K. Moon,³⁰ R. Mussa,²⁴ E. Nakano,⁵³ M. Nakao,^{15,12} T. Nanut,²⁵ K. J. Nath,¹⁸ M. Nayak,¹⁹ K. Negishi,⁶⁸ M. Niiyama,³¹ N. K. Nisar,^{64,1} S. Nishida,^{15,12} S. Ogawa,⁶⁷ S. Okuno,²⁶ S. L. Olsen,⁵⁷ G. Pakhlova,⁴¹ B. Pal,⁶ C. W. Park,⁶¹ H. Park,³² T. K. Pedlar,³⁶ R. Pestotnik,²⁵ M. Petrič,²⁵ L. E. Piilonen,⁷⁴ C. Pulvermacher,²⁷ J. Rauch,⁶⁶ M. Ritter,³⁵ A. Rostomyan,⁷ S. Ryu,⁵⁷ H. Sahoo,¹⁴ Y. Sakai,^{15,12} S. Sandilya,⁶⁴ L. Santelj,¹⁵ T. Sanuki,⁶⁸ Y. Sato,⁴² V. Savinov,⁵⁵ T. Schlüter,³⁵ O. Schneider,³³ G. Schnell,^{2,16} C. Schwanda,²² A. J. Schwartz,⁶ Y. Seino,⁵⁰ K. Senyo,⁷⁶ O. Seon,⁴² I. S. Seong,¹⁴ M. E. Sevier,³⁹ V. Shebalin,^{3,52} T.-A. Shibata,⁷¹ J.-G. Shiu,⁴⁷ B. Shwartz,^{3,52} F. Simon,^{38,65} Y.-S. Sohn,⁷⁷ A. Sokolov,²³ S. Stanič,⁵¹ M. Starič,²⁵ J. Stypula,⁴⁸ M. Sumihama,¹⁰ T. Sumiyoshi,⁷² M. Takizawa,⁵⁸ U. Tamponi,^{24,73} Y. Teramoto,⁵³ K. Trabelsi,^{15,12} V. Trusov,²⁷ M. Uchida,⁷¹ T. Uglov,⁴¹ Y. Unno,¹³ S. Uno,^{15,12} P. Urquijo,³⁹ Y. Usov,^{3,52} P. Vanhoefer,³⁸ G. Varner,¹⁴ K. E. Varvell,⁶² A. Vinokurova,^{3,52} A. Vossen,²⁰ M. N. Wagner,⁹ C. H. Wang,⁴⁶ M.-Z. Wang,⁴⁷ P. Wang,²¹ X. L. Wang,⁷⁴ Y. Watanabe,²⁶ K. M. Williams,⁷⁴ E. Won,³⁰ J. Yamaoka,⁵⁴ S. Yashchenko,⁷ H. Ye,⁷ J. Yelton,⁸ C. Z. Yuan,²¹ Y. Yusa,⁵⁰ Z. P. Zhang,⁵⁶ V. Zhilich,^{3,52} V. Zhulanov,^{3,52} and A. Zupanc^{34,25}

(The Belle Collaboration)

¹Aligarh Muslim University, Aligarh 202002

²University of the Basque Country UPV/EHU, 48080 Bilbao

³Budker Institute of Nuclear Physics SB RAS, Novosibirsk 630090

⁴Faculty of Mathematics and Physics, Charles University, 121 16 Prague

⁵Chonnam National University, Kwangju 660-701

⁶University of Cincinnati, Cincinnati, Ohio 45221

⁷Deutsches Elektronen-Synchrotron, 22607 Hamburg

⁸University of Florida, Gainesville, Florida 32611

⁹Justus-Liebig-Universität Gießen, 35392 Gießen

¹⁰Gifu University, Gifu 501-1193

¹¹II. Physikalisches Institut, Georg-August-Universität Göttingen, 37073 Göttingen

¹²SOKENDAI (The Graduate University for Advanced Studies), Hayama 240-0193

¹³Hanyang University, Seoul 133-791

¹⁴University of Hawaii, Honolulu, Hawaii 96822

¹⁵High Energy Accelerator Research Organization (KEK), Tsukuba 305-0801

¹⁶IKERBASQUE, Basque Foundation for Science, 48013 Bilbao

¹⁷Indian Institute of Technology Bhubaneswar, Satya Nagar 751007

¹⁸Indian Institute of Technology Guwahati, Assam 781039

¹⁹Indian Institute of Technology Madras, Chennai 600036

²⁰Indiana University, Bloomington, Indiana 47408

²¹Institute of High Energy Physics, Chinese Academy of Sciences, Beijing 100049

²²Institute of High Energy Physics, Vienna 1050

²³Institute for High Energy Physics, Protvino 142281

- ²⁴INFN - Sezione di Torino, 10125 Torino
²⁵J. Stefan Institute, 1000 Ljubljana
²⁶Kanagawa University, Yokohama 221-8686
²⁷Institut für Experimentelle Kernphysik, Karlsruher Institut für Technologie, 76131 Karlsruhe
²⁸King Abdulaziz City for Science and Technology, Riyadh 11442
²⁹Korea Institute of Science and Technology Information, Daejeon 305-806
³⁰Korea University, Seoul 136-713
³¹Kyoto University, Kyoto 606-8502
³²Kyungpook National University, Daegu 702-701
³³École Polytechnique Fédérale de Lausanne (EPFL), Lausanne 1015
³⁴Faculty of Mathematics and Physics, University of Ljubljana, 1000 Ljubljana
³⁵Ludwig Maximilians University, 80539 Munich
³⁶Luther College, Decorah, Iowa 52101
³⁷University of Maribor, 2000 Maribor
³⁸Max-Planck-Institut für Physik, 80805 München
³⁹School of Physics, University of Melbourne, Victoria 3010
⁴⁰Moscow Physical Engineering Institute, Moscow 115409
⁴¹Moscow Institute of Physics and Technology, Moscow Region 141700
⁴²Graduate School of Science, Nagoya University, Nagoya 464-8602
⁴³Kobayashi-Maskawa Institute, Nagoya University, Nagoya 464-8602
⁴⁴Nara Women's University, Nara 630-8506
⁴⁵National Central University, Chung-li 32054
⁴⁶National United University, Miao Li 36003
⁴⁷Department of Physics, National Taiwan University, Taipei 10617
⁴⁸H. Niewodniczanski Institute of Nuclear Physics, Krakow 31-342
⁴⁹Nippon Dental University, Niigata 951-8580
⁵⁰Niigata University, Niigata 950-2181
⁵¹University of Nova Gorica, 5000 Nova Gorica
⁵²Novosibirsk State University, Novosibirsk 630090
⁵³Osaka City University, Osaka 558-8585
⁵⁴Pacific Northwest National Laboratory, Richland, Washington 99352
⁵⁵University of Pittsburgh, Pittsburgh, Pennsylvania 15260
⁵⁶University of Science and Technology of China, Hefei 230026
⁵⁷Seoul National University, Seoul 151-742
⁵⁸Showa Pharmaceutical University, Tokyo 194-8543
⁵⁹Soongsil University, Seoul 156-743
⁶⁰University of South Carolina, Columbia, South Carolina 29208
⁶¹Sungkyunkwan University, Suwon 440-746
⁶²School of Physics, University of Sydney, NSW 2006
⁶³Department of Physics, Faculty of Science, University of Tabuk, Tabuk 71451
⁶⁴Tata Institute of Fundamental Research, Mumbai 400005
⁶⁵Excellence Cluster Universe, Technische Universität München, 85748 Garching
⁶⁶Department of Physics, Technische Universität München, 85748 Garching
⁶⁷Toho University, Funabashi 274-8510
⁶⁸Department of Physics, Tohoku University, Sendai 980-8578
⁶⁹Earthquake Research Institute, University of Tokyo, Tokyo 113-0032
⁷⁰Department of Physics, University of Tokyo, Tokyo 113-0033
⁷¹Tokyo Institute of Technology, Tokyo 152-8550
⁷²Tokyo Metropolitan University, Tokyo 192-0397
⁷³University of Torino, 10124 Torino
⁷⁴CNP, Virginia Polytechnic Institute and State University, Blacksburg, Virginia 24061
⁷⁵Wayne State University, Detroit, Michigan 48202
⁷⁶Yamagata University, Yamagata 990-8560
⁷⁷Yonsei University, Seoul 120-749

We report the first observation of the decay $\Lambda_c^+ \rightarrow pK^+\pi^-$ using a 980 fb^{-1} data sample collected by the Belle detector at the KEKB asymmetric-energy e^+e^- collider. This is the first observation of a doubly Cabibbo-suppressed decay of a charmed baryon. We measure the branching ratio of this decay with respect to its Cabibbo-favored counterpart to be $\mathcal{B}(\Lambda_c^+ \rightarrow pK^+\pi^-)/\mathcal{B}(\Lambda_c^+ \rightarrow pK^-\pi^+) = (2.35 \pm 0.27 \pm 0.21) \times 10^{-3}$, where the uncertainties are statistical and systematic, respectively.

PACS numbers: 13.30.Eg, 14.20.Lq

Several doubly Cabibbo-suppressed (DCS) decays of charmed mesons have been observed [1–4]. Their mea-

sured branching ratios with respect to the corresponding Cabibbo-favored (CF) decays play an important role in constraining models of the decay of charmed hadrons and in the study of flavor- $SU(3)$ symmetry [1, 3–6]. On the other hand, because of the smaller production cross sections for charmed baryons, DCS decays of charmed baryons have not yet been observed; only an upper limit, $\mathcal{B}(\Lambda_c^+ \rightarrow pK^+\pi^-)/\mathcal{B}(\Lambda_c^+ \rightarrow pK^-\pi^+) < 0.46\%$ with 90% confidence level, has been reported by the FOCUS Collaboration [7]. Theoretical calculations of DCS decays of charmed baryons have been very few and limited to two-body decay modes [8, 9].

In this letter, we report the first observation of the DCS decay $\Lambda_c^+ \rightarrow pK^+\pi^-$ and the measurement of its branching ratio with respect to its counterpart CF decay $\Lambda_c^+ \rightarrow pK^-\pi^+$ [10]. Typical decay diagrams of DCS and CF decays are shown in Fig. 1. In brief, the diagrams are categorized as external W -emission, internal W -emission, and W -exchange processes. Since W exchange is allowed in $\Lambda_c^+ \rightarrow pK^-\pi^+$ as shown in Fig. 1(e) but absent in $\Lambda_c^+ \rightarrow pK^+\pi^-$, the ratio $\mathcal{B}(\Lambda_c^+ \rightarrow pK^+\pi^-)/\mathcal{B}(\Lambda_c^+ \rightarrow pK^-\pi^+)$ may be smaller than the naïve expectation [7] of $\tan^4 \theta_c$ (0.285%), where θ_c is the Cabibbo mixing angle [11] and $\sin \theta_c = 0.225 \pm 0.001$ [12]. We can also compare the ratio $\mathcal{B}(\Lambda_c^+ \rightarrow pK^+\pi^-)/\mathcal{B}(\Lambda_c^+ \rightarrow$

$pK^-\pi^+)$ with similar ratios in charmed meson decays, such as $\sqrt{\frac{\mathcal{B}(D^+ \rightarrow K^+\pi^+\pi^-)}{\mathcal{B}(D^+ \rightarrow K^-\pi^+\pi^-)}} \frac{\mathcal{B}(D_s^+ \rightarrow K^+K^+\pi^-)}{\mathcal{B}(D_s^+ \rightarrow K^+K^-\pi^+)} = (1.25 \pm 0.08) \tan^4 \theta_c$ [1] or $\mathcal{B}(D^0 \rightarrow K^+\pi^-)/\mathcal{B}(D^0 \rightarrow K^-\pi^+) = (1.24 \pm 0.05) \tan^4 \theta_c$ [2]. By doing so, similarities and differences between charmed meson and baryon decays can provide additional insight into flavor- $SU(3)$ symmetry and QCD. For example, flavor- $SU(3)$ symmetry breaking in Λ_c^+ decay may affect the ratio as is the case in D meson decay.

We analyze data taken at or near the $\Upsilon(1S)$, $\Upsilon(2S)$, $\Upsilon(3S)$, $\Upsilon(4S)$, and $\Upsilon(5S)$ resonances collected by the Belle detector at the KEKB asymmetric-energy e^+e^- collider [13]. The integrated luminosity of the data sample is 980 fb^{-1} . The Belle detector is a large-solid-angle magnetic spectrometer comprising a silicon vertex detector (SVD) [14], a central drift chamber (CDC), an array of aerogel threshold Cherenkov counters (ACC), a barrel-like arrangement of time-of-flight scintillation counters (TOF), and an electromagnetic calorimeter comprised of CsI(Tl) crystals (ECL) located inside a superconducting solenoid coil that provides a 1.5 T magnetic field. The detector is described in detail elsewhere [15]. The combined particle identification (PID) likelihoods, $\mathcal{L}(h)$ ($h = p, K$, or π), are derived from ACC and TOF measurements and dE/dx measurements in CDC. The discriminant $\mathcal{R}(h|h')$, defined as $\mathcal{L}(h)/(\mathcal{L}(h) + \mathcal{L}(h'))$, is the ratio of likelihoods for h and h' identification. The electron likelihood ratio, $\mathcal{R}(e)$, for e and h identification is derived from ACC, CDC, and ECL measurements [16]. We use samples of $e^+e^- \rightarrow c\bar{c}$ Monte Carlo (MC) events, which are generated with PYTHIA [17] and EvtGen [18] and propagated by GEANT3 [19] to simulate the detector performance, to estimate reconstruction efficiencies and to study backgrounds.

In this analysis, our selection criteria follow mostly those typically used in other charmed hadron studies at Belle (for example, Ref. [1, 20, 21]). However, our final criteria, described in the next paragraph, are determined by a figure-of-merit (FoM) study performed using a control sample of the CF decay ($\Lambda_c^+ \rightarrow pK^-\pi^+$) in real data, together with sidebands to the DCS signal region. We use this blinded study to optimize the FoM, defined as $n_{\text{sig}}/\sqrt{n_{\text{sig}} + n_{\text{bkg}}}$, where n_{sig} is the fitted yield of the control sample multiplied by the presumed ratio of the DCS and CF decays (0.0025), and n_{bkg} is the number of background events from the sideband region in the DCS decay.

A Λ_c^+ candidate is reconstructed from the three charged hadrons, and all charged tracks are required to have a distance of closest approach to the interaction point (DOCA) less than 2.0 cm and 0.1 cm in the beam direction (z) and in the transverse (r - ϕ) direction, respectively. The number of SVD hits is also required to be at least one, both in the z and r - ϕ directions, for each of three charged particles. The charged particles




FIG. 1. Typical external [internal] W -emission diagrams for (a) [(c)] $\Lambda_c^+ \rightarrow pK^+\pi^-$ and (b) [(d)] $\Lambda_c^+ \rightarrow pK^-\pi^+$, and (e) a typical W -exchange diagram of $\Lambda_c^+ \rightarrow pK^-\pi^+$.




FIG. 2. Distribution of $M(pK^-\pi^+)$. The curves indicate the fit result: the full fit model (solid) and the combinatorial background only (dashed).

are identified by the PID measurements: $\mathcal{R}(p|h) > 0.9$ for both $h = \pi$ and K is required for charged protons, $\mathcal{R}(K|p) > 0.4$ and $\mathcal{R}(K|\pi) > 0.9$ are required for charged kaons, $\mathcal{R}(\pi|p) > 0.4$ and $\mathcal{R}(\pi|K) > 0.4$ are required for charged pions, and $\mathcal{R}(e) < 0.9$ is required for all charged particles. The identification efficiencies of p , K , and π are 75%, 75%, and 95%, respectively, for the typical momentum range of the decays. Probabilities of misidentifying h as h' , $P(h \rightarrow h')$, are estimated by using data and MC samples of the CF decay to be 8% ($P(p \rightarrow K)$), 5% ($P(p \rightarrow \pi)$), 11% ($P(K \rightarrow \pi)$), 2% ($P(K \rightarrow p)$), 2% ($P(\pi \rightarrow K)$), and less than 1% ($P(\pi \rightarrow p)$) for the typical momentum range. To suppress combinatorial backgrounds, especially from B meson decays, we place a requirement on the scaled momentum: $x_p > 0.53$, where x_p is defined as $p^*/\sqrt{E_{\text{cm}}^2/4 - M^2}$; here, E_{cm} is the total center-of-mass energy, p^* is the momentum in the center-of-mass frame, and M is the mass of the Λ_c^+ candidate. In addition, the χ^2 value from the common vertex fit of the charged tracks must be less than 40.

Figures 2 and 3 show invariant mass distributions, $M(pK^-\pi^+)$ (CF) and $M(pK^+\pi^-)$ (DCS), with the final selection criteria. DCS decay events are clearly observed in $M(pK^+\pi^-)$. We perform a binned least- χ^2 fit to the two distributions from 2.15 GeV/c^2 to 2.42 GeV/c^2 with 0.01 MeV/c^2 bin width, and the figures are drawn with merged bins. The probability density functions (PDFs) for the fits are the sum of two Gaussian distributions, with a common central value, to represent the signals,




FIG. 3. Distribution of $M(pK^+\pi^-)$ (top) and residuals of data with respect to the fitted combinatorial background (bottom). Curves are drawn as described in Fig. 2.

and polynomials of fifth and third order for the combinatorial backgrounds in the $M(pK^-\pi^+)$ and $M(pK^+\pi^-)$ distributions, respectively. In the fit to $M(pK^+\pi^-)$, the resolution and central value of the signal function are fixed to be the same as those found from the fit to $M(pK^-\pi^+)$. The equality of these quantities is expected from first principles and is confirmed using the MC simulation. The reduced χ^2 values ($\chi^2/d.o.f$) of the fits are 1.03 (27749/26989) and 1.01 (27131/26995) for the CF and DCS decays, respectively. From the fit results, the signal yields of $\Lambda_c^+ \rightarrow pK^-\pi^+$ and $\Lambda_c^+ \rightarrow pK^+\pi^-$ decays are determined to be $(1.452 \pm 0.015) \times 10^6$ events and 3587 ± 380 events, respectively, where the uncertainties are statistical. There is a small excess above background on the right side of the Λ_c^+ peak (around 2.297 GeV/c^2) in the DCS spectrum of Fig. 3. We attribute this to a statistical fluctuation as no known process would make such a narrow feature at this position even when possible particle misidentification, such as the misidentification of both the K and the π , is taken into account.

The DCS decay has a peaking background from the SCS decay $\Lambda_c^+ \rightarrow \Lambda K^+$ with $\Lambda \rightarrow p\pi^-$, which has the same final state topology. However, because of the long Λ lifetime, many of the Λ vertexes are displaced by several centimeters from the main vertex so the DOCA and χ^2 requirements suppress most of this background. The remaining SCS-decay yield is included in the signal yield

of $\Lambda_c^+ \rightarrow pK^+\pi^-$ decay and is estimated via the relation

$$\mathcal{N}(SCS; \Lambda \rightarrow p\pi^-) = \frac{\epsilon(SCS; \Lambda \rightarrow p\pi^-)}{\epsilon(CF)} \frac{\mathcal{B}(SCS; \Lambda \rightarrow p\pi^-)}{\mathcal{B}(CF)} \mathcal{N}(CF), \quad (1)$$

where $\mathcal{N}(CF)$ is the signal yield of the CF decay, $\mathcal{B}(SCS; \Lambda \rightarrow p\pi^-)/\mathcal{B}(CF) = (0.61 \pm 0.13)\%$ is the branching ratio [12], and $\epsilon(SCS; \Lambda \rightarrow p\pi^-)/\epsilon(CF) = 0.023$ is the relative efficiency found using MC samples. This calculation gives a yield of 208 ± 78 events from this source, where the uncertainty is estimated by comparing the signal yields from this calculation and a fit to $M(pK^+\pi^-)$ with loosened selection criteria for the vertex point and Λ selection in $M(p\pi^-)$. After subtraction of this SCS component, the signal yield of the DCS decay is $3379 \pm 380 \pm 78$, where the first uncertainty is statistical and the second is systematic due to this subtraction.

To estimate the statistical significance of the DCS signal, we exclude the SCS signal by vetoing events with $1.1127 \text{ GeV}/c^2 < M(p\pi^-) < 1.1187 \text{ GeV}/c^2$. The significance is estimated as $\sqrt{-2 \ln(\mathcal{L}_0/\mathcal{L})}$, where \mathcal{L}_0 and \mathcal{L} are the maximum likelihood values from binned maximum likelihood fits with the signal yield fixed to zero and allowed to float, respectively. The calculated significance corresponds to 9.4σ .

We calculate the reconstruction efficiency using a mixture of subchannels weighted with their corresponding branching fractions. For the CF decay, the subchannels and their branching fractions are taken from the Ref. [12] and the estimated efficiency of the CF decay is $(13.83 \pm 0.05)\%$, where the uncertainty is from MC statistics. To estimate the uncertainty arising from the mix of intermediate states in the CF decay, the reconstruction efficiency is calculated using the efficiency of each bin of the $M^2(K^-\pi^+)$ vs. $M^2(pK^-)$ Dalitz distribution [22], shown in Fig. 4, and weighting them by the number of events in the bin of the real data. The relative difference between the reconstruction efficiencies, before and after this weighting, is 3.0%. For the DCS decay, we use the $pK^*(892)^0$, $\Delta(1232)^0 K^+$, and non-resonant subchannels with branching fractions of 0.23, 0.18, and 0.59, respectively. These values represent the branching fractions for the corresponding subchannels of the CF decay, adjusted for the fact that $\Lambda(1520)$ cannot be produced in the DCS decay. With the assumed subchannels and their branching fractions, the reconstruction efficiency of the DCS decay is estimated to be $(13.71 \pm 0.05)\%$, where the uncertainty is from MC statistics. Due to the low signal-to-background ratio in the DCS signal peak, the uncertainty from the assumed mixture of intermediate states cannot be estimated using the method used for the CF decay. Therefore, the largest difference between the efficiency of a subchannel and the overall reconstruction efficiency is taken as the efficiency uncertainty; the largest relative difference is 4.5% from $\Delta(1232)^0 K^+$ sub-




FIG. 4. Invariant mass squared of $K^-\pi^+$ versus pK^- within $2.2746 \text{ GeV}/c^2 < M(pK^-\pi^+) < 2.2986 \text{ GeV}/c^2$ in real data (top) and estimated efficiency using the MC (bottom). The bin widths of x and y axes are $0.016 \text{ GeV}^2/c^4$ and $0.027 \text{ GeV}^2/c^4$, respectively.

channel. The relative efficiency of the CF and DCS decays is 1.01 ± 0.05 , where the uncertainty is due to the uncertainty in the composition of the intermediate states as described above.

The branching ratio, $\mathcal{B}(\Lambda_c^+ \rightarrow pK^+\pi^-)/\mathcal{B}(\Lambda_c^+ \rightarrow pK^-\pi^+)$, is $(2.35 \pm 0.27 \pm 0.21) \times 10^{-3}$, where the uncertainties are statistical and systematic, respectively. Sources of the systematic uncertainty and their values are listed in Table I. The uncertainty from the binning and range of the fits is estimated by changing the bin

TABLE I. Systematic uncertainties and sources.

| Source | Uncertainty (%) |
|-----------------------------|-----------------|
| Background from SCS signal | ± 2.3 |
| Intermediate states | ± 5.4 |
| Binning and fit range (DCS) | ± 5.5 |
| Binning and fit range (CF) | ± 0.6 |
| PDF shape (DCS) | ± 2.6 |
| PDF shape (CF) | ± 1.4 |
| MC statistics | ± 0.4 |
| PID | ± 2.2 |
| Charge-conjugate mode | ± 1.8 |
| Total | ± 9.0 |

width to $3 \text{ MeV}/c^2$ and adjusting the fitted range of the invariant mass distributions. The uncertainty due to the PDF shapes is estimated by changing the order of the polynomial background function, by changing the signal function to the sum of three Gaussian distributions, and by fixing the resolution of the signal function to the MC-derived resolution value. The PID uncertainty is determined by data-MC comparison of several control samples. We treat the relative efficiency difference between charge-conjugate modes as a systematic uncertainty.

The branching fraction of the CF decay, $(6.84 \pm 0.24^{+0.21}_{-0.27}) \times 10^{-2}$, was already well-measured in a previous Belle analysis [23]. Combining that with our measurement, we determine the absolute branching fraction of the DCS decay to $(1.61 \pm 0.23^{+0.07}_{-0.08}) \times 10^{-4}$, where the first uncertainty is the total uncertainty of the branching ratio and the second is uncertainty of the branching fraction of CF decay. This measured branching ratio corresponds to $(0.82 \pm 0.12) \tan^4 \theta_c$, where the uncertainty is the total.

The branching ratio suggests a slightly smaller decay width than the naïve expectation, although the significance is only 1.5σ . This is consistent with the expectation that the Δ isobar, in addition to $\Lambda^*(1520)$, does not contribute to the DCS decay [9]. Omitting those two contributions, which are $(25 \pm 4)\%$ [12], from the CF decay rate, the ratio becomes $(1.10 \pm 0.17) \tan^4 \theta_c$ which is consistent with $\tan^4 \theta_c$ within 1σ . This result suggests that W -exchange effects are modest in the decay $\Lambda_c^+ \rightarrow pK^-\pi^+$, except possibly for the sub-mode with an intermediate Δ . In addition, we note that the observed DCS/CF ratio for charmed baryons is not significantly different from the measured ratio for charmed meson decay.

In conclusion, the first DCS decay of a charmed baryon, $\Lambda_c^+ \rightarrow pK^+\pi^-$, is observed with a statistical significance of 9.4σ . The branching ratio relative to its counterpart CF decay is $(2.35 \pm 0.27 \pm 0.21) \times 10^{-3}$, which corresponds to $(0.82 \pm 0.12) \tan^4 \theta_c$. This result sheds new light on charmed hadron decays, and such DCS measurements are important ingredients in modeling the non-

leptonic decays of hadrons. However, the current experimental precision on the strengths of DCS modes and the level of detail of the available theoretical results are not sufficient to constrain the relative importance of the different subprocesses shown in Fig. 1. Future progress in this field will require more precise experimental measurements as well as more refined theoretical calculations.

We thank the KEKB group for excellent operation of the accelerator; the KEK cryogenics group for efficient solenoid operations; and the KEK computer group, the NII, and PNNL/EMSL for valuable computing and SINET4 network support. We acknowledge support from MEXT, JSPS and Nagoya's TLPSC (Japan); ARC and DIISR (Australia); FWF (Austria); NSFC and CCEPP (China); MSMT (Czechia); CZF, DFG, and VS (Germany); DST (India); INFN (Italy); MOE, MSIP, NRF, GSDC of KISTI, and BK21Plus (Korea); MNiSW and NCN (Poland); MES and RFAAE (Russia); ARRS (Slovenia); IKERBASQUE and UPV/EHU (Spain); SNSF (Switzerland); NSC and MOE (Taiwan); and DOE and NSF (USA).

-
- [1] B.R. Ko *et al.* (Belle Collaboration), Phys. Rev. Lett. **102**, 221802 (2009).
 - [2] R. Aaij *et al.* (LHCb Collaboration), Phys. Rev. Lett. **110**, 101802 (2013).
 - [3] E. Won *et al.* (Belle Collaboration), Phys. Rev. Lett. **107**, 221801 (2011).
 - [4] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. D **74**, 011107(R) (2006).
 - [5] D.N. Gao, Phys. Lett. B **645**, 59 (2007).
 - [6] H.J. Lipkin, Nucl. Phys. B, Proc. Suppl. **115**, 117 (2003).
 - [7] J.M. Link *et al.* (FOCUS Collaboration), Phys. Lett. B **624**, 166 (2005).
 - [8] K.K. Sharma and R.C. Verma, Phys. Rev. D **55**, 7067 (1997).
 - [9] T. Uppal, R.C. Verma, and M.P. Khanna, Phys. Rev. D **49**, 3417 (1994).
 - [10] Unless stated otherwise, charge-conjugate modes are implied throughout this paper.
 - [11] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963).
 - [12] K. A. Olive *et al.* (Particle Data Group), Chin. Phys. C **38**, 090001 (2014).
 - [13] S. Kurokawa and E. Kikutani, Nucl. Instrum. Methods Phys. Res., Sect. A **499**, 1 (2003), and other papers included in this volume; T.Abe *et al.*, Prog. Theor. Exp. Phys. **2013**, 03A001 (2013) and references therein.
 - [14] Z. Natkaniec *et al.* (Belle SVD2 Group), Nucl. Instrum. Methods Phys. Res., Sect. A **560**, 1 (2006); Y. Ushiroda (Belle SVD2 Group), Nucl. Instrum. Methods Phys. Res., Sect. A **511**, 6 (2003).
 - [15] A. Abashian *et al.* (Belle Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A **479**, 117 (2002); also see detector section in J.Brodzicka *et al.*, Prog. Theor. Exp. Phys. **2012**, 04D001 (2012).
 - [16] H. Hanagaki *et al.*, Nucl. Instrum. Methods Phys. Res., Sect. A **485**, 490 (2002)

- [17] T. Sjöstrand, S. Mrenna, and P. Skands, J. High Energy Phys. **05**, 026 (2006).
- [18] D. Lange, Nucl. Instrum. Methods Phys. Res., Sect. A **462**, 152 (2001); T. Sjöstrand, P. Edén, C. Friberg, L. Lönnblad, G. Miu, S. Mrenna, and E. Norrbin, Comput. Phys. Commun. **135**, 238 (2001).
- [19] R. Brun *et al.*, GEANT 3.21, CERN Report DD/EE/84-1, 1984.
- [20] K. Abe *et al.* (Belle Collaboration), Phys. Lett. B **524**, 33 (2002).
- [21] R. Chistov *et al.* (Belle Collaboration), Phys. Rev. D **88**, 071103(R) (2013).
- [22] R.H. Dalitz, Philos. Mag. **44**, 1068 (1953).
- [23] A. Zupanc *et al.* (Belle Collaboration), Phys. Rev. Lett. **113**, 042002 (2014).