



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Anomalous Thermalization in Ergodic Systems

David J. Luitz and Yevgeny Bar Lev

Phys. Rev. Lett. **117**, 170404 — Published 21 October 2016

DOI: [10.1103/PhysRevLett.117.170404](https://doi.org/10.1103/PhysRevLett.117.170404)

Anomalous thermalization in ergodic systems

David J. Luitz¹ and Yevgeny Bar Lev²

¹*Department of Physics and Institute for Condensed Matter Theory,
University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA**

²*Department of Chemistry, Columbia University,
3000 Broadway, New York, New York 10027, USA*

It is commonly believed that quantum isolated systems satisfying the eigenstate thermalization hypothesis (ETH) are diffusive. We show that this assumption is too restrictive, since there are systems that are asymptotically in a thermal state, yet exhibit anomalous, subdiffusive thermalization. We show that such systems satisfy a modified version of the ETH ansatz and derive a general connection between the scaling of the variance of the offdiagonal matrix elements of local operators, written in the eigenbasis of the Hamiltonian, and the dynamical exponent. We find that for subdiffusively thermalizing systems the variance scales more slowly with system size than expected for diffusive systems. We corroborate our findings by numerically studying the distribution of the coefficients of the eigenfunctions and the offdiagonal matrix elements of local operators of the random field Heisenberg chain, which has anomalous transport in its thermal phase. Surprisingly, this system also has non-Gaussian distributions of the eigenfunctions, thus directly violating Berry's conjecture.

PACS numbers: 05.30.-d, 05.70.Ln, 75.10.Pq

Recently, the long standing question of thermalization in closed quantum systems [1] has regained importance due to advances in cold atoms experiments [2], as well as the theoretical prediction of a dynamical phase transition, known as the many-body localization (MBL) transition between ergodic and nonergodic phases [3–7]. Thermalization in classical systems is normally associated with their underlying ergodicity, a property which is one of the basic assumptions of statistical mechanics. The situation for quantum systems is more delicate, since the evolution of any eigenstate amounts to a time dependent global phase (see recent reviews [8–10]). Major progress was achieved by Berry who conjectured [11] that the coefficients of high energy eigenstates of a quantum system in a generic basis corresponding to a chaotic classical system are independent Gaussian variables, similarly to the distribution of the eigenstates in the corresponding random matrix ensemble [12]. The connection between random matrix theory and realistic systems was made in Deutsch's seminal paper [13], showing that perturbing the Hamiltonian with a random matrix leads to thermalization. Later, it was shown by Srednicki for a gas of hard core particles that if Berry's conjecture is satisfied, the distribution of the velocities of the particles approaches the Maxwell-Boltzmann distribution for large systems. It was therefore concluded that the validity of Berry's conjecture is *required* for thermalization in quantum systems [14]. Building on this intuition, and the analogy to random-matrix theory, Srednicki proposed that an ergodic isolated quantum system should satisfy the Eigenstate Thermalization Hypothesis (ETH) ansatz [15],

$$\langle \alpha | \hat{O} | \beta \rangle = \bar{O}(E) \delta_{\alpha\beta} + e^{-S(E)/2} f(E, \omega) R_{\alpha\beta}, \quad (1)$$

where α, β are the eigenstates, \hat{O} is a generic operator, $S(E)$ is the microcanonical entropy, $\bar{O}(E)$, $f(E, \omega)$ are smooth functions of their arguments, and $E = (E_\alpha + E_\beta)/2$ and $\omega = E_\beta - E_\alpha$. Here, the normal distribution with zero mean and unit variance of the random term $R_{\alpha\beta}$ is justified through Berry's conjecture. The first, diagonal term in the ETH ansatz is equal to the micro-canonical expectation value of the corresponding observable, thus representing a static thermodynamic quantity. This relation was numerically verified by Rigol et al. for certain generic quantum systems [16]. The exponential decay with system size of the second term, as well as the validity of the Gaussian distribution of the noise ($R_{\alpha\beta}$), was subsequently verified for a number of generic quantum systems [17–24]. In the present work we show that there is a class of ergodic systems which exhibit anomalous (non-diffusive) relaxation to equilibrium while still satisfying a modified ETH ansatz, such that the off-diagonal elements in (1) include a power law correction to their scaling with the system size. To characterize the approach to equilibrium, we follow the derivations in Refs. [25, 26] and [9] (Sec. 6.8), and use the correlator,

$$C_\alpha(t) = \langle \alpha | \hat{O}(t) \hat{O}(0) | \alpha \rangle = \sum_{\beta \neq \alpha} \left| \langle \alpha | \hat{O} | \beta \rangle \right|^2 e^{i(E_\alpha - E_\beta)t}. \quad (2)$$

where $|\alpha\rangle, |\beta\rangle$ are eigenstates and in the last step we have subtracted the element $\beta = \alpha$ (assuming a generic system with no degeneracy), to have a correlator with a vanishing infinite time average. Using (1) we have,

$$C_\alpha(t) = \sum_{\beta \neq \alpha} e^{-S(E_\alpha + \frac{\omega}{2})} \left| f\left(E_\alpha + \frac{\omega}{2}, \omega\right) \right|^2 |R_{\alpha\beta}|^2 e^{-i\omega t}. \quad (3)$$

For further simplification we replace the sum over eigenstates by an integral over the density of states, which we write as $e^{S(E)}$:

$$\sum_{\beta \neq \alpha} \rightarrow \int dE_\beta e^{S(E_\beta)} = \int d\omega e^{S(E_\alpha + \omega)}. \quad (4)$$

The Fourier transform to frequency space yields

$$C_\alpha(\omega) = 2\pi \exp \left[S(E_\alpha + \omega) - S \left(E_\alpha + \frac{\omega}{2} \right) \right] \times \left| f \left(E_\alpha + \frac{\omega}{2}, \omega \right) \right|^2 |R_{E_\alpha, E_\alpha + \omega}|^2. \quad (5)$$

Assuming that $S(E)$ and $f(E, \omega)$ are smooth functions of energy and frequency we can expand,

$$S(E_\alpha + \omega) - S \left(E_\alpha + \frac{\omega}{2} \right) = \frac{\partial S}{\partial E} \omega - \frac{\partial S}{\partial E} \frac{\omega}{2} = \frac{\omega}{2T} + \mathcal{O}(\omega^2), \quad (6)$$

where we used $\partial S / \partial E = 1/T$, where T is the micro-canonical temperature, and we set the Boltzmann constant to one. Expanding the other term gives

$$f \left(E_\alpha + \frac{\omega}{2}, \omega \right) = f(E_\alpha, \omega) + \frac{\omega}{2} \left. \frac{\partial f(E, \omega)}{\partial E} \right|_{E=E_\alpha} + \mathcal{O}(\omega^2). \quad (7)$$

Therefore to the leading order in ω we get,

$$C_\alpha(\omega) = 2\pi e^{\omega/(2T)} \left[|f(E_\alpha, \omega)|^2 + \frac{\omega}{2} \left. \frac{\partial |f(E, \omega)|^2}{\partial E} \right|_{E=E_\alpha} \right]. \quad (8)$$

For an Hermitian operator, \hat{O} we have $f(E_\alpha, \omega) = f(E_\alpha, -\omega)$, yielding,

$$|f(E_\alpha, \omega)|^2 = \frac{1}{4\pi} \left[e^{-\omega/(2T)} C_\alpha(\omega) + e^{\omega/(2T)} C_\alpha(-\omega) \right]. \quad (9)$$

In the limit of small frequencies $\omega/T \rightarrow 0$, we have,

$$|f(E_\alpha, \omega)|^2 = \int_{-\infty}^{\infty} dt \langle \alpha | \{ \hat{O}(t), \hat{O}(0) \} | \alpha \rangle e^{i\omega t}, \quad (10)$$

where $\{.,.\}$ is an anti-commutator. We now *assume* that $\hat{O}(t)$ is a conserved quantity which exhibits anomalous transport,

$$\langle \Psi | \{ \hat{O}(t), \hat{O}(0) \} | \Psi \rangle \asymp t^{-\gamma}. \quad (11)$$

For such a decay of the correlation function $|f(E_\alpha, \omega)|^2$ is given by,

$$|f(E_\alpha, \omega)|^2 \propto \int_{-\infty}^{\infty} dt |t|^{-\gamma} e^{i\omega t} \propto |\omega|^{-(1-\gamma)}. \quad (12)$$

For a finite system of size L , saturation will occur after time $t_c \approx L^{1/\gamma}$, analogous to the Thouless time [27]. This follows from the relation between the return probability exponent γ , and the mean-square displacement exponent, which is valid for one dimensional systems [28].

The power-law dependence will be therefore cut-off for frequencies, $\omega < t_c^{-1} = L^{-1/\gamma}$, and $|f(E_\alpha, \omega)|^2$ will become structureless [9],

$$|f(E_\alpha, \omega)|^2 \approx t_c^{1-\gamma} = L^{(1-\gamma)/\gamma}, \quad \omega < L^{-1/\gamma} \quad (13)$$

Then, the offdiagonal elements should scale with system size as,

$$O_{\alpha\beta} \propto e^{-Ls(E)/2} L^{(1-\gamma)/(2\gamma)} R_{\alpha\beta}, \quad |E_\alpha - E_\beta| < L^{-1/\gamma} \quad (14)$$

where we write the micro-canonical entropy density as $s(E) = S(E)/L$, to make the dependence on system size explicit. Note that we keep the general form of the ETH ansatz and assume that the distribution of the random numbers $R_{\alpha\beta}$ has zero mean and unit variance. The scaling with system size of the standard deviation of the off-diagonal matrix elements after the dominant exponential factor has been removed is therefore given by,

$$\text{std} \left(O_{\alpha\beta} e^{Ls(E)/2} \right) \asymp L^\delta, \quad |E_\alpha - E_\beta| < L^{-1/\gamma} \quad (15)$$

where, $\delta \equiv (1 - \gamma) / (2\gamma)$. A special case of this relation was established in Ref. [9] for diffusive one-dimensional systems, where $\delta = 1/2$ and $\gamma = 1/2$. We note in passing, that the scaling of $\langle |O_{\alpha\beta}|^2 \rangle$ with system size was computed in Ref. [21] for generic clean systems and in Ref. [29] for a generic disordered system. In both works, departure from exponential dependence on system size is observed when ω is taken to be small. Our results suggest that the cause of this discrepancy is the logarithmic correction resulting from (15).

To show that (14) holds for systems with anomalous transport, we numerically study the spin- $\frac{1}{2}$ Heisenberg chain in a random magnetic field,

$$\hat{H} = J \sum_i \vec{S}_i \cdot \vec{S}_{i+1} + \sum_i h_i \hat{S}_i^z, \quad (16)$$

where J is the spin-spin coupling, which we will set to 1, and $h_i \in [-W, W]$ are random fields drawn from a uniform distribution. Previous studies [24, 30–36], have established that the ergodic phase of this model is characterized by anomalous transport with a continuously varying dynamical exponent $\gamma(W) \lesssim 1/2$, as a function of the disorder strength W . The dynamical exponent vanishes at the many body localization transition, as the system no longer thermalizes in the MBL phase. In general, exact numerical studies of high energy many-body eigenstates are a formidable task and full diagonalization becomes very expensive for systems of size $L \gtrsim 18$. Since we strive to access systems that are as large as possible, we use the shift-invert technique, which transforms the spectrum of the Hamiltonian such that the states of interest are moved to the lowest (highest) energies in the transformed spectrum and become tractable




Figure 1. *Left column:* Distribution of the off-diagonal elements ($\alpha \neq \beta$) of the operator \hat{S}_i^z , written in the eigenstate basis of the Hamiltonian (16), for different disorder strengths, $W = 0.4$ and 2.0 and system sizes, $L \in [12, 22]$. Darker tones correspond to larger system sizes. The eigenstates correspond to 50 closest eigenvalues to the middle of the many-body spectrum and the distributions have been sampled from roughly 1000 disorder realizations, except for $L = 22$, where we only used 100 realizations. *Right column:* Distributions rescaled to have a unit variance. At $W = 0.4$ the distribution is very close to Gaussian (dashed line).

by Krylov space methods. The most commonly used spectral transformation for this purpose is $(H - \sigma I)^{-1}$, where the explicit inversion of the shifted Hamiltonian can be avoided and replaced by a repeated solution of a set of linear equations. We use the massively parallel MUMPS library [37, 38] for this purpose and are able to obtain exact mid-spectrum eigenstates for system sizes up to $L = 22$. For all system sizes, we calculate a fixed number $k = 50$ of eigenstates and eigenvalues in the middle of the spectrum. For these energy densities the transition to the MBL phase occurs at a critical disorder strength of $W_c \approx 3.7$ [39]. In what follows, we will focus on the limit of small disorder, $W < W_c$, where the system is ergodic and the diagonal elements of local operators were shown to satisfy ETH, although with non-Gaussian distributions [24]. We will show that the *offdiagonal* elements satisfy our scaling prediction (14). Since the many-body density of states scales exponentially with the system size, for a fixed number of states around some energy the assumption on the energy difference, $\omega = E_\alpha - E_\beta = k \exp(-Ls(E)) < L^{-1/\gamma}$, in (14) is always satisfied for sufficiently large systems.

For every pair $|\alpha\rangle, |\beta\rangle$ of these eigenstates with $\alpha \neq \beta$,




Figure 2. *Left column:* Distributions of the eigenfunction elements in the basis of the local magnetization for small ($W = 0.4$, top) and intermediate ($W = 1.6$, bottom) disorder strengths for various system sizes. *Right column:* Same distribution as in the left column, rescaled such that the variance is equal to one. Darker tones correspond to larger system sizes. Clearly, the distribution differs strongly from a Gaussian distribution at intermediate disorder. At weak disorder, the difference from Gaussian (dashed line) is visible mostly in the tails and the excess of weight at zero.

we calculate the matrix elements $\langle \alpha | \hat{S}_i^z | \beta \rangle$ of the local \hat{S}_i^z operator for all sites i in the chain using periodic boundary conditions. In the left column of Fig. 1 we present the probability distribution of the off-diagonal elements computed for different disorder strengths and system sizes, the right panel shows the same distributions, renormalized by their standard deviation σ , in order to compare the shapes of the distributions across system sizes. This normalization procedure allows us to directly extract $R_{\alpha\beta}$, since the resulting distribution has a unit variance. The shape of the rescaled distribution is Gaussian deep in the ergodic phase (for weak disorder) and thus corresponds to the general expectation of the ETH ansatz [18, 21]. Closer to the MBL transition the shape of the distribution is clearly non-Gaussian, which hints on the violation of the Berry's conjecture. To directly test the validity of Berry's conjecture we calculate the distribution of the coefficients $\langle i | \alpha \rangle$ of the eigenfunctions $|\alpha\rangle$ in the spin basis $|i\rangle$. Surprisingly even for the smallest disorder we study ($W = 0.4$), Berry's conjecture is clearly violated.

To verify that the exponent obtained from rescaling according to (14) is indeed linked to the dynamical exponent γ , we study the behavior of the corre-

lation function $\langle \psi | \{ \hat{S}_i^z(t), \hat{S}_i^z \} | \psi \rangle$. As it is very difficult for large systems to obtain high energy eigenstates, we use random states with an average energy density of $\epsilon = 0.5$, corresponding to the energy $\langle \psi | H | \psi \rangle = E_{\frac{1}{2}} := (E_{\max} + E_{\min})/2$ and a small variance of the energy $(\langle H^2 \rangle - \langle H \rangle^2) / \langle H \rangle^2 \ll 1$. We generate such typical high energy states starting from a random state $|\psi_0\rangle$ and using the power method for the *folded* Hamiltonian $(H - E_{\frac{1}{2}})^2$ to iteratively reduce the uncertainty in the energy around $E_{\frac{1}{2}}$. Typically, a few hundred iterations suffice to reduce the standard deviation of the energy to a few percent of the bandwidth. We then use the resulting *energy squeezed states* in the calculation of the correlation function, which is obtained using exact time evolution by a Krylov space method [32, 34, 40]. After a short time transient, this function decays as a power law superposed by oscillations as observed in previous studies for similar quantities [30–32, 41]. We find that the most reliable way of extracting the dynamical exponent γ is by using open boundary conditions (OBC) and studying the correlation function on one of the boundaries. This yields the same result as the bulk, but the effect of the other boundary is delayed compared to other setups, which gives access to longer times for which bulk transport is observed. For smaller system sizes we have verified that using the eigenstates as the initial condition $|\psi\rangle$ points to similar results. To reliably extract the dynamical exponent γ it is crucial to fit also the transient behavior which includes decaying oscillations superimposed onto the power law decay. For this purpose we use the ansatz proposed in Ref. [32],

$$C(t) = ae^{-t/\tau} \cos(\omega_1 t + \phi), \quad (17)$$

$$+ bt^{-\gamma} [1 + ct^{-\eta} \sin(\omega_2 t + \phi)]$$

yielding excellent fits. In Fig. 3 we present the dynamical exponent γ calculated from (17), together with the exponent γ , obtained from the exponent δ (see (15)). The left panel of Fig. 3 shows the LHS of Eq. (15) as a function of system size for various disorder strengths on a log-log scale, demonstrating that it indeed follows a power law. Here, we have estimated the density of states e^S from the energy interval, in which we find k eigenvalues. Note that approaching the MBL transition, visible deviations from power law behavior appear, signaling the violation of the scaling (14). However, sufficiently far from the MBL transition the agreement of the two exponents is remarkable. Surprisingly, while Berry’s conjecture is violated, the excellent collapse between the two exponents as predicted by (14) suggests that the ETH ansatz (1) still applies, just with *non-Gaussian* fluctuations and with a modified scaling of the offdiagonal elements with the system size.

In summary, we have shown that there are systems which are thermal and exhibit anomalous transport of conserved quantities, but still satisfy ETH, though in a




Figure 3. *Left panel*: Extraction of the exponent from the scaling relation (14) for various disorder strengths after the dominant exponential scaling term was eliminated. *Right panel*: Exponent extracted from the scaling relation (black circles) versus direct computation of the dynamical exponent γ from the correlation function using energy squeezed states.

modified form. We have derived the dependence of the standard deviation of offdiagonal matrix elements of local operators (written in the basis of the eigenstates of the Hamiltonian) on the system size for systems with both normal and *anomalous* transport. This dependence includes power law corrections to the customary exponential ETH term. We have derived a scaling relation between the exponent δ of this power law, and the dynamical transport exponent γ , and thoroughly tested the validity of this scaling using extensive numerical calculations on the random field Heisenberg model in its thermal phase. The scaling relation works perfectly for low to indeterminate disorder strengths sufficiently far from the MBL transition. Our numerical results also show that the distributions of the offdiagonal matrix elements are Gaussian at weak disorder, where the dynamics is roughly diffusive and become strongly non-Gaussian for stronger disorder, when the system becomes subdiffusive. These pathological distributions are accompanied by a violation of Berry’s conjecture, as the distributions of the wave function coefficients deviate strongly from Gaussian distributions. It would be interesting to explore the possible connection between anomalous transport and the violation of Berry’s conjecture in future works. In our analysis we have relied only on the second moment of the distributions of offdiagonal matrix elements, thus ignoring additional information encoded in its shape, which will show up in the relation between their moments. A number of previous studies discussed the existence of an intermediate phase with multifractal eigenstates [39, 42–47] and multifractal offdiagonal matrix elements of local operators [48]. While the non-Gaussian form of the obtained distributions is consistent with these studies, we leave the detailed exploration of this connection to a subsequent work.

Since the exponential dependence on the system size of the offdiagonal elements stems from the randomness assumption of the eigenfunctions coefficients, we speculate that the derived power law corrections follow from residual correlations between these coefficients induced by the conservation laws of the underlying system. It would be therefore interesting to see how the obtained corrections are affected by the number of conserved quantities in the system, a question which we leave for future studies.

Acknowledgments. – We thank Achileas Lazarides for inspiring discussions. YBL would like to thank David R. Reichman for valuable discussions and pointing out the connection to the Berry’s conjecture. DJL thanks Eduardo Fradkin, Anatoli Polkovnikov and Marcos Rigol for useful comments and Fabien Alet, Bryan Clark, Nicolas Laflorencie and Xiongjie Yu for related collaborations.

This work was supported in part by the Gordon and Betty Moore Foundation’s EPIQS Initiative through Grant No. GBMF4305 at the University of Illinois and the French ANR program ANR-11-IS04-005-01. The code is based on the PETSc [49–51], SLEPc[52] and MUMPS[37, 38] libraries and calculations were partly performed using HPC resources from CALMIP (grant 2015-P0677) as well as on Blue Waters. This research is part of the Blue Waters sustained-petascale computing project, which is supported by the National Science Foundation (awards OCI-0725070 and ACI-1238993) and the state of Illinois. Blue Waters is a joint effort of the University of Illinois at Urbana-Champaign and its National Center for Supercomputing Applications.

* dluitz@illinois.edu

- [1] J. v. Neumann, *Zeitschrift für Physik* **57**, 30 (1929)EPJ H **35**, 201 (2010)
- [2] I. Bloch, J. Dalibard, and W. Zwerger, *Rev. Mod. Phys.* **80**, 885 (2008)
- [3] P. W. Anderson, *Phys. Rev.* **109**, 1492 (1958)
- [4] D. M. Basko, I. L. Aleiner, and B. L. Altshuler, *Annals of Physics* **321**, 1126 (2006)
- [5] R. Nandkishore and D. A. Huse, *Annual Review of Condensed Matter Physics* **6**, 15 (2015)
- [6] E. Altman and R. Vosk, *Annual Review of Condensed Matter Physics* **6**, 383 (2015)
- [7] R. Vasseur and J. E. Moore, “Nonequilibrium quantum dynamics and transport: from integrability to many-body localization,” (2016), arXiv:1603.06618
- [8] V. Yukalov, *Laser Physics Letters* **8**, 485 (2011)
- [9] L. D’Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol, “From Quantum Chaos and Eigenstate Thermalization to Statistical Mechanics and Thermodynamics,” (2015), arXiv:1509.06411
- [10] F. Borgonovi, F. M. Izrailev, L. F. Santos, and V. G. Zelevinsky, *Physics Reports* **626**, 1 (2016)
- [11] M. V. Berry, *J. Phys. A: Math. Gen.* **10**, 2083 (1977)
- [12] M. L. Mehta, *Random Matrices* (Academic Press, New York, 1991)
- [13] J. M. Deutsch, *Phys. Rev. A* **43**, 2046 (1991)
- [14] M. Srednicki, *Phys. Rev. E* **50**, 888 (1994)
- [15] M. Srednicki, *J. Phys. A: Math. Gen.* **29**, L75 (1996)
- [16] M. Rigol, V. Dunjko, and M. Olshanii, *Nature* **452**, 854 (2008)
- [17] M. Rigol and M. Srednicki, *Phys. Rev. Lett.* **108**, 110601 (2012)
- [18] R. Steinigeweg, J. Herbrych, and P. Prelovšek, *Phys. Rev. E* **87**, 012118 (2013)
- [19] T. N. Ikeda, Y. Watanabe, and M. Ueda, *Phys. Rev. E* **87**, 012125 (2013)
- [20] W. Beugeling, R. Moessner, and M. Haque, *Phys. Rev. E* **89**, 042112 (2014)
- [21] W. Beugeling, R. Moessner, and M. Haque, *Phys. Rev. E* **91**, 012144 (2015)
- [22] V. Alba, *Phys. Rev. B* **91**, 155123 (2015)
- [23] R. Mondaini, K. R. Fratus, M. Srednicki, and M. Rigol, *Phys. Rev. E* **93**, 032104 (2016)
- [24] D. J. Luitz, *Phys. Rev. B* **93**, 134201 (2016)
- [25] M. Srednicki, *J. Phys. A: Math. Gen.* **32**, 1163 (1999)
- [26] E. Khatami, G. Pupillo, M. Srednicki, and M. Rigol, *Phys. Rev. Lett.* **111**, 050403 (2013)
- [27] J. T. Edwards and D. J. Thouless, *J. Phys. C: Solid State Phys.* **5**, 807 (1972)
- [28] S. Alexander, J. Bernasconi, W. R. Schneider, and R. Orbach, *Rev. Mod. Phys.* **53**, 175 (1981)
- [29] M. Serbyn, Z. Papić, and D. A. Abanin, *Phys. Rev. X* **5**, 041047 (2015)
- [30] Y. Bar Lev, G. Cohen, and D. R. Reichman, *Phys. Rev. Lett.* **114**, 100601 (2015)
- [31] K. Agarwal, S. Gopalakrishnan, M. Knap, M. Müller, and E. Demler, *Phys. Rev. Lett.* **114**, 160401 (2015)
- [32] D. J. Luitz, N. Laflorencie, and F. Alet, *Phys. Rev. B* **93**, 060201 (2016)
- [33] S. Gopalakrishnan, K. Agarwal, E. A. Demler, D. A. Huse, and M. Knap, *Phys. Rev. B* **93**, 134206 (2016)
- [34] V. K. Varma, A. Leroze, F. Pietracaprina, J. Gould, and A. Scardicchio, “Energy diffusion in the ergodic phase of a many body localizable spin chain,” (2015), arXiv:1511.09144
- [35] M. Znidaric, A. Scardicchio, and V. K. Varma, “Diffusive and subdiffusive spin transport in the ergodic phase of a many-body localizable system,” (2016), arXiv:1604.08567
- [36] I. Khait, S. Gazit, N. Y. Yao, and A. Auerbach, *Phys. Rev. B* **93**, 224205 (2016), 1603.06588
- [37] P. R. Amestoy, I. S. Duff, J. Koster, and J.-Y. L’Excellent, *SIAM J. Matrix Anal. Appl.* **23**, 15 (2001)
- [38] P. R. Amestoy, A. Guermouche, J.-Y. L’Excellent, and S. Pralet, *Parallel Computing* **32**, 136 (2006)
- [39] D. J. Luitz, N. Laflorencie, and F. Alet, *Phys. Rev. B* **91**, 081103 (2015)
- [40] A. Nauts and R. E. Wyatt, *Phys. Rev. Lett.* **51**, 2238 (1983)
- [41] K. Fabricius and B. M. McCoy, *Phys. Rev. B* **57**, 8340 (1998)
- [42] A. De Luca, B. L. Altshuler, V. E. Kravtsov, and A. Scardicchio, *Phys. Rev. Lett.* **113**, 046806 (2014)
- [43] A. D. Luca and A. Scardicchio, *EPL* **101**, 37003 (2013)
- [44] X. Chen, X. Yu, G. Y. Cho, B. K. Clark, and E. Fradkin, *Phys. Rev. B* **92**, 214204 (2015)
- [45] E. J. Torres-Herrera and L. F. Santos, *Phys. Rev. B* **92**, 014208 (2015)
- [46] B. L. Altshuler, E. Cuevas, L. B. Ioffe, and V. E.

- Kravtsov, “Non-ergodic phases in strongly disordered random regular graphs,” (2016), [arXiv:1605.02295](https://arxiv.org/abs/1605.02295)
- [47] K. S. Tikhonov, A. D. Mirlin, and M. A. Skvortsov, “Anderson localization on random regular graphs,” (2016), [arXiv:1604.05353](https://arxiv.org/abs/1604.05353)
- [48] C. Monthus, “Many-Body-Localization Transition : strong multifractality spectrum for matrix elements of local operators,” (2016), [arXiv:1603.04701](https://arxiv.org/abs/1603.04701)
- [49] S. Balay, W. D. Gropp, L. C. McInnes, and B. F. Smith, in *Modern Software Tools in Scientific Computing*, edited by E. Arge, A. M. Bruaset, and H. P. Langtangen (Birkhäuser Press, 1997) pp. 163–202
- [50] S. Balay, S. Abhyankar, M. F. Adams, J. Brown, P. Brune, K. Buschelman, V. Eijkhout, W. D. Gropp, D. Kaushik, M. G. Knepley, L. C. McInnes, K. Rupp, B. F. Smith, and H. Zhang, *PETSc Users Manual*, Tech. Rep. ANL-95/11 - Revision 3.5 (Argonne National Laboratory, 2014)
- [51] S. Balay, S. Abhyankar, M. F. Adams, J. Brown, P. Brune, K. Buschelman, V. Eijkhout, W. D. Gropp, D. Kaushik, M. G. Knepley, L. C. McInnes, K. Rupp, B. F. Smith, and H. Zhang, “PETSc Web page,” <http://www.mcs.anl.gov/petsc> (2014)
- [52] V. Hernandez, J. E. Roman, and V. Vidal, *ACM Trans. Math. Softw.* **31**, 351 (2005)