

This is the accepted manuscript made available via CHORUS. The article has been published as:

Ringed Structures of the HD 163296 Protoplanetary Disk Revealed by ALMA

Andrea Isella, Greta Guidi, Leonardo Testi, Shangfei Liu, Hui Li, Shengtai Li, Erik Weaver, Yann Boehler, John M. Carperter, Itziar De Gregorio-Monsalvo, Carlo F. Manara, Antonella Natta, Laura M. Pérez, Luca Ricci, Anneila Sargent, Marco Tazzari, and Neal Turner

Phys. Rev. Lett. **117**, 251101 — Published 12 December 2016

DOI: [10.1103/PhysRevLett.117.251101](https://doi.org/10.1103/PhysRevLett.117.251101)

Ringed structure of the HD 163296 disk revealed by ALMA

Andrea Isella,¹ Greta Guidi,² Leonardo Testi,^{2,3} Shangfei Liu,^{1,4} Hui Li,⁴
Shengtai Li,⁴ Erik Weaver,¹ Yann Boehler,¹ John M. Carperter,⁵ Itziar
De Gregorio-Monsalvo,⁵ Carlo F. Manara,⁶ Antonella Natta,^{7,2} Laura M.
Pérez,⁸ Luca Ricci,⁹ Anneila Sargent,¹⁰ Marco Tazzari,³ and Neal Turner¹¹

¹*Department of Physics and Astronomy, Rice University,
6100 Main St., 77005, MS-108, Houston, Texas**

²*INAF/Osservatorio Astrofisico di Arcetri,
Largo E. Fermi 5, I-50125 Firenze, Italy*

³*ESO, Karl Schwarzschild str. 2, 85748 Garching bei Munchen, Germany*

⁴*Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA*

⁵*Joint ALMA Observatory (JAO), Alonso de
Cordova 3107 Vitacura - Santiago de Chile, Chile*

⁶*Scientific Support Office, Directorate of Science,
European Space Research and Technology Centre (ESA/ESTEC),
Keplerlaan 1, 2201 AZ Noordwijk, The Netherlands*

⁷*School of Cosmic Physics, Dublin Institute for Advanced Studies,
31 Fitzwilliams Place, 2 Dublin, Ireland*

⁸*Max-Planck-Institut fr Radioastronomie Bonn,
auf dem Hgel 69, D-53121 Bonn, Germany*

⁹*Harvard-Smithsonian Center for Astrophysics,
60 Garden Street, Cambridge, MA 02138, USA*

¹⁰*Division of Physics, Mathematics and Astronomy,
California Institute of Technology, Pasadena, CA 91125, USA*

¹¹*Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109, USA*

(Dated: October 28, 2016)

Abstract

We present Atacama Large Millimeter and sub-millimeter Array (ALMA) observations of the protoplanetary disk around the Herbig Ae star HD 163296 that trace the spatial distribution of millimeter-sized particles and cold molecular gas on spatial scales as small as 25 astronomical units (au). The image of the disk recorded in the 1.3 mm continuum emission reveals three dark concentric rings which indicate the presence of dust depleted gaps at about 60 au, 100 au, and 160 au from the central star. The maps of the ^{12}CO , ^{13}CO , and C^{18}O J=2-1 emission do not show such structures but reveal a change in the slope of the radial intensity profile across the positions of the dark rings in the continuum image. By comparing the observations with theoretical models for the disk emission, we find that the density of CO molecules is reduced inside the middle and outer dust gaps. However, in the inner ring there is no evidence of CO depletion. From the measurements of the dust and gas densities, we deduce that the gas-to-dust ratio varies across the disk and, in particular, it increases by at least a factor 5 within the inner dust gap compared to adjacent regions of the disk. The depletion of both dust and gas suggests that the middle and outer rings could be due to the gravitational torque exerted by two Saturn-mass planets orbiting at 100 au and 160 au from the star. On the other hand, the inner dust gap could result from dust accumulation at the edge of a Magneto-Rotational instability (MRI) dead zone, or from dust opacity variations at the edge of the CO frost line. Observations of the dust emission at higher angular resolution and of molecules that probe dense gas are required to establish more precisely the origins of the dark rings observed in the HD 163296 disk.

INTRODUCTION

After centuries of speculation about planets orbiting stars other than our Sun, the discovery of an extrasolar planet with Jupiter-like-mass came in 1995 [1]. To date, the most fruitful methods of planetary searches have been transit and radial velocity surveys [2, 3], especially *Kepler*'s transit surveys [4]. However, the discoveries of more than 2000 exoplanet candidates [5], mostly gas giants and super-Earths, have raised even more challenges to understanding such systems. For example, while giant planets are believed to form at 0.1-10 au from the central star [6], the newly-discovered orbital radii cover a much broader range, 0.01-100 au. This suggests either a surprisingly large mobility of planets or a fundamental flaw in current planetary formation models.

Recently, the unprecedented angular resolution and sensitivity of Atacama Large Millimeter and sub-millimeter Array (ALMA) observations have led to the discovery of ring-shaped structures in the mm continuum emission from the HL Tau and TW Hya protoplanetary disks [7, 8]. While the origin of these structures is still debated, it is likely that they are related to the formation of planets, and hence may provide long-awaited observational constraints on the processes responsible for planetary system assembly and evolution.

A number of explanations for the presence of rings in the continuum emission in protoplanetary disks have been put forward. These include the aggregation of solids at the outer edge of disk regions characterized by low turbulence [e.g., 9–14], variations in the dust opacity at the frost line of volatile elements [15, 16], and gravitational perturbations caused by yet unseen giant planets [17]. In the first two cases, the observed structures might aid the formation of planetesimals by concentrating solid particles and preventing them from accreting onto the star [18, 19]. In the last case, a circular gap is expected when a forming planet is orbiting in a disk. The orbital radius and mass of the planet can be inferred from the location and shape of this gap [17]. For HL Tau, it has been suggested that the observed ringed structure is due to the presence of Saturn mass planets orbiting at several tens of au from the central star [20]. This implies that giant planets can form on much shorter timescales, and at much larger distances from the central star, than predicted by current planet formation models [6].

The initial results of ALMA observations are promising but key questions remain unanswered. In particular it is not clear if ring-like structures are common to all protoplanetary

disks. ALMA high angular resolution observations targeted the most luminous disks first, and these may not be representative of the bulk of the disk population. Furthermore, dust continuum images do not probe the distribution of the circumstellar gas and thus cannot distinguish unambiguously between different formation processes. Addressing such questions will require a multi-pronged approach involving both large continuum surveys of nearby protoplanetary disks with ALMA and in-depth multi-wavelength studies of the most relevant structures revealed.

In this Letter, we report on the structure of the HD 163296 disk as revealed by ALMA observations of the dust continuum emission and also line emission from three isotopologues of carbon monoxide, ^{12}CO , ^{13}CO , and C^{18}O . HD 163296 is a young (5 Myr) intermediate mass ($2.3 M_{\odot}$) Herbig Ae star located at 122^{+17}_{-13} pc from Earth [21]. The star is surrounded by a gaseous Keplerian disk with a radius of about 550 au. By contrast, mm-wave continuum emission arising from solid particles is confined within 250 au from the star [22, 23]. Previous ALMA observations showed that the continuum emission has a local maximum at about 90 au [24], suggesting a dusty ring at a distance that corresponds to the frost line for CO molecules [25]. Recently, we have imaged the HD 163296 disk with ALMA at $0.2''$ angular resolution, a factor of 3 higher than previous observations and corresponding to a physical scale of 25 au at the distance of the star. The observations reveal rings in the dust continuum emission similar to those observed toward HL Tau. They also reveal the morphology of the ^{12}CO , ^{13}CO , and C^{18}O emission on the same angular scales as the continuum. Here we show that the combination of continuum and molecular line observations offers stringent tests for several physical processes related to planet formation and planet-disk interaction.

OBSERVATIONS

Figure 1 shows the images of the HD 163296 continuum emission together with those of the continuum-subtracted, spectrally integrated ^{12}CO , ^{13}CO , and C^{18}O J=2-1 lines. The acquisition and calibration of the data are described in the Supplementary Material. Also displayed are the radial intensity profiles across the disk calculated by averaging the flux density over elliptical annuli characterized by a position angle of 132° and an eccentricity of 0.7 to correct for the disk inclination and orientation [22, 25].

The continuum emission is in first approximation centrosymmetric and confined to an




FIG. 1. ALMA images of the HD 163296 disk emission recorded in 1.3 mm dust continuum (top left), ^{12}CO (top middle), ^{13}CO (bottom left), and C^{18}O (bottom middle) $J=2-1$ line emission. The angular resolution of the observations, $0.22'' \times 0.14''$, is indicated by the filled white ellipse in the continuum image. The dashed ellipses in the CO maps indicate the locations of the dark rings seen in the continuum map. Azimuthally averaged profiles, normalized to the peak intensities, are shown in the right panels. They are calculated by averaging on elliptical annuli with a position angle of 132° , an eccentricity of 0.7, and a width equal to $1/4$ of the angular resolution of the observations. The error bars are calculated by dividing the root mean square noise of the observations (see Table 1 in the Supplemental Material) by the square root of the number of independent beams in each annulus. The vertical lines indicate the position of the dark (D) and bright (B) rings observed in the continuum map. The horizontal bar in the top right panel indicates the angular resolution of the observations.

angular radius of about $2''$ from the central star. **It is also characterized by three pairs of low and high luminosity elliptical rings, labeled as (D1, B1), (D2, B2), and (D3, B3), where (D1,B1) and (D3, B3) are the innermost and outermost pairs, respectively.** The aspect ratios of the rings correspond to an intrinsically circular disk inclined by 42° with respect to the line of sight; 90° corresponds to an edge-on orientation. The inclination agrees with previous measurements based on CO observations [22, 25]. The radial profile of the dust continuum emission decreases by a factor of 1000 within $2''$ of the star. The ratios of the continuum intensity for each high/low luminosity ring pair are 1.6 (B1/D1), 1.3 (B2/D2), and 1.08 (B3/D3). In first approximation, the azimuthally averaged profile of the continuum flux density can be modeled by a combination of Gaussian functions, $f(\theta) = g(\theta) - g_1(\theta) - g_2(\theta) - g_3(\theta)$, where $g(\theta) = ae^{-(\theta/b)^2}$ and $g_n(r) = a_n e^{-((\theta-\theta_n)/b_n)^2}$. This profile mimics that of a disk with gaps in the continuum flux density characterized by radius θ_n and width b_n . A nonlinear least-squares Marquardt-Levenberg fit of the observations returns $\theta_1 = (0.44 \pm 0.01)''$, $\theta_2 = (0.81 \pm 0.01)''$, $\theta_3 = (1.13 \pm 0.02)''$, and $b = (0.76 \pm 0.02)''$, $b_1 = (0.23 \pm 0.01)''$, $b_2 = (0.15 \pm 0.01)''$, $b_3 = (0.28 \pm 0.02)''$, with a reduced $\chi^2 \sim 2$. **A table with the best fit value of the gaussian modeling is provided in the Supplemental Material.**

Molecular emission is detected in all three CO isotopologues, and differs from the continuum in three main aspects. First, the CO intensity maps do not show any prominent ringed structure and, differently from the continuum, their radial profiles cannot be approximated by a simple mathematical function. Second, the ^{13}CO and C^{18}O line intensities have local minima at the center of the disk, while both the continuum and the ^{12}CO line are centrally peaked. We ascribe the ^{13}CO or C^{18}O central minima to an over subtraction of the continuum emission, and not to a physical decrement of the gas density or temperature. This point is discussed in more details in the Supplemental material. Third, the molecular emission is much more extended and decreases with the orbital radius more slowly than the continuum. In particular, the ^{12}CO emission decreases only to about 15% of its peak value at radius $2''$. Zooming into the central disk regions, as in the bottom right panel on Figure 1, indicates that only mild changes in the CO emission occurs at the location of the continuum rings. A bump in ^{12}CO emission is observed near D1 and a flattening of the slope of the emission can be seen near B2 for all three isotologues, with the C^{18}O emission increasing slightly between $1''$ and $1.2''$.

DATA ANALYSIS

In this section, we investigate the ringed structure observed in the continuum and the aforementioned morphological differences between continuum and molecular line emission. We do this by modeling the continuum and spectrally integrated line emission maps presented in Figure 1. The analysis of the spectrally resolved line emission will be presented in a future work.

The continuum and line emission of a protoplanetary disks depend on the three dimensional density and temperature distribution of dust and gas, the gas velocity, the dust opacity, the disk inclination, and the distance of the source. In recent years, the HD 163296 disk has been the subject of several studies whose results provide a starting point for our investigation. Following the methodology described in [23] and [26], we describe the surface density of gas and dust as

$$\Sigma(r) = \Sigma_c (r/r_c)^{-\gamma} \exp[-(r/r_c)^{2-\gamma}], \quad (1)$$

and the disk temperature as

$$T(r, z) = \begin{cases} T_a(r, z) + [T_m(r) - T_a(r, z)] \left[\cos \frac{\pi z}{2z_q(r)} \right]^{2d(r)}, & \text{if } |z| < z_q(r) \\ T_a(r, z), & \text{otherwise} \end{cases} \quad (2)$$

In the latter equation, $T_m(r) = T_{m,0}(r/r_0)^{-q_m}$ and $T_a(r, z) = T_{a,0}(\sqrt{r^2 + z^2}/r_0)^{-q_a}$ are the temperature of the disk midplane and surface layer, respectively. The temperatures of these two layers are smoothly connected through the functions $d(r) = d_0(r - r_0) + c$ and $z_q(r) = z_{q,1}(r/r_1)^{q_z} e^{-(r/r_2)^2}$. This rather complex parameterization of the disk temperature mimics the thermal structure of an externally illuminated disk and enable us to generate synthetic models for the dust and gas emission avoiding calculating the disk temperature using time consuming radiative transfer models.

Using Equation 1 and 2, and the procedure described in the Supplemental Material, we find that the dust emission measured at angular radii less than $0.3''$ and larger than $1.8''$ is consistent with a dust surface density characterized by $\gamma = 0.1$, $r_c = 90$ au, $\Sigma_c = 0.42$ g cm⁻². **The low value of γ implies that the dust density within about 90 au from the star scales with the radius approximately as a Gaussian. This result is consistent with the parametric fitting of the dust continuum emission described in the previous section.**




FIG. 2. Constraints on the dust and gas surface density (left and middle panels, respectively, and on the gas-to-dust ratio (right panel), derived by comparing ALMA observations of the HD163296’s disk with synthetic models for the disk emission. The red solid curve indicates the prediction for a disk model perturbed by three planets with masses of 0.1, 0.3, and 0.3 M_J orbiting at 62, 105, and 160 au from the central star.

Applying the same procedure to the ^{12}CO , ^{13}CO and C^{18}O emission, we find gas surface densities characterized by $\gamma = 0.8$, $r_c = 165$ au, $\Sigma_c(^{12}\text{CO}) = 1.6 \times 10^{-3} \text{ g cm}^{-2}$, $\Sigma_c(^{13}\text{CO}) = 2.3 \times 10^{-5} \text{ g cm}^{-2}$, and $\Sigma_c(\text{C}^{18}\text{O}) = 2.9 \times 10^{-6} \text{ g cm}^{-2}$. Our modeling suggests that (i) the $^{12}\text{CO}/^{13}\text{CO}$ and $^{12}\text{CO}/\text{C}^{18}\text{O}$ density ratios are radially constant, (ii) the CO/dust ratio changes with the orbital radius, and (iii) that the disk thermal structure is characterized by $T_m(r) = 24\text{K}(r/100\text{au})^{-0.5}$ and $T_a(r, z) = 68\text{K}(\sqrt{r^2 + z^2}/100\text{au})^{-0.6}$.

However, disk models characterized by a smooth density profile predict too much emission in the region where rings are seen in dust continuum. To correct for this excess, we introduce circular gaps in the dust and gas surface density. Each gap is described by three parameters: the distance between the center of the gap and the star (R), the width of the gap (W), and the ratio between the surface density inside and outside the gap (Δ). We calculate synthetic images for the dust continuum, ^{12}CO , ^{13}CO , and C^{18}O J=2-1 emission for different values of the gap parameters (R, W, Δ), and compare them to the observations. Figure 2 shows the set of models that better reproduce the observations, while the modeling procedure is described in the Supplemental Material.

The colored area in left panel of Figure 2 shows the values of the dust surface density compatible with the observations. The dark rings observed in the continuum map are compatible with the presence of three partially dust depleted gaps centered at $R_1 = 60$ au,

$R_2 = 100$ au, and $R_3 = 160$ au. Given the finite angular resolution of the observations, there is a degeneracy between the gap width W and dust depletion Δ^d . We find that the narrowest gaps compatible with the observations have widths of $W_1 = 25$ au, $W_2 = 22$ au, and $W_3 = 45$ au. The dust depletion factors inside the gaps are $\Delta_1^d > 100$, $\Delta_2^d = 70$, and $\Delta_3^d = 6$. The widest gaps compatible with the observations have widths of $W_1 = 33$ au, $W_2 = 26$ au, and $W_3 = 55$ au, and depletion factors $\Delta_1^d = 13$, $\Delta_2^d = 7$, and $\Delta_3^d = 3.6$. The dust surface density inside the inner gap is $\lesssim 3 \times 10^{-2}$ g cm $^{-2}$, while it varies between 3×10^{-3} - 3×10^{-2} g cm $^{-2}$ and 4×10^{-3} - 7×10^{-3} g cm $^{-2}$ in the middle and outer gaps, respectively. The ratios between the half-width and the radius of the gaps are between 0.21-0.28 for the innermost gap, 0.1-13 for the middle gap, and 0.14-0.17 for the outermost gap.

The central panel of Figure 2 shows the constraints on the gas (H_2) surface density derived from the CO observations adopting a cosmic molecular abundance $n(^{12}\text{CO}/\text{H}_2) = 5 \times 10^{-5}$ [27]. In our analysis, we assume that the gaps in the dust and CO have the same width and radius, and explore different values for the depletion factor Δ^{CO} . Inside the inner dust gap, we measure Δ_1^{CO} between 0-2.5. This implies that CO molecules are at least 5 times less depleted than the dust. In the middle and outer gaps, we measure Δ_2^{CO} between 3.5-70 and Δ_3^{CO} between 1.8-6, respectively. These latter values are similar to those inferred for the dust.

Finally, the right panel of Figure 2 shows the gas-to-dust ratio obtained by combining the results for the dust and gas surface density. The gas-to-dust ratio increases from about 40 at 100 au to more than 1000 for $r > 250$ au. The gas-to-dust ratio inside the first dust gap is at least 5 times larger than outside the gap. Such an increase results from the discrepancy between the depletion in dust and CO. Inside the middle and outer dust gaps the gas-to-dust ratio varies by less than a factor of 2. Our model also suggests that the gas-to-dust ratio might increase for $r < 50$ au. However, this result should be taken with caution because both the dust and CO emission inside this radius are optically thick and the corresponding surface densities are affected by a large uncertainty.

DISCUSSION

To date, the HD 163296 disk is the third system known to host multiple rings in the dust continuum emission, but it is the only one for which we have succeeded in mapping

the molecular gas emission at the same angular resolution of the continuum. As mentioned above, such ringed structures are thought to be related to the formation of planets, but it is unclear whether they trace an early stage of planet assembly or whether they result from the interaction with already formed planets. In this section we discuss how measuring the spatial distribution of both gas and dust is key to address this question.

Disk-planet interaction: The interaction between disks and planets is based on the notion that the gravitational field of a planet exerts a tidal torque on the circumstellar material, which allows the transfer of angular momentum between the planet and the disk [see, e.g., 28]. The consequences are twofold: one an hand, the planet is predicted to migrate; on the other hand, the density waves launched by the planet are expected to alter the disk surface density. If the planet gravity exceeds the stellar gravity across the vertical extent of the disk, then the planet opens a circular gap in the disk surface density [17]. The radius, width, and level of gas and dust depletion within the gap carry information about the planet and disk properties.

Numerical simulations show that the width W of a gap opened by a planet should depend on the planet mass M_p and orbital radius R_p following the relation

$$W \sim C \times R_p \left(\frac{M_p}{3M_\star} \right)^{1/3}, \quad (3)$$

where the proportionality constant C varies between 4-8 [17, 29, 30]. In the case of HD 163296, the gap widths derived in the previous section would imply planet masses between 0.5-2 M_J for the innermost planet, 0.05-0.3 M_J for the middle planet, and 0.15-0.5 M_J for the outer planet. In addition, the disk viscosity can affect both the gap width and depth [31]. Various models have predicted that, at equilibrium, the depth of a gap opened by a planet in the gas surface density should depend on the planet mass through the relation [32]

$$\frac{M_p}{M_\star} = 5 \times 10^{-4} (\Delta - 1)^{1/2} \left(\frac{h}{0.1} \right)^{5/2} \left(\frac{\alpha}{10^{-3}} \right)^{1/2}, \quad (4)$$

where h is the gas disk aspect ratio, Δ is the gas depletion ratio, and α is the Shakura-Sunyaev viscosity parameter [33]. This relation indicates that generally a lower disk viscosity and aspect ratio would result in a deeper gap for a fixed planet mass.

In the case of HD 163296, we use our disk temperature model derived from the observations to calculate a disk aspect ratio of 0.05 at 60 au, 0.065 at 100 au, and 0.07 at

160 au. By combining Equation 3 and Equation 4 we estimate the viscosity parameter α . We obtain $\alpha \gtrsim 2 \times 10^{-2}$ in the inner gap, $10^{-6} < \alpha < 5 \times 10^{-3}$ in the middle gap, and $10^{-4} < \alpha < 7 \times 10^{-3}$ in the outer gap. The larger α of the inner gap results from the fact that low or no gas depletion is observed despite a dust gap width that is larger than those of the other two dust gaps. These heuristic considerations work reasonably well for the middle and outer gaps but the inner gap presents additional challenges. The gas-to-dust ratio is the largest for this gap, which for a given planet mass, would require a relatively low viscosity [e.g., 19], in contrast to the value derived above. This discrepancy suggests that additional processes might be happening around the first gap, including the possibility of having multiple planets.

To better understand the effect of planet perturbations on the structure of the HD 163296 disk, we have compared the gas and dust surface density derived from ALMA observations with hydrodynamic simulations of the planet-disk interaction. The setup of the simulations is described in the Supplemental Material. The red line in Figure 2 shows the dust and gas surface density for a disk perturbed by three planets with masses of $0.1 M_J$, $0.3 M_J$, and $0.3 M_J$ orbiting at 62 au, 105 au, and 160 au from the star. The surface density profiles were extracted after 2000 orbits of the innermost planet corresponding 7×10^5 yr.

We find that the gas and dust surface density measured across the middle and outer gaps are compatible with perturbations from Saturn mass planets and a disk viscosity parameter $10^{-3} < \alpha < 10^{-2}$. The planets carve rather shallow and axisymmetric gaps in the gas surface density, whereas the gaps in the dust are deeper due to migration of solid particles toward the edges of the gas gaps. As a consequence, the gas-to-dust ratio inside the gaps formed by the planets increases with respect to the surrounding disk regions.

Our simulations show that perturbations from a single planet cannot explain the properties of the inner gap. The width of the gap generated by a $0.1 M_J$ planet is narrower than that suggested by the observations. A more massive planet would make the gap wider, but the gap would be too depleted in gas. Increasing both the planet mass and the disk viscosity would make the gap wider and shallower, but, a larger viscosity would cause the gas-to-dust ratio inside the gap to drop well below 5. As a possible solution, we argue that the inner gap might be created by multiple planets between 50-70 au. For example, two Saturn mass planets orbiting around 55 au and 65 au might create a joint gap consistent with the observations though the stability of such a configuration remains to be investigated

[see, e.g., 34]. However, observations with higher angular resolution would be required to test this hypothesis.

MRI “dead-zone”: An ionized Keplerian disk embedded in a weak magnetic field is predicted to be subject to the magneto-rotational instability [MRI, 35]. MRI should generate turbulent motions capable of transporting angular momentum outward, therefore allowing the circumstellar material to accrete onto the star. If the gas ionization varies with the orbital radius, then the mass flow rate throughout the disk is not constant and gas and dust can pile up forming ringed structures.

There are two main transitions in the gas ionization in a disk characterized by a monotonically decreasing surface density. The first happens at a fraction of au from the star where the gas temperature exceed 1000 K and the thermal ionization is strong. The ionized region is predicted to have high turbulence velocities and quickly accrete onto the central star, possibly leading to the formation of a large central hole [36]. At larger distances the ionization is non-thermal and caused mostly by stellar ultraviolet and X-ray photons and interstellar cosmic rays, which are absorbed over columns of 0.1, 10 and 100 g cm⁻², respectively. [37–39]. The ionized regions experience MRI which leads to turbulent speeds that increase with height above the disk midplane in proportion to the Alfvén speed, reaching the local sound speed above several disk scale heights [40, 41]. At surface densities above about 10 g cm⁻², the disk midplane is expected to be mostly neutral and quiescent. In the outermost disk regions, for $\Sigma < 10$ g cm², the disk midplane returns to be partially ionized. The sharp gradient in turbulence, and henceforth in mass flow rate, at the interface between the MRI “dead-zone” and the MRI active midplane is predicted to lead to a pile up of dust and gas, which might appear as an emission ring once observed with ALMA [42].

MHD simulations of protoplanetary disks predict that the gas surface density at the outer edge of an MRI dead zone might increase by a factor of a few compared to the surrounding regions [42]. The increase in gas density creates a gas pressure maxima which might trap dust particles, therefore augmenting the radial variation in the dust surface density. Our measurement of the gas surface density of the HD 163296 disk (central panel of Figure 2) indicates that the MRI “dead zone” might extend to about 60-100 au from the central star. This distance is similar to the radius of the first bright continuum ring, suggesting that this process might perhaps be involved in the ring formation. Furthermore, the large gas-to-dust ratio measured inside the inner dust gap would imply low viscosity. This is consistent with

the presence of a MRI “dead-zone”. The middle and outer dust gaps at 100 au and 160 au from the star do not seem to be compatible with this scenario though.

Phase transition of volatile elements: The phase transition from solid to gas of molecules such as H_2O and CO is predicted to alter the coagulation and fragmentation properties of dust grains, possibly leading to the accumulation of dust particles in ring-like structures. Dust grains located in the cold regions of protoplanetary disks are expected to be covered by ice mantles, which make them more sticky compared to bare grains and more prone to growth in size as result of collisions. As the size of a dust grain increases, so does the drag caused by the aerodynamic friction with the circumstellar gas. As a result of this process, larger icy grains are predicted to drift inward toward warmer disk regions. Eventually they will cross the frost lines relative to the molecules that form their ice mantles, which will then evaporate. When this happens, the grain size shrinks affecting both its opacity and the gas drag. The combined result is an increase in the dust opacity right behind the frost line, which might result in the formation of bright emission rings in the microwave continuum emission [15, 43].

A second effect of the evaporation of volatile elements from dust grains is described by the so called “sintering” model [16] which postulates that the grains are fractal aggregates of smaller particles. Outside each volatile’s frost line, the aggregates are covered by a layer of ice. As inward drifting particles approach the frost line, the ice molecules become mobile and migrate to the necks between the component particles, making rolling and sliding difficult. This reduces the internal degrees of freedom available to take up excess energy when aggregates collide. The aggregates become hard but brittle, and break apart easily in collisions. The smaller fragments drift more slowly, enhancing the dust surface density near the snow line and causing radial variations in the dust opacity [16].

In general, the snow lines of H_2O and CO should be the easiest to observed due to the large abundance of these molecules. At the typical densities of protoplanetary disks, water and CO condensate at temperature between 130-150 K and 18-28 K respectively [15, 25]. At the temperatures of the HD 163296 disk (Figure 3), the water snow line occurs within 3 au from the star and is not resolved by our observations. Instead, the CO snow line might occur between 40-180 au and might overlap with the ringed structures observed in the continuum.

A key aspect of the sintering and frost line models is that the formation of continuum emission rings by evaporation of volatile elements does not require and might not produce

any variation in the gas distribution. Actually, these models require a constant radial inflow of particles which will not happen if the gas density has bumps in the radial direction. Therefore, the depletion of CO molecules measured inside the middle and outer gaps is inconsistent with the evaporation of volatile elements. However, this process might play a role in shaping the disk regions around the innermost dust gap at 59 au, where the CO observations do not show any evidence of gas depletion. Future ALMA observation of molecules capable of tracing denser gas, e.g., HCO+ [44], are required to set more stringent constraints on the nature of this inner gap and on the role of the evaporation of volatile elements.

CONCLUSIONS

We have presented ALMA observations of the HD 163296 circumstellar disk that reveal three dark rings in the dust continuum emission. HD 163296 is the third system known to host multiple rings in the dust continuum emission, but it is the only one for which we have succeeded in mapping the molecular gas emission at the same angular resolution of the continuum. By comparing the observations with radiative transfer models, we find that the continuum rings can be explained with the presence of three circular dust depleted gaps with radii of 60 au, 100 au, and 160 au. Furthermore, we find that the density of CO molecules inside the middle and outer dust gaps is also reduced compared to the surrounding regions.

We argue that the morphology of these two gaps is consistent with gravitational perturbations caused by two Saturn-mass planets orbiting at about 100 au and 160 au. Such large orbital radii and the young age of this disk, would imply that giant planets might form much farther away from the central star and much more rapidly than previously thought. Finally, we note that the presented planet configuration might not be unique. A detailed analysis of the HD 163296 ALMA observations in the framework of planet-disk interaction models will be presented in Liu et al. (in preparation).

We do not find any evidence of gas depletion inside the inner dust gap although, due to the large optical depth of the observed CO lines, we cannot rule out the presence of a shallow gas gap. The morphology of the inner dust gap is not well described by perturbations caused by a single planet. Instead we suggest that it might have formed as the result of opacity variations at the outer edge of a MRI “dead-zone” or at the CO frost line.




FIG. 3. Temperature of the HD 163296 disk inferred from the observations of the CO emission. The white solid lines correspond to temperatures of 18 K, 23 K, and 28 K, and indicate the range of condensation temperatures for CO molecules. The vertical dashed line indicate the location of the dust gaps. The green and yellow curves indicate the pressure scale height of gas (h_{gas}) and of large dust grains (h_{dust}), respectively.

The results presented in this Letter highlight the importance of imaging protoplanetary disks in both the dust and molecular line emission at high angular resolution. Thanks to the large flux and radial extent, the HD 163296 disk is a perfect target for future ALMA followup observations. Furthermore, differently from HL Tau, the relative isolation of the

HD 163296 disk enables optical and infrared observations to map the morphology of the disk surface and, perhaps, directly detect forming planets.

We thank Munetake Momose, Misato Fukagawa, and Giovanni Rosotti for the helpful conversation. This paper makes use of the following ALMA data: ADS/JAO.ALMA#2013.1.00601.S. ALMA is a partnership of ESO (representing its member states), NSF (USA) and NINS (Japan), together with NRC (Canada) and NSC and ASIAA (Taiwan) and KASI (Republic of Korea), in cooperation with the Republic of Chile. The Joint ALMA Observatory is operated by ESO, auI/NRAO and NAOJ. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. The National Radio Astronomy Observatory is a facility of the National Science Foundation operated under cooperative agreement by Associated Universities, Inc. A. Isella and Y. Boehler acknowledge support from the NSF award AST-1535809 and the NASA award NNX15AB06G. E. Weaver acknowledge support from the NRAO Student Observing Support award AST-0836064. H. Li and S. Li acknowledge the support from the LDRD program at LANL. C.F. Manara gratefully acknowledges an ESA Research Fellowship. Part of this research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration.

* isella@rice.edu

- [1] M. Mayor and D. Queloz, *Nature* (London) **378**, 355 (1995).
- [2] A. W. Howard, G. W. Marcy, J. A. Johnson, D. A. Fischer, J. T. Wright, H. Isaacson, J. A. Valenti, J. Anderson, D. N. C. Lin, and S. Ida, *Science* **330**, 653 (2010), arXiv:1011.0143 [astro-ph.EP].
- [3] M. Mayor, M. Marmier, C. Lovis, S. Udry, D. Ségransan, F. Pepe, W. Benz, J. . Bertaux, F. Bouchy, X. Dumusque, G. Lo Curto, C. Mordasini, D. Queloz, and N. C. Santos, *ArXiv e-prints* (2011), arXiv:1109.2497 [astro-ph.EP].
- [4] W. J. Borucki, D. G. Koch, G. Basri, N. Batalha, T. M. Brown, S. T. Bryson, D. Caldwell, J. Christensen-Dalsgaard, W. D. Cochran, E. DeVore, E. W. Dunham, T. N. Gautier, III, J. C. Geary, R. Gilliland, A. Gould, S. B. Howell, J. M. Jenkins, D. W. Latham, J. J. Lissauer,

- G. W. Marcy, J. Rowe, D. Sasselov, A. Boss, D. Charbonneau, D. Ciardi, L. Doyle, A. K. Dupree, E. B. Ford, J. Fortney, M. J. Holman, S. Seager, J. H. Steffen, J. Tarter, W. F. Welsh, C. Allen, L. A. Buchhave, J. L. Christiansen, B. D. Clarke, S. Das, J.-M. Désert, M. Endl, D. Fabrycky, F. Fressin, M. Haas, E. Horch, A. Howard, H. Isaacson, H. Kjeldsen, J. Kolodziejczak, C. Kulesa, J. Li, P. W. Lucas, P. Machalek, D. McCarthy, P. MacQueen, S. Meibom, T. Miquel, A. Prsa, S. N. Quinn, E. V. Quintana, D. Ragozzine, W. Sherry, A. Shporer, P. Tenenbaum, G. Torres, J. D. Twicken, J. Van Cleve, L. Walkowicz, F. C. Witteborn, and M. Still, *Astrophys. J.* **736**, 19 (2011), arXiv:1102.0541 [astro-ph.EP].
- [5] T. D. Morton, S. T. Bryson, J. L. Coughlin, J. F. Rowe, G. Ravichandran, E. A. Petigura, M. R. Haas, and N. M. Batalha, *Astrophys. J.* **822**, 86 (2016), arXiv:1605.02825 [astro-ph.EP].
- [6] R. Helled, P. Bodenheimer, M. Podolak, A. Boley, F. Meru, S. Nayakshin, J. J. Fortney, L. Mayer, Y. Alibert, and A. P. Boss, *Protostars and Planets VI*, 643 (2014), arXiv:1311.1142 [astro-ph.EP].
- [7] ALMA Partnership, C. L. Brogan, L. M. Pérez, T. R. Hunter, W. R. F. Dent, A. S. Hales, R. E. Hills, S. Corder, E. B. Fomalont, C. Vlahakis, Y. Asaki, D. Barkats, A. Hirota, J. A. Hodge, C. M. V. Impellizzeri, R. Kneissl, E. Liuzzo, R. Lucas, N. Marcelino, S. Matsushita, K. Nakanishi, N. Phillips, A. M. S. Richards, I. Toledo, R. Aladro, D. Brogiere, J. R. Cortes, P. C. Cortes, D. Espada, F. Galarza, D. Garcia-Appadoo, L. Guzman-Ramirez, E. M. Humphreys, T. Jung, S. Kamenoi, R. A. Laing, S. Leon, G. Marconi, A. Mignano, B. Nikolic, L.-A. Nyman, M. Radiszcz, A. Remijan, J. A. Rodón, T. Sawada, S. Takahashi, R. P. J. Tilanus, B. Vila Vilaro, L. C. Watson, T. Wiklind, E. Akiyama, E. Chapillon, I. de Gregorio-Monsalvo, J. Di Francesco, F. Gueth, A. Kawamura, C.-F. Lee, Q. Nguyen Luong, J. Mangum, V. Pietu, P. Sanhueza, K. Saigo, S. Takakuwa, C. Ubach, T. van Kempen, A. Wootten, A. Castro-Carrizo, H. Francke, J. Gallardo, J. Garcia, S. Gonzalez, T. Hill, T. Kaminski, Y. Kurono, H.-Y. Liu, C. Lopez, F. Morales, K. Plarre, G. Schieven, L. Testi, L. Videla, E. Villard, P. Andreani, J. E. Hibbard, and K. Tatematsu, *Astrophys. J. Lett.* **808**, L3 (2015), arXiv:1503.02649 [astro-ph.SR].
- [8] S. M. Andrews, D. J. Wilner, Z. Zhu, T. Birnstiel, J. M. Carpenter, L. M. Pérez, X.-N. Bai, K. I. Öberg, A. M. Hughes, A. Isella, and L. Ricci, *Astrophys. J. Lett.* **820**, L40 (2016), arXiv:1603.09352 [astro-ph.EP].
- [9] P. Barge and J. Sommeria, *Astron. Astrophys.* **295**, L1 (1995), arXiv:astro-ph/9501050.

- [10] H. H. Klahr and T. Henning, *Icarus* **128**, 213 (1997).
- [11] H. Li, S. A. Colgate, B. Wendroff, and R. Liska, *Astrophys. J.* **551**, 874 (2001), arXiv:astro-ph/0012479.
- [12] W. Lyra, A. Johansen, H. Klahr, and N. Piskunov, *Astron. Astrophys.* **491**, L41 (2008), arXiv:0807.2622.
- [13] W. Lyra and M.-K. Lin, *Astrophys. J.* **775**, 17 (2013), arXiv:1307.3770 [astro-ph.EP].
- [14] W. Fu, H. Li, S. Lubow, S. Li, and E. Liang, *Astrophys. J. Lett.* **795**, L39 (2014), arXiv:1410.4196 [astro-ph.EP].
- [15] K. Zhang, G. A. Blake, and E. A. Bergin, *Astrophys. J. Lett.* **806**, L7 (2015), arXiv:1505.00882 [astro-ph.EP].
- [16] S. Okuzumi, M. Momose, S.-i. Sirono, H. Kobayashi, and H. Tanaka, *Astrophys. J.* **821**, 82 (2016), arXiv:1510.03556 [astro-ph.SR].
- [17] G. Bryden, X. Chen, D. N. C. Lin, R. P. Nelson, and J. C. B. Papaloizou, *Astrophys. J.* **514**, 344 (1999).
- [18] P. Pinilla, M. Benisty, and T. Birnstiel, *Astron. Astrophys.* **545**, A81 (2012), arXiv:1207.6485 [astro-ph.EP].
- [19] Z. Zhu, R. P. Nelson, R. Dong, C. Espaillat, and L. Hartmann, *Astrophys. J.* **755**, 6 (2012), arXiv:1205.5042 [astro-ph.SR].
- [20] S. Jin, S. Li, A. Isella, H. Li, and J. Ji, *Astrophys. J.* **818**, 76 (2016), arXiv:1601.00358 [astro-ph.EP].
- [21] M. E. van den Ancker, P. S. The, H. R. E. Tjin A Djie, C. Catala, D. de Winter, P. F. C. Blondel, and L. B. F. M. Waters, *Astron. Astrophys.* **324**, L33 (1997).
- [22] A. Isella, L. Testi, A. Natta, R. Neri, D. Wilner, and C. Qi, *Astron. Astrophys.* **469**, 213 (2007), arXiv:0704.0616.
- [23] I. de Gregorio-Monsalvo, F. Ménard, W. Dent, C. Pinte, C. López, P. Klaassen, A. Hales, P. Cortés, M. G. Rawlings, K. Tachihara, L. Testi, S. Takahashi, E. Chapillon, G. Mathews, A. Juhasz, E. Akiyama, A. E. Higuchi, M. Saito, L.-Å. Nyman, N. Phillips, J. Rodón, S. Corder, and T. Van Kempen, *Astron. Astrophys.* **557**, A133 (2013), arXiv:1307.1357 [astro-ph.SR].
- [24] G. Guidi, M. Tazzari, L. Testi, I. de Gregorio-Monsalvo, C. J. Chandler, L. Pérez, A. Isella, A. Natta, S. Ortolani, T. Henning, S. Corder, H. Linz, S. Andrews, D. Wilner, L. Ricci,

- J. Carpenter, A. Sargent, L. Mundy, S. Storm, N. Calvet, C. Dullemond, J. Greaves, J. Lazio, A. Deller, and W. Kwon, *Astron. Astrophys.* **588**, A112 (2016), arXiv:1601.07542 [astro-ph.SR].
- [25] C. Qi, K. I. Öberg, S. M. Andrews, D. J. Wilner, E. A. Bergin, A. M. Hughes, M. Hogherheijde, and P. D’Alessio, *Astrophys. J.* **813**, 128 (2015), arXiv:1510.00968 [astro-ph.SR].
- [26] K. A. Rosenfeld, S. M. Andrews, A. M. Hughes, D. J. Wilner, and C. Qi, *Astrophys. J.* **774**, 16 (2013), arXiv:1306.6475 [astro-ph.SR].
- [27] T. L. Wilson, *Reports on Progress in Physics* **62**, 143 (1999).
- [28] J. C. B. Papaloizou, R. P. Nelson, W. Kley, F. S. Masset, and P. Artymowicz, *Protostars and Planets V*, 655 (2007), astro-ph/0603196.
- [29] G. P. Rosotti, A. Juhasz, R. A. Booth, and C. J. Clarke, *Mon. Not. R. Astron. Soc* **459**, 2790 (2016), arXiv:1603.02141 [astro-ph.EP].
- [30] S. Wolf, A. Moro-Martín, and G. D’Angelo, *Planet. Space Sci.* **55**, 569 (2007).
- [31] K. D. Kanagawa, T. Muto, H. Tanaka, T. Tanigawa, T. Takeuchi, T. Tsukagoshi, and M. Momose, *Pub. Astron. Soc. Jap.* **68**, 43 (2016), arXiv:1603.03853 [astro-ph.EP].
- [32] K. D. Kanagawa, T. Muto, H. Tanaka, T. Tanigawa, T. Takeuchi, T. Tsukagoshi, and M. Momose, *Astrophys. J. Lett.* **806**, L15 (2015), arXiv:1505.04482 [astro-ph.EP].
- [33] N. I. Shakura and R. A. Sunyaev, *Astron. Astrophys.* **24**, 337 (1973).
- [34] D. Tamayo, A. H. M. J. Triaud, K. Menou, and H. Rein, *Astrophys. J.* **805**, 100 (2015), arXiv:1502.05099 [astro-ph.EP].
- [35] S. A. Balbus and J. F. Hawley, *Astrophys. J.* **376**, 214 (1991).
- [36] E. Chiang and R. Murray-Clay, *Nature Physics* **3**, 604 (2007), arXiv:0706.1241.
- [37] J. Igea and A. E. Glassgold, *Astrophys. J.* **518**, 848 (1999).
- [38] D. Perez-Becker and E. Chiang, *Astrophys. J.* **735**, 8 (2011), arXiv:1104.2320 [astro-ph.EP].
- [39] T. Umebayashi and T. Nakano, *Pub. Astron. Soc. Jap.* **33**, 617 (1981).
- [40] K. A. Miller and J. M. Stone, *Astrophys. J.* **534**, 398 (2000), astro-ph/9912135.
- [41] J. B. Simon, P. J. Armitage, and K. Beckwith, *Astrophys. J.* **743**, 17 (2011), arXiv:1107.3561 [astro-ph.SR].
- [42] M. Flock, J. P. Ruge, N. Dzyurkevich, T. Henning, H. Klahr, and S. Wolf, *Astron. Astrophys.* **574**, A68 (2015), arXiv:1411.2736 [astro-ph.EP].
- [43] A. Banzatti, P. Pinilla, L. Ricci, K. M. Pontoppidan, T. Birnstiel, and F. Ciesla, *Astrophys.*

- J. Lett. **815**, L15 (2015), arXiv:1511.06762 [astro-ph.EP].
- [44] H.-W. Yen, H. B. Liu, P.-G. Gu, N. Hirano, C.-F. Lee, E. Puspitaningrum, and S. Takakuwa, Astrophys. J. Lett. **820**, L25 (2016), arXiv:1603.01378 [astro-ph.EP].