



This is the accepted manuscript made available via CHORUS. The article has been published as:

## Universal Bell Correlations Do Not Exist

Cole A. Graham and William M. Hoza

Phys. Rev. Lett. **119**, 050402 — Published 1 August 2017

DOI: [10.1103/PhysRevLett.119.050402](https://doi.org/10.1103/PhysRevLett.119.050402)

# Universal Bell correlations do not exist

Cole A. Graham\*

Department of Mathematics, Stanford University, Stanford, CA 94305

William M. Hoza†

Department of Computer Science, University of Texas, Austin, TX 78712

(Dated: June 8, 2017)

We prove that there is no finite-alphabet nonlocal box that generates exactly those correlations that can be generated using a maximally entangled pair of qubits. More generally, we prove that if some finite-alphabet nonlocal box is strong enough to simulate arbitrary local projective measurements of a maximally entangled pair of qubits, then that nonlocal box cannot itself be simulated using *any* finite amount of entanglement. We also give a quantitative version of this theorem for approximate simulations, along with a corresponding positive result.

A classic goal of quantum information research is to understand *the power of quantum nonlocality*: Which nonlocal tasks can be performed using quantum entanglement? For example, quantum entanglement is helpful for playing certain nonlocal games, but it cannot be used to achieve faster-than-light communication. The notion of a *correlation box* serves as a conceptual tool for reasoning about this question. A correlation box is a hypothetical randomized “channel” through which two separated parties, Alice and Bob, can interact. Mathematically, a correlation box is a function  $C : X \times Y \rightarrow \Delta(A \times B)$ , where  $X, Y, A, B$  are countable (finite or countably infinite) alphabets and  $\Delta(A \times B)$  denotes the set of all probability distributions over  $A \times B$ . We imagine that Alice chooses  $x \in X$  and Bob chooses  $y \in Y$ . A sample  $(a, b)$  is drawn from  $C(x, y)$ , and Alice is given  $a$  and Bob is given  $b$ . We will abuse notation and write  $C : X \times Y \rightarrow A \times B$ .

The canonical example [1, 2] is the *Popescu-Rohrlich box*  $C_{\text{PR}} : \{0, 1\} \times \{0, 1\} \rightarrow \{0, 1\} \times \{0, 1\}$ , defined by

$$C_{\text{PR}}(x, y) = \begin{cases} (0, xy) & \text{with probability } 1/2 \\ (1, 1 - xy) & \text{with probability } 1/2. \end{cases} \quad (1)$$

PR boxes cannot be used to communicate, since the marginal distributions of  $a$  and  $b$  are uniform regardless of  $x$  and  $y$ . However,  $C_{\text{PR}}$  is a *nonlocal* box: given access to a PR box, Alice and Bob can perform tasks that would be impossible if they were isolated in a classical world. The standard example is winning the CHSH game [3] with certainty. As usual, we model the classical scenario by assuming that each player’s behavior is a function of his or her private input and a random variable shared between the two players.

For any correlation box  $C$ , rather than analyzing the capabilities of two parties with access to  $C$ , we can instead analyze the *problem of simulating*  $C$ . That is, Alice is given  $x \in X$  and Bob is given  $y \in Y$ . Alice is supposed to output  $a \in A$  and Bob is supposed to output  $b \in B$  such that  $(a, b)$  has distribution  $C(x, y)$ . Let  $\mathbf{Q}$  be the class of all correlation boxes that can be simulated if Alice and Bob have unlimited shared randomness

and an arbitrary but finite amount of entanglement. The question at the beginning of this letter can now be sharpened: Which correlation boxes are in  $\mathbf{Q}$ ? For example, the Tsirelson bound [4] implies that  $C_{\text{PR}} \notin \mathbf{Q}$ .

Let  $\mathbf{B}$  be the class of all correlation boxes that can be simulated if Alice and Bob have unlimited shared randomness, each holds one of a pair of maximally entangled qubits, and they are only allowed to make projective measurements. Clearly,  $\mathbf{B} \subseteq \mathbf{Q}$ ; understanding  $\mathbf{B}$  is a good first step toward understanding  $\mathbf{Q}$ . Bell [5] famously showed that there are correlation boxes in  $\mathbf{B}$  that cannot be simulated using only shared randomness.

A long line of work [6–13] investigated the problem of simulating Bell correlations using *classical communication*, culminating in a protocol by Toner and Bacon [13] for simulating any correlation box in  $\mathbf{B}$  using shared randomness and a single classical bit of one-way communication. Cerf et al. [14] improved on the Toner-Bacon theorem by showing that instead of a bit of communication, it suffices to have a single PR box. (Taking a cue from quantum mechanics, we think of correlation boxes as “single use only”.)

In general, if every correlation box in  $\mathbf{B}$  can be simulated using a particular correlation box  $C$ , then  $C$  can be interpreted as an *upper bound* on the power of  $\mathbf{B}$ . Part of what makes the result by Cerf et al. so appealing is that  $C_{\text{PR}}$  has *finite alphabets*, making it an extremely *explicit* upper bound on the power of  $\mathbf{B}$ : a PR box is a “discrete” device. On the other hand, a “defect” of the result by Cerf et al. is that  $C_{\text{PR}} \notin \mathbf{B}$ , and hence  $C_{\text{PR}}$  is a *loose* upper bound. Local projective measurements of a Bell pair can be simulated using a PR box, but not vice versa.

It is natural, therefore, to hope to construct some finite-alphabet correlation box  $C_*$  such that every correlation box in  $\mathbf{B}$  can be simulated using  $C_*$ , and  $C_*$  is in  $\mathbf{B}$ . Such a correlation box  $C_*$  would *exactly characterize*  $\mathbf{B}$ , greatly clarifying the power of quantum nonlocality. Unfortunately, our main result is that *no such correlation box*  $C_*$  *exists*.

Actually, the situation is even worse in several respects. The simulations we have discussed so far are *one-query*

*reductions.* In general, if  $C_1$  and  $C_2$  are correlation boxes, a  $k$ -query reduction from  $C_1$  to  $C_2$  is a protocol for simulating  $C_1$  in which Alice and Bob have unlimited shared randomness and  $k$  copies of  $C_2$  that they use in a pre-specified order. We say that  $C_1$  *reduces to*  $C_2$  if there is a  $k$ -query reduction from  $C_1$  to  $C_2$  for some finite  $k$ . We prove that  $C_*$  does not exist even if we allow *countably infinite* input alphabets, we allow an *arbitrary* finite amount of entanglement when simulating  $C_*$ , we consider *general* reductions, and we only try to simulate correlation boxes in  $\mathbf{B}$  with *binary alphabets*:

**Theorem 1.** *Suppose  $C_* \in \mathbf{Q}$  has countable input alphabets and finite output alphabets. Then there is some correlation box  $C : \{0, 1\} \times \{0, 1\} \rightarrow \{0, 1\} \times \{0, 1\}$  such that  $C \in \mathbf{B}$ , but  $C$  does not reduce to  $C_*$ .*

Adopting terminology from computational complexity theory, if  $C$  is a correlation box and  $\mathbf{C}$  is a class of correlation boxes, we say that  $C$  is  $\mathbf{C}$ -complete if  $C \in \mathbf{C}$  and every correlation box in  $\mathbf{C}$  reduces to  $C$ .

**Corollary 1.** *There does not exist a finite-alphabet  $\mathbf{B}$ -complete correlation box.*

**Corollary 2.** *There does not exist a finite-alphabet  $\mathbf{Q}$ -complete correlation box.*

Our results can be thought of as “bad news” for the project of understanding the power of quantum nonlocality. In this respect, our results are in the same spirit as previous work showing that certain information-theoretic conditions fail to exactly characterize quantum correlations [15, 16]. Our results should also be considered in the context of the body of research [17–23] investigating the power of correlation boxes in their own right, apart from quantum entanglement. In particular, Barrett and Pironio [17] gave a reduction from any non-signaling correlation box with binary output alphabets to  $C_{\text{PR}}$ . Our result shows that there is no corresponding phenomenon for  $\mathbf{B}$ . Furthermore, Dupuis et al. [19] showed that no finite-alphabet correlation box is complete for the class of non-signaling correlation boxes; our result is analogous.

We now sketch a proof of a weaker version of Theorem 1. We will consider a certain class of nonlocal games, parameterized by a real value  $p$ . Lawson, Linden, and Popescu [24] showed that the optimal entangled success probability for these games depends nonlinearly on  $p$ . On the other hand, if  $C_*$  is a finite-alphabet correlation box, the optimal success probability of a strategy using shared randomness and a single query to  $C_*$  can be shown to be some piecewise-linear function of  $p$ . Hence,  $C_*$  is not  $\mathbf{Q}$ -complete with respect to one-query reductions. A more careful analysis will justify the stronger claim expressed by Theorem 1. We remark that our argument is similar in spirit to Bell’s original proof [5] of his namesake theorem.

We also give a quantitative version of our result for approximate simulations. An  $\varepsilon$ -error reduction is defined like an ordinary reduction except that we allow  $\varepsilon$  total variation error.

**Theorem 2.** *Suppose  $C_* : X \times Y \rightarrow A \times B$  is a finite-alphabet correlation box in  $\mathbf{Q}$ . Then there is some correlation box  $C : \{0, 1\} \times \{0, 1\} \rightarrow \{0, 1\} \times \{0, 1\}$  in  $\mathbf{B}$  such that for every  $k$ , if there is a  $k$ -query  $\varepsilon$ -error reduction from  $C$  to  $C_*$ , then*

$$k^4 \cdot (2|X|)^{4|A|^k} \cdot (2|Y|)^{4|B|^k} \geq \Omega(1/\varepsilon). \quad (2)$$

Conversely, for any  $\varepsilon > 0$ , we give a simple construction of  $C_* : X \times Y \rightarrow \{1, -1\} \times \{1, -1\}$  with  $|X| = |Y| \leq O(1/\varepsilon^2)$  such that  $C_* \in \mathbf{B}$  and every correlation box in  $\mathbf{B}$  reduces to  $C_*$  via a one-query  $\varepsilon$ -error reduction. Note that for  $|A| = |B| = 2, k = 1$ , Theorem 2 implies that  $|X| \cdot |Y|$  must be at least  $1/\varepsilon^{\Omega(1)}$ . On the other hand, when  $|A|, |B|, k$  are large, Theorem 2 might be very far from tight.

## PROOFS OF NEGATIVE RESULTS

For  $0 \leq p, q \leq 1$ , the *biased CHSH game*  $\text{CHSH}[p, q]$  is defined as follows [24]: The referee picks  $x, y \in \{0, 1\}$  independently at random, with  $\Pr[x = 1] = p$ ,  $\Pr[y = 1] = q$ . Alice gets  $x$  and Bob gets  $y$ . Alice outputs  $a \in \{0, 1\}$  and Bob outputs  $b \in \{0, 1\}$ . They win if  $a + b = xy \pmod{2}$ . Alice and Bob know  $p$  and  $q$ . The standard CHSH game [3] is the case  $p = q = \frac{1}{2}$ .

We can think of a correlation box  $C : \{0, 1\} \times \{0, 1\} \rightarrow \{0, 1\} \times \{0, 1\}$  as a *strategy* for the biased CHSH game. The probability that  $C$  wins  $\text{CHSH}[p, q]$  is the probability that  $a + b = xy \pmod{2}$ , where  $(a, b) = C(x, y)$  and the probability is over both the internal randomness of  $C$  and the inputs  $x, y$ . (The inputs  $(x, y)$  are independent of the internal randomness of  $C$ .) Lawson et al. computed the optimal success probability for entangled strategies:

**Lemma 1** ([24]). *If  $\frac{1}{2} \leq q \leq \frac{1}{2p} \leq 1$ , then there exists  $C_{p,q} \in \mathbf{B}$  that wins  $\text{CHSH}[p, q]$  with probability  $\frac{1}{2} + \frac{1}{2}\sqrt{2}\sqrt{q^2 + (1-q)^2}\sqrt{p^2 + (1-p)^2}$ . Furthermore, no correlation box in  $\mathbf{Q}$  achieves higher win probability.*

Throughout the rest of the paper, we define

$$\omega(p) = \frac{1}{2} + \frac{1}{2}\sqrt{p^2 + (1-p)^2}, \quad (3)$$

which is the bound of Lemma 1 for  $q = 1/2$ . We now prove that playing  $\text{CHSH}[p, 1/2]$  near-optimally using some  $C_* \in \mathbf{Q}$  requires approximating  $\omega$  by a small set of *linear* functions (i.e. polynomials in  $p$  of degree one). Roughly, the idea is that there are only so many things one can do with  $C_*$ , and each of them gives rise to a strategy for the biased CHSH game with a success probability that depends linearly on  $p$ .

**Lemma 2.** Suppose  $C_* : X \times Y \rightarrow A \times B$  is in  $\mathbf{Q}$  and  $k \in \mathbb{N}$ . For  $p \in [1/2, 1]$ , let  $C_{p,1/2}$  be the box of Lemma 1. There is a set  $L_{C_*,k}$  of linear functions  $\mathbb{R} \rightarrow \mathbb{R}$  such that:

1. For every  $p \in [1/2, 1]$  and every  $\varepsilon > 0$ , if there exists a  $k$ -query  $\varepsilon$ -error reduction from  $C_{p,1/2}$  to  $C_*$ , then there exists  $\ell \in L_{C_*,k}$  such that  $|\ell(p) - \omega(p)| \leq \varepsilon$ .
2. If  $X, Y, A, B$  are all finite, then  $|L_{C_*,k}|$  is at most  $(2|X|)^{2|A|^k} \cdot (2|Y|)^{2|B|^k}$ . If  $X, Y$  are countable and  $A, B$  are finite, then  $L_{C_*,k}$  is countable.

*Proof.* A deterministic  $k$ -query  $C_*$ -protocol is a protocol that uses  $k$  copies of  $C_*$  in a deterministic way. (The output of such a protocol is random, but only because of the internal randomness of  $C_*$ .) For each such protocol  $\Pi$ , let  $\ell_\Pi(p)$  be the probability that  $\Pi$  wins CHSH $[p, 1/2]$ . Then  $\ell_\Pi$  is a linear function, since

$$\ell_\Pi(p) = \frac{1-p}{2}P_{00} + \frac{1-p}{2}P_{01} + \frac{p}{2}P_{10} + \frac{p}{2}P_{11}, \quad (4)$$

where  $P_{xy}$  is the probability that  $a + b = xy \pmod{2}$  where  $(a, b) = \Pi(x, y)$ . Let  $L_{C_*,k}$  be the set of all  $\ell_\Pi$ .

To prove the first item, let  $\Lambda$  be a  $k$ -query  $\varepsilon$ -error reduction from  $C_{p,1/2}$  to  $C_*$ . We can think of  $\Lambda$  as a *distribution over deterministic  $C_*$ -protocols*  $\Pi$ . By the correctness of the reduction,  $|\mathbb{E}_{\Pi \sim \Lambda}[\ell_\Pi(p)] - \omega(p)| \leq \varepsilon$ . The best case is at least as good as the average case, so there is a deterministic protocol  $\Pi_*$  such that  $\ell_{\Pi_*}(p) \geq \omega(p) - \varepsilon$ . Since  $C_* \in \mathbf{Q}$  and  $\mathbf{Q}$  is closed under reductions,  $\Pi_*$  implements a correlation box in  $\mathbf{Q}$ . Therefore, by the optimality clause of Lemma 1,  $\ell_{\Pi_*}(p) \leq \omega(p)$ , and hence  $|\ell_{\Pi_*}(p) - \omega(p)| \leq \varepsilon$ .

To prove the second item, we bound the number of deterministic  $k$ -query  $C_*$ -protocols. Such a protocol can be specified by:

- Functions  $q_i : \{0, 1\} \times A^{i-1} \rightarrow X$  for each  $1 \leq i \leq k$ , telling the  $i$ th query that Alice makes as a function of her input and the query responses she has seen so far, and corresponding functions  $r_i : \{0, 1\} \times B^{i-1} \rightarrow Y$  for Bob.
- A function  $s : \{0, 1\} \times A^k \rightarrow \{0, 1\}$ , telling the output Alice gives as a function of her input and all query responses, and a corresponding function  $t : \{0, 1\} \times B^k \rightarrow \{0, 1\}$  for Bob.

If  $X, Y$  are countable and  $A, B$  are finite, then there are only countably many possibilities for each of these functions, so there are countably many such protocols. Suppose now that  $X, Y, A, B$  are all finite and  $|A|, |B| \geq 2$ . The number of possibilities for  $q_i$  is  $|X|^{2|A|^{i-1}}$ , and similarly for  $r_i$ . The number of possibilities for  $s$  is  $2^{|A|^k}$ ,

and similarly for  $t$ . Therefore,  $|L_{C_*,k}|$  is bounded by

$$\left( \prod_{i=1}^k |X|^{2|A|^{i-1}} \right) \left( \prod_{i=1}^k |Y|^{2|B|^{i-1}} \right) \cdot 2^{2|A|^k} \cdot 2^{2|B|^k} \quad (5)$$

$$= |X|^{2 \sum_i |A|^{i-1}} \cdot |Y|^{2 \sum_i |B|^{i-1}} \cdot 2^{2|A|^k} \cdot 2^{2|B|^k} \quad (6)$$

$$\leq |X|^{2|A|^k} \cdot |Y|^{2|B|^k} \cdot 2^{2|A|^k} \cdot 2^{2|B|^k} \quad (7)$$

$$= (2|X|)^{2|A|^k} \cdot (2|Y|)^{2|B|^k}. \quad (8)$$

Finally, if  $A$  is a singleton set, the step above where we bounded  $\sum_i |A|^{i-1}$  by  $|A|^k$  was not valid, but in this case  $q_1, \dots, q_k$  do not need to be specified anyway, so the final bound still holds. Similarly if  $B$  is a singleton set.  $\square$

Next, we show that  $\omega$  *cannot* be well-approximated by a small set of linear functions. To prove Theorem 1, the following trivial fact suffices.

**Lemma 3.** Suppose  $L$  is a countable set of linear functions  $\mathbb{R} \rightarrow \mathbb{R}$ . Then there is some  $p \in [1/2, 1]$  such that for every  $\ell \in L$ ,  $\ell(p) \neq \omega(p)$ .

*Proof.* Suppose  $\ell(p) = \omega(p)$ , where  $\ell \in L$ . Rearranging,

$$p^2 + (1-p)^2 = r(p)^2, \quad (9)$$

where  $r(p)$  is another linear function. The quadratic expression on the left hand side of (9) is not a square (e.g. the discriminant is  $-4 \neq 0$ ). Therefore, (9) is a non-degenerate quadratic equation, so it has at most two solutions  $p$ . So  $\ell$  intersects  $\omega$  at most twice, and hence  $L$  intersects  $\omega$  in countably many places.  $\square$

*Proof of Theorem 1.* Fix  $C_* : X \times Y \rightarrow A \times B$ , where  $X, Y$  are countable,  $A, B$  are finite, and  $C_* \in \mathbf{Q}$ . We will show that there is some choice of  $p$  so that there is no reduction from  $C_{p,1/2}$  to  $C_*$ ; since  $C_{p,1/2} \in \mathbf{B}$  and  $C_{p,1/2}$  has binary alphabets, this will complete the proof.

For each  $k \in \mathbb{N}$ , let  $L_{C_*,k}$  be the set of linear functions given by Lemma 2. The alphabet bounds for  $C_*$  imply that  $L_{C_*,k}$  is countable. Let  $L = \bigcup_{k \in \mathbb{N}} L_{C_*,k}$ , so that  $L$  is still countable. By Lemma 3, choose  $p \in [1/2, 1]$  so that for every  $\ell \in L$ ,  $\ell(p) \neq \omega(p)$ . Then  $C_{p,1/2}$  does not reduce to  $C_*$ , because if there were a  $k$ -query (0-error) reduction for some  $k$ , Lemma 2 would imply that there was some  $\ell \in L$  with  $\ell(p) = \omega(p)$ .  $\square$

To prove Theorem 2, we need a *quantitative* lower bound on the error of any approximation of  $\omega$  by linear functions:

**Lemma 4.** Pick  $p \in [1/2, 1]$  uniformly at random. Then for any linear function  $\ell : \mathbb{R} \rightarrow \mathbb{R}$  and any  $\varepsilon > 0$ ,

$$\Pr[|\ell(p) - \omega(p)| \leq \varepsilon] \leq O(\sqrt{\varepsilon}). \quad (10)$$

*Proof.* Let  $I = [1/2, 1]$ . We first compute

$$\omega''(x) = \frac{1}{2} \left[ 2 \left( x - \frac{1}{2} \right)^2 + \frac{1}{2} \right]^{-3/2} \geq \frac{1}{2} \quad \text{on } I. \quad (11)$$

Hence  $\omega$  is uniformly convex on  $I$ .

Without loss of generality we can assume that the graph of  $\ell$  intersects the graph of  $\omega$  twice (with a point of tangency counted as a double intersection). After all, if  $\ell < \omega$  on  $I$ , translate  $\ell$  up until the first moment of equality with  $\omega$ , thus decreasing the pointwise error between  $\ell$  and  $\omega$  at every  $x \in I$ . If  $\ell$  is then tangent to  $\omega$ , we are done. Otherwise,  $\ell$  intersects  $\omega$  at an endpoint, so rotate  $\ell$  up about this point until it is tangent to  $\omega$  (no other intersections occur because  $\omega$  is uniformly convex). Again, pointwise errors do not increase under this rotation. Similar considerations hold if initially  $\ell > \omega$  or  $\ell$  intersects  $\omega$  at one point.

Therefore, suppose  $\ell$  linearly interpolates  $\omega$  at the (potentially coincident) points  $x_1, x_2 \in I$ . By a standard argument in interpolation theory, for all  $x \in I$ , there exists  $\xi_x \in I$  such that

$$\omega(x) - \ell(x) = \frac{\omega''(\xi_x)}{2} (x - x_1)(x - x_2). \quad (12)$$

By (11),  $|\omega(x) - \ell(x)| \geq \frac{1}{4}|x - x_1||x - x_2|$ . In particular, when  $\min\{|x - x_1|, |x - x_2|\} > 2\sqrt{\varepsilon}$ ,  $|\omega(x) - \ell(x)| > \varepsilon$ . The probability that  $p$  is within  $2\sqrt{\varepsilon}$  of either  $x_1$  or  $x_2$  is  $O(\sqrt{\varepsilon})$ .  $\square$

*Proof of Theorem 2.* Fix  $C_* : X \times Y \rightarrow A \times B$ , where  $X, Y, A, B$  are finite and  $C_* \in \mathbf{Q}$ . Let  $L_{C_*,k}$  be the set of linear functions given by Lemma 2. Pick  $p \in [1/2, 1]$  uniformly at random. By Lemma 4 and the union bound, for any  $\varepsilon_k > 0$ , the probability that some  $\ell \in L_{C_*,k}$  satisfies  $|\ell(p) - \omega(p)| \leq \varepsilon_k$  is at most  $O(\sqrt{\varepsilon_k} \cdot |L_{C_*,k}|)$ . Therefore, by the union bound over  $k$ , the probability that there is such an  $\ell$  for *any*  $k$  is

$$O\left(\sum_{k=1}^{\infty} \sqrt{\varepsilon_k} \cdot |L_{C_*,k}|\right). \quad (13)$$

Choose  $\varepsilon_k$  so that  $\sqrt{\varepsilon_k} \cdot |L_{C_*,k}| = c/k^2$ , where  $c$  is a sufficiently small constant so that the bound in (13) is strictly less than 1. (Such a  $c$  exists because  $\sum_k 1/k^2$  converges.) This implies  $\varepsilon_k \geq \Omega(k^{-4}|L_{C_*,k}|^{-2})$ . Hence, by Lemma 2,

$$k^4 \cdot (2|X|)^{4|A|^k} \cdot (2|Y|)^{4|B|^k} \geq \Omega(1/\varepsilon_k). \quad (14)$$

By our choice of  $\varepsilon_k$ , there exists some  $p$  so that for every  $k$ , for every  $\ell \in L_{C_*,k}$ ,  $|\ell(p) - \omega(p)| > \varepsilon_k$ . Choose  $C = C_{p,1/2}$ . By Lemma 2, if there is a  $k$ -query  $\varepsilon$ -error reduction from  $C$  to  $C_*$ , then  $\varepsilon > \varepsilon_k$ .  $\square$

We end this section with two simple consequences of our main results. We say that  $C_1 \leq C_2$  if there is a reduction from  $C_1$  to  $C_2$ . We say that  $C_1 < C_2$  (“simulating  $C_1$  is strictly easier than simulating  $C_2$ ”) if  $C_1 \leq C_2$  and  $C_2 \not\leq C_1$ .

**Theorem 3.** *For any finite-alphabet correlation box  $C_1 \in \mathbf{B}$ , there is another finite-alphabet correlation box  $C_2 \in \mathbf{B}$  such that  $C_1 < C_2$ .*

*Proof.* By Theorem 1, there is a correlation box  $C \in \mathbf{B}$  with binary alphabets such that  $C \not\leq C_1$ . Write  $C_1 : X_1 \times Y_1 \rightarrow A_1 \times B_1$ . By relabeling if necessary, we can assume that  $0, 1 \notin X_1, Y_1$ . Define  $X_2 = X_1 \cup \{0, 1\}$ ,  $Y_2 = Y_1 \cup \{0, 1\}$ ,  $A_2 = A_1 \cup \{0, 1\}$ ,  $B_2 = B_1 \cup \{0, 1\}$ . Define  $C_2 : X_2 \times Y_2 \rightarrow A_2 \times B_2$  by the following  $\mathbf{B}$  protocol: If  $x \in X_1$ , then Alice does what she would have done in the protocol witnessing  $C_1 \in \mathbf{B}$ . Otherwise, if  $x \in \{0, 1\}$ , she does what she would have done in the protocol witnessing  $C \in \mathbf{B}$ . Bob acts similarly. By construction,  $C_1 \leq C_2$  and  $C \leq C_2$ , so by transitivity,  $C_1 < C_2$ .  $\square$

The same technique used to prove Theorem 3 can also be used to generalize Theorem 1 as follows. Suppose we have a finite set of correlation boxes  $\{C_*^{(1)}, C_*^{(2)}, \dots, C_*^{(t)}\} \subseteq \mathbf{Q}$ , each with countable input alphabets and finite output alphabets. As in the proof of Theorem 3, we can construct a single correlation box  $C_* \in \mathbf{Q}$  with countable input alphabets and finite output alphabets such that every  $C_*^{(i)}$  reduces to  $C_*$ . Hence, by Theorem 1, there is a correlation box  $C : \{0, 1\} \times \{0, 1\} \rightarrow \{0, 1\} \times \{0, 1\}$  such that  $C \in \mathbf{B}$ , but it is impossible to simulate  $C$  using shared randomness and any finite number of copies of each  $C_*^{(i)}$ .

## POSITIVE RESULTS

We now show how to construct a finite-alphabet correlation box that is approximately complete for  $\mathbf{B}$ . The construction is simple, and essentially consists of a discretization of the Bloch sphere [25].

**Theorem 4.** *For every  $\varepsilon > 0$ , there exists  $C_* : \{1, \dots, T\} \times \{1, \dots, T\} \rightarrow \{1, -1\} \times \{1, -1\}$  with  $T \leq O(1/\varepsilon^2)$  such that  $C_* \in \mathbf{B}$ , and for every  $C \in \mathbf{B}$ , there is a one-query  $\varepsilon$ -error reduction from  $C$  to  $C_*$ .*

*Proof.* Let  $S^2$  denote the unit sphere in  $\mathbb{R}^3$ . A pair of local projective measurements as in the definition of  $\mathbf{B}$  can be described by  $x \in S^2$  chosen by Alice and  $y \in S^2$  chosen by Bob; the  $(\pm 1)$ -valued outcomes  $a, b$  satisfy  $E[a] = E[b] = 0$  and  $E[ab] = x \cdot y$ . (For example,  $x$  and  $y$  might specify spin axes along which Alice and Bob are measuring.)

Let  $c_1, c_2, \dots, c_T \in S^2$  be points such that every point in  $S^2$  is within  $\varepsilon$  of some  $c_i$  in  $\ell_2$  distance. Such a collection of points exists with  $T \leq O(1/\varepsilon^2)$ . The correlation



box  $C_*$  simply makes the measurements described by  $c_i$  and  $c_j$ , where  $i$  is the input to Alice and  $j$  is the input to Bob. By construction,  $C_* \in \mathbf{B}$ .

For the reduction, when Alice and Bob need to measure according to  $x, y \in S^2$ , they simulate the measurement by inputting  $i, j$  to  $C_*$ , where  $i$  and  $j$  minimize  $\|c_i - x\|_2$  and  $\|c_j - y\|_2$ . Correctness follows from the fact that

$$|x \cdot y - c_i \cdot c_j| \leq |x \cdot c_j - c_i \cdot c_j| + |c_i \cdot y - c_i \cdot c_j| \quad (15)$$

$$\leq \|x - c_i\|_2 + \|y - c_j\|_2. \quad (16)$$

□

**Proposition 1.** *There exists  $C_* : \mathbb{N} \times \mathbb{N} \rightarrow \{0, 1\} \times \{0, 1\}$  such that  $C_* \in \mathbf{B}$ , and for every  $C \in \mathbf{B}, \varepsilon > 0$ , there is a one-query  $\varepsilon$ -error reduction from  $C$  to  $C_*$ .*

*Proof sketch.* Use a countable dense subset of  $S^2$ . □

## CONCLUSION

To better understand quantum entanglement, it is desirable to characterize what can and cannot be achieved using quantum nonlocality. In this letter, we have *ruled out* a natural type of characterization, even if attention is restricted to projective measurements of a single Bell pair. We hope that this letter inspires future researchers to *circumvent* our results by formulating a different type of characterization of the power of quantum nonlocality.

We thank Scott Aaronson and Ronald de Wolf for helpful comments and encouragement. We thank referees for numerous suggestions. This material is based upon work supported by the National Science Foundation Graduate Research Fellowship under Grant No. DGE-1610403. Cole Graham gratefully acknowledges the support of the Fannie and John Hertz Foundation.

---

\* grahamca@stanford.edu

† whoza@utexas.edu

- [1] L. A. Khalfin and B. S. Tsirelson, in *Symposium on the foundations of modern physics*, Vol. 85 (Singapore: World Scientific, 1985) p. 441.
- [2] S. Popescu and D. Rohrlich, *Foundations of Physics* **24**, 379 (1994).
- [3] J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, *Physical Review Letters* **23**, 880 (1969).
- [4] B. S. Cirel'son, *Letters in Mathematical Physics* **4**, 93 (1980).
- [5] J. Bell, *Physics* **1**, 195 (1964).
- [6] T. Maudlin, in *PSA: Proceedings of the Biennial Meeting of the Philosophy of Science Association* (JSTOR, 1992) pp. 404–417.
- [7] G. Brassard, R. Cleve, and A. Tapp, *Physical Review Letters* **83**, 1874 (1999).
- [8] M. Steiner, *Physics Letters A* **270**, 239 (2000).
- [9] N. J. Cerf, N. Gisin, and S. Massar, *Physical Review Letters* **84**, 2521 (2000).
- [10] J. A. Csirik, *Physical Review A* **66**, 014302 (2002).
- [11] A. Coates, arXiv preprint quant-ph/0203112 (2002).
- [12] D. Bacon and B. F. Toner, *Physical Review Letters* **90**, 157904 (2003).
- [13] B. F. Toner and D. Bacon, *Physical Review Letters* **91**, 187904 (2003).
- [14] N. J. Cerf, N. Gisin, S. Massar, and S. Popescu, *Physical Review Letters* **94**, 220403 (2005).
- [15] R. Gallego, L. E. Würflinger, A. Acín, and M. Navascués, *Physical Review Letters* **107**, 210403 (2011).
- [16] M. Navascués, Y. Guryanova, M. J. Hoban, and A. Acín, *Nature communications* **6** (2015).
- [17] J. Barrett and S. Pironio, *Physical Review Letters* **95**, 140401 (2005).
- [18] J. Barrett, N. Linden, S. Massar, S. Pironio, S. Popescu, and D. Roberts, *Physical Review A* **71**, 022101 (2005).
- [19] F. Dupuis, N. Gisin, A. Hasidim, A. A. Méthot, and H. Pilpel, *Journal of mathematical physics* **48**, 082107 (2007), note that the article claims to prove this theorem more generally for correlation boxes with finite *output* alphabets and arbitrary input alphabets, but the authors agree that the proof is flawed (F. Dupuis, personal communication).
- [20] G. Brassard, H. Buhrman, N. Linden, A. A. Méthot, A. Tapp, and F. Unger, *Physical Review Letters* **96**, 250401 (2006).
- [21] M. Forster, S. Winkler, and S. Wolf, *Physical Review Letters* **102**, 120401 (2009).
- [22] N. Brunner and P. Skrzypczyk, *Physical Review Letters* **102**, 160403 (2009).
- [23] W. van Dam, *Natural Computing* **12**, 9 (2013).
- [24] T. Lawson, N. Linden, and S. Popescu, arXiv preprint arXiv:1011.6245 (2010).
- [25] F. Bloch, *Physical review* **70**, 460 (1946).