



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Primordial Black Holes and r-Process Nucleosynthesis

George M. Fuller, Alexander Kusenko, and Volodymyr Takhistov

Phys. Rev. Lett. **119**, 061101 — Published 7 August 2017

DOI: [10.1103/PhysRevLett.119.061101](https://doi.org/10.1103/PhysRevLett.119.061101)

Primordial Black Holes and r -Process Nucleosynthesis

George M. Fuller,^{1,*} Alexander Kusenko,^{2,3,†} and Volodymyr Takhistov^{2,‡}

¹*Department of Physics, University of California, San Diego
La Jolla, California 92093-0354, USA*

²*Department of Physics and Astronomy, University of California, Los Angeles
Los Angeles, CA 90095-1547, USA*

³*Kavli Institute for the Physics and Mathematics of the Universe (WPI), UTIAS
The University of Tokyo, Kashiwa, Chiba 277-8583, Japan*

(Dated: June 21, 2017)

We show that some or all of the inventory of r -process nucleosynthesis can be produced in interactions of primordial black holes (PBHs) with neutron stars (NSs) if PBHs with masses $10^{-14} M_{\odot} < M_{\text{PBH}} < 10^{-8} M_{\odot}$ make up a few percent or more of the dark matter. A PBH captured by a neutron star (NS) sinks to the center of the NS and consumes it from the inside. When this occurs in a rotating millisecond-period NS, the resulting spin-up ejects $\sim 0.1 - 0.5 M_{\odot}$ of relatively cold neutron-rich material. This ejection process and the accompanying decompression and decay of nuclear matter can produce electromagnetic transients, such as a kilonova-type afterglow and fast radio bursts. These transients are not accompanied by significant gravitational radiation or neutrinos, allowing such events to be differentiated from compact object mergers occurring within the distance sensitivity limits of gravitational wave observatories. The PBH-NS destruction scenario is consistent with pulsar and NS statistics, the dark matter content and spatial distributions in the Galaxy and Ultra Faint Dwarfs (UFD), as well as with the r -process content and evolution histories in these sites. Ejected matter is heated by beta decay, which leads to emission of positrons in an amount consistent with the observed 511-keV line from the Galactic Center.

PACS numbers: 95.35.d, 26.30.k, 97.60.Gb, 97.60.Lf

Primordial black holes (PBHs) can account for all or part of the dark matter (DM) [1–13]. If a PBH is captured by a neutron star (NS), it settles into the center and grows until the supply of nuclear matter is exhausted by accretion and ejection.

In this *Letter* we show that NS disruptions by PBHs in DM-rich environments, such as the Galactic center (GC) and dwarf spheroidal galaxies, provide a viable site for r -process nucleosynthesis, thus offering a solution to a long-standing puzzle [14–18]. The transients accompanying NS disruption events and the positrons produced in these events are consistent with present observations, and they offer a way of testing the NS–PBH scenario in the future.

We will demonstrate that, when a PBH accretes matter inside a rapidly rotating millisecond pulsar (MSP), the resulting pulsar spin-up causes $\sim 0.1 - 0.5 M_{\odot}$ of neutron-rich material to be ejected without significant heating and only modest neutrino emission. This provides a favorable setting for r -process nucleosynthesis, occurring on the Galactic time-scales, which can evade several problems that have challenged the leading proposed r -process production sites, such as neutrino-heated winds from core collapse supernovae or binary compact object mergers (COM) [19, 20]. The unusual distribution of r -process abundances within the ultra-faint dwarf spheroidal galaxies [21, 22] (UFDs) is naturally explained by the rates of PBH capture in these systems. The rates are also consistent with the paucity of pulsars in the GC [23]. A similar distribution of r -process material in UFDs can be expected from NS disruptions due to black holes produced

in the NS interiors by accretion of particle dark matter onto the NS [24], although the rates and the implications for dark matter properties are, of course, different. The probability of PBH capture depends on both the PBH and the NS densities. DM-rich environments, such as the GC and dwarf spheroidal galaxies, are not known to host NSs. An exception is the young magnetar [25, 26], whose age is small compared to the time scales of PBH capture. NSs are found in the disk and the halo, as well as in the globular clusters, where the dark matter density [27, 28] is too low to cause a substantial decrease in the pulsar population. Positrons emitted from the heated neutron-rich ejecta can account for the observed 511-keV line from the GC [29, 30]. The final stages of neutron star demise can be the origin [31, 32] of some of the recently observed [33] fast radio bursts (FRBs), as well as X-ray and γ -ray transients. A kilonova-type [34–41] afterglow can accompany the decompressing nuclear matter ejecta, but unlike COM, these events are not associated with a significant release of neutrinos or gravitational radiation. Therefore, future observations of gravitational waves and kilonovae will be able to distinguish between r -process scenarios.

Millisecond pulsars are responsible for the predominant contribution to the nucleosynthesis initiated by PBH-induced centrifugal ejection of neutron-rich material, since MSPs have the highest angular velocities at the time of PBH capture. The most prominent sites of r -process production must have a high density of MSPs as well as PBHs. The latter trace the DM spatial distri-

bution. The DM density is high in the GC and in the UFDs. On the other hand, the MSP density is high in molecular clouds, including the Central Molecular Zone (CMZ), and the globular clusters. While the CMZ is located within the GC with an extremely high DM density, observations imply that the DM content of globular clusters is fairly low [27, 28]. The product of the DM density and the MSP density is still sufficient to allow for some r -process nucleosynthesis in both the UFDs and the globular clusters, but we estimate that CMZ accounts for 10% to 50% of the total Galactic production. We include contributions from the GC (CMZ) and the rest of the halo (which may be comparable, within uncertainties) [42, 43].

The CMZ has an approximate size of ~ 200 pc and is located near the GC, where supernova rates are the highest. Since the DM density peaks at the GC, the CMZ is a site of frequent PBH-MSP interactions. The pulsar formation rate [44] in the CMZ is $1.5 \times 10^{-3} \text{yr}^{-1}$ (7% of the Galactic formation rate), consistent with the GeV γ -ray flux observed [45] from the GC by the *Fermi* Large Area Telescope. Hence, we expect $N_p^{\text{GC}} \approx 1.5 \times 10^7$ NSs to be produced at the GC during the lifetime of the Galaxy, $t_G \sim 10^{10}$ yr. Roughly, 30%-50% of these NSs become MSPs [46], and the number of MSPs with a particular rotation period can then be estimated from a population model [46–48].

Simulations and observations of UFDs imply [21] that ~ 2000 core-collapse supernovae have occurred in 10 UFDs during $t_{\text{UFD}} \sim 5 \times 10^8$ yr. Hence, we expect that $N_p^{\text{UFD}} \sim 10^2$ pulsars have been produced in each of these systems, on average, and we estimate the fraction of fast-rotating MSPs using a population model [47].

The *black hole capture rate* can be calculated using the initial PBH mass (m_{PBH}), DM density and velocity dispersion (assuming a Maxwellian distribution). With the base MW and UFD capture rates denoted as F_0^{MW} and F_0^{UFD} , one obtains [49] the full PBH-NS capture rates $F = (\Omega_{\text{PBH}}/\Omega_{\text{DM}})F_0^{\text{MW}}$ and $F = (\Omega_{\text{PBH}}/\Omega_{\text{DM}})F_0^{\text{UFD}}$ in the MW and UFD, respectively. The number of PBHs captured by a NS is $F \cdot t$, where the time t is either t_G or t_{UFD} for the MW and the UFD, respectively. For our analysis we consider a typical NS with mass $M_{\text{NS}} = 1.5M_\odot$ and radius $R_{\text{NS}} = 10$ km. The base NS-PBH capture rate F_0 is given by [49]

$$F_0 = \sqrt{6\pi} \frac{\rho_{\text{DM}}}{m_{\text{PBH}}} \left[\frac{R_{\text{NS}} R_s}{\bar{v}(1 - R_s/R_{\text{NS}})} \right] \left(1 - e^{-3E_{\text{loss}}/(m_{\text{PBH}}\bar{v}^2)} \right), \quad (1)$$

where \bar{v} is the DM velocity dispersion, $R_S = 2GM_{\text{NS}}$ and R_{NS} are the BH Schwarzschild radius and the NS radius, respectively. E_{loss} is the energy loss associated with PBH-NS interaction. BH capture can occur when $E_{\text{loss}} > m_{\text{PBH}}v_0^2/2$, with v_0 being the asymptotic velocity of the PBH. Taking a uniform flux of PBHs across the star, the average energy loss for a typical NS is found to be $E_{\text{loss}} \simeq 58.8G^2m_{\text{PBH}}^2M_{\text{NS}}/R_{\text{NS}}^2$. Since MSPs origi-

nate from binaries, a higher binary gravitational potential causes an increase in the capture rate. We, therefore, assume that the capture rate for MSPs is a factor 2 higher than for isolated NSs. For example, for typical values of parameters and $m_{\text{PBH}} = 10^{19}$ g, one obtains

$$\begin{cases} F_0^{\text{MW}} = 1.5 \times 10^{-11}/\text{yr} \\ F_0^{\text{UFD}} = 6.0 \times 10^{-10}/\text{yr} \end{cases} \quad (2)$$

For the MW we have used a velocity dispersion of 48 km/s and 105 km/s for NS and DM, respectively, as well as DM density $8.8 \times 10^2 \text{GeV}/\text{cm}^3$. The pulsar and DM velocity dispersions are simultaneously taken into account for the MW as described below. For UFD we have used DM velocity dispersion 2.5 km/s and DM density $10 \text{GeV}/\text{cm}^3$.

A PBH could also be captured by a NS progenitor prior to supernova core collapse [50], but this does not increase the capture rates significantly.

Natal pulsar kicks can enable pulsars to escape from the region of interest. We include this effect in our calculations (see Supplemental Material [51]).

Pulsar lifetimes in the presence of PBHs with a given number density can be estimated as $\langle t_{\text{NS}} \rangle = 1/F + t_{\text{loss}} + t_{\text{con}}$, where the first term describes the mean BH capture time, t_{loss} is the time for the PBH to be brought within the NS once it is gravitationally captured, and t_{con} is the time for the black hole to consume the NS. For a typical NS one finds [49] that $t_{\text{loss}} \simeq 4.1 \times 10^4 (m_{\text{PBH}}/10^{22} \text{g})^{-3/2} \text{yr}$. The spherical accretion rate of NS matter onto the PBH is described by the Bondi equation $dm_{\text{BH}}/dt = 4\pi\lambda_s G^2 m_{\text{BH}}^2 \rho_c / v_s^3 = C_0 m_{\text{BH}}^2$, where $m_{\text{BH}}(t)$ is the time-dependent mass of the central black hole, v_s is the sound speed, ρ_c is the central density and λ_s is a density profile parameter. For typical NS values of [52] $v_s = 0.17$, $\rho_c = 10^{15} \text{g}/\text{cm}^3$ and $\lambda_s = 0.707$ (for a star described by an $n = 3$ polytrope) we obtain that $t_{\text{con}} = 10 (10^{19} \text{g}/m_{\text{PBH}}) \text{yr}$. If PBHs make up all of the DM, we calculate that $\langle t_{\text{NS}} \rangle < 10^{12} \text{yr}$ for $10^{17} \text{g} < m_{\text{PBH}} < 10^{25} \text{g}$, implying that a $\mathcal{O}(1-10)\%$ fraction of pulsars should have been consumed in the age of the Galaxy. This is consistent with observations [23] suggesting an under abundance of MSPs near the central Galactic black hole, Sgr A*. A recently discovered young, 4×10^4 yr old, magnetar J1745-2900 located just 0.1 pc from the GC [25, 26] is also consistent with our results, since this magnetar's age is shorter than $\langle t_{\text{NS}} \rangle$. The unusual surface temperature [53] and X-ray luminosity of J1745-2900 warrants scrutiny, as this activity might be consistent with PBH destruction in progress.

Angular momentum transfer determines the dynamics of a NS spin-up. As the captured PBH starts to grow and consume the spinning pulsar from the inside, the radius of the neutron star decreases and angular momentum conservation forces a spin-up. As the star contracts, the fractional change in radius is greater for accreted matter in the inner regions than it is for material further out.

This could lead to differential rotation. However, if angular momentum can be efficiently transferred outward, the star can maintain rigid-body rotation. Viscosity [54, 55] and magnetic stresses [55] can prevent differential rotation from developing. It can be shown (see Supplemental Material [51]) that angular momentum is transferred efficiently on the relevant time scales and that Bondi accretion proceeds nearly uninterrupted throughout the BH evolution.

Ejected mass originates from the star spin-up when matter at the equator exceeds the escape velocity. Using polytropic NS density runs with different indices [52], we have calculated analytically (see end of “Ejected Mass” section in Supplemental Material [51]) the amount of ejected material. The results are shown in Figure 1 for NS victims with a range of initial rotation periods. Based on our estimates discussed in Supplemental Materials, NS with periods of a few ms can eject more than $10^{-1}M_{\odot}$ of material. A detailed calculation taking into account general relativistic effects [56, 57] is needed to improve understanding of the ejected mass.

The number of MSPs in the disk with periods greater than P is described by a power-law distributed population model [47]. We assume that the distribution in the CMZ is the same, and we normalize the total to the number of neutron stars produced in supernova explosions. According to the population model [47], $N_{\text{MSP}} \simeq 1.6 \times 10^4 (1.56 \text{ ms}/P)$. Using the differential distribution $d(N_{\text{MSP}})/dP$, we obtain the population-averaged ejected mass:

$$\langle M_{\text{ej}} \rangle = \frac{\int_{P_{\text{min}}}^{\infty} \left(\frac{dN_{\text{MSP}}}{dP} \right) M_{\text{ej}}(P) dP}{\int_{P_{\text{min}}}^{\infty} \left(\frac{dN_{\text{MSP}}}{dP} \right) dP}, \quad (3)$$

where P_{min} is the minimal MSP period in the population, and $M_{\text{ej}}(P)$ is the ejected mass function interpolated from the distribution shown in Figure 1. We find that the population-averaged ejected mass is $\langle M_{\text{ej}} \rangle = 0.18M_{\odot}$ and $0.1M_{\odot}$ if we take the shortest period to be $P_{\text{min}} = 0.7$ ms (theoretically predicted) and $P_{\text{min}} = 1.56$ ms (observed), respectively. Since realistic nuclear matter equations of state suggest flatter NS density profiles than our polytropic approximations, our estimate is conservative and $\langle M_{\text{ej}} \rangle$ can be up to a factor of few larger. Alternative population models [46], such as those based on [48], do not significantly alter the results.

Nucleosynthesis takes place in the ejecta. Heating accompanying the growth of a BH inside a NS results in a temperature increase near the event horizon that is only a factor of few higher than the NS surface temperature [54]: $T_{\text{h}}/T_{\text{surf}} \sim 3$. Consequently, neutrino emission is negligible and ejected material does not suffer significant heating or exposure to neutrinos.

Decompression of the centrifugally-ejected, relatively




FIG. 1: Total ejected mass (M_{ej}) from a MSP with initial rotation period P disrupted by a PBH. $n = 3$ polytrope (red) and $n = 1.5$ polytrope (orange) NS density profiles are shown. The black line indicates ejection of $0.1 M_{\odot}$. The MSP period–population distribution [47] is displayed with a dashed blue line.

low entropy and very low electron fraction nuclear matter in this scenario could be expected to result in a significant mass fraction of this material participating in r -process nucleosynthesis [58–64]. The large neutron excess in this scenario, relatively unmolested by neutrino charged current capture-induced reprocessing of the neutron-to-proton ratio, could lead to fission cycling [64, 65], thereby tying together the nuclear mass number $A = 130$ and $A = 195$ r -process abundance peaks. Unlike COM r -process ejecta, which will have a wide range of neutrino exposures, entropy, and electron fraction and thereby can reproduce the solar system r -process abundance pattern [66], the PBH scenario may be challenged in producing the low mass, $A < 100$, r -process material.

The material ejected in the PBH-NS destruction process is heated by beta decay and fission, resulting in thermodynamic conditions and abundances closely akin to those in the COM-induced “tidal tail” nuclear matter decompression that gives rise to kilonova-like electromagnetic signatures [34–41]. This could be a more luminous and longer duration transient compared to the classic COM-generated kilonovae, as the ejecta in the PBH scenario can have more mass than the tidal tails of COM.

The total amount of ejected r -process material in the PBH-NS destruction process can be estimated via $M_{\text{tot}}^r = FtN_{\text{MSP}}\langle M_{\text{ej}} \rangle$, assuming that the bulk of the ejecta undergoes r -process nucleosynthesis. The overall mass of r -process material in the Galaxy is $M_{\text{tot}}^{r,\text{MW}} \sim 10^4 M_{\odot}$. The required fraction of dark matter in the form of PBHs is $(\Omega_{\text{PBH}}/\Omega_{\text{DM}}) = M_{\text{tot}}^{r,\text{MW}} / (F_0^{\text{MW}} t_G N_{\text{MSP}}^{\text{GC}} \langle M_{\text{ej}} \rangle)$. If the mass of ejected r -process material in a single event is $0.1 - 0.5 M_{\odot}$, the PBH capture rate $10^{-5} - 10^{-6} \text{ Mpc}^{-3} \text{ yr}^{-1}$ can account for *all* of the r -process in the Galaxy. At

this rate, 10^5 NS disruption events have occurred in the lifetime of the Galaxy.

This rate of NS disruptions in UFDs is also consistent with the observationally inferred UFD r -process content and with the uneven distribution of this material among the observed UFD. Observations imply that one in ten of UFDs have been a host to r -process nucleosynthesis events, which must, therefore, be rare [21, 22, 67]. The rate F_0^{UFD} implies that the probability of a NS disruption in a single UFD is about 0.1, which explains the uneven distribution. The amount of r -process material supplied by a single event, $\sim 0.1 M_\odot$, is more than sufficient to explain the observations [21, 22, 67]. Only a small fraction $\sim (\bar{v}^{\text{UFD}}/v_\infty) \sim 10^{-3}$ of the produced r -process material is likely to remain in the shallow gravitational potential well of a UFD because it is produced with a velocity $v_\infty \sim 0.1 v_{\text{esc}}$. The observations are consistent with this: the required $10^{-4} M_\odot$ of r -process material is consistent with a 0.1% fraction of the $0.1 M_\odot$ produced in a single event.

We have separately fit to the r -process abundances for the MW and UFD, accounting for uncertainties in various quantities as described below. The combined requirements result in the allowed region of parameter space shown on Figure 2, along with the current constraints for PBH contribution to the DM abundance. The region denoted “all r -process” shows parameter space for which r -process observations are fully explained simultaneously in the MW and in UFDs. For our fit we have varied the input parameters over a broad range, covering significant parameter space (see Supplemental Material [51]). The enclosed region above the line can be interpreted as a constraint of r -process material over-production from PBH-NS interactions, subject to large uncertainties in astrophysical input parameters.

We note that COM-produced r -process, with an event rate of $10^{-4} - 10^{-5} \text{ Mpc}^{-3} \text{ yr}^{-1}$, could also be consistent with this analysis [68]. However, COM simulations suggest an ejecta mass of $\sim 0.01 M_\odot$. This would imply a COM rate near the upper end of the allowed range, if COM are to explain all of the r -process. Such a rate is still marginally consistent with the current Advanced LIGO (aLIGO) limits, but readily verifiable or refutable when aLIGO reaches its design sensitivity [69] in a few years. Sensitivity similar to aLIGO is expected in the upcoming Advanced Virgo [70] (aVirgo) and KAGRA [71] experiments. A recent analysis of kilonova [72] also exhibits tension with observations and highlights the need for an extremely efficient ejection of r -process material in COM scenario.

Positron emission from ejecta can explain the observed 511 keV emission line from the Galactic central region [79], which is consistent with the e^+e^- annihilation line via positronium formation. The origin of the positrons remains unknown [29]. The 511 keV line flux in the bulge component is [80] $\sim 10^{-3}$ pho-




FIG. 2: Parameter space where PBHs can account for all or partial r -process element production in the Milky Way and the UFDs simultaneously. Constraints from extragalactic γ -rays from BH evaporation [73] (EG γ), femto-lensing [74] (FL), white dwarf abundance [75] (WD), Kepler star milli/micro-lensing [76] (K), Subaru HSC micro-lensing [77] (HSC) and MACHO/EROS/OGLE micro-lensing [78] (ML) are displayed.

tons $\text{cm}^{-2} \text{ s}^{-1}$. The line can be explained through electron-positron annihilations that occur at a rate of $\Gamma(e^+e^- \rightarrow \gamma\gamma) \sim 10^{50} \text{ yr}^{-1}$. Ejected cold nuclear matter expands on a dynamical time scale of $\tau_e \sim \alpha/\sqrt{G_N \rho} = 446\alpha(\rho/\text{g cm}^{-3})^{-1/2} \text{ s}$, where $\alpha = 0.01 - 10$ is a model-dependent parameter [58, 59, 68]. At the same time, beta decays and fission raise the temperature to $T \sim 0.1 \text{ MeV}$ [58, 59, 68]. This temperature is high enough to generate a sizable equilibrium density of positrons, which leak through the surface of each clump. Taking the radius of each clump as $R \sim 0.1 \text{ km}$ and the density as $\rho \sim 10^8 \text{ g/cm}^3$, the total surface area of $0.1 M_\odot$ of ejected material is $A \sim 4\pi R^2(0.1 M_\odot/\rho)/(4\pi R^3/3) \sim 10^{20} \text{ cm}^2$. The number of positrons emitted in a single event, during the time τ_e , while the temperature $T \sim 0.1 \text{ MeV}$ is maintained, can be estimated as $N_{e^+} \sim Av\tau_e \times 2(m_e T/2\pi)^{3/2} \exp(-m_e/T)$, where v is the average speed of positrons emitted with a relativistic γ factor $\gamma \sim (3T/m_e)$. If the neutron star disruption events occur in the GC on the time scale of $\tau_d \sim 10^5 \text{ yr}$, the average rate of positron production is

$$R_{e^+} = N_{e^+}/\tau_d \sim 10^{50} \text{ yr}^{-1}. \quad (4)$$

Since the average positron energy $E_{e^+} \approx 3T$ is below 3 MeV, the positrons do not annihilate in flight in the interstellar medium [81].

Fast radio bursts, kilonovae and other signatures are expected from the PBH capture-induced NS demise. During the final stages of the event, described by dynamical time scales of the order of a few to tens of milliseconds, $10^{41} - 10^{43}$ ergs of energy stored in the magnetic field are released. Inside the cold NS, at temperatures below 0.4

MeV, the nuclear matter is a Type II superconductor and magnetic field is concentrated in flux tubes. A consequence of the rapid rearrangement of nuclear matter accompanying ejection is a prodigious release of electromagnetic radiation from magnetic field reconnection and decay. The resulting bursts of radio waves [31, 32] with duration of a few milliseconds can account for some of the observed [33] FRBs. One FRB is known to be a repeater, while the others appear to be one-time events. The FRB energy of 10^{41} erg is consistent with observations [82]. If 1–10% of the magnetic field energy is converted to radio waves, an FRB could accompany an NS destruction event. The rapidly evolving magnetic field can also accelerate charged particles leading to X-ray and γ -ray emission.

Detection of an “orphaned” kilonova (macronova) within the aLIGO, aVirgo and KAGRA sensitivity distance (~ 200 Mpc) that is *not* accompanied by a binary compact object in-spiral gravitational wave signal or a short γ -ray burst, but possibly associated with an FRB, would constitute an indirect argument that NS disruptions via PBH capture occur and could account for a significant fraction of the r -process. Sophisticated numerical simulations of PBH-induced NS collapse and of the accompanying nucleosynthesis and electromagnetic emission (including FRB) could help enable feasible observational search strategies. The search can be further assisted by detailed mapping of chemical abundances that will be made possible by the future Hitomi-2 detector. The stakes are high, as finding evidence for PBH-NS destruction could have profound implications for our understanding of the origin of the heavy elements and for the source and composition of dark matter.

Acknowledgments. We thank B. Carr, Y. Inoue, S. Nagataki, R. Rothschild, and E. Wright for helpful discussions. The work of G.M.F. was supported in part by National Science Foundation Grants No. PHY-1307372 and PHY-1614864. The work of A.K. and V.T. was supported by the U.S. Department of Energy Grant No. DE-SC0009937. A.K. was also supported by the World Premier International Research Center Initiative (WPI), MEXT, Japan.

* gfuller@ucsd.edu

† kusenko@ucla.edu

‡ vtakhist@physics.ucla.edu

- [1] Y. B. Zel’dovich and I. D. Novikov, *Sov. Astron.* **10**, 602 (1967).
- [2] S. Hawking, *Mon. Not. Roy. Astron. Soc.* **152**, 75 (1971).
- [3] B. J. Carr and S. W. Hawking, *Mon. Not. Roy. Astron. Soc.* **168**, 399 (1974).
- [4] J. Garcia-Bellido, A. D. Linde, and D. Wands, *Phys. Rev. D* **54**, 6040 (1996), arXiv:astro-ph/9605094 [astro-ph].
- [5] M. Yu. Khlopov, *Res. Astron. Astrophys.* **10**, 495 (2010), arXiv:0801.0116 [astro-ph].
- [6] P. H. Frampton, M. Kawasaki, F. Takahashi, and T. T. Yanagida, *JCAP* **1004**, 023 (2010), arXiv:1001.2308 [hep-ph].
- [7] M. Kawasaki, A. Kusenko, and T. T. Yanagida, *Phys. Lett.* **B711**, 1 (2012), arXiv:1202.3848 [astro-ph.CO].
- [8] M. Kawasaki, A. Kusenko, Y. Tada, and T. T. Yanagida, *Phys. Rev. D* **94**, 083523 (2016), arXiv:1606.07631 [astro-ph.CO].
- [9] E. Cotner and A. Kusenko, (2016), arXiv:1612.02529 [astro-ph.CO].
- [10] B. Carr, F. Kuhnel, and M. Sandstad, *Phys. Rev. D* **94**, 083504 (2016), arXiv:1607.06077 [astro-ph.CO].
- [11] K. Inomata, M. Kawasaki, K. Mukaida, Y. Tada, and T. T. Yanagida, (2016), arXiv:1611.06130 [astro-ph.CO].
- [12] K. Inomata, M. Kawasaki, K. Mukaida, Y. Tada, and T. T. Yanagida, (2017), arXiv:1701.02544 [astro-ph.CO].
- [13] J. Georg and S. Watson, (2017), arXiv:1703.04825 [astro-ph.CO].
- [14] M. E. Burbidge, G. R. Burbidge, W. A. Fowler, and F. Hoyle, *Rev. Mod. Phys.* **29**, 547 (1957).
- [15] M. Arnould, S. Goriely, and K. Takahashi, *Phys. Rept.* **450**, 97 (2007), arXiv:0705.4512 [astro-ph].
- [16] A. G. W. Cameron, *Proc. Astron. Soc. Pac.* **69**, 201 (1957).
- [17] J. W. Truran, J. J. Cowan, C. A. Pilachowski, and C. Sneden, *Publ. Astron. Soc. Pac.* **114**, 1293 (2002), arXiv:astro-ph/0209308 [astro-ph].
- [18] Y. Z. Qian and G. J. Wasserburg, *Phys. Rept.* **442**, 237 (2007), arXiv:0708.1767 [astro-ph].
- [19] D. Argast, M. Samland, F. K. Thielemann, and Y. Z. Qian, *Astron. Astrophys.* **416**, 997 (2004), arXiv:astro-ph/0309237 [astro-ph].
- [20] C. Freiburghaus, S. Rosswog, and F.-K. Thielemann, *Astrophys. J. Lett.* **525**, L121 (1999).
- [21] A. P. Ji, A. Frebel, A. Chiti, and J. D. Simon, *Nature* **531**, 610 (2016), arXiv:1512.01558.
- [22] P. Beniamini, K. Hotokezaka, and T. Piran, *Astrophys. J.* **832**, 149 (2016), arXiv:1608.08650 [astro-ph.HE].
- [23] J. Dexter and R. M. O’Leary, *Astrophys. J.* **783**, L7 (2014), arXiv:1310.7022 [astro-ph.GA].
- [24] J. Bramante and T. Linden, *Astrophys. J.* **826**, 57 (2016), arXiv:1601.06784 [astro-ph.HE].
- [25] K. Mori et al., *Astrophys. J.* **770**, L23 (2013), arXiv:1305.1945 [astro-ph.HE].
- [26] J. A. Kennea et al., *Astrophys. J.* **770**, L24 (2013), arXiv:1305.2128 [astro-ph.HE].
- [27] J. D. Bradford et al., *Astrophys. J.* **743**, 167 (2011), [Erratum: *Astrophys. J.* 778,85(2013)], arXiv:1110.0484 [astro-ph.CO].
- [28] R. Ibata et al., *Mon. Not. Roy. Astron. Soc.* **428**, 3648 (2013), arXiv:1210.7787 [astro-ph.CO].
- [29] N. Prantzos et al., *Rev. Mod. Phys.* **83**, 1001 (2011), arXiv:1009.4620 [astro-ph.HE].
- [30] R. E. Lingens, J. C. Hildon, and R. E. Rothschild, *Phys. Rev. Lett.* **103**, 031301 (2009), arXiv:0904.1025 [astro-ph.HE].
- [31] J. Fuller and C. Ott, *Mon. Not. Roy. Astron. Soc.* **450**, L71 (2015), arXiv:1412.6119 [astro-ph.HE].
- [32] Z. Shand, A. Ouyed, N. Koning, and R. Ouyed, *Res. Astron. Astrophys.* **16**, 080 (2016), arXiv:1505.08147 [astro-ph.HE].
- [33] D. R. Lorimer, M. Bailes, M. A. McLaughlin, D. J.

- Narkevic, and F. Crawford, *Science* **318**, 777 (2007), arXiv:0709.4301 [astro-ph].
- [34] D. Kasen, R. Fernandez, and B. Metzger, *Mon. Not. Roy. Astron. Soc.* **450**, 1777 (2015), arXiv:1411.3726 [astro-ph.HE].
- [35] L.-X. Li and B. Paczynski, *Astrophys. J.* **507**, L59 (1998), arXiv:astro-ph/9807272 [astro-ph].
- [36] B. D. Metzger et al., *Mon. Not. Roy. Astron. Soc.* **406**, 2650 (2010), arXiv:1001.5029 [astro-ph.HE].
- [37] L. F. Roberts, D. Kasen, W. H. Lee, and E. Ramirez-Ruiz, *Astrophys. J.* **736**, L21 (2011), arXiv:1104.5504 [astro-ph.HE].
- [38] K. Hotokezaka and T. Piran, *Mon. Not. Roy. Astron. Soc.* **450**, 1430 (2015), arXiv:1501.01986 [astro-ph.HE].
- [39] T. Piran, E. Nakar, and S. Rosswog, *Mon. Not. Roy. Astron. Soc.* **430**, 2121 (2013), arXiv:1204.6242 [astro-ph.HE].
- [40] J. Barnes and D. Kasen, *Astrophys. J.* **775**, 18 (2013), arXiv:1303.5787 [astro-ph.HE].
- [41] D. Martin et al., *Astrophys. J.* **813**, 2 (2015), arXiv:1506.05048 [astro-ph.SR].
- [42] J. I. Read, L. Mayer, A. M. Brooks, F. Governato, and G. Lake, *Mon. Not. Roy. Astron. Soc.* **397**, 44 (2009), arXiv:0902.0009 [astro-ph.GA].
- [43] J. I. Read, G. Lake, O. Agertz, and V. P. Debattista, *Mon. Not. Roy. Astron. Soc.* **389**, 1041 (2008), arXiv:0803.2714 [astro-ph].
- [44] R. M. O’Leary, M. D. Kistler, M. Kerr, and J. Dexter, (2016), arXiv:1601.05797 [astro-ph.HE].
- [45] M. Ajello et al. (Fermi-LAT), *Astrophys. J.* **819**, 44 (2016), arXiv:1511.02938 [astro-ph.HE].
- [46] D. R. Lorimer, *Proc., IAU Symp.* **291**, 237 (2013), arXiv:1210.2746 [astro-ph.GA].
- [47] J. M. Cordes and D. F. Chernoff, *Astrophys. J.* **482**, 971 (1997), arXiv:astro-ph/9706162 [astro-ph].
- [48] C.-A. Faucher-Giguere and V. M. Kaspi, *Astrophys. J.* **643**, 332 (2006), arXiv:astro-ph/0512585 [astro-ph].
- [49] F. Capela, M. Pshirkov, and P. Tinyakov, *Phys. Rev.* **D87**, 123524 (2013), arXiv:1301.4984 [astro-ph.CO].
- [50] F. Capela, M. Pshirkov, and P. Tinyakov, *Phys. Rev.* **D90**, 083507 (2014), arXiv:1403.7098 [astro-ph.CO].
- [51] See Supplemental Material [url], which includes Ref. [24, 42, 43, 47, 52, 54, 55, 83–88], for further details regarding pulsar kicks, viscosity and magnetic differential rotation breaking, as well as variation of various input parameters for the fit.
- [52] S. L. Shapiro and S. A. Teukolsky, *Black holes, white dwarfs, and neutron stars* (1983).
- [53] F. C. Zelati et al., *Mon. Not. Roy. Astron. Soc.* **449**, 2685 (2015), arXiv:1503.01307 [astro-ph.HE].
- [54] C. Kouvaris and P. Tinyakov, *Phys. Rev.* **D90**, 043512 (2014), arXiv:1312.3764 [astro-ph.SR].
- [55] D. Markovic, *Mon. Not. Roy. Astron. Soc.* **277**, 25 (1995).
- [56] N. Andersson and K. D. Kokkotas, *Int. J. Mod. Phys.* **D10**, 381 (2001), arXiv:gr-qc/0010102 [gr-qc].
- [57] B. J. Owen, L. Lindblom, C. Cutler, B. F. Schutz, A. Vecchio, and N. Andersson, *Phys. Rev.* **D58**, 084020 (1998), arXiv:gr-qc/9804044 [gr-qc].
- [58] J. M. Lattimer, F. Mackie, D. G. Ravenhall, and D. N. Schramm, *Astrophys. J.* **213**, 225 (1977).
- [59] B. S. Meyer, *Astrophys. J.* **343**, 254 (1989).
- [60] J. Lippuner and L. F. Roberts, *Astrophys. J.* **815**, 82 (2015), arXiv:1508.03133 [astro-ph.HE].
- [61] P. Jaikumar, B. S. Meyer, K. Otsuki, and R. Ouyed, *Astron. Astrophys.* **471**, 227 (2007), arXiv:nucl-th/0610013 [nucl-th].
- [62] S. Goriely, A. Bauswein, and H. T. Janka, *Astrophys. J.* **738**, L32 (2011), arXiv:1107.0899 [astro-ph.SR].
- [63] S. Goriely et al., *J. Phys. Conf. Ser.*, **665**, 012052 (2016).
- [64] M. Eichler et al., *Astrophys. J.* **808**, 30 (2015), arXiv:1411.0974 [astro-ph.HE].
- [65] M. R. Mumpower, G. C. McLaughlin, R. Surman, and A. W. Steiner, *Astrophys. J.* **833**, 282 (2016), arXiv:1603.02600 [nucl-th].
- [66] S. Wanajo, Y. Sekiguchi, N. Nishimura, K. Kiuchi, K. Kyutoku, and M. Shibata, *Astrophys. J.* **789**, L39 (2014), arXiv:1402.7317 [astro-ph.SR].
- [67] Y. Hirai et al., *Astrophys. J.* **814**, 41 (2015), arXiv:1509.08934 [astro-ph.GA].
- [68] S. Goriely, N. Chamel, H. T. Janka, and J. M. Pearson, *Astron. Astrophys.* **531**, A78 (2011), arXiv:1105.2453 [astro-ph.SR].
- [69] B. Cote et al., *Astrophys. J.* **836**, 230 (2017), arXiv:1610.02405 [astro-ph.GA].
- [70] F. Acernese et al. (VIRGO), *Class. Quant. Grav.* **32**, 024001 (2015), arXiv:1408.3978 [gr-qc].
- [71] Y. Aso et al. (KAGRA), *Phys. Rev.* **D88**, 043007 (2013), arXiv:1306.6747 [gr-qc].
- [72] Z.-P. Jin, K. Hotokezaka, X. Li, M. Tanaka, P. D’Avanzo, Y.-Z. Fan, S. Covino, D.-M. Wei, and T. Piran, *Nature Commun.* **7**, 2898 (2016), arXiv:1603.07869 [astro-ph.HE].
- [73] B. J. Carr, K. Kohri, Y. Sendouda, and J. Yokoyama, *Phys. Rev.* **D81**, 104019 (2010), arXiv:0912.5297 [astro-ph.CO].
- [74] A. Barnacka, J. F. Glicenstein, and R. Moderski, *Phys. Rev.* **D86**, 043001 (2012), arXiv:1204.2056 [astro-ph.CO].
- [75] P. W. Graham, S. Rajendran, and J. Varela, *Phys. Rev.* **D92**, 063007 (2015), arXiv:1505.04444 [hep-ph].
- [76] K. Griest, A. M. Cieplak, and M. J. Lehner, *Astrophys. J.* **786**, 158 (2014), arXiv:1307.5798 [astro-ph.CO].
- [77] H. Niikura et al., (2017), arXiv:1701.02151 [astro-ph.CO].
- [78] P. Tisserand et al. (EROS-2), *Astron. Astrophys.* **469**, 387 (2007), arXiv:astro-ph/0607207 [astro-ph].
- [79] P. Jean et al., *Astron. Astrophys.* **407**, L55 (2003), arXiv:astro-ph/0309484 [astro-ph].
- [80] J. Knodlseder et al., *Astron. Astrophys.* **411**, L457 (2003), arXiv:astro-ph/0309442 [astro-ph].
- [81] J. F. Beacom and H. Yuksel, *Phys. Rev. Lett.* **97**, 071102 (2006), arXiv:astro-ph/0512411 [astro-ph].
- [82] K. Dolag, B. M. Gaensler, A. M. Beck, and M. C. Beck, *Mon. Not. Roy. Astron. Soc.* **451**, 4277 (2015), arXiv:1412.4829 [astro-ph.CO].
- [83] S. L. Shapiro and M. Shibata, *Astrophys. J.* **577**, 904 (2002), arXiv:astro-ph/0209251 [astro-ph].
- [84] M. Kaplinghat, R. E. Keeley, T. Linden, and H.-B. Yu, *Phys. Rev. Lett.* **113**, 021302 (2014), arXiv:1311.6524 [astro-ph.CO].
- [85] J. F. Navarro, C. S. Frenk, and S. D. M. White, *Astrophys. J.* **490**, 493 (1997), arXiv:astro-ph/9611107 [astro-ph].
- [86] S. E. Kopusov et al., *Astrophys. J.* **811**, 62 (2015), arXiv:1504.07916 [astro-ph.GA].
- [87] A. Burkert, *Proc., IAU Symp.* **171**, 175 (1996), [*Astrophys. J.*447,L25(1995)], arXiv:astro-ph/9504041 [astro-ph].

[88] A. G. Lyne et al., Mon. Not. Roy. Astron. Soc. **295**, 743 (1998).