

This is the accepted manuscript made available via CHORUS. The article has been published as:

Proton-Proton Fusion and Tritium β Decay from Lattice Quantum Chromodynamics

Martin J. Savage, Phiala E. Shanahan, Brian C. Tiburzi, Michael L. Wagman, Frank Winter, Silas R. Beane, Emmanuel Chang, Zohreh Davoudi, William Detmold, and Kostas Orginos
(NPLQCD Collaboration)

Phys. Rev. Lett. **119**, 062002 — Published 10 August 2017

DOI: [10.1103/PhysRevLett.119.062002](https://doi.org/10.1103/PhysRevLett.119.062002)



Proton-proton fusion and tritium β -decay from lattice quantum chromodynamics

Martin J. Savage,^{1,2} Phiala E. Shanahan,^{3,2} Brian C. Tiburzi,^{4,5,6,2} Michael L. Wagman,^{7,2} Frank Winter,⁸
 Silas R. Beane,^{7,2} Emmanuel Chang,¹ Zohreh Davoudi,^{3,2} William Detmold,^{3,2} and Kostas Orginos^{9,8}

(NPLQCD Collaboration)

¹*Institute for Nuclear Theory, University of Washington, Seattle, WA 98195-1550, USA*

²*Kavli Institute for Theoretical Physics, University of California, Santa Barbara, CA 93106, USA*

³*Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA*

⁴*Department of Physics, The City College of New York, New York, NY 10031, USA*

⁵*Graduate School and University Center, The City University of New York, New York, NY 10016, USA*

⁶*RIKEN BNL Research Center, Brookhaven National Laboratory, Upton, NY 11973, USA*

⁷*Department of Physics, University of Washington, Box 351560, Seattle, WA 98195, USA*

⁸*Jefferson Laboratory, 12000 Jefferson Avenue, Newport News, VA 23606, USA*

⁹*Department of Physics, College of William and Mary, Williamsburg, VA 23187-8795, USA*

(Dated: July 6, 2017)

The nuclear matrix element determining the $pp \rightarrow de^+\nu$ fusion cross section and the Gamow-Teller matrix element contributing to tritium β -decay are calculated with lattice Quantum Chromodynamics (QCD) for the first time. Using a new implementation of the background field method, these quantities are calculated at the SU(3)-flavor-symmetric value of the quark masses, corresponding to a pion mass of $m_\pi \sim 806$ MeV. The Gamow-Teller matrix element in tritium is found to be $0.979(03)(10)$ at these quark masses, which is within 2σ of the experimental value. Assuming that the short-distance correlated two-nucleon contributions to the matrix element (meson-exchange currents) depend only mildly on the quark masses, as seen for the analogous magnetic interactions, the calculated $pp \rightarrow de^+\nu$ transition matrix element leads to a fusion cross section at the physical quark masses that is consistent with its currently accepted value. Moreover, the leading two-nucleon axial counterterm of pionless effective field theory is determined to be $L_{1,A} = 3.9(0.2)(1.0)(0.4)(0.9) \text{ fm}^3$ at a renormalization scale set by the physical pion mass, also in agreement with the accepted phenomenological range. This work concretely demonstrates that weak transition amplitudes in few-nucleon systems can be studied directly from the fundamental quark and gluon degrees of freedom and opens the way for subsequent investigations of many important quantities in nuclear physics.

PACS numbers: 11.15.Ha, 12.38.Gc, 13.40.Gp

Weak nuclear processes play a central role in many settings, from the instability of the neutron to the dynamics of core-collapse supernova. In this work, the results of the first lattice Quantum Chromodynamics (LQCD) calculations of two such processes are presented, namely the $pp \rightarrow de^+\nu_e$ fusion process and tritium β -decay. The $pp \rightarrow de^+\nu$ process is centrally important in astrophysics as it is primarily responsible for initiating the proton-proton fusion chain reaction that provides the dominant energy production mechanism in stars like the Sun. Significant theoretical effort has been expended in refining calculations of the $pp \rightarrow de^+\nu$ cross section at the energies relevant to solar burning, and progress continues to be made with a range of techniques [1–10], as summarized in Ref. [11]. This process is related to the $\bar{\nu}d \rightarrow nne^+$ neutrino breakup reaction [12–14], relevant to the measurement of neutrino oscillations at the Sudbury Neutrino Observatory (SNO) [15, 16], and to the muon capture reaction, $\mu^-d \rightarrow nn\nu_\mu$, which is the focus of current investigation in the MuSun experiment [17]. The

second process studied in this work, tritium β -decay, is a powerful tool for investigating the weak interactions of the Standard model and plays an important role in the search for new physics. The super-allowed process ${}^3\text{H} \rightarrow {}^3\text{He} e^- \bar{\nu}$ is theoretically clean and is the simplest semileptonic weak decay of a nuclear system. In contrast to pp fusion, this decay has been very precisely studied in the laboratory (see Ref. [18] for a review) and provides important constraints on the antineutrino mass [19]. Tritium β -decay is also potentially sensitive to sterile neutrinos [20, 21] and to interactions not present in the Standard Model [21–24]. Although the dominant contributions to the decay rate are under theoretical control as this is a super-allowed process, the Gamow-Teller (GT) contribution (axial current) is somewhat more challenging to address than the Fermi (F) contribution (vector current). Improved constraints on multi-body contributions to the GT matrix element will translate into reduced uncertainties in predictions for decay rates of larger nuclei and are a first step towards understanding

the quenching of g_A in nuclei [25–27], a long standing problem in nuclear theory.

In this Letter, LQCD is used to study the $pp \rightarrow de^+\nu_e$ fusion process and the Gamow-Teller matrix element contributing to tritium β -decay for the first time, albeit at unphysically large values of the light quark masses and neglecting the effects of isospin-breaking and electromagnetism. This is accomplished using a new algorithm for implementing background fields, which is superior to existing methods. Further, the quantities of interest are extracted at high precision using a refined analysis strategy made possible by this new approach. For $pp \rightarrow de^+\nu_e$, the deviations from the single-nucleon contributions are small but are well resolved with the new technique. The leading two-nucleon axial counterterm of pionless effective field theory ($\not\equiv$ EFT), $L_{1,A}$, is determined. The axial coupling of ${}^3\text{H}$ that determines the matrix element for ${}^3\text{H} \rightarrow {}^3\text{He} e^- \bar{\nu}$ in the isospin limit is found to be slightly smaller than that of the proton and is consistent with previous phenomenological estimates [6].

Background Axial Fields: Background field techniques were first used in LQCD in the pioneering works of Ref. [28] and Refs. [29, 30] in the cases of axial and magnetic fields, respectively. Significant effort has been applied to using background electromagnetic fields to extract magnetic moments and electromagnetic polarizabilities of hadrons [31–35] and nuclei [36–38], as well as the magnetic transition amplitude for the $np \rightarrow d\gamma$ process [39]. Very recently, axial background fields have been employed to extract the axial charge of the proton [40, 41], and generalizations to nonzero momentum transfer [42–44] have been used [45] to access the axial form factor of the nucleon.

In this work, a new method is used to generate hadronic correlation functions order-by-order in the background field. In the standard approach, correlation functions are constructed from the contraction of quark propagators that are modified by the presence of a background field. The same effect can be achieved by directly constructing propagators that include explicit current insertions, then using such propagators to construct correlation functions. For the quantities studied in this work only a single insertion of the background axial field is required. To this end, the *compound propagator*

$$S_{\lambda_q; \Gamma}^{(q)}(x, y) = S^{(q)}(x, y) + \lambda_q \int dz S^{(q)}(x, z) \Gamma S^{(q)}(z, y) \quad (1)$$

is constructed for $\Gamma = \gamma_3 \gamma_5$ and flavors $q = \{u, d\}$, where $S^{(q)}(x, y)$ is the quark propagator of flavor q and λ_q is a constant (a similar approach is implemented in Ref. [46] in a different context). The second term in this expression is computed using standard sequential source techniques and the procedure can be repeated to produce propagators with higher-order couplings. These compound propagators are sufficient to construct the isovec-

tor axial matrix elements for zero momentum insertion in any hadronic or nuclear system.¹ This work focuses on zero-momentum-projected correlation functions,

$$C_{\lambda_u; \lambda_d}^{(h)}(t) = \sum_{\mathbf{x}} \langle 0 | \chi_h(\mathbf{x}, t) \chi_h^\dagger(0) | 0 \rangle_{\lambda_u; \lambda_d}, \quad (2)$$

where $\langle \dots \rangle_{\lambda_u; \lambda_d}$ denotes the expectation value determined using the compound propagators. The interpolating operators for hadrons and nuclei, χ_h , are those previously used to study spectroscopy of these systems [47, 48]. By construction, $C_{\lambda_u; \lambda_d}^{(h)}(t)$ is a polynomial of maximum order $\lambda_u^{N_u} \lambda_d^{N_d}$ in the field strengths, where $N_{u(d)}$ is the number of up (down) quarks in the particular interpolating operator.

Details of the LQCD Calculation: The calculations presented below used an ensemble of gauge-field configurations generated with a clover-improved fermion action [49] and a Lüscher-Weisz gauge action [50]. The ensemble was generated with $N_f = 3$ degenerate light-quark flavors with masses tuned to the physical strange quark mass, producing a pion of mass $m_\pi \sim 806$ MeV, with a volume of $L^3 \times T = 32^3 \times 48$ and a lattice spacing of $a \sim 0.12$ fm. For these calculations, 437 configurations, with a spacing of 10 trajectories between configurations, were used. Correlation functions were computed for $h = \{p, n, d, nn, np({}^1S_0), pp, {}^3\text{H}, {}^3\text{He}\}$ from propagators generated from a smeared source and either a smeared (SS) or point (SP) sink. Sixteen different source locations were averaged over on each configuration. Compound propagators and correlation functions were calculated at six different values of the background field strength parameter $\lambda = \{\pm 0.05, \pm 0.1, \pm 0.2\}$. The axial current renormalization factor $Z_A = 0.867(43)$ was determined from computations of the vector current in the proton, noting that $Z_A = Z_V + \mathcal{O}(a)$ and assigning a 5% systematic uncertainty associated with lattice-spacing artifacts (statistical uncertainties are negligible).²

The Proton Axial Charge: The simplest matrix element of the isovector axial current determines the axial charge of the proton. The correlation function $C_{\lambda_u; \lambda_d}^{(p)}(t)$ is at most quadratic in λ_u and linear in λ_d when constructed from the compound propagators $S_{\lambda_u; \gamma_3 \gamma_5}^{(u)}(x, y)$ and $S_{\lambda_d; \gamma_3 \gamma_5}^{(d)}(x, y)$, as the proton has two valence up quarks and one valence down quark. Consequently, using at least one (two) nonzero value(s) of $\lambda_{d(u)}$ enables extraction of the axial current matrix element as the linear

¹ This work does not address isoscalar responses which also involve insertions on the sea-quark propagators.

² A determination that removes the leading lattice-spacing artifacts leads to $Z_A = 0.8623(01)(71)$ [51, 52] at a pion mass of $m_\pi \sim 317$ MeV.




FIG. 1. The ratios of correlation functions that determine the unrenormalized isovector axial charge of the proton. The orange diamonds (blue circles) correspond to the SS (SP) effective correlator ratios, $\bar{R}_p(t)$, defined in Eq. (4), and the band corresponds to a constant fit to the plateau interval of both SS and SP.

response by using suitable polynomial fits. The difference of the up-quark and down-quark matrix elements can be used to construct the desired three-point function containing the isovector axial current. This can then be combined with the zero-field two-point function to form a ratio that asymptotes to the desired axial charge at late times, namely

$$R_p(t) = \frac{C_{\lambda_u; \lambda_d=0}^{(p)}(t) \big|_{\mathcal{O}(\lambda_u)} - C_{\lambda_u=0; \lambda_d}^{(p)}(t) \big|_{\mathcal{O}(\lambda_d)}}{C_{\lambda_u=0; \lambda_d=0}^{(p)}(t)}, \quad (3)$$

where the ratios are spin-weighted averages, and “ $\big|_{\mathcal{O}(\lambda_q)}$ ” extracts the coefficient of λ_q in the preceding expression. Then,

$$\bar{R}_p(t) \equiv R_p(t+1) - R_p(t) \xrightarrow{t \rightarrow \infty} \frac{g_A}{Z_A}, \quad (4)$$

where corrections to this relation from backwards propagating states originating from the finite extent of the time direction are suppressed by at least $e^{-2m_\pi T/3} \sim 10^{-7}$ in the signal region in the present set of calculations. The *effective- g_A plots* resulting from the correlator differences are shown in Fig. 1, along with the result of a combined constant fit to the SS and SP ratios that extracts g_A/Z_A from the late-time asymptote. The extracted value is $g_A/Z_A = 1.298(2)(7)$, where the first uncertainty is statistical (determined from a bootstrap analysis) and the second is systematic (arising from choices of fit ranges in both the field strengths and temporal separation as well as from differences in analysis techniques). Including the renormalization factor yields an axial charge of $g_A = 1.13(2)(7)$, which is consistent with previous determinations from standard three-point func-

tion techniques at this pion mass [53, 54].

The GT Matrix Element for Tritium β -decay: The half-life of tritium, $t_{1/2}$, is related to the F and GT matrix elements by [1]

$$\frac{(1 + \delta_R)f_V}{K/G_V^2} t_{1/2} = \frac{1}{\langle \mathbf{F} \rangle^2 + f_A/f_V g_A^2 \langle \mathbf{GT} \rangle^2}, \quad (5)$$

where the factors on the left-hand side are known precisely from theory or experiment. On the right-hand side, $f_{A,V}$ denote known Fermi functions [55] and $\langle \mathbf{F} \rangle$ and $\langle \mathbf{GT} \rangle$ are the F and GT reduced matrix elements, respectively. The Ademollo-Gatto theorem [56] implies $\langle \mathbf{F} \rangle \sim 1$, modified only by second-order isospin-breaking and by electromagnetic corrections. However, $\langle {}^3\text{He} | \bar{q} \gamma_k \gamma_5 \tau^+ q | {}^3\text{H} \rangle = \bar{u} \gamma_k \gamma_5 \tau^+ u g_A \langle \mathbf{GT} \rangle$ (assuming vanishing electron mass and at vanishing lepton momentum) is less constrained, and its evaluation is the focus of this section.

By isospin symmetry, the spin-averaged GT matrix element for ${}^3\text{H} \rightarrow {}^3\text{He} \ e^- \bar{\nu}$ is related to the axial charge of the triton, $g_A({}^3\text{H})$, when the light quarks are degenerate and in the absence of electromagnetism. Analogous to $R_p(t)$ in Eq. (3), the ratio $R_{3\text{H}}(t)$ of correlation functions in background fields is constructed such that, analogous to Eq. (4), $\bar{R}_{3\text{H}}(t) \rightarrow g_A({}^3\text{H})/Z_A$ in the large-time limit. The analysis of these correlation functions is more complex than for the proton because the triton has four up quarks and five down quarks and the correlators are thus quartic and quintic polynomials in $\lambda_{u,d}$, respectively. Polynomial fits to the calculated correlation functions are sufficient to extract the terms linear in $\lambda_{u,d}$. Results for $\bar{R}_{3\text{H}}(t)$ are shown in Fig. 2 along with a constant fit to the asymptotic value $g_A({}^3\text{H})/Z_A$. Also shown in Fig. 2 is $\langle \mathbf{GT} \rangle(t) = \bar{R}_{3\text{H}}(t)/\bar{R}_p(t)$, which is independent of Z_A , and the fit to its asymptotic value, $g_A({}^3\text{H})/g_A$. Analyses of these ratios lead to

$$\frac{g_A({}^3\text{H})}{Z_A} = 1.272(6)(22), \quad \frac{g_A({}^3\text{H})}{g_A} = 0.979(3)(10), \quad (6)$$

where the first uncertainties are statistical and the second arise from systematics as described for g_A . The result for $g_A({}^3\text{H})/g_A$ is quite close to the precise, experimentally-determined value of $\langle \mathbf{GT} \rangle = 0.9511(13)$ [6] at the physical quark masses. In the context of $\not{E}\text{FT}$, the short-distance two-nucleon axial-vector operator, with coefficient $L_{1,A}$ [4], is expected to give the leading contribution to the difference of this ratio from unity [57].

The Low-Energy Proton-Proton Fusion Cross Section: The low-energy cross section for $pp \rightarrow de^+ \nu$ is dictated by the matrix element

$$|\langle d; j | A_k^- | pp \rangle| \equiv g_A C_\eta \sqrt{\frac{32\pi}{\gamma^3}} \Lambda(p) \delta_{jk}, \quad (7)$$

where $A_k^a(x)$ is the axial current with isospin and spin components a and k respectively, j is the deuteron spin




FIG. 2. The ratios of correlation functions that determine the unrenormalized isovector axial matrix element in ^3H (upper panel), and the ratio of the isovector axial matrix element in ^3H to that in the proton (lower panel). The orange diamonds (blue circles) correspond to the SS (SP) effective correlator ratios and the bands correspond to constant fits to the asymptotic behavior.

index, C_η is the Sommerfeld factor and γ is the deuteron binding momentum. The quantity $\Lambda(p)$ has been calculated at threshold in $\not\epsilon\text{EFT}$ to N^2LO [3] and N^4LO [4] and later with a dibaryon approach [10, 57] and in pionful EFT [58]. With the approach of Ref. [4], resumming all of the effective range contributions [10, 59, 60], $\Lambda(0)$ at N^2LO is related to the renormalization-scale independent short-distance quantity $L_{1,A}^{sd-2b}$ that is a solely two-body contribution, along with scattering parameters and Coulomb corrections:

$$\Lambda(0) = \frac{1}{\sqrt{1-\gamma\rho}} \{e^\chi - \gamma a_{pp} [1 - \chi e^\chi \Gamma(0, \chi)] + \frac{1}{2} \gamma^2 a_{pp} \sqrt{r_1 \rho}\} - \frac{1}{2g_A} \gamma a_{pp} \sqrt{1-\gamma\rho} L_{1,A}^{sd-2b}. \quad (8)$$

Here $\chi = \alpha M_p / \gamma$, where α is the QED fine-structure constant and M_p is the mass of the proton. The pp scattering length is a_{pp} , r_1 and ρ are the effective ranges in the $^1\text{S}_0$ and $^3\text{S}_1$ channels, respectively, and $\Gamma(0, \chi)$ is the incomplete gamma function. A determination of $L_{1,A}^{sd-2b}$, or equivalently of the $\not\epsilon\text{EFT}$ coupling $L_{1,A}$ which is determined from the scale-independent constant

$$\bar{L}_{1,A} = \frac{1}{2g_A} \frac{1-\gamma\rho}{\gamma} L_{1,A}^{sd-2b} - \frac{1}{2} \sqrt{r_1 \rho} \quad (9)$$

as shown explicitly in Ref. [4], is a goal of the present LQCD calculations.

A background isovector axial-vector field mixes the $J_z = I_z = 0$ components of the $^3\text{S}_1$ and $^1\text{S}_0$ two-nucleon channels, enabling the pp -fusion matrix element to be accessed. Using the new background field construction, the relevant off-diagonal matrix element $C_{\lambda_u; \lambda_d}^{(^3\text{S}_1, ^1\text{S}_0)}(t)$ is a cubic polynomial in both λ_u and λ_d . In Ref. [39],

the analogous mixing between the two-nucleon channels induced by an isovector magnetic field was treated by diagonalizing a (channel-space) matrix of correlators and determining the splittings between energy eigenvalues. This provided access to the matrix element dictating $np \rightarrow d\gamma$ at low energies, as was proposed in Ref. [61]. Such a method can also be used for the axial field, but the improved approach implemented here makes use of the finite-order polynomial structure to access the matrix element directly. For a background field that couples to the u quarks,

$$C_{\lambda_u; \lambda_d=0}^{(^3\text{S}_1, ^1\text{S}_0)}(t) = \lambda_u \sum_{\tau=0}^t \sum_{\mathbf{x}, \mathbf{y}} \langle 0 | \chi_{^3\text{S}_1}^3(\mathbf{x}, t) A_3^u(\mathbf{y}, \tau) \chi_{^1\text{S}_0}^\dagger(0) | 0 \rangle + c_2 \lambda_u^2 + c_3 \lambda_u^3, \quad (10)$$

where $\chi_{^3\text{S}_1}^3(\chi_{^1\text{S}_0})$ is an interpolating field for the $J_z = 0$ ($I_z = 0$) component of the $^3\text{S}_1$ ($^1\text{S}_0$) channel, $A_3^u = \bar{u}\gamma_3\gamma_5 u$, and $c_{2,3}$ are irrelevant terms. Calculations of the background field correlators at three or more values of λ_u allow for the extraction of the term that is linear in λ_u . A similar procedure yields the term that is linear in λ_d from background fields coupling to the d quark. Taking the difference of the ratios of these terms to the corresponding zero-field two-point functions determines the transition matrix element in the finite lattice volume;

$$R_{^3\text{S}_1, ^1\text{S}_0}(t) = \frac{C_{\lambda_u, \lambda_d=0}^{(^3\text{S}_1, ^1\text{S}_0)}(t) \Big|_{\mathcal{O}(\lambda_u)} - C_{\lambda_u=0, \lambda_d}^{(^3\text{S}_1, ^1\text{S}_0)}(t) \Big|_{\mathcal{O}(\lambda_d)}}{\sqrt{C_{\lambda_u=0, \lambda_d=0}^{(^3\text{S}_1, ^3\text{S}_1)}(t)} C_{\lambda_u=0, \lambda_d=0}^{(^1\text{S}_0, ^1\text{S}_0)}(t)}}. \quad (11)$$

Consequently, the difference between ratios at neighboring timeslices determines the isovector matrix element;

$$\bar{R}_{^3\text{S}_1, ^1\text{S}_0}(t) \equiv R_{^3\text{S}_1, ^1\text{S}_0}(t+1) - R_{^3\text{S}_1, ^1\text{S}_0}(t) \xrightarrow{t \rightarrow \infty} \frac{\langle ^3\text{S}_1; J_z=0 | A_3^3 | ^1\text{S}_0; I_z=0 \rangle}{Z_A}, \quad (12)$$

in the limit where $\Delta E = E_d - E_{pp}$ is small (as is the case with the quark masses used in this calculation [47]), and when the contributions from excited states are suppressed. This quantity, measured with both SS and SP correlators, is shown in Fig. 3, along with the extracted value of the axial matrix element, $\langle ^3\text{S}_1; J_z=0 | A_3^3 | ^1\text{S}_0; I_z=0 \rangle / Z_A = 2.568(5)(31)$, where the first uncertainty is statistical and the second is a systematic encompassing choices of fit ranges in time, field strength and variations in analysis techniques. The latter includes an estimate of the violation of Wigner's $\text{SU}(4)$ symmetry, contributing an uncertainty $\mathcal{O}(1/N_c^4) \sim 1\%$ to the extraction of the matrix element in the large- N_c limit. At the pion mass of this study, the initial and final two-nucleon states are deeply bound [47] and the finite-volume effects in the matrix elements are negligible [62, 63]. At lighter values of the quark masses, where the $np(^1\text{S}_0)$ system and/or the deuteron are unbound or only weakly bound, the connection between finite-volume




FIG. 3. The ratios of correlation functions that determine the unrenormalized isovector axial matrix element in the $J_z = I_z = 0$ coupled two-nucleon system (upper panel), and the unrenormalized difference between the axial matrix element in this channel and $2g_A$ (lower panel). The orange diamonds (blue circles) correspond to the SS (SP) effective correlator ratios and the bands correspond to fits to the asymptotic plateau behavior and include only the statistical and fitting systematic uncertainties (the additional 1% uncertainty from Wigner symmetry breaking is not represented in the bands).

matrix elements and transition amplitudes requires the framework developed in Refs. [62, 63].

To isolate the two-body contribution, the combination $L_{1,A}^{sd-2b}(t)/Z_A = [\bar{R}_{3S_1,1S_0}(t) - 2\bar{R}_p(t)]/2$ is formed as shown in the lower panel of Fig. 3. Taking advantage of the near-degeneracy of the 3S_1 and 1S_0 two-nucleon channels at the quark masses used in this calculation, it is straightforward to show that this correlated difference leads directly to the short-distance two-nucleon quantity, $L_{1,A}^{sd-2b}$. Fitting a constant to the late-time behavior of this quantity leads to

$$\frac{L_{1,A}^{sd-2b}}{Z_A} = \frac{\langle ^3S_1; J_z = 0 | A_3^3 | ^1S_0; I_z = 0 \rangle - 2g_A}{2Z_A} = -0.011(01)(15), \quad (13)$$

where the first uncertainty is statistical and the second encompasses fitting and analysis systematics.

In light of the mild quark-mass dependence of the analogous short-distance, two-body quantity contributing to $np \rightarrow d\gamma$ [39], $L_{1,A}^{sd-2b}$ is likely to be largely insensitive to the pion mass between $m_\pi \sim 806$ MeV and its physical value. This approximate independence and the associated systematic uncertainty will need to be refined in subsequent calculations. Based on this expectation, the result obtained here at $m_\pi \sim 806$ MeV is used to estimate the value of $L_{1,A}^{sd-2b}$ at the physical pion mass by including an additional 50% additive uncertainty. Propagating this uncertainty through Eq. (8), the threshold value of $\Lambda(p)$ in this system at the physical quark masses is determined to be $\Lambda(0) = 2.659(2)(9)(5)$, where the uncertainties are statistical, fitting and analysis systematic, and

quark-mass extrapolation systematic, respectively. Uncertainties in the scattering parameters and other physical mass inputs are also propagated and included in the systematic uncertainty. This result is remarkably close to the currently accepted, precise phenomenological value, $\Lambda(0) = 2.65(1)$ [11] (see also Ref. [57]). The N²LO relation of Ref. [4], when enhanced by the summation of the effective ranges to all orders using the dibaryon field approach [10, 59, 60], gives $\Lambda(0) = 2.62(1) + 0.0105(1)L_{1,A}$, enabling a determination of the \not{A} EFT coupling

$$L_{1,A} = 3.9(0.2)(1.0)(0.4)(0.9) \text{ fm}^3, \quad (14)$$

at a renormalization scale $\mu = m_\pi$. The uncertainties are statistical, fitting and analysis systematic, mass extrapolation systematic, and a power-counting estimate of higher order corrections in \not{A} EFT, respectively. This value is also very close to previous phenomenological estimates, as summarized in Refs. [11, 14].

Summary: The primary results of this work are the isovector axial-current matrix elements in two and three-nucleon systems calculated directly from the underlying theory of the strong interactions using lattice QCD.³ These matrix elements determine the cross section for the pp fusion process $pp \rightarrow de^+\nu$ and the Gamow-Teller contribution to tritium β -decay, $^3\text{H} \rightarrow ^3\text{He} e^-\bar{\nu}$. While the calculations are performed at unphysical quark masses corresponding to $m_\pi \sim 806$ MeV and at a single lattice spacing and volume, the mild mass dependence of the analogous short-distance quantity in the $np \rightarrow d\gamma$ magnetic transition enables an estimate of the $pp \rightarrow de^+\nu$ matrix element at the physical point, and the results are found to agree within uncertainties with phenomenology. Future LQCD calculations including electromagnetic effects beyond Coulomb at lighter quark masses with isospin splittings, larger volumes, and finer lattice spacings, making use of the new techniques that are introduced here, will enable extractions of these axial matrix elements with fully quantified uncertainties and will be of great importance in phenomenology, providing increasingly precise values for the pp -fusion cross section and GT matrix element in tritium β -decay.

Beyond the current study, background axial-field calculations also allow the extraction of second-order, as well as momentum-dependent, responses to axial fields. Second-order responses are important for determining nuclear $\beta\beta$ -decay matrix elements, both with and without (for a light Majorana neutrino) the emission of associated neutrinos. Momentum-dependent axial background fields will allow the determination of nuclear effects in neutrino-nucleus scattering. In both cases, LQCD calcu-

³ See Supplemental Material for additional discussion of technical aspects of these calculations, which includes Refs. [64–68].

lations of these quantities in light nuclei will provide vital input with which to constrain the nuclear many-body methods that are used to determine the matrix elements for these processes in heavy nuclei.

Acknowledgments: We would like to thank Jiunn-Wei Chen and Peter Kammel for several interesting discussions. This research was supported in part by the National Science Foundation under grant number NSF PHY11-25915 and SRB, ZD, WD, MJS, PES, BCT and MLW acknowledge the Kavli Institute for Theoretical Physics for hospitality during completion of this work. Calculations were performed using computational resources provided by NERSC (supported by U.S. Department of Energy grant number DE-AC02-05CH11231), and by the USQCD collaboration. This research used resources of the Oak Ridge Leadership Computing Facility at the Oak Ridge National Laboratory, which is supported by the Office of Science of the U.S. Department of Energy under Contract number DE-AC05-00OR22725. The PRACE Research Infrastructure resources at the Très Grand Centre de Calcul and Barcelona Supercomputing Center were also used. Parts of the calculations used the Chroma software suite [69]. SRB was partially supported by NSF continuing grant number PHY1206498 and by the U.S. Department of Energy through grant number DE-SC001347. EC was supported in part by the USQCD SciDAC project, the U.S. Department of Energy through grant number DE-SC00-10337, and by U.S. Department of Energy grant number DE-FG02-00ER41132. ZD, WD and PES were partly supported by U.S. Department of Energy Early Career Research Award DE-SC0010495 and grant number DE-SC0011090. KO was partially supported by the U.S. Department of Energy through grant number DE-FG02-04ER41302 and through contract number DE-AC05-06OR23177 under which JSA operates the Thomas Jefferson National Accelerator Facility. MJS was supported by DOE grant number DE-FG02-00ER41132, and in part by the USQCD SciDAC project, the U.S. Department of Energy through grant number DE-SC00-10337. BCT was supported in part by a joint City College of New York-RIKEN/Brookhaven Research Center fellowship, and by the U.S. National Science Foundation, under grant number PHY15-15738. MLW was supported in part by DOE grant number DE-FG02-00ER41132. FW was partially supported through the USQCD Scientific Discovery through Advanced Computing (SciDAC) project funded by U.S. Department of Energy, Office of Science, Offices of Advanced Scientific Computing Research, Nuclear Physics and High Energy Physics and by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics under contract DE-AC05-06OR23177.

-
- [1] R. Schiavilla *et al.*, Phys. Rev. **C58**, 1263 (1998), arXiv:nucl-th/9808010 [nucl-th].
 - [2] X. Kong and F. Ravndal, Nucl. Phys. **A656**, 421 (1999), arXiv:nucl-th/9902064 [nucl-th].
 - [3] X. Kong and F. Ravndal, Phys. Rev. **C64**, 044002 (2001), arXiv:nucl-th/0004038 [nucl-th].
 - [4] M. Butler and J.-W. Chen, Phys. Lett. **B520**, 87 (2001), arXiv:nucl-th/0101017 [nucl-th].
 - [5] A. Baroni, L. Girlanda, S. Pastore, R. Schiavilla, and M. Viviani, Phys. Rev. **C93**, 015501 (2016), [Erratum: Phys. Rev. **C93**, no.4, 049902(2016)], arXiv:1509.07039 [nucl-th].
 - [6] A. Baroni, L. Girlanda, A. Kievsky, L. E. Marcucci, R. Schiavilla, and M. Viviani, Phys. Rev. **C94**, 024003 (2016), arXiv:1605.01620 [nucl-th].
 - [7] T.-S. Park, D.-P. Min, and M. Rho, Phys. Rept. **233**, 341 (1993), arXiv:hep-ph/9301295 [hep-ph].
 - [8] T.-S. Park, K. Kubodera, D.-P. Min, and M. Rho, Nucl. Phys. **A684**, 101 (2001), arXiv:nucl-th/0005069 [nucl-th].
 - [9] T. S. Park, L. E. Marcucci, R. Schiavilla, M. Viviani, A. Kievsky, S. Rosati, K. Kubodera, D. P. Min, and M. Rho, Phys. Rev. **C67**, 055206 (2003), arXiv:nucl-th/0208055 [nucl-th].
 - [10] S. Ando, J. W. Shin, C. H. Hyun, S. W. Hong, and K. Kubodera, Phys. Lett. **B668**, 187 (2008), arXiv:0801.4330 [nucl-th].
 - [11] E. G. Adelberger *et al.*, Rev. Mod. Phys. **83**, 195 (2011), arXiv:1004.2318 [nucl-ex].
 - [12] M. Butler and J.-W. Chen, Nucl. Phys. **A675**, 575 (2000), arXiv:nucl-th/9905059 [nucl-th].
 - [13] M. Butler, J.-W. Chen, and X. Kong, Phys. Rev. **C63**, 035501 (2001), arXiv:nucl-th/0008032 [nucl-th].
 - [14] M. Butler, J.-W. Chen, and P. Vogel, Phys. Lett. **B549**, 26 (2002), arXiv:nucl-th/0206026 [nucl-th].
 - [15] Q. R. Ahmad *et al.* (SNO), Phys. Rev. Lett. **89**, 011301 (2002), arXiv:nucl-ex/0204008.
 - [16] J.-W. Chen, K. M. Heeger, and R. G. H. Robertson, Phys. Rev. **C67**, 025801 (2003), arXiv:nucl-th/0210073 [nucl-th].
 - [17] V. A. Andreev *et al.* (MuSun), (2010), arXiv:1004.1754 [nucl-ex].
 - [18] E. W. Otten and C. Weinheimer, Rept. Prog. Phys. **71**, 086201 (2008), arXiv:0909.2104 [hep-ex].
 - [19] G. Drexlin, V. Hannen, S. Mertens, and C. Weinheimer, Adv. High Energy Phys. **2013**, 293986 (2013), arXiv:1307.0101 [physics.ins-det].
 - [20] S. Mertens, T. Lasserre, S. Groh, G. Drexlin, F. Glück, A. Huber, A. W. P. Poon, M. Steidl, N. Steinbrink, and C. Weinheimer, JCAP **1502**, 020 (2015), arXiv:1409.0920 [physics.ins-det].
 - [21] J. Barry, J. Heeck, and W. Rodejohann, JHEP **07**, 081 (2014), arXiv:1404.5955 [hep-ph].
 - [22] G. J. Stephenson, Jr., J. T. Goldman, and B. H. J. McKellar, Phys. Rev. **D62**, 093013 (2000), arXiv:hep-ph/0006095 [hep-ph].
 - [23] J. Bonn, K. Eitel, F. Glück, D. Sevilla-Sanchez, and N. Titov, Phys. Lett. **B703**, 310 (2011), arXiv:0704.3930 [hep-ph].
 - [24] P. O. Ludl and W. Rodejohann, JHEP **06**, 040 (2016), arXiv:1603.08690 [hep-ph].

- [25] B. Buck and S. Perez, Phys.Rev.Lett. **50**, 1975 (1983).
- [26] D. Krofcheck, E. Sugarbaker, J. Rapaport, D. Wang, R. Byrd, C. Foster, C. Goodman, I. V. Heerden, T. Taddeucci, J. N. Bahcall, C. Gaarde, S. Larsen, and D. Horen, Phys.Rev.Lett. **55**, 1051 (1985).
- [27] W.-T. Chou, E. Warburton, and B. Brown, Phys.Rev. **C47**, 163 (1993).
- [28] F. Fucito, G. Parisi, and S. Petrarca, Phys. Lett. **B115**, 148 (1982).
- [29] G. Martinelli, G. Parisi, R. Petronzio, and F. Rapuano, Phys.Lett. **B116**, 434 (1982).
- [30] C. W. Bernard, T. Draper, K. Olynyk, and M. Rushton, Phys.Rev.Lett. **49**, 1076 (1982).
- [31] F. X. Lee, L. Zhou, W. Wilcox, and J. C. Christensen, Phys.Rev. **D73**, 034503 (2006), arXiv:hep-lat/0509065 [hep-lat].
- [32] W. Detmold, B. Tiburzi, and A. Walker-Loud, Phys.Rev. **D73**, 114505 (2006), arXiv:hep-lat/0603026 [hep-lat].
- [33] W. Detmold, B. C. Tiburzi, and A. Walker-Loud, Phys.Rev. **D79**, 094505 (2009), arXiv:0904.1586 [hep-lat].
- [34] W. Detmold, B. Tiburzi, and A. Walker-Loud, Phys.Rev. **D81**, 054502 (2010), arXiv:1001.1131 [hep-lat].
- [35] A. Parreño, M. J. Savage, B. C. Tiburzi, J. Wilhelm, E. Chang, W. Detmold, and K. Orginos, (2016), arXiv:1609.03985 [hep-lat].
- [36] S. Beane, E. Chang, S. Cohen, W. Detmold, H. Lin, *et al.*, Phys.Rev.Lett. **113**, 252001 (2014), arXiv:1409.3556 [hep-lat].
- [37] E. Chang, W. Detmold, K. Orginos, A. Parreño, M. J. Savage, B. C. Tiburzi, and S. R. Beane (NPLQCD), Phys. Rev. **D92**, 114502 (2015), arXiv:1506.05518 [hep-lat].
- [38] W. Detmold, K. Orginos, A. Parreño, M. J. Savage, B. C. Tiburzi, S. R. Beane, and E. Chang, Phys. Rev. Lett. **116**, 112301 (2016), arXiv:1508.05884 [hep-lat].
- [39] S. R. Beane, E. Chang, W. Detmold, K. Orginos, A. Parreño, M. J. Savage, and B. C. Tiburzi (NPLQCD), Phys. Rev. Lett. **115**, 132001 (2015), arXiv:1505.02422 [hep-lat].
- [40] A. J. Chambers *et al.* (QCDSF/UKQCD, CSSM), Phys. Rev. **D90**, 014510 (2014), arXiv:1405.3019 [hep-lat].
- [41] A. J. Chambers *et al.*, Phys. Rev. **D92**, 114517 (2015), arXiv:1508.06856 [hep-lat].
- [42] W. Detmold, Phys. Rev. **D71**, 054506 (2005), arXiv:hep-lat/0410011 [hep-lat].
- [43] Z. Davoudi and W. Detmold, Phys. Rev. **D92**, 074506 (2015), arXiv:1507.01908 [hep-lat].
- [44] G. Bali and G. Endrodi, Phys. Rev. **D92**, 054506 (2015), arXiv:1506.08638 [hep-lat].
- [45] A. J. Chambers *et al.*, PoS **LATTICE2015**, 125 (2016), arXiv:1511.07090 [hep-lat].
- [46] G. M. de Divitiis, R. Petronzio, and N. Tantalo, Phys. Lett. **B718**, 589 (2012), arXiv:1208.5914 [hep-lat].
- [47] S. R. Beane, E. Chang, S. D. Cohen, W. Detmold, H. W. Lin, T. C. Luu, K. Orginos, A. Parreno, M. J. Savage, and A. Walker-Loud (NPLQCD), Phys. Rev. **D87**, 034506 (2013), arXiv:1206.5219 [hep-lat].
- [48] S. R. Beane *et al.* (NPLQCD), Phys. Rev. **C88**, 024003 (2013), arXiv:1301.5790 [hep-lat].
- [49] B. Sheikholeslami and R. Wohlert, Nucl.Phys. **B259**, 572 (1985).
- [50] M. Lüscher and P. Weisz, Commun.Math.Phys. **97**, 59 (1985).
- [51] J. Green, “Light and strange axial form factors of the nucleon at pion mass 317 meV,” (2016), the 34th International Symposium on Lattice Field Theory.
- [52] B. Yoon *et al.*, Phys. Rev. **D95**, 074508 (2017), arXiv:1611.07452 [hep-lat].
- [53] B. Yoon *et al.*, Phys. Rev. **D93**, 114506 (2016), arXiv:1602.07737 [hep-lat].
- [54] H.-W. Lin, (2011), arXiv:1112.2435 [hep-lat].
- [55] J. J. Simpson, Phys. Rev. **C35**, 752 (1987).
- [56] M. Ademollo and R. Gatto, Phys. Rev. Lett. **13**, 264 (1964).
- [57] H. De-Leon, L. Platter, and D. Gazit, (2016), arXiv:1611.10004 [nucl-th].
- [58] B. Acharya, B. D. Carlsson, A. Ekström, C. Forssén, and L. Platter, Phys. Lett. **B760**, 584 (2016), arXiv:1603.01593 [nucl-th].
- [59] S. R. Beane and M. J. Savage, Nucl.Phys. **A694**, 511 (2001), arXiv:nucl-th/0011067 [nucl-th].
- [60] D. R. Phillips, G. Rupak, and M. J. Savage, Phys. Lett. **B473**, 209 (2000), arXiv:nucl-th/9908054 [nucl-th].
- [61] W. Detmold and M. J. Savage, Nucl.Phys. **A743**, 170 (2004), arXiv:hep-lat/0403005 [hep-lat].
- [62] R. A. Briceño and Z. Davoudi, Phys. Rev. **D88**, 094507 (2013), arXiv:1204.1110 [hep-lat].
- [63] R. A. Briceño and M. T. Hansen, Phys. Rev. **D94**, 013008 (2016), arXiv:1509.08507 [hep-lat].
- [64] S. R. Beane *et al.*, (2017), arXiv:1705.09239 [hep-lat].
- [65] E. Berkowitz, T. Kurth, A. Nicholson, B. Joo, E. Rinaldi, M. Strother, P. M. Vranas, and A. Walker-Loud, Phys. Lett. **B765**, 285 (2017), arXiv:1508.00886 [hep-lat].
- [66] B. C. Tiburzi, M. L. Wagman, F. Winter, E. Chang, Z. Davoudi, W. Detmold, K. Orginos, M. J. Savage, and P. E. Shanahan, (2017), arXiv:1702.02929 [hep-lat].
- [67] J.-W. Chen, G. Rupak, and M. J. Savage, Nucl. Phys. **A653**, 386 (1999), arXiv:nucl-th/9902056 [nucl-th].
- [68] T. Iritani, S. Aoki, T. Doi, T. Hatsuda, Y. Ikeda, T. Inoue, N. Ishii, H. Nemura, and K. Sasaki, (2017), arXiv:1703.07210 [hep-lat].
- [69] R. G. Edwards and B. Joo (SciDAC Collaboration, LHPC Collaboration, UKQCD Collaboration), Nucl.Phys.Proc.Suppl. **140**, 832 (2005), arXiv:hep-lat/0409003 [hep-lat].