

This is the accepted manuscript made available via CHORUS. The article has been published as:

## Exposing the QCD Splitting Function with CMS Open Data

Andrew Larkoski, Simone Marzani, Jesse Thaler, Aashish Tripathee, and Wei Xue

Phys. Rev. Lett. **119**, 132003 — Published 26 September 2017

DOI: [10.1103/PhysRevLett.119.132003](https://doi.org/10.1103/PhysRevLett.119.132003)

# Exposing the QCD Splitting Function with CMS Open Data

Andrew Larkoski,<sup>1,\*</sup> Simone Marzani,<sup>2,†</sup> Jesse Thaler,<sup>3,‡</sup> Aashish Tripathy,<sup>3,§</sup> and Wei Xue<sup>3,¶</sup>

<sup>1</sup>*Physics Department, Reed College, Portland, OR 97202, USA*

<sup>2</sup>*University at Buffalo, The State University of New York, Buffalo, NY 14260-1500, USA*

<sup>3</sup>*Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA*

The splitting function is a universal property of quantum chromodynamics (QCD) which describes how energy is shared between partons. Despite its ubiquitous appearance in many QCD calculations, the splitting function cannot be measured directly since it always appears multiplied by a collinear singularity factor. Recently, however, a new jet substructure observable was introduced which asymptotes to the splitting function for sufficiently high jet energies. This provides a way to expose the splitting function through jet substructure measurements at the Large Hadron Collider. In this letter, we use public data released by the CMS experiment to study the 2-prong substructure of jets and test the  $1 \rightarrow 2$  splitting function of QCD. To our knowledge, this is the first ever physics analysis based on the CMS Open Data.

Quantum chromodynamics (QCD), like any weakly-coupled gauge theory, exhibits universal behavior in the small angle limit. When two partons become collinear in QCD, the cross section for a  $2 \rightarrow n$  scattering process factorizes into a  $2 \rightarrow n-1$  scattering cross section multiplied by a universal  $1 \rightarrow 2$  splitting probability, with corrections suppressed by the degree of collinearity. Collinear universality is a fundamental property of QCD and appears in many applications, most famously in deriving the DGLAP evolution equations [1–3] (see also [4–13]), and it is at the heart of the factorization theorem in hadron-hadron collisions [14, 15]. In addition, parton shower generators are based on recursively applying  $1 \rightarrow 2$  splittings [16–18], fixed-order subtraction schemes utilize the  $1 \rightarrow 2$  splitting function [19–21], and the  $k_t$  jet clustering metric is based on  $2 \rightarrow 1$  recombination [22–24]. Collinear universality can be extended to multi-parton splittings at tree level and beyond [25–41]; however its all-orders validity [42, 43] is spoiled in the presence of Glauber modes [44–47]. More recently, jet substructure techniques [48–52] have been introduced to distinguish  $1 \rightarrow n$  decays of heavy particles from  $1 \rightarrow n$  splittings in QCD in order to enhance the search for new physics at the Large Hadron Collider (LHC) [53–56].

Despite its ubiquity, however, the  $1 \rightarrow 2$  splitting function cannot be directly measured at a collider, since collinear universality is inseparable from the existence of collinear singularities and closely related non-perturbative fragmentation functions. Specifically, when two partons are separated by an angle  $\theta$ , the  $1 \rightarrow 2$  splitting probability takes the form

$$dP_{i \rightarrow jk} = \frac{d\theta}{\theta} dz P_{i \rightarrow jk}(z), \quad (1)$$

where the  $P_{i \rightarrow jk}$  are the Altarelli-Parisi QCD splitting functions [3] which depend on the momentum fraction  $z$  and the parton flavors  $i, j$ , and  $k$ . Crucially, this expression has a real emission singularity in the  $\theta \rightarrow 0$  limit, as required to cancel corresponding virtual singularities from loop diagrams. In this sense, there is no way to

directly measure the splitting function  $P_{i \rightarrow jk}(z)$  in data, though there is of course overwhelming indirect evidence that  $P_{i \rightarrow jk}(z)$  is a universal function from the many successes of QCD in describing high-energy scattering (see e.g. [57–67]).

In this letter, we present a semi-direct method to test the  $1 \rightarrow 2$  splitting function in QCD by studying the 2-prong substructure of jets. Our method is based on soft drop declustering [68] (see also [52, 69, 70]), which recursively removes soft radiation from a jet until hard 2-prong substructure is found. When applied to ordinary quark- and gluon-initiated jets with no intrinsic substructure, soft drop exposes the collinear core of the jet. As shown in ref. [71], the momentum sharing between the two prongs (denoted  $z_g$ ) is closely related to the momentum fraction  $z$  appearing in eq. (1), and the cross section for  $z_g$  asymptotes to the QCD splitting function in the high-energy limit. While variants of  $z_g$  have appeared in many jet substructure studies (notably the  $\sqrt{y}$  parameter in [52, 72]), to the best of our knowledge, no published  $z_g$  distribution has ever been presented using actual collider data, though there are preliminary  $z_g$  results from CMS [73], STAR [74], and ALICE [75]. Here, we present the first analysis of  $z_g$  using LHC data, taking advantage for the first time of public data released by the CMS experiment [76].

The CMS Open Data is derived from 7 TeV center-of-mass proton-proton collisions recorded in 2010 and released to the public on the CERN Open Data Portal in November 2014 [77]. The data is provided in AOD (Analysis Object Data) format, which is a CMS-specific data scheme based on the ROOT framework [78]. Crucially for the purposes of studying jet substructure, the AOD format contains all of the particle flow candidates (PFCs) [79, 80] used for jet finding within CMS [81], and we can apply jet substructure techniques directly on the PFCs themselves. The AOD files have an associated conditions database which include jet energy correction (JEC) factors and recommended jet quality cuts, though no specific calibration tools for jet substructure studies. The main




FIG. 1. Jet  $p_T$  spectrum from the CMS Open Data compared to three parton shower generators. Indicated is the  $p_T > 150$  GeV cut used in later analyses.

limitation of the 2010 CMS Open Data release is that it does not come accompanied by detector-simulated Monte Carlo samples, though this issue has been partially addressed in the 2011 CMS Open Data release [82]. Even without a detector simulation, we can improve the robustness of our analysis by using a charged-particle subset of PFCs with better angular resolution. Overall, this study highlights the fantastic performance of CMS’s particle flow algorithm and the exciting physics opportunities made possible by this public data release.

Our analysis is based on  $31.8 \text{ pb}^{-1}$  [83, 84] of data collected using the Jet Primary Dataset [76], which contains events selected by single-jet triggers, di-jet triggers, as well as some quad-jet and  $H_T$  triggers. We use the HLT\_Jet30U/50U/70U/100U/140U triggers for this analysis, which gives us near 100% efficiency to select single jets with transverse momentum  $p_T > 85$  GeV. All jets in our analysis are clustered using the anti- $k_t$  jet clustering algorithm [85] with radius parameter  $R = 0.5$ ; we validated that the anti- $k_t$  jets reported by CMS in the AOD format agree with those found by directly clustering the PFCs with FASTJET 3.1.3 [86]. To gain a more transparent understanding of the CMS data, we converted the AOD file format into our own text-based MIT Open Data (MOD) file format. Information about the MOD format as well as a broader suite of jet substructure analyses will be presented in a companion paper [87]. The substructure results shown here use the RecursiveTools 1.0.0 package from FASTJET CONTRIB 1.019 [88].

To validate initial jet reconstruction, Fig. 1 shows the  $p_T$  spectrum of the hardest jet in the event, with a pseudorapidity cut of  $|\eta| < 2.4$  and transverse momentum cut of  $p_T > 85$  GeV. This spectrum is obtained after applying the “loose” jet quality criteria provided by CMS as well as rescaling the jet  $p_T$  by the provided JEC factors.




FIG. 2. Schematic of the soft drop algorithm, which removes angular-ordered branches whose momentum fraction  $z$  is below  $z_{\text{cut}}\theta^\beta$ . The final groomed kinematics are indicated by the  $g$  subscript.

For comparison, we show the same spectrum obtained from three parton shower generators with their default settings: PYTHIA 8.219 [89], HERWIG 7.0.3 [90], and SHERPA 2.2.1 [91]. The qualitative agreement between all four samples is excellent. Note that this spectrum is obtained after combining five different CMS triggers with prescale factors that changed over the course of the 2010 run. No kinks are observed at the transitions between the various triggers, giving us confidence that we can derive jet spectra using the trigger and prescale values provided in the AOD files.

We now turn to an analysis of the 2-prong substructure of the hardest jet, imposing a further cut of  $p_T > 150$  GeV in order to avoid the large prescale factors present in the HLT\_Jet30U/50U triggers. To partially account for the finite energy resolution and efficiency of the CMS detector, we only consider PFCs within the hardest jet above  $p_T^{\text{min}} = 1$  GeV. Moreover, because charged particles have better angular resolution than neutral ones, our analysis will be only based on charged particles with associated tracks; we refer the reader to ref. [87] for substructure analyses with both charged and neutral PFCs. The charged PFCs are reclustered with the Cambridge/Aachen (C/A) algorithm [92, 93] to form an angular-ordered clustering tree. We then apply the soft drop declustering procedure [68] in Fig. 2, which recursively declusters the C/A tree, removing the softer  $p_T$  branch until 2-prong substructure is found which satisfies

$$z > z_{\text{cut}}\theta^\beta, \quad z \equiv \frac{\min[p_{T1}, p_{T2}]}{p_{T1} + p_{T2}}, \quad \theta = \frac{R_{12}}{R}. \quad (2)$$

Here,  $p_{T1}$  and  $p_{T2}$  are the transverse momenta of the two branches of the C/A tree, and  $R_{12} = \sqrt{(y_{12})^2 + (\phi_{12})^2}$  is their relative rapidity-azimuth distance. Throughout our analysis, we take the momentum fraction cut and angular exponent to be

$$z_{\text{cut}} = 0.1, \quad \beta = 0, \quad (3)$$

such that soft drop acts like the modified mass drop tagger (mMDT) [69] with  $\mu = 1$ . The values of  $z$  and  $\theta$  obtained after soft drop are referred to as  $z_g$  and  $\theta_g$ ,




FIG. 3. Track jet mass spectra before and after the soft drop procedure with  $\beta = 0$  (i.e. mMDT with  $\mu = 1$ ), comparing the CMS Open Data to PYTHIA.

where the  $g$  subscript is a reminder that these values were obtained after jet grooming. These two observables encode information about the two non-trivial kinematic variables in the unpolarized  $1 \rightarrow 2$  QCD splitting function from eq. (1). Note that  $z_g$  is a ratio of  $p_T$  scales, so not affected by the JEC factor applied to the jet  $p_T$  as a whole. Similarly, as a dimensionless quantity,  $z_g$  is relatively insensitive to the absolute energy scale of the PFCs, and is only mildly affected by the  $p_T^{\min} = 1$  GeV restriction.

The key observable used in jet substructure analyses at ATLAS and CMS is the jet invariant mass [94–96]. The track-only jet mass spectrum before and after soft drop is shown in Fig. 3 and compared to predictions from PYTHIA. There is reasonable qualitative agreement between the CMS Open Data and PYTHIA for  $m > 10$  GeV; below 10 GeV one expects deviations from the finite detector resolution of CMS and the fact that the PFCs do not include full hadron mass information. We emphasize that no additional corrections have been applied to the CMS Open Data, apart from the JEC factor needed to impose the  $p_T > 150$  GeV criteria and the  $p_T^{\min} = 1$  GeV PFC restriction needed to account for finite energy resolution and efficiency. Similarly, we are showing particle-level predictions from PYTHIA using the default tune with no detector simulation (but the same restriction to charged hadrons with  $p_T^{\min} = 1$  GeV). Because we do not have access to detector-simulated Monte Carlo samples, and because there is insufficient information in the AOD format to estimate systematic uncertainties, the error bars shown only include statistical uncertainties.

To see the 2-prong structure revealed by soft drop, Fig. 4 shows the double-differential track  $(z_g, \theta_g)$  spectrum seen in the CMS Open Data. The peak towards small values of  $z_g$  and  $\theta_g$  reflects the double-logarithmic




FIG. 4. Double-differential distribution of track  $z_g$  versus  $\theta_g$  in the CMS Open Data, i.e. the dimensionless probability density  $p(z_g, \theta_g)$  whose integral is 1.

structure in eq. (1), since soft gluon emission from a hard quark or gluon is approximated by

$$dP_{i \rightarrow ig} \simeq \frac{2\alpha_s C_i}{\pi} \frac{d\theta}{\theta} \frac{dz}{z}, \quad (4)$$

where  $\alpha_s$  is the strong coupling constant and  $C_i$  is the Casimir factor (4/3 for quarks, 3 for gluons). The  $z_g$  distribution is cut off by  $z_{\text{cut}}$ , which regulates the soft singularity of QCD. In principle, the  $\theta_g$  distribution could extend all the way to zero, but it is cut off both by the angular resolution of the CMS detector and by non-perturbative QCD effects which are relevant for  $\theta_g \simeq \Lambda_{\text{QCD}}/(z_{\text{cut}} p_T R) \simeq 10^{-1}$ . In addition, the perturbative  $\theta_g \rightarrow 0$  singularity in eq. (1) is regulated by a single-logarithmic form factor [68], which we now exploit to perform analytic calculations of the  $z_g$  distribution.

In perturbative QCD,  $z_g$  with  $\beta = 0$  is a collinear-unsafe observable and therefore not calculable order-by-order in an expansion in the strong coupling constant  $\alpha_s$ . In particular,  $z_g$  is ambiguous for a jet containing a single parton, and therefore real emission singularities associated with 2 partons (where  $z_g$  is well-defined) cannot cancel against virtual emission singularities associated with 1 parton (where  $z_g$  is ill-defined). That said, we can follow the strategy outlined in refs. [71, 97] and express the normalized  $z_g$  probability distribution  $p(z_g)$  as

$$p(z_g) = \int d\theta_g p(\theta_g) p(z_g|\theta_g), \quad (5)$$

where  $p(\theta_g)$  is the probability distribution for  $\theta_g$ , and  $p(z_g|\theta_g)$  is the conditional probability distribution for  $z_g$  given a fixed value of  $\theta_g$ . While  $z_g$  is collinear unsafe, the conditional probability distribution  $p(z_g|\theta_g)$  is calculable as a perturbative expansion, since any finite value

of  $\theta_g$  will remove the 1 parton region of phase space. By resumming the  $p(\theta_g)$  distribution to all orders in  $\alpha_s$ , the  $\theta_g \rightarrow 0$  limit is regulated, and the integral in eq. (5) yields a finite distribution for  $p(z_g)$ . In this way,  $z_g$  is a collinear unsafe but “Sudakov safe” observable [97].

Remarkably, to lowest non-trivial order, the probability distribution for  $p(z_g)$  can be directly expressed in terms of the QCD splitting function as [71]

$$p(z_g) = \sum_i f_i p_i(z_g), \quad (6)$$

where  $f_i$  is the fraction of the event sample composed of jets initiated by partons of flavor  $i$  (i.e. quarks or gluons), and

$$p_i(z) = \frac{\bar{P}_i(z)}{\int_{z_{\text{cut}}}^{1/2} dz' \bar{P}_i(z')} \Theta(z > z_{\text{cut}}) + \mathcal{O}(\alpha_s), \quad (7)$$

where

$$\bar{P}_i(z) = \sum_{j,k} [P_{i \rightarrow jk}(z) + P_{i \rightarrow jk}(1-z)]. \quad (8)$$

The  $z_g$  distribution is a flavor-averaged,  $z$ -symmetrized,  $z_{\text{cut}}$ -truncated, and normalized version of the QCD splitting function. Due to a supersymmetric relationship between the quark and gluon splitting functions [98, 99],  $\bar{P}_i$  is the same for quarks and gluons to an excellent approximation, such that

$$p(z_g) \simeq \frac{2 \frac{z_g}{1-z_g} + 2 \frac{1-z_g}{z_g} + 1}{\frac{3}{2} (2z_{\text{cut}} - 1) + 2 \log \frac{1-z_{\text{cut}}}{z_{\text{cut}}}}, \quad (9)$$

and the probability distribution for  $z_g$  is independent of  $\alpha_s$  at leading order. In this way, measuring  $z_g$  exposes the QCD splitting function. The predicted  $z_g$  distribution can be refined by performing higher-order calculations. As in ref. [71], we calculate  $p(\theta_g)$  to modified leading-logarithmic (MLL) accuracy, which includes running coupling effects and subleading terms in the splitting functions. We also calculate  $p(z_g|\theta_g)$  to leading fixed order in the collinear approximation and obtain an analytic prediction for  $p(z_g)$  using eq. (5). While not shown below, the theoretical uncertainties on  $p(z_g)$  can be estimated by varying the different renormalization scales that enter the calculation [87].

In Fig. 5, we show the  $z_g$  distribution for our jet selection, comparing the analytic expression in eq. (5) (which extends eq. (9) to MLL accuracy), three parton shower generators, and the CMS Open Data. Strictly speaking, the theoretical calculation described above should be modified [100, 101] to account for the fact that the current analysis is based only on charged particles; for this reason, we show  $p(z_g)$  without its uncertainty band to emphasize its qualitative nature. Notwithstanding the above, the CMS Open Data agrees very well with the




FIG. 5. Distribution of  $z_g$  from mMDT/soft drop. The theory distribution is from an all-particle prediction, yet agrees very well with the track-based distributions.

theory calculation as well as with the Monte Carlo parton showers, and the characteristic  $1/z$  behavior expected from the QCD splitting function is seen in all distributions. The one point where there is a noticeable (but expected) difference between the open data and the parton showers is at  $z_g = 0$ , which corresponds to jets that have only one constituent after soft drop. Because close-by particles can be reconstructed as a single PFC due to finite angular resolution, the CMS Open Data is expected to have more “one particle” jets than the parton shower generators. We have evidence that the small difference between the parton showers and the theory distribution at  $z_g \simeq z_{\text{cut}}$  is due to growing logarithms of  $z_g$  that are not resummed in our MLL approach. We verified that these discrepancies are suppressed for  $z_{\text{cut}} = 0.2$  and enhanced for  $z_{\text{cut}} = 0.05$ , consistent with this expectation.

The CMS Open Data represents a new chapter in particle physics, since for the first time, high-quality collider data has been released to scientists not affiliated with an experimental collaboration. In this paper, we applied state-of-the-art jet substructure techniques on the CMS Open Data and exposed the QCD splitting function, which encodes the universal behavior of gauge theories in the collinear limit. This was only possible because of theoretical advances on Sudakov safe observables, which allowed us to predict the  $z_g$  distribution from first principles, and the fantastic experimental performance of the CMS detector, which allowed us to perform a detailed study of the substructure of jets. We hope this letter inspires scientists outside of the LHC collaborations to incorporate CMS Open Data into their research and motivates the LHC collaborations to continue their support of open data initiatives.

We applaud CERN for the historic launch of the Open Data Portal, and we congratulate the CMS collaboration for the fantastic performance of their detector and the high quality of the resulting public data set. We thank Alexis Romero for collaboration in the early stages of this work. We are indebted to Salvatore Rappoccio and Kati Lassila-Perini for helping us navigate the CMS software framework. We benefitted from code and encouragement from Tim Andeen, Matt Bellis, Andy Buckley, Kyle Cranmer, Sarah Demers, Guenther Dissertori, Javier Duarte, Peter Fisher, Achim Geiser, Giacomo Govi, Phil Harris, Beate Heinemann, Harri Hirvonsalo, Markus Klute, Greg Landsberg, Yen-Jie Lee, Elliot Lipeles, Peter Loch, Marcello Maggi, David Miller, Ben Nachman, Christoph Paus, Alexx Perloff, Andreas Pfeiffer, Maurizio Pierini, Ana Rodriguez, Gunther Roland, Ariel Schwartzman, Liz Sexton-Kennedy, Maria Spiropulu, Nhan Tran, Ana Trisovic, Chris Tully, Marta Verweij, Mikko Voutilainen, and Mike Williams. This work is supported by the MIT Charles E. Reed Faculty Initiatives Fund. The work of JT, AT, and WX is supported by the U.S. Department of Energy (DOE) under grant contract numbers DE-SC-00012567 and DE-SC-00015476. The work of AL was supported by the U.S. National Science Foundation, under grant PHY-1419008, the LHC Theory Initiative. SM is supported by the U.S. National Science Foundation, under grants PHY-0969510 (LHC Theory Initiative) and PHY-1619867. AT is also supported by the MIT Undergraduate Research Opportunities Program.

---

\* larkoski@reed.edu

† smarzani@buffalo.edu

‡ jthaler@mit.edu

§ aashisht@mit.edu

¶ weixue@mit.edu

- [1] V.N. Gribov and L.N. Lipatov, “Deep inelastic  $e p$  scattering in perturbation theory,” *Sov.J.Nucl.Phys.* **15**, 438–450 (1972).
- [2] Yuri L. Dokshitzer, “Calculation of the Structure Functions for Deep Inelastic Scattering and  $e^+ e^-$  Annihilation by Perturbation Theory in Quantum Chromodynamics,” *Sov.Phys.JETP* **46**, 641–653 (1977).
- [3] Guido Altarelli and G. Parisi, “Asymptotic Freedom in Parton Language,” *Nucl.Phys.* **B126**, 298 (1977).
- [4] E. G. Floratos, D. A. Ross, and Christopher T. Sachrajda, “Higher Order Effects in Asymptotically Free Gauge Theories: The Anomalous Dimensions of Wilson Operators,” *Nucl. Phys.* **B129**, 66–88 (1977), [Erratum: *Nucl. Phys.* **B139**, 545 (1978)].
- [5] E. G. Floratos, D. A. Ross, and Christopher T. Sachrajda, “Higher Order Effects in Asymptotically Free Gauge Theories. 2. Flavor Singlet Wilson Operators and Coefficient Functions,” *Nucl. Phys.* **B152**, 493–520 (1979).
- [6] Antonio Gonzalez-Arroyo, C. Lopez, and F. J. Yndurain, “Second Order Contributions to the Structure Functions in Deep Inelastic Scattering. 1. Theoretical Calculations,” *Nucl. Phys.* **B153**, 161–186 (1979).
- [7] Antonio Gonzalez-Arroyo and C. Lopez, “Second Order Contributions to the Structure Functions in Deep Inelastic Scattering. 3. The Singlet Case,” *Nucl. Phys.* **B166**, 429–459 (1980).
- [8] G. Curci, W. Furmanski, and R. Petronzio, “Evolution of Parton Densities Beyond Leading Order: The Nonsinglet Case,” *Nucl. Phys.* **B175**, 27–92 (1980).
- [9] W. Furmanski and R. Petronzio, “Singlet Parton Densities Beyond Leading Order,” *Phys. Lett.* **B97**, 437–442 (1980).
- [10] E. G. Floratos, C. Kounnas, and R. Lacaze, “Higher Order QCD Effects in Inclusive Annihilation and Deep Inelastic Scattering,” *Nucl. Phys.* **B192**, 417–462 (1981).
- [11] R. Hamberg and W. L. van Neerven, “The Correct renormalization of the gluon operator in a covariant gauge,” *Nucl. Phys.* **B379**, 143–171 (1992).
- [12] A. Vogt, S. Moch, and J. A. M. Vermaseren, “The Three-loop splitting functions in QCD: The Singlet case,” *Nucl. Phys.* **B691**, 129–181 (2004), arXiv:hep-ph/0404111 [hep-ph].
- [13] S. Moch, J. A. M. Vermaseren, and A. Vogt, “The Three loop splitting functions in QCD: The Nonsinglet case,” *Nucl. Phys.* **B688**, 101–134 (2004), arXiv:hep-ph/0403192 [hep-ph].
- [14] John C. Collins, Davison E. Soper, and George F. Sterman, “All Order Factorization for Drell-Yan Cross-sections,” *Phys. Lett.* **B134**, 263 (1984).
- [15] John C. Collins, Davison E. Soper, and George F. Sterman, “Factorization for Short Distance Hadron - Hadron Scattering,” *Nucl. Phys.* **B261**, 104–142 (1985).
- [16] P. Mazzanti and R. Odorico, “A Monte Carlo Program for QCD Event Simulation in  $e^+e^-$  Annihilation at LEP Energies,” *Z. Phys.* **C7**, 61 (1980).
- [17] Torbjorn Sjostrand, “The Lund Monte Carlo for Jet Fragmentation,” *Comput. Phys. Commun.* **27**, 243 (1982).
- [18] G. Marchesini and B. R. Webber, “Simulation of QCD Jets Including Soft Gluon Interference,” *Nucl. Phys.* **B238**, 1–29 (1984).
- [19] R. Keith Ellis, D. A. Ross, and A. E. Terrano, “The Perturbative Calculation of Jet Structure in  $e^+ e^-$  Annihilation,” *Nucl. Phys.* **B178**, 421 (1981).
- [20] K. Fabricius, I. Schmitt, G. Kramer, and G. Schierholz, “Higher Order Perturbative QCD Calculation of Jet Cross-Sections in  $e^+ e^-$  Annihilation,” *Z. Phys.* **C11**, 315 (1981).
- [21] S. Catani and M. H. Seymour, “A general algorithm for calculating jet cross sections in NLO QCD,” *Nucl. Phys.* **B485**, 291–419 (1997), arXiv:hep-ph/9605323.
- [22] S. Catani, Yuri L. Dokshitzer, M. Olsson, G. Turnock, and B. R. Webber, “New clustering algorithm for multi-jet cross-sections in  $e^+ e^-$  annihilation,” *Phys. Lett.* **B269**, 432–438 (1991).
- [23] S. Catani, Yuri L. Dokshitzer, M.H. Seymour, and B.R. Webber, “Longitudinally invariant  $K_t$  clustering algorithms for hadron hadron collisions,” *Nucl.Phys.* **B406**, 187–224 (1993).
- [24] Stephen D. Ellis and Davison E. Soper, “Successive combination jet algorithm for hadron collisions,” *Phys.Rev.* **D48**, 3160–3166 (1993), arXiv:hep-ph/9305266 [hep-ph].

- [25] Stefano Catani and Massimiliano Grazzini, “Collinear factorization and splitting functions for next-to-next-to-leading order QCD calculations,” *Phys. Lett.* **B446**, 143–152 (1999), arXiv:hep-ph/9810389 [hep-ph].
- [26] Stefano Catani and Massimiliano Grazzini, “Infrared factorization of tree level QCD amplitudes at the next-to-next-to-leading order and beyond,” *Nucl. Phys.* **B570**, 287–325 (2000), arXiv:hep-ph/9908523 [hep-ph].
- [27] Zvi Bern, Vittorio Del Duca, and Carl R. Schmidt, “The Infrared behavior of one loop gluon amplitudes at next-to-next-to-leading order,” *Phys. Lett.* **B445**, 168–177 (1998), arXiv:hep-ph/9810409 [hep-ph].
- [28] Zvi Bern, Vittorio Del Duca, William B. Kilgore, and Carl R. Schmidt, “The infrared behavior of one loop QCD amplitudes at next-to-next-to leading order,” *Phys. Rev.* **D60**, 116001 (1999), arXiv:hep-ph/9903516 [hep-ph].
- [29] S. D. Badger and E. W. Nigel Glover, “Two loop splitting functions in QCD,” *JHEP* **07**, 040 (2004), arXiv:hep-ph/0405236 [hep-ph].
- [30] Frits A. Berends and W. T. Giele, “Recursive Calculations for Processes with  $n$  Gluons,” *Nucl. Phys.* **B306**, 759–808 (1988).
- [31] Michelangelo L. Mangano and Stephen J. Parke, “Multiparton amplitudes in gauge theories,” *Phys. Rept.* **200**, 301–367 (1991), arXiv:hep-th/0509223 [hep-th].
- [32] John M. Campbell and E. W. Nigel Glover, “Double unresolved approximations to multiparton scattering amplitudes,” *Nucl. Phys.* **B527**, 264–288 (1998), arXiv:hep-ph/9710255 [hep-ph].
- [33] Vittorio Del Duca, Alberto Frizzo, and Fabio Maltoni, “Factorization of tree QCD amplitudes in the high-energy limit and in the collinear limit,” *Nucl. Phys.* **B568**, 211–262 (2000), arXiv:hep-ph/9909464 [hep-ph].
- [34] T. G. Birthwright, E. W. Nigel Glover, V. V. Khoze, and P. Marquard, “Multi-gluon collinear limits from MHV diagrams,” *JHEP* **05**, 013 (2005), arXiv:hep-ph/0503063 [hep-ph].
- [35] T. G. Birthwright, E. W. Nigel Glover, V. V. Khoze, and P. Marquard, “Collinear limits in QCD from MHV rules,” *JHEP* **07**, 068 (2005), arXiv:hep-ph/0505219 [hep-ph].
- [36] Zvi Bern, Gordon Chalmers, Lance J. Dixon, and David A. Kosower, “One loop  $N$  gluon amplitudes with maximal helicity violation via collinear limits,” *Phys. Rev. Lett.* **72**, 2134–2137 (1994), arXiv:hep-ph/9312333 [hep-ph].
- [37] Zvi Bern, Lance J. Dixon, David C. Dunbar, and David A. Kosower, “One loop  $n$  point gauge theory amplitudes, unitarity and collinear limits,” *Nucl. Phys.* **B425**, 217–260 (1994), arXiv:hep-ph/9403226 [hep-ph].
- [38] Zvi Bern and Gordon Chalmers, “Factorization in one loop gauge theory,” *Nucl. Phys.* **B447**, 465–518 (1995), arXiv:hep-ph/9503236 [hep-ph].
- [39] David A. Kosower and Peter Uwer, “One loop splitting amplitudes in gauge theory,” *Nucl. Phys.* **B563**, 477–505 (1999), arXiv:hep-ph/9903515 [hep-ph].
- [40] Stefano Catani, Daniel de Florian, and German Rodrigo, “The Triple collinear limit of one loop QCD amplitudes,” *Phys. Lett.* **B586**, 323–331 (2004), arXiv:hep-ph/0312067 [hep-ph].
- [41] Zvi Bern, Lance J. Dixon, and David A. Kosower, “Two-loop  $g \rightarrow \bar{g}g$  splitting amplitudes in QCD,” *JHEP* **08**, 012 (2004), arXiv:hep-ph/0404293 [hep-ph].
- [42] David A. Kosower, “All order collinear behavior in gauge theories,” *Nucl. Phys.* **B552**, 319–336 (1999), arXiv:hep-ph/9901201 [hep-ph].
- [43] Ilya Feige and Matthew D. Schwartz, “Hard-Soft-Collinear Factorization to All Orders,” *Phys. Rev.* **D90**, 105020 (2014), arXiv:1403.6472 [hep-ph].
- [44] Stefano Catani, Daniel de Florian, and German Rodrigo, “Space-like (versus time-like) collinear limits in QCD: Is factorization violated?” *JHEP* **07**, 026 (2012), arXiv:1112.4405 [hep-ph].
- [45] Jeffrey R. Forshaw, Michael H. Seymour, and Andrzej Siodmok, “On the Breaking of Collinear Factorization in QCD,” *JHEP* **11**, 066 (2012), arXiv:1206.6363 [hep-ph].
- [46] Ira Z. Rothstein and Iain W. Stewart, “An Effective Field Theory for Forward Scattering and Factorization Violation,” *JHEP* **08**, 025 (2016), arXiv:1601.04695 [hep-ph].
- [47] Matthew D. Schwartz, Kai Yan, and Hua Xing Zhu, “Collinear factorization violation and effective field theory,” (2017), arXiv:1703.08572 [hep-ph].
- [48] M. H. Seymour, “Tagging a heavy Higgs boson,” in *ECFA Large Hadron Collider (LHC) Workshop: Physics and Instrumentation Aachen, Germany, October 4-9, 1990* (1991).
- [49] Michael H. Seymour, “Searches for new particles using cone and cluster jet algorithms: A Comparative study,” *Z. Phys.* **C62**, 127–138 (1994).
- [50] J. M. Butterworth, B. E. Cox, and Jeffrey R. Forshaw, “ $WW$  scattering at the CERN LHC,” *Phys. Rev.* **D65**, 096014 (2002), arXiv:hep-ph/0201098 [hep-ph].
- [51] J. M. Butterworth, John R. Ellis, and A. R. Raklev, “Reconstructing sparticle mass spectra using hadronic decays,” *JHEP* **05**, 033 (2007), arXiv:hep-ph/0702150 [HEP-PH].
- [52] Jonathan M. Butterworth, Adam R. Davison, Mathieu Rubin, and Gavin P. Salam, “Jet substructure as a new Higgs search channel at the LHC,” *Phys. Rev. Lett.* **100**, 242001 (2008), arXiv:0802.2470 [hep-ph].
- [53] A. Abdesselam, E. Bergeaas Kuutmann, U. Bitenc, G. Brooijmans, J. Butterworth, *et al.*, “Boosted objects: A Probe of beyond the Standard Model physics,” *Eur.Phys.J.* **C71**, 1661 (2011), arXiv:1012.5412 [hep-ph].
- [54] A. Altheimer, S. Arora, L. Asquith, G. Brooijmans, J. Butterworth, *et al.*, “Jet Substructure at the Tevatron and LHC: New results, new tools, new benchmarks,” *J.Phys.* **G39**, 063001 (2012), arXiv:1201.0008 [hep-ph].
- [55] A. Altheimer *et al.*, “Boosted objects and jet substructure at the LHC. Report of BOOST2012, held at IFIC Valencia, 23rd-27th of July 2012,” *BOOST 2012 Valencia, Spain, July 23-27, 2012*, *Eur. Phys. J.* **C74**, 2792 (2014), arXiv:1311.2708 [hep-ex].
- [56] D. Adams *et al.*, “Towards an Understanding of the Correlations in Jet Substructure,” *Eur. Phys. J.* **C75**, 409 (2015), arXiv:1504.00679 [hep-ph].
- [57] R. Brandelik *et al.* (TASSO), “Evidence for Planar Events in  $e^+e^-$  Annihilation at High-Energies,” *Phys. Lett.* **B86**, 243–249 (1979).
- [58] D. P. Barber *et al.*, “Discovery of Three Jet Events and a Test of Quantum Chromodynamics at PETRA Energies,” *Phys. Rev. Lett.* **43**, 830 (1979).
- [59] Christoph Berger *et al.* (PLUTO), “Evidence for Gluon Bremsstrahlung in  $e^+e^-$  Annihilations at High-



- Energies,” Phys. Lett. **B86**, 418–425 (1979).
- [60] W. Bartel *et al.* (JADE), “Observation of Planar Three Jet Events in  $e^+e^-$  Annihilation and Evidence for Gluon Bremsstrahlung,” Phys. Lett. **B91**, 142–147 (1980).
- [61] P. Abreu *et al.* (DELPHI), “Measurement of the gluon fragmentation function and a comparison of the scaling violation in gluon and quark jets,” Eur. Phys. J. **C13**, 573–589 (2000).
- [62] M. Z. Akrawy *et al.* (OPAL), “A Study of coherence of soft gluons in hadron jets,” Phys. Lett. **B247**, 617–628 (1990).
- [63] G. Abbiendi *et al.* (OPAL), “A Simultaneous measurement of the QCD color factors and the strong coupling,” Eur. Phys. J. **C20**, 601–615 (2001), arXiv:hep-ex/0101044 [hep-ex].
- [64] A. Heister *et al.* (ALEPH), “Studies of QCD at  $e^+e^-$  centre-of-mass energies between 91-GeV and 209-GeV,” Eur. Phys. J. **C35**, 457–486 (2004).
- [65] J. Abdallah *et al.* (DELPHI), “A study of the energy evolution of event shape distributions and their means with the DELPHI detector at LEP,” Eur. Phys. J. **C29**, 285–312 (2003), arXiv:hep-ex/0307048.
- [66] G. Abbiendi *et al.* (OPAL), “Measurement of event shape distributions and moments in  $e^+e^- \rightarrow j$  hadrons at 91-GeV - 209-GeV and a determination of  $\alpha(s)$ ,” Eur. Phys. J. **C40**, 287–316 (2005), arXiv:hep-ex/0503051.
- [67] P. Achard *et al.* (L3), “Studies of hadronic event structure in  $e^+e^-$  annihilation from 30-GeV to 209-GeV with the L3 detector,” Phys. Rept. **399**, 71–174 (2004), arXiv:hep-ex/0406049.
- [68] Andrew J. Larkoski, Simone Marzani, Gregory Soyez, and Jesse Thaler, “Soft Drop,” JHEP **1405**, 146 (2014), arXiv:1402.2657 [hep-ph].
- [69] Mrinal Dasgupta, Alessandro Fregoso, Simone Marzani, and Gavin P. Salam, “Towards an understanding of jet substructure,” JHEP **09**, 029 (2013), arXiv:1307.0007 [hep-ph].
- [70] Mrinal Dasgupta, Alessandro Fregoso, Simone Marzani, and Alexander Powling, “Jet substructure with analytical methods,” Eur. Phys. J. **C73**, 2623 (2013), arXiv:1307.0013 [hep-ph].
- [71] Andrew J. Larkoski, Simone Marzani, and Jesse Thaler, “Sudakov Safety in Perturbative QCD,” Phys. Rev. **D91**, 111501 (2015), arXiv:1502.01719 [hep-ph].
- [72] Georges Aad *et al.* (ATLAS), “Search for high-mass diboson resonances with boson-tagged jets in proton-proton collisions at  $\sqrt{s} = 8$  TeV with the ATLAS detector,” JHEP **12**, 055 (2015), arXiv:1506.00962 [hep-ex].
- [73] *Splitting function in pp and PbPb collisions at 5.02 TeV*, Tech. Rep. CMS-PAS-HIN-16-006 (2016).
- [74] Kolja Kauder (STAR collaboration), “Measurement of the shared momentum fraction  $z_g$  using jet reconstruction in p+p and au+au collisions with star,” Conference talk at Hard Probes 2016.
- [75] Kirill Lapidus (ALICE collaboration), “Hard substructure of jets probed in p-pb collisions,” Poster at Quark Matter 2017.
- [76] “Jet primary dataset in AOD format from RunB of 2010 (/Jet/Run2010B-Apr21ReReco-v1/AOD),” CMS Collaboration, CERN Open Data Portal (), 10.7483/OPENDATA.CMS.3S7F.2E9W.
- [77] “CERN Open Data Portal,” <http://opendata.cern.ch>.
- [78] R. Brun and F. Rademakers, “ROOT: An object oriented data analysis framework,” *New computing techniques in physics research V. Proceedings, 5th International Workshop, AIHENP '96, Lausanne, Switzerland, September 2-6, 1996*, Nucl. Instrum. Meth. **A389**, 81–86 (1997).
- [79] *Particle-Flow Event Reconstruction in CMS and Performance for Jets, Taus, and MET*, Tech. Rep. CMS-PAS-PFT-09-001 (CERN, 2009. Geneva, 2009).
- [80] *Commissioning of the Particle-flow Event Reconstruction with the first LHC collisions recorded in the CMS detector*, Tech. Rep. CMS-PAS-PFT-10-001 (2010).
- [81] Vardan Khachatryan *et al.* (CMS), “Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV,” JINST **12**, P02014 (2017), arXiv:1607.03663 [hep-ex].
- [82] “Jet primary dataset in AOD format from RunA of 2011 (/Jet/Run2011A-12Oct2013-v1/AOD),” CMS Collaboration, CERN Open Data Portal (), 10.7483/OPENDATA.CMS.UP77.P6PQ.
- [83] Michele De Gruttola, *The  $Z \rightarrow \mu^+\mu^-$  decay channel in the CMS experiment at LHC: from cross-section measurement with the 2010 7 TeV collision dataset to offline machine luminosity monitor*, Ph.D. thesis, Naples U. (2010).
- [84] *Measurement of CMS Luminosity*, Tech. Rep. CMS-PAS-EWK-10-004 (CERN, 1900. Geneva, 1900).
- [85] Matteo Cacciari, Gavin P. Salam, and Gregory Soyez, “The Anti-k(t) jet clustering algorithm,” JHEP **0804**, 063 (2008), arXiv:0802.1189 [hep-ph].
- [86] Matteo Cacciari, Gavin P. Salam, and Gregory Soyez, “FastJet User Manual,” Eur.Phys.J. **C72**, 1896 (2012), arXiv:1111.6097 [hep-ph].
- [87] Aashish Tripathy, Wei Xue, Andrew Larkoski, Simone Marzani, and Jesse Thaler, “Jet Substructure Studies with CMS Open Data,” (2017), arXiv:1704.05842 [hep-ph].
- [88] “Fastjet contrib,” <http://fastjet.hepforge.org/contrib/>.
- [89] Torbjorn Sjostrand, Stephen Mrenna, and Peter Skands, “A Brief Introduction to PYTHIA 8.1,” Comput.Phys.Commun.178:852-867,2008 (2007), 10.1016/j.cpc.2008.01.036, arXiv:0710.3820v1 [hep-ph].
- [90] Johannes Bellm *et al.*, “Herwig 7.0/Herwig++ 3.0 release note,” Eur. Phys. J. **C76**, 196 (2016), arXiv:1512.01178 [hep-ph].
- [91] T. Gleisberg, Stefan. Hoeche, F. Krauss, M. Schonherr, S. Schumann, F. Siegert, and J. Winter, “Event generation with SHERPA 1.1,” JHEP **02**, 007 (2009), arXiv:0811.4622 [hep-ph].
- [92] M. Wobisch and T. Wengler, “Hadronization corrections to jet cross-sections in deep inelastic scattering,” (1998), arXiv:hep-ph/9907280 [hep-ph].
- [93] Yuri L. Dokshitzer, G.D. Leder, S. Moretti, and B.R. Webber, “Better jet clustering algorithms,” JHEP **9708**, 001 (1997), arXiv:hep-ph/9707323 [hep-ph].
- [94] Georges Aad *et al.* (ATLAS Collaboration), “Jet mass and substructure of inclusive jets in  $\sqrt{s} = 7$  TeV pp collisions with the ATLAS experiment,” JHEP **1205**, 128 (2012), arXiv:1203.4606 [hep-ex].
- [95] Serguei Chatrchyan *et al.* (CMS Collaboration), “Studies of jet mass in dijet and W/Z + jet events,” JHEP **1305**, 090 (2013), arXiv:1303.4811 [hep-ex].
- [96] T. Aaltonen *et al.* (CDF), “Study of Substructure of High Transverse Momentum Jets Produced in Proton-



- Antiproton Collisions at  $\sqrt{s} = 1.96$  TeV,” Phys. Rev. **D85**, 091101 (2012), arXiv:1106.5952 [hep-ex].
- [97] Andrew J. Larkoski and Jesse Thaler, “Unsafe but Calculable: Ratios of Angularities in Perturbative QCD,” JHEP **1309**, 137 (2013), arXiv:1307.1699 [hep-ph].
  - [98] Yuri L. Dokshitzer, Valery A. Khoze, Alfred H. Mueller, and S. I. Troian, *Basics of perturbative QCD* (1991).
  - [99] M.H. Seymour, “Jet shapes in hadron collisions: Higher orders, resummation and hadronization,” Nucl.Phys. **B513**, 269–300 (1998), arXiv:hep-ph/9707338 [hep-ph].
  - [100] Hsi-Ming Chang, Massimiliano Procura, Jesse Thaler, and Wouter J. Waalewijn, “Calculating Track-Based Observables for the LHC,” Phys. Rev. Lett. **111**, 102002 (2013), arXiv:1303.6637 [hep-ph].
  - [101] Hsi-Ming Chang, Massimiliano Procura, Jesse Thaler, and Wouter J. Waalewijn, “Calculating Track Thrust with Track Functions,” Phys. Rev. **D88**, 034030 (2013), arXiv:1306.6630 [hep-ph].