

This is the accepted manuscript made available via CHORUS. The article has been published as:

Exact Solutions for Nonlinear Development of a Kelvin-Helmholtz Instability for the Counterflow of Superfluid and Normal Components of Helium II

Pavel M. Lushnikov and Nikolay M. Zubarev

Phys. Rev. Lett. **120**, 204504 — Published 17 May 2018

DOI: [10.1103/PhysRevLett.120.204504](https://doi.org/10.1103/PhysRevLett.120.204504)

Exact solutions for nonlinear development of Kelvin-Helmholtz instability for counterflow of superfluid and normal components of Helium II

Pavel M. Lushnikov^{1,2,*} and Nikolay M. Zubarev^{3,4,†}

¹*Department on Mathematics and Statistics, University of New Mexico, New Mexico 87131, USA*

²*Landau Institute for Theoretical Physics, 2 Kosygin St., Moscow 119334, Russia*

³*Institute for Electrophysics, Ural Branch, Russian Academy of Sciences, Yekaterinburg, 620016 Russia*

⁴*Lebedev Physical Institute, Russian Academy of Sciences, Moscow, 119991 Russia*

(Dated: March 13, 2018)

A relative motion of the normal and superfluid components of Helium II results in quantum Kelvin-Helmholtz instability (KHI) at their common free surface. We found the integrability and exact growing solutions for the nonlinear stage of the development of that instability. Contrary to the usual KHI of the interface between two classical fluids, the dynamics of Helium II free surface allows reduction to the Laplace growth equation which has an infinite number of exact solutions including the generic formation of sharp cusps at the free surface in a finite time.




FIG. 1: (Color online) A schematic of counterflow in superfluid ^4He . Heater results in the flux of heat \mathbf{Q} which is carried by the normal fluid component with velocity \mathbf{v}_n while superfluid component moves in the opposite direction with the velocity \mathbf{v}_s . Both components coexist in the same volume of fluid and share the same free surface.

Kelvin-Helmholtz instability (KHI) is perhaps the most important hydrodynamic instability which commonly occurs either at the interface between two fluids moving with different velocities or in the presence of the tangential velocity jump/shear flow in the same fluid [1]. Recently KHI attracted significant experimental and theoretical attention in superfluids. KHI was studied either for interface between different phases of ^3He [2–6], which has many similarities with KHI in classical fluids, or KHI from relative motion of components of ^4He [7–11] which has no classical analog thus we refer to it as quantum KHI. We focus on the second case, i.e. on quantum KHI of the free surface of ^4He in the superfluid phase (He-II state) in the presence of counterflow of superfluid and normal fluid components [12, 13]. Principle difference here from KHI of classical fluids is that relative fluid motion in quantum KHI occurs not from different sides of interface but from the same side of the He-II free surface with fluids components coexisting in the same volume which is purely quantum effect. A counterflow is achieved in experiment by the action of a stationary heat flow within the liquid in the direction tangent to the free surface as shown in Fig. 1.

Linear analysis of both classical KHI and quantum KHI results in the exponential growth of surface perturbations [1, 12, 13]. As these initial perturbations reach amplitudes comparable with their wavelength, nonlinear

effects must be considered. Weak nonlinearity approximation takes into account the leading order nonlinear correction over the small parameter which is the typical slope of surface. Weakly nonlinear equations for development of KHI of classical fluids results in a finite time singularity [14] which means that solutions become strongly nonlinear beyond the perturbation theory. Two-dimensional (2D) dynamics of interface between two fluids in weak nonlinearity approximation can be reduced to the motion of complex singularities through the analytical continuation into complex plane from the interface [15, 16]. Approach of singularity to the interface always means a formation of its geometric singularity. Other examples of analysis of weakly nonlinear 2D dynamics through motion of singularities include the interface between ideal fluid and light highly viscous fluid [17], and vortex sheet in ideal fluid [18]. Extending weakly nonlinear solutions into strongly nonlinear solution is challenging and mostly was done for the particular case of free surface hydrodynamics (i.e. the density of the second fluid turns into zero) [19–24] including drops pinchoff [25, 26]. Another exception is the ideal fluid pushed through viscous fluid in a narrow gap between two parallel plates (Hele-Shaw flow) which can be approximately reduced to the Laplace growth equation (LGE) admitting an infinite set of exact solutions [27–32].

We use a key property of quantum KHI that both fluid components share the same volume. It allows to find the exact strongly nonlinear solutions and moreover, general integrability of growing solutions. This is achieved through the exact reduction of quantum KHI dynamics to LGE for arbitrary level of nonlinearity. These new solutions, in particular, describe the formation of cusps (dimples) on the He-II free surface in a finite time with both a surface curvature and velocities of components of He-II diverging at singular points. We expect that these singularities will be possible to observe in He-II experiment which is different from weaker singularities of the Moore’s type (were identified from approximate analysis

in Refs. [15, 16, 18]) and predicts smooth surface with jump only in second derivative. LGE is integrable in a sense of the existence of infinite number of integrals of motion and relation to the dispersionless limit of the integrable Toda hierarchy [33]. We suggest that the obtained reduction of quantum KHI to LGE is important to the general problem of integrability of surface dynamics [34]. It provides a very rare example of integrable physical system.

Superfluid component of He-II necessary has quantized vortices if counterflow velocity exceeds several mm/s with their density growing with that velocity [35]. Here we consider the dynamics of He-II at macroscopic scale where we can average over vortices. We neglect average vorticity from such averaging as well as ignore vorticity of normal component similar to Ref. [12, 13] which refers to that approximation as non-dissipative two-fluid description. In that approximation the dynamics of both fluid components is potential one, i.e. $\mathbf{v}_s = \nabla\Phi_s$ and $\mathbf{v}_n = \nabla\Phi_n$, where \mathbf{v}_s , \mathbf{v}_n are velocities of superfluid and normal components with Φ_s and Φ_n being the corresponding velocity potentials. We assume that both components are incompressible with densities $\rho_s \equiv \text{const}$, $\rho_n \equiv \text{const}$ and the total density $\rho \equiv \rho_s + \rho_n$. Incompressibility implies Laplace equation for each component, $\nabla^2\Phi_{n,s} = 0$. We focus on 2D flow $\mathbf{r} \equiv (x, y)$, where x and y are horizontal and vertical coordinates, respectively. We assume that both fluids occupy the region $-\infty < y \leq \eta(x, t)$, where $y = \eta(x, t)$ is the free surface elevation with the unperturbed surface given by $\eta(x, t) \equiv 0$. The flow of both components deep inside He-II ($y \rightarrow -\infty$) as well as at $|x| \rightarrow \infty$ is assumed to be uniform following x direction, which implies $\Phi_{n,s} \rightarrow V_{n,s}x$, where $V_{n,s}$ are the corresponding horizontal velocities. We use the reference frame of the center of mass such that $\rho_n V_n + \rho_s V_s = 0$ and introduce the relative velocity $V = V_s - V_n > 0$ between fluid components meaning that $V_{n,s} = \mp \rho_{s,n} V / \rho$.

The dynamic boundary condition (BC) at the free surface ($y = \eta$) follows from the generalization of Bernoulli Eq. into two fluid components, see e.g. Chap. 140 of Ref. [1] and Refs. [12, 13])

$$\rho_n \left(\frac{\partial\Phi_n}{\partial t} + \frac{(\nabla\Phi_n)^2}{2} \right) + \rho_s \left(\frac{\partial\Phi_s}{\partial t} + \frac{(\nabla\Phi_s)^2}{2} \right) \Big|_{y=\eta} = \Gamma - P_\alpha - P_g, \quad (1)$$

where $P_\alpha = -\alpha \frac{\partial}{\partial x} [\eta_x (1 + \eta_x^2)^{-1/2}]$ is the pressure jump at the free surface due to the surface tension α (the pressure is zero outside the fluid assuming that there is a vacuum there), $\eta_x \equiv \partial\eta/\partial x$, $P_g = \rho g \eta$ is the gravity pressure (the contribution of the acceleration due to gravity g) and $\Gamma = \rho_n \rho_s V^2 / (2\rho)$ is the Bernoulli constant which ensures that Eq. (1) is satisfied at $|x| \rightarrow \infty$.

The kinematic BCs at the free surface are given by

$$\eta_t (1 + \eta_x^2)^{-1/2} = \partial_n \Phi_n|_{y=\eta} = \partial_n \Phi_s|_{y=\eta}, \quad (2)$$

where $\eta_t \equiv \partial\eta/\partial t$, $\partial_n \equiv \mathbf{n} \cdot \nabla$ is the outward normal derivative to the free surface with $\mathbf{n} = (-\eta_x, 1)(1 + \eta_x^2)^{-1/2}$. Eqs. (1) and (2) together with $\nabla^2\Phi_n = \nabla^2\Phi_s = 0$ and BC at infinity form a closed set of equations of two-fluid hydrodynamics for KHI problem.

We introduce the average velocity $\mathbf{v} = (\rho_n \mathbf{v}_n + \rho_s \mathbf{v}_s) / \rho$ and the auxiliary potentials $\Phi = (\rho_n \Phi_n + \rho_s \Phi_s) / \rho$, $\phi = \sqrt{\rho_n \rho_s} (\Phi_n - \Phi_s) / \rho$ which are linear combinations of Φ_n and Φ_s thus satisfying Laplace equation together with $\nabla\Phi = \mathbf{v}$. BCs at either $y \rightarrow -\infty$ or $|x| \rightarrow \infty$ are reduced to

$$\Phi \rightarrow 0 \text{ and } \phi \rightarrow -Vx\sqrt{\rho_n \rho_s} / \rho. \quad (3)$$

Eq. (1) turns into

$$\frac{\partial\Phi}{\partial t} + \frac{(\nabla\Phi)^2}{2} + \frac{(\nabla\phi)^2}{2} \Big|_{y=\eta} = \frac{c^2}{2} - \frac{P_\alpha + P_g}{\rho}, \quad (4)$$

where $c = \sqrt{2\Gamma/\rho}$ is the constant which has the dimension of velocity. Eqs. (2) are reduced to

$$\eta_t (1 + \eta_x^2)^{-1/2} = \partial_n \Phi|_{y=\eta} \quad (5)$$

and

$$\partial_n \phi|_{y=\eta} = 0. \quad (6)$$

We replace ϕ by its harmonic conjugate ψ such that Cauchy-Riemann equations $\phi_x = \psi_y$ and $\phi_y = -\psi_x$ are valid. BC (6) for Laplace Eq.

$$\nabla^2\psi = 0 \quad (7)$$

at the free surface reduces to vanishing of tangential derivatives $\partial_\tau \psi|_{y=\eta} = 0$ because $\partial_\tau \psi|_{y=\eta} = -\partial_n \phi|_{y=\eta}$. Without the loss of generality we set

$$\psi|_{y=\eta} = 0. \quad (8)$$

BC at either $y \rightarrow -\infty$ or $|x| \rightarrow \infty$ are reduced to

$$\psi \rightarrow -Vy\sqrt{\rho_n \rho_s} / \rho = -cy. \quad (9)$$

If we introduce the stream functions $\Psi_{n,s}$ for the components of He-II (they satisfy Cauchy-Riemann equations $\partial_x \Phi_{n,s} = \partial_y \Psi_{n,s}$ and $\partial_y \Phi_{n,s} = -\partial_x \Psi_{n,s}$), then $\psi = (\Psi_n - \Psi_s) \sqrt{\rho_n \rho_s} / \rho$. ψ is fully determined by $\eta(x, t)$ from (8) and (9) while being independent on Φ . Dynamic BC (4) in terms of Φ and ψ is given by

$$\frac{\partial\Phi}{\partial t} + \frac{(\nabla\Phi)^2}{2} + \frac{(\nabla\psi)^2}{2} \Big|_{y=\eta} = \frac{c^2}{2} - \frac{P_\alpha + P_g}{\rho}. \quad (10)$$

Eqs. (5), (8) and (10) together with $\nabla^2\Phi = \nabla^2\psi = 0$ and BCs (3) and (9) at infinity form a closed set of equations equivalent (through harmonic conjugation) to solving two-fluid He-II hydrodynamics for KHI problem. It

is remarkable that this set is equivalent (up to trivial change of constants) to the problem of 2D dynamics of charged surface of ideal fluid in the limit when surface charges fully screen the electric field above the fluid free surface. This limit was realized experimentally for the He-II (with negligible ρ_n) free surface charged by electrons [36]. In that case Φ has the meaning of the only (ideal) fluid component and ψ represents (up to the multiplication on constant) the electrostatic potential in the ideal fluid. The term $\propto (\nabla\psi)^2$ in Eq. (10) corresponds to the electrostatic pressure.

Refs. [37, 38] found exact time-dependent solutions for this problem of the dynamics of charged surface of superfluid He-II in the limit of zero surface tension and gravity as well as for the limit of zero temperature (i.e. neglecting the normal component of He-II). We apply that approach for full (nonlinear) KHI problem with finite temperature. We set $\alpha = g = 0$ in right-hand side of Eq. (10). Below we provide estimates of the applicability of such neglect of surface tension and gravity for two-component dynamics of He-II.

Our goal is to reduce Eqs. (5), (8) and (10) together with $\nabla^2\Phi = \nabla^2\psi = 0$ and BCs (3) and (9) to solving LGE. Differentiation of Eq. (8) over t and x results in

$$\eta_t = -\psi_t/\psi_y|_{y=\eta}, \quad \eta_x = -\psi_x/\psi_y|_{y=\eta}$$

respectively. Using these expressions in kinematic BC (5) rewritten in the equivalent form $\eta_t = \Phi_y - \eta_x\Phi_x|_{y=\eta}$, allows to obtain that

$$\psi_t + \nabla\psi \cdot \nabla\Phi|_{y=\eta} = 0. \quad (11)$$

The sum and difference of Eqs. (10) and (11) (with $P_\alpha = P_g = 0$) result in

$$F_t^{(\pm)} \mp cF_y^{(\pm)} + (\nabla F^{(\pm)})^2|_{y=\eta} = 0, \quad (12)$$

where we introduced the harmonics potentials

$$F^{(\pm)} = (\Phi \pm \psi \pm cy)/2, \quad (13)$$

which satisfy the Laplace Eqs.

$$\nabla^2 F^{(\pm)} = 0, \quad F^{(\pm)} \rightarrow 0 \quad \text{for } y \rightarrow -\infty \text{ or } |x| \rightarrow \infty. \quad (14)$$

According to Eqs. (8) and (13), the motion of the free surface is determined by the implicit expression

$$c\eta = F^{(+)} - F^{(-)}|_{y=\eta}. \quad (15)$$

Returning to physical $\Phi_{n,s}$ and $\Psi_{n,s}$, we find that

$$2\rho F^{(\pm)} = \rho_n\Phi_n + \rho_s\Phi_s \pm \sqrt{\rho_n\rho_s}(\Psi_n - \Psi_s + Vy). \quad (16)$$

Eqs. (14) together with (12), and (15) is equivalent to the KHI problem. It is crucial that nonlinear Eqs. (12) decouple into separate Eqs. for $F^{(+)}$ and $F^{(-)}$.

We note that such decoupling does not occur for the classical KHI problem (the interface between two fluids) where the velocity potentials and stream functions of each of two fluids are defined in physically distinct regions ($y < \eta$ and $y > \eta$) thus making impossible a superposition of the type (16). (Decoupling is however possible by other method in small angle approximation with leading quadratic nonlinearity in perturbation series for classical KHI [16].)

The full set of equations (12), (14)-(15) is still generally coupled through Eq. (15). But an additional assumption (reduction) that either

$$F^{(+)} = 0 \text{ or } F^{(-)} = 0 \quad (17)$$

ensures the closed Eqs. which have a wide family of exact nontrivial solutions described below. That assumption remains valid as time evolves. It follows from Eqs. (13) that (17) ensure the relations between $\Phi_{n,s}$ and $\Psi_{n,s}$ as

$$\mp \sqrt{\rho_n\rho_s}(\Psi_n - \Psi_s + Vy) = \rho_n\Phi_n + \rho_s\Phi_s.$$

We look at the physical meaning of our reductions (17), based on the particular limit of small amplitude surface waves. We neglect the nonlinear term in Eqs. (12) resulting in the linear system which we solve in the form of plane waves

$$F^{(\pm)} = a^{(\pm)} \exp(ikx + ky - i\omega^{(\pm)}t), \quad \eta = b^{(+)} \times \exp(ikx - i\omega^{(+)}t) + b^{(-)} \exp(ikx - i\omega^{(-)}t), \quad (18)$$

where $a^{(\pm)}$ and $b^{(\pm)}$ are small constants, $\omega^{(\pm)}$ are frequencies and k is the wavenumber. First Eq. in (18) ensures the exact solution of Eqs. (14) with decaying BCs at $y \rightarrow -\infty$. Substitution of (18) into Eq. (15) and the linearization of (12) results in the relations

$$\omega^{(\pm)} = \pm ick, \quad cb^{(\pm)} = \pm a^{(\pm)}. \quad (19)$$

which are two branches of the dispersion relation of KHI with $g = \alpha = 0$ [1, 12, 13]. Superscripts “+” and “-” correspond to exponentially growing and decaying perturbations of the flat free surface, respectively. Eqs. (17) choose one of these two branches. Thus Eqs. (12), (14)-(15) together with (17) represent the fully nonlinear stage of such separation into two branches.

The generic initial conditions include both unstable and stable part (18) with the unstable part dominates as time evolves. Also it was shown in Refs. [38, 39] that small perturbation of $F^{(-)}$ on the background of large $F^{(+)}$ decays to zero. Thus the choice of the reduction $F^{(-)} = 0$ (which is assumed below) in Eq. (17) is the natural one to address the nonlinear stage of KHI. Then Eqs. (13) imply that $F^{(+)} = \Phi = \psi + cy$, i.e. Φ is determined by ψ . The boundary value problem (BVP) (7)-(9) solves for ψ at each t . The motion of the free surface is determined by Eq. (5) as

$$(\eta_t - c)(1 + \eta_x^2)^{-1/2} = \partial_n\psi|_{y=\eta}. \quad (20)$$

To solve BVP (7)-(9) we consider the conformal map $z = z(w, t)$ [40] from the lower complex half-plane $-\infty < v \leq 0$, $-\infty < u < +\infty$ of the complex variable $w = u + iv$ into the area $-\infty < y \leq \eta(x, t)$ occupied by the fluid in the physical plane $z = x + iy$ with the real line $v = 0$ mapped into fluid free surface. Then the free surface is given in the parametric form $y = Y(u, t) \equiv \text{Im } z(u, t)$ and $x = X(u, t) \equiv \text{Re } z(u, t)$. Solutions of both BVP (7)-(9) and the harmonically conjugated BVP $\nabla^2 \phi = 0$, (3), (6) in (u, v) variables are given by $\phi + i\psi = -c(u + iv)$. It means that the conformal variables u and v have a simple physical meaning: $u = -\phi/c$ and $v = -\psi/c$ corresponding (up to multiplication to the constant $-1/c$) to the harmonically conjugated potentials ϕ and ψ .

We consider w as independent variable while $z(w, t)$ as the unknown function. Eq. (20) is given by $Y_t X_u - Y_u X_t = cX_u - c$, which can be rewritten as

$$\text{Im}(\bar{G}_t G_u) = c. \quad (21)$$

where $G(u, t) = z(u, t) - ict$. Eq. (21) has the exact form of LGE which has the infinite number of exact solutions often involving logarithms (see e.g. Refs.[29–32]). We look at a periodic solution [31] with the wavenumber k ,

$$z = w - ikA^2(t)/2 - iA(t) \exp[-ikw], \quad (22)$$

where $A(t)$ is the amplitude of the free surface surface perturbation satisfying a nonlinear ordinary differential equation $dA/dt = ckA(1 - k^2 A^2)^{-1}$ which develops a finite-time singularity in dA/dt at the time $t = t_c$ with $A(t_c) = 1/k$. As t approaches t_c , a leading order solution is given by $A = 1/k - \sqrt{c\tau/k} + O(\tau)$, where $\tau = t_c - t$. Singularities of the conformal map (22) are determined by a condition $z_w = 0$ implying that they approach the real line $v = 0$ from above with the increase of t . That line is reached at $\tau = 0$ and $u = 2\pi n/k$, $n = 0, \pm 1, \pm 2, \dots$. In particular, choosing $n = 0$, expanding at $u = 0$ and assuming $\tau \rightarrow 0$ we obtain that

$$\begin{aligned} X &= u\sqrt{ck\tau} + k^2 u^3/6 + O(u\tau + u^3\tau^{1/2}), \\ Y &= -3/2k + 2\sqrt{ck\tau/k} + ku^2/2 + O(\tau + u^2\tau^{1/2}). \end{aligned} \quad (23)$$

Fig. 2 shows an example of such solution at different t . It follows from Eq. (23) that a cusp pointing downward (a dimple) $y + 3/2k \propto |x|^{2/3}$ is formed at the free surface at $t = t_c$ (i.e. $\tau = 0$) with the vertical velocity diverging as $\tau^{-1/2}$ at the tip of the cusp [29, 31].

Near the singularity (the tip of the cusp) one has to take into account the surface tension and the finite viscosity of the normal component to regularize the singularity. Surface tension near the singularity is given by $P_\alpha \approx \alpha/r$, where r is the radius of curvature of the free surface. It follows from Eq. (23) that $r \approx c\tau$, which implies that $P_\alpha \approx \alpha/c\tau$. The dynamic pressure P_v , which determines the development of KHI in LGE reduction, is given by $P_v = \rho v^2/2 \equiv \rho[(\nabla\Phi)^2 + (\nabla\psi)^2]/2$, where




FIG. 2: Evolution of an initial periodic perturbation of the free surface $y(x)$ for Eq. (22) with $kA(0) \approx 0.15$. The surface shape is shown over one spatial period for the times $ckt = 0, 0.8, 1.2, 1.4$ until the cusp singularity is formed. The dashed line shows the unperturbed free surface, $y \equiv 0$.

v is the typical velocity. Near singularity $v \simeq \sqrt{2c/k\tau}$ and $P_v = \rho c/k\tau$. Thus both P_v and $P_\alpha \propto \tau^{-1}$. Surface tension effect is small if the Weber number $We = P_v/P_\alpha$, the ratio of dynamic and surface tension pressures, is well above 1. Using $We \approx \rho c^2/\alpha k = \rho_n \rho_s V^2/(\rho \alpha k)$, and assuming $We \gtrsim 1$ for applicability of LGE regime, we obtain the condition for the wavelength $\lambda = 2\pi/k \gtrsim 2\pi \rho \alpha/(\rho_n \rho_s V^2)$. He-II at the temperature 1.5 K has $\rho_n = 0.016 \text{ g/cm}^3$, $\rho_s = 0.129 \text{ g/cm}^3$ and $\alpha = 0.332 \text{ dyn/cm}$ [41]. E.g. if $V = 15 \text{ cm/s}$ then $\lambda \gtrsim 0.64 \text{ cm}$.

The relative strength of inertial and viscous forces near the singularity is determined by the Reynolds number $Re = vr/\nu$, where ν is kinematic viscosity of He-II. Using that $v \approx \sqrt{2c/k\tau}$ and $r \approx c\tau$, implies that $Re \approx cv^{-1}\sqrt{2r/k}$, i.e. Re turns small for $r \rightarrow 0$ and viscosity has to be taken into account. A typical scale r_ν below which the flow of the normal component cannot be considered as potential one is estimated by setting $Re \approx 1$ which gives $r_\nu \approx kv^2/2c^2$. For the temperature 1.5 K, we use $\nu = 9.27 \cdot 10^{-5} \text{ cm}^2/\text{s}$ [41]. Then $r_\nu \approx 1.8 \cdot 10^{-10} \text{ cm}$, i.e. $r_\nu \ll \lambda$ thus the viscous effect is much less than the surface tension. The influence of gravity, which is determined by the Froude number $Fr = P_v/P_g$, is small near the singularity because the gravity pressure $P_g \simeq \rho g y$ is finite while P_v diverges as τ^{-1} implying divergence of Fr .

We conclude that we reduced fully nonlinear quantum KHI dynamics to LGE which has the infinite set of exact solutions with the generic formation of cusps at the free surface in a finite time. The key is the exact transform from two-fluid description into the effective single-fluid description of Eq. (10). It suggests a roadmap for efficient use of conformal map to include gravity and capillarity into dynamics. Adding capillarity would ensure singularity regularization at small spatial scales. Conformal map can be used for electro-hydrodynamic instability [36, 39] and Faraday waves [42] of He-II. Viscosity can

be taken into account through conformal map in Stokes flow regime of normal component which would go beyond weakly nonlinear result [17].

Free surface represents vortex sheet which results in the additional generation of quantized vortices at the nonlinear stage of KHI. It is expected to push quantum turbulence states T1 towards T2/T3 states [35].

The authors thank L.P. Mezhov-Deglin for helpful discussions. The work of P.L. was partially supported by the National Science Foundation DMS-1412140. The work of P.L. on KHI was supported by the Russian Scientific Foundation, Grant No. 14-22-00259. The work of N.Z. was performed under the state assignment no. 0389-2015-0023 and supported in part by the Russian Foundation for Basic Research (grant nos. 16-08-00228 and 17-08-00430), by the Presidium of RAS (program no. 2), and by the Presidium of the Ural Branch of RAS (grant no. 18-2-2-15).

* Electronic address: plushnik@math.unm.edu

† Electronic address: nick@iep.uran.ru

- [1] L. D. Landau and E. M. Lifshitz, *Fluid Mechanics, Third Edition: Volume 6* (Pergamon, New York, 1989).
- [2] G. E. Volovik, JETP Letters **75**, 418 (2002).
- [3] R. Blaauwgeers, V. B. Eltsov, G. Eska, A. P. Finne, R. P. Haley, M. Krusius, J. J. Ruohio, L. Skrbek, and G. E. Volovik, Phys. Rev. Lett. **89**, 155301 (2002).
- [4] G. E. Volovik, *The Universe in a Helium Droplet* (Oxford University Press, Oxford, 2003).
- [5] A. P. Finne, V. B. Eltsov, R. Hanninen, N. B. Kopnin, J. Kopu, M. Krusius, M. Tsubota, and G. E. Volovik, Rep. Progr. Phys. **69**, 3157 (2006).
- [6] G. E. Volovik, Uspekhi Fizicheskikh Nauk **185**, 970 (2015).
- [7] R. Hanninen and A. W. Baggaley, PNAS **111**, 4667 (2014).
- [8] I. A. Remizov, A. A. Levchenko, and L. P. Mezhov-Deglin, J. Low Temp. Phys. **185**, 324 (2016).
- [9] S. Babuin, V. S. Lvov, A. Pomyalov, L. Skrbek, and E. Varga, Phys. Rev. B. **94**, 174504 (2016).
- [10] J. Gao, W. Guo, V. S. L'vov, A. Pomyalov, L. Skrbek, E. Varga, and W. F. Vinen, JETP Lett **103**, 648 (2016).
- [11] J. Gao, E. Varga, W. Guo, and W. F. Vinen, J. Low Temp. Phys. **187**, 490 (2017).
- [12] S. E. Korshunov, Europhys. Letters **16**, 673 (1991).
- [13] S. E. Korshunov, JETP Letters **75**, 496 (2002).
- [14] E. A. Kuznetsov and P. M. Lushnikov, J. Exp. Theor. Phys. **81**, 332 (1995).
- [15] E. Kuznetsov, M. Spector, and V. Zakharov, Physics Letters A **182**, 387 (1993).
- [16] N. M. Zubarev and E. A. Kuznetsov, J. Exp. Theor. Phys. **119**, 169 (2014).
- [17] P. M. Lushnikov, Physics Letters A **329**, 49 (2004).
- [18] D. W. Moore, Proc. R. Soc. London, Ser. A **365**, 105 (1979).
- [19] S. Tanveer, Proc. R. Soc. Lond. A **435**, 137 (1991).
- [20] S. Tanveer, Proc. R. Soc. Lond. A **441**, 501 (1993).
- [21] S. J. Cowley, G. R. Baker, and S. Tanveer, J. Fluid Mech. **378**, 233 (1999).
- [22] S. A. Dyachenko, P. M. Lushnikov, and A. O. Korotkevich, JETP Letters **98**, 675 (2013).
- [23] E. A. Karabut and E. N. Zhuravleva, J. Fluid Mech. **754**, 308 (2014).
- [24] P. M. Lushnikov, Journal of Fluid Mechanics **800**, 557 (2016).
- [25] J. Eggers, M. A. Fontelos, D. Leppinen, and J. H. Snoeijer, Phys. Rev. Lett. **98**, 094502 (2007).
- [26] K. S. Turitsyn, L. Lai, and W. W. Zhang, Phys. Rev. Lett. **103**, 124501 (2009).
- [27] P. Polubarinova-Kochina, Dokl. Akad. Nauk SSSR **47**, 254 (1945).
- [28] L. A. Galin, Dokl. Akad. Nauk SSSR **47**, 246 (1945).
- [29] B. I. Shraiman and D. Bensimon, Phys. Rev. A **30**, 2840 (1984).
- [30] S. D. Howison, SIAM J. Appl. Math. **46**, 20 (1986).
- [31] D. Bensimon, L. P. Kadanoff, S. Liang, B. I. Shraiman, and C. Tang, Rev. Mod. Phys. **58**, 977 (1986).
- [32] M. B. Mineev-Weinstein and S. P. Dawson, Phys. Rev. E **50**, R24 (1994).
- [33] M. Mineev-Weinstein, P. B. Wiegmann, and A. Zabrodin, Phys. Rev. Lett. **84**, 5106 (2000).
- [34] A. I. Dyachenko and V. E. Zakharov, Phys. Lett. A **190**, 144 (1994).
- [35] D. J. Melotte and C. F. Barenghi, Physical Review Letters **80**, 4181 (1998).
- [36] V. S. Edelman, Sov. Phys. Usp. **23**, 227 (1980).
- [37] N. M. Zubarev, JETP Lett. **71**, 367 (2000).
- [38] N. M. Zubarev, J. Exp. Theor. Phys. **94**, 534 (2002).
- [39] N. M. Zubarev, J. Exp. Theor. Phys. **107**, 668 (2008).
- [40] A. I. Dyachenko, E. A. Kuznetsov, M. D. Spector, and V. E. Zakharov, Phys. Lett. A **221**, 73 (1996).
- [41] R. J. Donnelly and C. F. Barenghi, Journal of Physical and Chemical Reference Data **27**, 1217 (1998).
- [42] A. A. Levchenko, L. P. Mezhov-Deglin, and A. A. Pel'menev, JETP Lett. **106**, 252 (2017).