



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Entropic Energy-Time Uncertainty Relation

Patrick J. Coles, Vishal Katariya, Seth Lloyd, Iman Marvian, and Mark M. Wilde

Phys. Rev. Lett. **122**, 100401 — Published 12 March 2019

DOI: [10.1103/PhysRevLett.122.100401](https://doi.org/10.1103/PhysRevLett.122.100401)

Entropic Energy-Time Uncertainty Relation

Patrick J. Coles,¹ Vishal Katariya,² Seth Lloyd,^{3,4} Iman Marvian,⁵ and Mark M. Wilde^{2,6}

¹*Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA*

²*Hearne Institute for Theoretical Physics, Department of Physics and Astronomy,*

Louisiana State University, Baton Rouge, Louisiana 70803, USA

³*Department of Mechanical Engineering, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

⁴*Research Laboratory of Electronics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

⁵*Departments of Physics & Electrical and Computer Engineering,*

Duke University, Durham, North Carolina 27708, USA

⁶*Center for Computation and Technology, Louisiana State University, Baton Rouge, Louisiana 70803, USA*

(Dated: February 20, 2019)

Energy-time uncertainty plays an important role in quantum foundations and technologies, and it was even discussed by the founders of quantum mechanics. However, standard approaches (e.g., Robertson’s uncertainty relation) do not apply to energy-time uncertainty because, in general, there is no Hermitian operator associated with time. Following previous approaches, we quantify time uncertainty by how well one can read off the time from a quantum clock. We then use entropy to quantify the information-theoretic distinguishability of the various time states of the clock. Our main result is an entropic energy-time uncertainty relation for general time-independent Hamiltonians, stated for both the discrete-time and continuous-time cases. Our uncertainty relation is strong, in the sense that it allows for a quantum memory to help reduce the uncertainty, and this formulation leads us to reinterpret it as a bound on the relative entropy of asymmetry. Due to the operational relevance of entropy, we anticipate that our uncertainty relation will have information-processing applications.

Introduction—The uncertainty principle is one of the most iconic implications of quantum mechanics, stating that there are pairs of observables that cannot be simultaneously known. It was first proposed by Heisenberg [1] for the position \hat{q} and momentum \hat{p} observables and then rigorously stated by Kennard [2] in the familiar form using standard deviations: $\Delta\hat{q}\Delta\hat{p} \geq \hbar/2$. Robertson [3] later formulated a similar relation for a different class of observables, namely, for pairs of bounded Hermitian observables \hat{X} and \hat{Z} (e.g., the Pauli spin operators), as $\Delta\hat{X}\Delta\hat{Z} \geq \frac{1}{2}|\langle[\hat{X}, \hat{Z}]\rangle|$. Since then, many alternative formulations have been proven for similar Hermitian operator pairs (e.g., [4, 5]).

Unfortunately, these relations do not apply to energy and time since time does not, in general, correspond to a Hermitian operator. In particular, Pauli’s theorem states that the semi-boundedness of a Hamiltonian precludes the existence of a Hermitian time operator, or in other words, if there was such an operator, then the Hamiltonian would be unbounded from below and thus unphysical [6]. Hence, formulating a general energy-time uncertainty relation is a nontrivial task. We point to [7] for an overview on time in quantum mechanics.

Nevertheless, the energy-time pair is of significant importance both fundamentally and technologically. Energy-time uncertainty was already discussed by the founders of quantum mechanics: Bohr, Heisenberg, Schrödinger, and Pauli (see [8] for a review). In the special case of the harmonic oscillator, this pair corresponds to number and phase, and number-phase uncertainty is relevant to metrology [9], e.g., phase estimation in interferometry. The energy-time pair is arguably the most general observable pair in the sense that it applies to all physical systems (i.e., all systems have a Hamiltonian).

Despite the lack of a Hermitian observable associated with time, relations with the feel of energy-time uncertainty relations have been formulated. Mandelstam and Tamm [10] re-

lated the energy standard deviation ΔE to the time τ that it takes for a state to move to an orthogonal state: $\tau\Delta E \geq \frac{\pi\hbar}{2}$. This relation can be thought of as a speed limit—a bound on how fast a quantum state can move—and other similar speed limits have been formulated [11]. Alternatively, it can be thought of as bounding how well a quantum system acts as a clock, since the time resolution of the clock is related to the time τ for the system to move to an orthogonal state.

In this work, we take the clock perspective on time uncertainty: one’s uncertainty about time corresponds to how well one can “read off” the time from measuring a quantum clock. A natural measure for this purpose is to consider the information-theoretic distinguishability of the various time states. As such, we propose using entropy to quantify time uncertainty, and our main result is an entropic energy-time uncertainty relation.

Entropy has been widely employed in uncertainty relations for position-momentum [12] and finite-dimensional observables [13, 14]—see [15] for a recent detailed review of entropic uncertainty relations. The key benefits of entropy as an uncertainty measure are its clear operational meaning and its relevance to information-processing applications. Indeed, entropic uncertainty relations form the cornerstone of security proofs for quantum key distribution and other quantum cryptographic tasks [15]. They furthermore allow one to recast the uncertainty principle in terms of a guessing game, as we do below for energy and time.

An entropic uncertainty relation for energy and time was previously given in [16] by constructing an almost-periodic time observable and using a so-called almost-periodic entropy for time. This approach was extended in [17], where the Holevo information bound was used to derive an entropic energy-time uncertainty relation. However, as indicated in [16], an almost-periodic time observable serves as a poor




FIG. 1. Guessing game for energy-time uncertainty. (1) Bob prepares a quantum clock in the state ρ_A and sends it to Alice. (2) Alice flips a coin and (3) either measures the clock’s energy or randomly sets the clock’s time (i.e., applies a time evolution e^{-iHt} with t randomly chosen from a predefined set). Bob’s goal is to, depending on Alice’s coin flip, guess the clock’s energy or guess t by reading the clock. Our uncertainty relations constrain Bob’s ability to win this game.

quantum clock for aperiodic systems. In [18], the entanglement between a system and a clock was used to derive an entropic energy-time uncertainty relation for a Hamiltonian with a uniformly spaced spectrum.

In this paper, we derive entropic energy-time uncertainty relations for general, time-independent Hamiltonians. We first derive a relation for discrete and arbitrarily spaced time, and then we extend this relation to infinitesimally closely spaced (i.e., continuous) time. Our results apply to systems with either finite- or infinite-dimensional Hamiltonians.

A novel aspect of our energy-time uncertainty relation is that it allows the observer to reduce their uncertainty through access to a quantum memory system, as was the case in prior uncertainty relations [19]. The two main benefits of allowing for quantum memory are that (1) it dramatically tightens the relation when the clock is in a mixed state, and (2) it makes the relation more relevant to cryptographic applications in which the eavesdropper may hold the memory system (e.g., see [19]). Furthermore, by allowing for quantum memory, we can reinterpret our uncertainty relation as a bound on the relative entropy of asymmetry [20], and we discuss below the implications of this reinterpretation.

The fact that our uncertainty relation is stated using operationally-relevant entropies implies that it should be useful for information processing applications. For example, if one can distinguish between the time states well, then it is possible to extract randomness by performing an energy measurement. True random bits are critical to the execution of secure protocols and numerical computations. In this case, the randomness of energy measurement outcomes is certified by our bound. Entropic uncertainty relations also find use in proving the security of quantum key distribution (QKD) protocols [21]. If one party is able to prepare states in both the phase and number bases of photons, and if another party is able to perform measurements in these two bases, then both parties can distill a secret key whose security is guaranteed by our relation. We provide more details regarding applications in the supplementary material (Appendix A) [34].

Uncertainty relations can be understood in the framework

of a guessing game involving two players, Alice and Bob [15, 19], and Figure 1 shows this game for the energy-time pair. Bob prepares system A in an arbitrary state ρ_A and sends it to Alice. Alice then flips a coin. If she gets heads, she performs an energy measurement, and Bob then must guess the outcome (possibly with the help of a memory system R that is initially correlated to A). If she gets tails, she applies a time evolution e^{-iHt} in which t is randomly chosen from some predefined set, and then sends A back to Bob, who then tries to guess which time t Alice applied. All of our uncertainty relations can be understood in terms of this guessing game and can be viewed as constraints on Bob’s probability of winning this game (i.e., guessing both the energy and time correctly). There are other variations of this energy-time uncertainty guessing game that are possible, one of which is discussed in the Supplementary Material (Appendix B).

In what follows, we give some necessary preliminaries before stating our main result for the Rényi entropy family in the discrete-time case, and then we extend to the continuous-time case for the von Neumann entropy. Finally, we apply our relation to an illustrative example of a spin-1/2 particle.

Preliminaries—We begin by considering a finite-dimensional Hamiltonian H that acts on a quantum system A , and suppose that it has $N_E \in \mathbb{Z}^+$ real energy eigenvalues taken from a set $\mathcal{E} \subset \mathbb{R}$. We thus write the Hamiltonian as $H_A = \sum_{\varepsilon \in \mathcal{E}} \varepsilon \Pi_A^\varepsilon$, where Π_A^ε denotes the projector onto the subspace spanned by energy eigenstates with eigenvalue ε . The projectors obey $\Pi^\varepsilon \Pi^{\varepsilon'} = \Pi^\varepsilon \delta_{\varepsilon, \varepsilon'}$, where $\delta_{\varepsilon, \varepsilon'} = 1$ if $\varepsilon = \varepsilon'$ and $\delta_{\varepsilon, \varepsilon'} = 0$ otherwise.

We now recall how to encode the classical state of a clock into a quantum system. Inspired by the Feynman-Kitaev history state formalism [22–24], as well as the quantum time proposal of [25], we introduce a register T for storing the time, which can be interpreted as a background reference clock. A measurement on the time register is treated in this framework as a time measurement. Let $\mathcal{T} = \{t_1, \dots, t_K\}$ denote a set of times, for integer $K \geq 2$, such that $t_k \in \mathbb{R}$ for all $k \in \{1, \dots, K\}$, and $t_1 \leq t_2 \leq \dots \leq t_K$. We suppose that the register T has a complete, discrete, and orthonormal basis $\{|t_k\rangle\}_{k=1}^K$. The time values need not be evenly spaced, which means that the basis for register T can include any combination of $|\mathcal{T}| = K$ distinct and orthonormal kets.

Now consider a clock system A that may initially be correlated to a memory system R , together in a joint state ρ_{AR} with $\rho_A = \text{Tr}_R(\rho_{AR})$. Let random variable E capture the outcomes of an energy measurement on the system A . The outcomes can be stored in a classical register, which we also denote without ambiguity by E in what follows. To quantify energy uncertainty, we employ the Rényi conditional entropy $S_\alpha(E|R)$ (defined below) of the classical-quantum state

$$\omega_{ER} \equiv \sum_{\varepsilon \in \mathcal{E}} |\varepsilon\rangle\langle\varepsilon|_E \otimes \text{Tr}_A\{\Pi_A^\varepsilon \rho_{AR}\}, \quad (1)$$

where the kets $\{|\varepsilon\rangle\}_{\varepsilon \in \mathcal{E}}$ are orthonormal, obeying $\langle\varepsilon'|\varepsilon\rangle = \delta_{\varepsilon', \varepsilon}$, and thus serve as classical labels for the energies of

the Hamiltonian. To quantify the time uncertainty, we employ the Rényi conditional entropy $S_\alpha(T|A)$ of the following classical-quantum state:

$$\kappa_{TA} \equiv \frac{1}{|\mathcal{T}|} \sum_{k=1}^K |t_k\rangle\langle t_k|_T \otimes e^{-iHt_k} \rho_A e^{iHt_k}. \quad (2)$$

In the above and henceforth, we set $\hbar = 1$. The state κ_{TA} can be interpreted as the joint state of system A (the local quantum clock) and the background reference clock T , at an unknown time $t_k \in \mathcal{T}$ chosen according to the uniform distribution. Equivalently, this state can be understood as a time-decohered version of the Feynman-Kitaev history state [22–24], the latter of which has the entire history of the state $\rho_A(t)$ encoded and entangled with a time register in superposition. The classical-quantum states in (1) and (2) are in one-to-one correspondence with the following labeled ensembles, respectively:

$$\{p(\varepsilon), |\varepsilon\rangle\langle\varepsilon|_E \otimes \text{Tr}_A\{\Pi_A^\varepsilon \rho_{AR}\}/p(\varepsilon)\}_{\varepsilon \in \mathcal{E}}, \\ \{1/|\mathcal{T}|, |t_k\rangle\langle t_k|_T \otimes e^{-iHt_k} \rho_A e^{iHt_k}\}_{t_k \in \mathcal{T}},$$

where $p(\varepsilon) = \text{Tr}\{\Pi_A^\varepsilon \rho_{AR}\}$.

Rényi entropies—For a probability distribution $\{p_j\}$, the Rényi entropies are defined for $\alpha \in (0, 1) \cup (1, \infty)$ by $S_\alpha(\{p_j\}) = \frac{1}{1-\alpha} \log_2 \sum_j p_j^\alpha$, and for $\alpha \in \{0, 1, \infty\}$ in the limit. This entropy family is generalized to quantum states via the sandwiched Rényi conditional entropy [26], defined for a bipartite state ρ_{AB} with $\alpha \in (0, \infty]$ as

$$S_\alpha(A|B)_\rho = -\inf_{\sigma_B} D_\alpha(\rho_{AB} \| I_A \otimes \sigma_B), \quad (3)$$

where the optimization is with respect to all density operators σ_B on system B . The quantity $S_\alpha(A|B)_\rho$ is in turn defined from the sandwiched Rényi relative entropy of a density operator ξ and a positive semi-definite operator ζ , which is defined for $\alpha \in (0, 1) \cup (1, \infty)$ as [26, 27]

$$D_\alpha(\xi \| \zeta) = \frac{1}{\alpha - 1} \log_2 \text{Tr}[(\zeta^{\frac{1-\alpha}{2\alpha}} \xi \zeta^{\frac{1-\alpha}{2\alpha}})^\alpha]. \quad (4)$$

If $\alpha > 1$ and the support of ξ is not contained in the support of ζ , then it is defined to be equal to $+\infty$. The quantity $D_\alpha(\xi \| \zeta)$ is defined for $\alpha \in \{1, \infty\}$ in the limit.

Entropic energy-time uncertainty relation—Let us now state our uncertainty relation for energy and time. For a pure state $\rho_A = |\psi\rangle\langle\psi|_A$ uncorrelated with a reference system R , it is as follows:

$$S_\alpha(T|A)_\kappa + S_\beta(\{p(\varepsilon)\}) \geq \log_2 |\mathcal{T}|, \quad (5)$$

holding for all $\alpha \in [1/2, \infty]$, with β satisfying $1/\alpha + 1/\beta = 2$, where $p(\varepsilon) = \langle\psi| \Pi_A^\varepsilon |\psi\rangle$. The above inequality (5) is saturated, e.g., when $|\psi\rangle$ is an energy eigenstate. Such states also maximize the time uncertainty, $S_\alpha(T|A)_\kappa = \log_2 |\mathcal{T}|$, since they are stationary states.

The concavity of entropy and concavity of conditional entropy [28] then directly imply that the same inequality in (5)

holds for a mixed state uncorrelated with a reference system R . However, if ρ_A is a maximally mixed state, the inequality in (5) yields a trivial bound on the total uncertainty. This is because the inequality does not capture the inherent uncertainty of the initial state.

One of our main results remedies this deficiency, capturing the inherent uncertainty mentioned above and holding non-trivially for mixed states:

$$S_\alpha(T|A)_\kappa + S_\beta(E|R)_\omega \geq \log_2 |\mathcal{T}|. \quad (6)$$

The entropic energy-time uncertainty relation in (6) holds for all $\alpha \in [1/2, \infty]$, where β satisfies $1/\alpha + 1/\beta = 2$, with the proof given in Appendix C. The quantity $S_\alpha(T|A)_\kappa$ represents the uncertainty about the time t_k from the perspective of someone holding the A system of the state κ_{TA} in (2). The quantity $S_\beta(E|R)_\omega$, which is determined by the state ρ_{AR} and the Hamiltonian H_A , represents the uncertainty about the outcome of an energy measurement from the perspective of someone who possesses the R system of the state ω_{ER} in (1). In the case that ρ_{AR} is pure, then the quantity $S_\beta(E|R)_\omega$ is determined by the reduced state ρ_A and the Hamiltonian H_A . According to (6), a good quantum clock state ρ_A , for which $S_\alpha(T|A)_\kappa \approx 0$, necessarily has a large uncertainty in the energy measurement, in the sense that $S_\beta(E|R)_\omega \gtrsim \log_2 |\mathcal{T}|$. Conversely, a state with a small uncertainty in the energy measurement, i.e., $S_\beta(E|R)_\omega \approx 0$, is necessarily a poor quantum clock state, i.e., $S_\alpha(T|A)_\kappa \approx \log_2 |\mathcal{T}|$.

Note that the uncertainties in (6) are entropic and hence do not quantify the uncertainties of time and energy in their units, but rather the amount of information (in bits) that we do not know about the respective quantities. For example, if a system can equally likely take on one of two energies E_1 and E_2 , then the entropic uncertainty in energy constitutes only one bit, and it does not depend on the magnitudes of E_1 or E_2 . Each entropy in (6) is analogous to a guessing probability, which quantifies how well one can guess the time t given the state $\rho_A(t)$, or the energy given the ability to measure a memory system R . In fact, $S_\alpha(A|B)$ converges to the negative logarithm of the guessing probability as $\alpha \rightarrow \infty$ [26, 29].

Considering the special case of $|\mathcal{T}| = 2$, one finds a simple, yet interesting corollary of (6): under the Hamiltonian H_A , a quantum state ρ_A can evolve to a perfectly distinguishable state, only if $S_\beta(E|R)_\omega \geq 1$ for $\beta \in [1/2, \infty]$. In other words, for $S_\beta(E|R)_\omega < 1$, the orthogonalization time τ in the Mandelstam-Tamm bound is infinite, which cannot be seen using Mandelstam-Tamm or other standard quantum speed limits.

By means of a quantum memory, one can also reduce the time uncertainty instead of only reducing the energy uncertainty. This can be accomplished by considering the memory system R to be a bipartite system $R_1 R_2$. One can then write the uncertainty relation in (6) as follows:

$$S_\alpha(T|AR_1)_\kappa + S_\beta(E|R_2)_\omega \geq \log_2 |\mathcal{T}|, \quad (7)$$

with full details given in Appendix D. This shows that the tightening of (5) to give (6) using quantum memory can re-

duce the uncertainties in both energy and time. We note that this rewriting is achieved only by relabeling systems, and is thus a consequence of our earlier result in (6).

An important special case of (6) is $\alpha = \beta = 1$ where both entropies are the von Neumann conditional entropy. This results in the following entropic uncertainty relation:

$$S(T|A)_\kappa + S(E|R)_\omega \geq \log_2 |\mathcal{T}|, \quad (8)$$

where the von Neumann conditional entropy of a bipartite state τ_{CD} can be written as $S(C|D)_\tau = -\text{Tr}[\tau_{CD} \log_2 \tau_{CD}] + \text{Tr}[\tau_D \log_2 \tau_D]$. In fact, we show in Appendix E of the Supplementary Material that the following equality holds for the von Neumann case when ρ_{AR} is pure:

$$S(T|A)_\kappa + S(E|R)_\omega = \log_2 |\mathcal{T}| + D(\kappa_A \| \sum_{\varepsilon} \Pi^{\varepsilon} \rho_A \Pi^{\varepsilon}).$$

As discussed in the Supplementary Material (Appendix E), when ρ_{AR} is pure, equality in (8) is achieved [equivalently, $D(\kappa_A \| \sum_{\varepsilon} \Pi^{\varepsilon} \rho_A \Pi^{\varepsilon}) = 0$] if and only if

$$\frac{1}{|\mathcal{T}|} \sum_{k=1}^K e^{-iHt_k} \rho_A e^{iHt_k} = \sum_{\varepsilon} \Pi^{\varepsilon} \rho_A \Pi^{\varepsilon}. \quad (9)$$

One way to satisfy (9) is if $[\rho_A, H] = 0$, and hence the relation is tight for states ρ_A that are diagonal in the energy eigenbasis. Another way to satisfy (9) is if $\frac{1}{|\mathcal{T}|} \sum_{k=1}^K e^{i(\varepsilon - \varepsilon')t_k} = \delta_{\varepsilon, \varepsilon'}$ for all combinations of $\varepsilon, \varepsilon'$. If the $|\mathcal{T}|$ times are equally spaced, this implies that $e^{i(\varepsilon - \varepsilon')t_K} = 1$ and $(\varepsilon - \varepsilon')t_K = 2\pi$. This can be understood as an exact inverse relationship between the conjugate variables, which is a signature of a saturated uncertainty relation.

Eq. (8) can be generalized to non-uniform probabilities for the various times. As shown in the Supplementary Material (Appendix E), the right-hand-side of (8) gets replaced by the entropy $S(T)_\kappa$ of the time distribution for this generalization.

Relative entropy of asymmetry formulation—As shown in the Supplementary Material (Appendix F), an alternative way of stating our main result in (6) is by employing the sandwiched Rényi relative entropy of asymmetry [30], which generalizes an asymmetry measure put forward in [20]:

$$S_\alpha(T|A)_\kappa + \inf_{\sigma: [H, \sigma]=0} D_\alpha(\rho \| \sigma) \geq \log_2 |\mathcal{T}|, \quad (10)$$

and holds for all $\alpha \in (0, \infty]$. The inequality in (10) delineates a trade-off, given the Hamiltonian H , between how well a state ρ_A can serve as a quantum clock and the asymmetry of ρ_A with respect to time translations. Moreover, this connection is exact for pure states.

In the limit $\alpha \rightarrow 1$, the quantity $\inf_{\sigma: [H, \sigma]=0} D_\alpha(\rho \| \sigma)$ reduces to the relative entropy of asymmetry [20]

$$\begin{aligned} \lim_{\alpha \rightarrow 1} \inf_{\sigma: [H, \sigma]=0} D_\alpha(\rho \| \sigma) &= \inf_{\sigma: [H, \sigma]=0} D(\rho \| \sigma) \\ &\equiv \Gamma_H(\rho) = S(\Delta(\rho)) - S(\rho), \end{aligned} \quad (11)$$




FIG. 2. Our uncertainty relations applied to a spin-1/2 particle in a magnetic field. For Hamiltonian $H = \kappa \sigma_z$ and $|\mathcal{T}| = 2$ (in the discrete-time case) or $T_F = 2$ (in the continuous-time case), the plot shows the variation in the uncertainties with θ , the angle the state makes with the z -axis of the Bloch sphere. The quantity $\inf_{\sigma: [H, \sigma]=0} D(\rho_A \| \sigma)$ is the energy uncertainty, while $S(T|A)$ and $s(T|A)$ are respectively the time uncertainties for discrete and continuous time. The black dotted line shows $\log_2 |\mathcal{T}| = \log_2 T_F$, i.e., our lower bounds on the total uncertainty from (6) and (12).

where the quantum relative entropy is defined as $D(\rho \| \sigma) \equiv \text{Tr}[\rho \log_2 \rho - \log_2 \sigma]$ [31] and $\Delta(\rho) = \sum_{\varepsilon \in \mathcal{E}} \Pi^{\varepsilon} \rho \Pi^{\varepsilon}$ (in the context of asymmetry, the function $S(\Delta(\rho)) - S(\rho)$ was first studied in [32]). Then the entropic uncertainty relation in (10) reduces to $S(T|A)_\kappa + \Gamma_H(\rho) \geq \log_2 |\mathcal{T}|$.

Extension to continuous time—We now extend the uncertainty relation in (6) so that it is applicable to continuous, as opposed to discrete, time, and to Hamiltonians with countable spectrum. From (6) and [33], we derive an inequality applicable to the von Neumann entropies. Full details are available in the Supplementary Material (Appendix G). Consider time to be continuous in the interval $[0, T_F]$. Given a state ρ_A and a Hamiltonian $H_A = \sum_{\varepsilon \in \mathcal{E}} \varepsilon \Pi_A^{\varepsilon}$, with \mathcal{E} countably infinite, we then have that

$$\inf_{\sigma: [H, \sigma]=0} D(\rho_A \| \sigma) + s(T|A) \geq \log_2 T_F. \quad (12)$$

For a continuously parametrized ensemble of states $\{p(x), \rho_B^x\}_{x \in \mathcal{X}}$, the differential conditional quantum entropy $s(X|B)$ is defined as $s(X|B) = -\int_{\mathcal{X}} dx D(p(x) \rho_B^x \| \rho_{\text{avg}})$, where $\rho_{\text{avg}} = \int_{\mathcal{X}} dx p(x) \rho_B^x$ [33]. For our case, this means

$$\begin{aligned} s(T|A) &= -\int_0^{T_F} dt D(\rho(t)/T_F \| \bar{\rho}), \\ \bar{\rho} &= \frac{1}{T_F} \int_0^{T_F} dt e^{-iHt} \rho_A e^{iHt}. \end{aligned}$$

Example: Spin in a magnetic field—Consider a spin-1/2 particle in a magnetic field $\mathbf{B} = B\hat{z}$. This is described by the Hamiltonian $\hat{H} = \kappa \sigma_z$, where κ is a constant proportional to B , and σ_z is the z -Pauli operator. Consider a pure state $\rho_A = |\psi(0)\rangle\langle\psi(0)|$ that makes an angle θ with the z -axis of the Bloch sphere, given by $|\psi(0)\rangle = \cos(\theta/2)|0\rangle + \sin(\theta/2)|1\rangle$.

After a time t , this state evolves to $|\psi(t)\rangle = e^{-iHt}|\psi(0)\rangle$. Figure 2 plots the variation of the uncertainty (time, energy, and total uncertainty) with θ for both our discrete- and continuous-time relations. For $\theta = \pi/2$, the energy uncertainty is maximal (one bit) while the time uncertainty is minimal (although still non-zero in this example). At the other extreme, for $\theta = 0$ or π , the energy uncertainty is zero while the time uncertainty is maximal (one bit), meaning that clock's time states cannot be distinguished. One can see in Figure 2 that our uncertainty relation is tight in this extreme case.

Discussion—We gave a conceptually clear and operational formulation of the energy-time uncertainty principle. We stated an entropic energy-time uncertainty relation for the Rényi entropies for discrete time sets. This relation was strengthened for mixed states by allowing the observer to possess a quantum memory, a feature that also allowed us to reinterpret our relation as a bound on the relative entropy of asymmetry. For the von Neumann entropy, we extended our uncertainty relation to continuous time sets. Our relation is saturated for all states ρ_A diagonal in the energy eigenbasis.

Expressed in terms of entropies, which are operationally important in information theory, our result should have uses in various tasks. Entropic uncertainty relations have been used previously to certify randomness and prove security of quantum cryptography protocols, and we believe our result will be an important tool used to develop such protocols further.

PJC acknowledges support from the Los Alamos National Laboratory ASC Beyond Moore's Law project. VK acknowledges support from the Department of Physics and Astronomy at Louisiana State University. SL was supported by IARPA under the QEO program, NSF, and ARO under the Blue Sky Initiative. MMW acknowledges support from the US National Science Foundation through grant no. 1714215. MMW is grateful to SL for hosting him for a research visit to University of Oxford during May 2018, during which some of this research was conducted.

[1] Werner Heisenberg. Über den anschaulichen inhalt der quantentheoretischen kinematik und mechanik. *Zeitschrift für Physik*, 43(3-4):172–198, March 1927.

[2] E. H. Kennard. Zur quantenmechanik einfacher bewegungstypen. *Zeitschrift für Physik*, 44(4-5):326–352, April 1927.

[3] H. P. Robertson. The uncertainty principle. *Physical Review*, 34(1):163–164, July 1929.

[4] Paul Busch, Pekka Lahti, and Reinhard F. Werner. Heisenberg uncertainty for qubit measurements. *Physical Review A*, 89(1):012129, January 2014. arXiv:1311.0837.

[5] Lorenzo Maccone and Arun K. Pati. Stronger uncertainty relations for all incompatible observables. *Physical Review Letters*, 113(26):260401, December 2014. arXiv:1407.0338.

[6] Wolfgang Pauli (ed. Norbert Straumann). *Die allgemeinen Prinzipien der Wellenmechanik*. Springer Berlin Heidelberg, 1990.

[7] Jeremy Butterfield. On time in quantum physics. In *A Com-*

panion to the Philosophy of Time (eds H. Dyke and A. Bardon), pages 220–241. John Wiley & Sons, Ltd, January 2013. arXiv:1406.4745.

[8] V. V. Dodonov and A. V. Dodonov. Energy–time and frequency–time uncertainty relations: exact inequalities. *Physica Scripta*, 90(7):074049, June 2015. arXiv:1504.00862.

[9] Vittorio Giovannetti, Seth Lloyd, and Lorenzo Maccone. Quantum-enhanced measurements: Beating the standard quantum limit. *Science*, 306(5700):1330–1336, November 2004. arXiv:quant-ph/0412078.

[10] L. Mandelstam and Ig. Tamm. The uncertainty relation between energy and time in non-relativistic quantum mechanics. In *Selected Papers*, pages 115–123. Springer Berlin Heidelberg, 1991.

[11] Norman Margolus and Lev B. Levitin. The maximum speed of dynamical evolution. *Physica D: Nonlinear Phenomena*, 120(1-2):188–195, September 1998. arXiv:quant-ph/9710043.

[12] I. I. Hirschman. A note on entropy. *American Journal of Mathematics*, 79(1):152–156, January 1957.

[13] David Deutsch. Uncertainty in quantum measurements. *Physical Review Letters*, 50(9):631–633, February 1983.

[14] Hans Maassen and J. B. M. Uffink. Generalized entropic uncertainty relations. *Physical Review Letters*, 60(12):1103–1106, March 1988.

[15] Patrick J. Coles, Mario Berta, Marco Tomamichel, and Stephanie Wehner. Entropic uncertainty relations and their applications. *Reviews of Modern Physics*, 89(1):015002, February 2017. arXiv:1511.04857.

[16] Michael J. W. Hall. Almost-periodic time observables for bound quantum systems. *Journal of Physics A*, 41(25):255301, May 2008. arXiv:0803.3721.

[17] Michael J. W. Hall. Entropic Heisenberg limits and uncertainty relations from the Holevo information bound. *Journal of Physics A*, 51(36):364001, September 2018. arXiv:1804.01343.

[18] A. Boette, R. Rossignoli, N. Gigena, and M. Cerezo. System-time entanglement in a discrete-time model. *Physical Review A*, 93(6):062127, June 2016. arXiv:1512.07313.

[19] Mario Berta, Matthias Christandl, Roger Colbeck, Joseph M. Renes, and Renato Renner. The uncertainty principle in the presence of quantum memory. *Nature Physics*, 6(9):659–662, July 2010. arXiv:0909.0950.

[20] Gilad Gour, Iman Marvian, and Robert W. Spekkens. Measuring the quality of a quantum reference frame: The relative entropy of frameness. *Physical Review A*, 80(1):012307, July 2009. arXiv:0901.0943.

[21] Charles H. Bennett and Gilles Brassard. Quantum cryptography: Public key distribution and coin tossing. In *Proceedings of IEEE International Conference on Computers Systems and Signal Processing*, pages 175–179, Bangalore, India, December 1984.

[22] Richard P. Feynman. Quantum mechanical computers. *Optics News*, 11(2):11–20, February 1985.

[23] Richard Feynman. Quantum mechanical computers. *Foundations of Physics*, 16(6):507–531, June 1986.

[24] A. Yu. Kitaev, A. H. Shen, and M. N. Vyalii. *Classical and Quantum Computation*. American Mathematical Society, Boston, MA, USA, 2002.

[25] Vittorio Giovannetti, Seth Lloyd, and Lorenzo Maccone. Quantum time. *Physical Review D*, 92(4):045033, August 2015. arXiv:1504.04215.

[26] Martin Müller-Lennert, Frédéric Dupuis, Oleg Szehr, Serge Fehr, and Marco Tomamichel. On quantum Rényi entropies: A new generalization and some properties. *Journal of Mathematical Physics*, 54(12):122203, December 2013. arXiv:1306.3142.

- [27] Mark M. Wilde, Andreas Winter, and Dong Yang. Strong converse for the classical capacity of entanglement-breaking and Hadamard channels via a sandwiched Rényi relative entropy. *Communications in Mathematical Physics*, 331(2):593–622, October 2014. arXiv:1306.1586.
- [28] Elliott H. Lieb and Mary Beth Ruskai. A fundamental property of quantum-mechanical entropy. *Physical Review Letters*, 30(10):434–436, March 1973.
- [29] Robert König, Renato Renner, and Christian Schaffner. The operational meaning of min- and max-entropy. *IEEE Transactions on Information Theory*, 55(9):4337–4347, September 2009. arXiv:0807.1338.
- [30] Li Gao, Marius Junge, and Nicholas LaRacuente. Strong subadditivity inequality and entropic uncertainty relations. October 2017. arXiv:1710.10038.
- [31] Hisaharu Umegaki. Conditional expectations in an operator algebra IV (entropy and information). *Kodai Mathematical Seminar Reports*, 14(2):59–85, 1962.
- [32] J. Vaccaro, F. Anselmi, H. Wiseman, and K. Jacobs. Tradeoff between extractable mechanical work, accessible entanglement, and ability to act as a reference system, under arbitrary superselection rules. *Physical Review A*, 77(3):032114, March 2008. arXiv:quant-ph/0501121.
- [33] Fabian Furrer, Mario Berta, Marco Tomamichel, Volkher B. Scholz, and Matthias Christandl. Position-momentum uncertainty relations in the presence of quantum memory. *Journal of Mathematical Physics*, 55(12):122205, December 2014. arXiv:1308.4527.
- [34] See Supplementary Material [URL] for these applications and mathematical details of some derivations, as well as Refs. [35–44].
- [35] R. Impagliazzo, L. A. Levin, and M. Luby. Pseudo-random generation from one-way functions. In *Proceedings of the twenty-first annual ACM symposium on Theory of computing - STOC '89*, pages 12–24. ACM Press, 1989.
- [36] Charles H. Bennett, Gilles Brassard, Claude Crépeau, and Ueli M. Maurer. Generalized privacy amplification. *IEEE Transactions on Information Theory*, 41(6):1915–1923, November 1995.
- [37] Giuseppe Vallone, Davide G. Marangon, Marco Tomasin, and Paolo Villoresi. Quantum randomness certified by the uncertainty principle. *Physical Review A*, 90(5):052327, November 2014. arXiv:1401.7917.
- [38] Davide G. Marangon, Giuseppe Vallone, and Paolo Villoresi. Source-device-independent ultrafast quantum random number generation. *Physical Review Letters*, 118(6):060503, February 2017. arXiv:1509.0739.
- [39] Zheshen Zhang, Jacob Mower, Dirk Englund, Franco N. C. Wong, and Jeffrey H. Shapiro. Unconditional security of time-energy entanglement quantum key distribution using dual-basis interferometry. *Physical Review Letters*, 112(12):120506, March 2014. arXiv:1311.0825.
- [40] Murphy Yuezhen Niu, Feihu Xu, Fabian Furrer, and Jeffrey H. Shapiro. Finite-key analysis for time-energy high-dimensional quantum key distribution. June 2016. arXiv:1606.08394.
- [41] Bing Qi. Single-photon continuous-variable quantum key distribution based on the energy-time uncertainty relation. *Optics Letters*, 31(18):2795–2797, August 2006. arXiv:quant-ph/0602158.
- [42] Patrick J. Coles. Unification of different views of decoherence and discord. *Physical Review A*, 85(4):042103, April 2012. arXiv:1110.1664.
- [43] Salman Beigi. Sandwiched Rényi divergence satisfies data processing inequality. *Journal of Mathematical Physics*, 54(12):122202, December 2013. arXiv:1306.5920.
- [44] Marius Junge, Renato Renner, David Sutter, Mark M. Wilde, and Andreas Winter. Universal recovery maps and approximate sufficiency of quantum relative entropy. *Annales Henri Poincaré*, 19(10):2955–2978, October 2018. arXiv:1509.07127.