



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Echo Technique to Distinguish Flavors of Astrophysical Neutrinos

Shirley Weishi Li, Mauricio Bustamante, and John F. Beacom

Phys. Rev. Lett. **122**, 151101 — Published 16 April 2019

DOI: [10.1103/PhysRevLett.122.151101](https://doi.org/10.1103/PhysRevLett.122.151101)

Echo Technique to Distinguish Flavors of Astrophysical Neutrinos

Shirley Weishi Li,^{1,2,*} Mauricio Bustamante,^{1,2,†} and John F. Beacom^{1,2,3,‡}

¹Center for Cosmology and AstroParticle Physics (CCAPP), Ohio State University, Columbus, OH 43210

²Department of Physics, Ohio State University, Columbus, OH 43210

³Department of Astronomy, Ohio State University, Columbus, OH 43210

(Dated: August 26, 2016)

The flavor composition of high-energy astrophysical neutrinos is a rich observable. However, present analyses cannot effectively distinguish particle showers induced by ν_e versus ν_τ . We show that this can be accomplished by measuring the intensities of the delayed, collective light emission from muon decays and neutron captures, which are, on average, greater for ν_τ than for ν_e . This new technique would significantly improve tests of the nature of astrophysical sources and of neutrino properties. We discuss the promising prospects for implementing it in IceCube and other detectors.

Introduction.— High-energy astrophysical neutrinos, long sought, were recently discovered by the IceCube Collaboration [1–6]. Their energy spectrum provides important clues about extreme astrophysical sources as well as neutrino properties at unexplored energies. However, pressing mysteries remain.

Exploiting the flavor composition — the ratios of the fluxes of $\nu_e + \bar{\nu}_e$, $\nu_\mu + \bar{\nu}_\mu$, and $\nu_\tau + \bar{\nu}_\tau$ to the total flux — offers crucial additional clues. In the nominal scenario, a composition of $(\frac{1}{3} : \frac{2}{3} : 0)_S$ at the source is transformed by neutrino vacuum mixing to $(\frac{1}{3} : \frac{1}{3} : \frac{1}{3})_\oplus$ at Earth [7, 8]. Even for arbitrary flavor composition at the source, the maximal range of flavor composition at Earth with only standard mixing is surprisingly narrow [9], making deviations sensitive indicators of new physics [10–39].

So far, IceCube measurements of the flavor composition mostly separate muon tracks — made primarily by charged-current (CC) $\nu_\mu + \bar{\nu}_\mu$ interactions — from particle showers — made by all other interactions. A significant limitation is their poor ability to distinguish between CC interactions of ν_e and ν_τ (unless noted, ν_l refers to $\nu_l + \bar{\nu}_l$).

Synopsis of the paper.— We propose a new technique to break this ν_e - ν_τ degeneracy, one that could work for a wider range of energies than existing ideas (Glashow resonance [40–42], double pulses [43], double bangs [7], and lollipops [44]).

We introduce two new shower observables. In showers, many low-energy muons and neutrons are produced; after delays, the muons decay and the neutrons capture. We call the collective Cherenkov emission from the many independent decays and captures the *muon echo* and the *neutron echo*. We show that the echoes are brighter for ν_τ -initiated than for ν_e -initiated showers, which could allow them to be distinguished on a statistical basis.

Our focus is pointing out new observables to help solve the important problem of flavor identification. The




FIG. 1. Flavor composition $f_{l,\oplus}$ ($l = e, \mu, \tau$) of astrophysical neutrinos at Earth. Each axis is read parallel to its ticks. Orange: the IceCube fit [5]. Blue: the expected precision of our proposed technique (for the case $f_{e,\oplus} = f_{\tau,\oplus}$), assuming 100 showers of 100 TeV with well-detected echoes. Results for other energies are similar, but collecting 100 showers with echoes will likely require a larger detector than IceCube. Green: the standard expectation [7, 8] and maximal range with standard mixing [9].

technical aspects of implementation require experimental expertise. Nevertheless, in a preliminary evaluation, grounded in the measured properties of IceCube, we find the detection prospects promising.

Figure 1 shows that the present ν_e - ν_τ degeneracy in IceCube elongates the contours of the measured flavor composition [5]. It also shows how detecting echoes could refine these measurements, probing the flavor composition better than the maximal range with standard mixing [9], which would lead to powerful conclusions.

High-energy neutrino signatures.— At present, IceCube identifies neutrino-initiated events only as tracks

* Now at SLAC National Accelerator Laboratory, shirleyl@slac.stanford.edu

† Now at Niels Bohr Institute, mbustamante@nbi.ku.dk

‡ beacom.7@osu.edu

and showers, for which the Cherenkov light appears to emanate from approximate lines and spheres. Tracks are caused by muons, which travel up to ~ 10 km in ice [2], due to their low interaction and decay rates. Showers are caused by all other neutrino-induced particles and extend only ~ 10 m in ice [2], due to the high interaction and decay rates of their constituent particles.

Neutrinos produce secondaries through deep-inelastic scattering [45–47]. A neutrino interacts with a nucleon N via the CC channel $\nu_l + N \rightarrow l + X$ or the neutral-current (NC) channel $\nu_l + N \rightarrow \nu_l + X$, where $l = e, \mu, \tau$, and X represents hadrons. A fraction $(1 - y)$ of the neutrino energy goes to the final-state lepton; the remaining fraction y goes to the final-state hadrons. The inelasticity distribution peaks at $y = 0$ and has an average $\langle y \rangle \approx 0.3$ at 100 TeV, for both ν and $\bar{\nu}$, CC and NC.

Tracks are produced by ν_μ CC interactions plus 17% of ν_τ CC interactions where the tau decays to a muon [48].

Showers are produced by all other neutrino interactions. For ν_e CC interactions, the electron- and hadron-initiated showers combine, and their sum energy equals the neutrino energy. For ν_τ CC interactions, the tau decays promptly, so again the showers combine (when the tau does not decay to a muon); the neutrino energy estimate is slightly biased because $\sim 25\%$ of its energy is lost to outgoing neutrinos from tau decay. For NC interactions of all flavors, the hadron-initiated shower carries a fraction y of the neutrino energy; because of the steeply falling neutrino spectrum, NC interactions are subdominant in the total shower spectrum [49]. (This is also true for mis-identified ν_μ CC interactions that appear to be a shower event because the track is missed [34].)

These points explain the basic features of the IceCube results in Fig. 1. Because there are track events, the ν_μ component of the flux must be nonzero; because there are shower events, the sum of the ν_e and ν_τ components must be nonzero. The similarity of ν_e - and ν_τ -initiated events makes the contours nearly horizontal; the degeneracy is weakly broken because increasing the ν_τ/ν_e fraction increases the number of tracks and decreases the shower energies. With present methods, improvement requires much larger exposure [9, 50, 51].

Electromagnetic versus hadronic showers.—

The key to our new method is understanding the low-energy physics underlying high-energy showers [52–59].

When showers are developing, particles multiply in number while decreasing in energy. An electromagnetic shower starts out with electrons, positrons, and gamma rays and stays composed predominantly of them; there is usually a small fraction of pions and nucleons produced by photonuclear processes. A hadronic shower starts out with pions and nucleons, and then builds up a progressively larger fraction of electromagnetic particles as prompt $\pi^0 \rightarrow \gamma\gamma$ decays deplete $\sim 1/3$ of the remaining hadronic energy with each shower generation.

Shower development ends when the average particle energy is low enough that the particle- and energy-loss rates exceed the particle-production rates. At that point,




FIG. 2. Time evolution of the light yield of a hadronic shower simulated with FLUKA, following injection of a 100-TeV charged pion. The shaded bands are exponentials with the respective timescales. For an electromagnetic shower of the same prompt energy, the echoes are ~ 10 times smaller.

the most abundant particles in all showers are ~ 100 -MeV electrons and positrons, which produce most of the prompt Cherenkov light. Pions carry only $\sim 10\%$ of the energy in hadronic showers and $\sim 1\%$ in electromagnetic showers. However, they are the key to separating electromagnetic and hadronic showers.

New shower observables.— At the end of shower development, charged pions come to rest by ionization; then π^- capture on nuclei and π^+ decay to μ^+ . The μ^+ decay with a lifetime of $2.2 \mu\text{s}$, producing e^+ with ~ 35 MeV. The collective Cherenkov light from these positrons is our first new observable: the *muon echo*.

Separately, neutrons lose energy by collisions until they reach thermal energy. They eventually capture on hydrogen, with a timescale of $\sim 200 \mu\text{s}$, producing 2.2 MeV gamma rays. (In seawater, 33% of neutrons capture on Cl; the emitted gamma rays have 8.6 MeV [60], making the neutron echoes much more visible.) The gamma rays Compton-scatter electrons to moderate energies, producing Cherenkov light. This collective emission is our second new observable: the *neutron echo*.

We simulate showers and subsequent echoes using the FLUKA Monte Carlo software (version 2011.2c-4) [61, 62]. We inject high-energy electrons or positrons to simulate electromagnetic showers and charged pions to simulate hadronic showers.

Figure 2 shows the averaged time profile of a 100-TeV hadronic shower. Because the features happen on very different timescales, it is appropriate to analyze their light yield L in bins of log time. Accordingly, we plot




FIG. 3. Numbers of muon decays and neutron captures (scaled down by 10) per shower, as a function of shower energy, for electromagnetic and hadronic showers simulated with FLUKA. The bands show 1σ intrinsic fluctuations.

$dL/d\log t \propto t dL/dt$; this makes the height of the curve proportional to its contribution to the integrated light yield. The echo shapes are exponentials with the respective timescales. The echoes are well-separated from the prompt shower and from each other.

Figure 2 also shows that the echoes have low intensities: the muon echo has $\sim 3 \times 10^{-3}$ of the prompt shower energy and the neutron echo has $\sim 6 \times 10^{-4}$. The first number results from the facts that 10% of hadronic shower energy goes to pions, 10% of those pions are π^+ that come to rest and decay, and 30% of the pion decay energy goes to positrons from muon decays. The second number results from the facts that there are about 10 times more neutron captures than muon decays, that the capture energy is about 20 times smaller, and that the Cherenkov efficiency is about 3 times smaller.

The points above carry over for electromagnetic showers, except for a crucial difference: the pions carry only $\sim 1\%$ of the shower energy as opposed to $\sim 10\%$. Thus the echo intensities are expected to be ~ 10 times higher in hadronic showers than in electromagnetic showers.

Figure 3 shows that there are indeed about 10 times as many muon decays and neutron captures in hadronic showers. This difference is much larger than the intrinsic fluctuations of these numbers. Because the number of decays and captures, and, therefore, the light coming from them, grows linearly with shower energy, this factor-of-10 difference between electromagnetic and hadronic showers is present at all energies. The yields may have an overall shift of up to a factor of 2 due to hadronic and nuclear uncertainties [57, 63–65], but this can be calibrated by




FIG. 4. Probability distributions of the numbers of muon decays per shower (of energy 100 TeV) for different neutrino interaction channels, each normalized separately.

external measurements [66–69] or *in situ*.

Separating ν_e and ν_τ .— We now examine how echoes can be used to help identify the flavors of neutrino-induced showers. In realistic neutrino interactions, the differences in the echoes are less stark than above.

Showers initiated by ν_e are mostly electromagnetic because the outgoing electron typically carries more energy than the final-state hadrons. But showers initiated by ν_τ are mostly hadronic because, in addition to the shower from the final-state hadrons, 67% of tau decays are hadronic. (NC showers are purely hadronic.)

We consider flavor separation at fixed shower energy, as opposed to fixed neutrino energy, to make contact with experiment. We simulate neutrino interactions with appropriate energies to give $E_{sh} = 100$ TeV, including 10% energy resolution [70]. For NC interactions, we mimic the final-state hadrons by directly injecting charged pions at the shower energy.

Figure 4 shows how the numbers of muon decays per shower are distributed for different neutrino interaction channels. As expected, ν_e CC showers produce fewer muons than ν_τ CC showers.

The basics of the distributions in Fig. 4 can be understood easily. For pure electromagnetic showers, the peak would be at ~ 500 decays; it would be narrow because most pions are produced late in the shower and the fluctuations are mostly Poissonian. For pure hadronic showers, the peak would be at ~ 8000 decays; it would be broad because there are large fluctuations in how much energy goes into π^0 in the first few shower generations. The shapes shown in Fig. 4 depend also on the y distributions for neutrino interactions. For ν_e CC events,

the distribution is substantially broadened because the differential cross section $d\sigma/dy$, while peaked at $y = 0$, has a substantial tail. For ν_τ CC events, there is a slight shift to the left, due to the 17% of tau decays to muons.

The results in Fig. 4 make it possible to distinguish ν_e and ν_τ on a statistical basis. We next estimate the sensitivity to flavor composition using the echoes from an ensemble of events, assuming well-detected echoes.

First, we use the results in Fig. 4 to generate the muon decay distributions for each flavor, assuming an equal flux of ν_l and $\bar{\nu}_l$, and NC to CC event ratios consistent with a power-law spectral index of 2.5 [5]. Next, for an assumed flavor composition, we randomly sample the number of muon decays for each shower in an ensemble of 100 showers of $E_{\text{sh}} = 100$ TeV. Then, we treat the flavor composition $f_{e,\oplus}$ and $f_{\tau,\oplus}$ as free parameters ($f_{\mu,\oplus} = 1 - f_{e,\oplus} - f_{\tau,\oplus}$) and use an unbinned maximum-likelihood procedure to find their best-fit values. We generate 10^3 different realizations of the shower ensemble, and find the average best-fit values and uncertainties of $f_{e,\oplus}$ and $f_{\tau,\oplus}$. Further details are in the Supplemental Material.

Figure 1 shows the predicted sensitivity on $f_{e,\oplus}$ and $f_{\tau,\oplus}$, assuming equal ν_e and ν_τ content, *i.e.*, a composition of the form $(x : 1 - 2x : x)_{\oplus}$, where x varies in $[0, 0.5]$. The vertical shape of the band shows that the sensitivity to $f_{e,\oplus}$ and $f_{\tau,\oplus}$ does not depend on the ν_μ content. (Because our method is only weakly sensitive to $f_{\mu,\oplus}$, we suppress its uncertainty in the plot.)

Our results are conservative. The sensitivity improves slightly with shower energy. Assuming well-detected echoes, the sensitivity is comparable whether we use muon echoes only, neutron echoes only, or both. It is also comparable, or better, for other choices of input parameters. See the Supplemental Material for details.

Observability of the echoes.— Echo detection depends on how the echo light yield compares to that from ambient backgrounds and detector transients. These quantities are detector-dependent, and we use IceCube as a concrete example, either for itself or to guide the design of upgrades or future detectors.

The echoes are faint, but they are well localized, which enhances their visibility. In space, like the parent shower, they are concentrated among only the few photomultiplier tubes (PMT) on a single string that are closest to the neutrino interaction vertex [71]. In time, they occur $\sim 2.2 \mu\text{s}$ and $\sim 200 \mu\text{s}$ after the prompt shower. These timescales require long-time data collection, made possible by the recent development of the Hit-Spooling technique, which can go to hours for infrequent events [72, 73]. In direction, the shower light is beamed forward but the echo light is isotropic. Light scattering makes the shower more isotropic and increases its duration [70], which could partially obscure the muon echo.

The total light yield of a shower in IceCube is ~ 100 detected photoelectrons (p.e.) per TeV [2]. For 100 TeV, the muon echo in a hadronic shower is expected to yield ~ 30 p.e. (300 GeV) and the neutron echo ~ 6 p.e. (60 GeV). At 30 p.e., IceCube can easily trigger on neu-

trino events of that yield [74]; the low efficiency at that energy shown in many analyses reflects the need to reject atmospheric muons. For echoes, because the place and time of the preceding shower is known, IceCube could restrict attention to a small number of PMTs, allowing triggering on echoes with as few as 4 p.e. [73, 75]. The minimum energy for our method is thus at least ~ 10 TeV.

Ambient backgrounds in IceCube do not eclipse the echoes. For an average p.e. noise rate of ~ 500 Hz per PMT [76], the expected backgrounds in $2 \mu\text{s}$ and $200 \mu\text{s}$ are only $\sim 10^{-3}$ and $\sim 10^{-1}$ p.e. per PMT, respectively. (Even with correlated noise, due to nuclear decays near the PMT, the backgrounds will be small in all but a few PMTs, and those will be identifiable [77].) And the cosmic-ray muon rate in IceCube is 3 kHz [2], so the probability of a muon lighting up several specific PMTs in the short time between shower and echo is small.

A serious concern is the detector transient called afterpulsing, where a PMT registers late p.e. with total charge proportional to the initial signal (the shower) and with a time profile characteristic to the PMT. For the IceCube PMTs, the muon echo will compete with an afterpulse feature of relative amplitude $\sim 10^{-2} E_{\text{sh}}$ near $2 \mu\text{s}$ [76]; though larger than the echo, it is not overwhelmingly so. Encouragingly, the neutron echo, though smaller, is late enough that afterpulsing seems to be negligible.

In summary, the prospects for observing echoes are promising, and they improve with shower energy. Doing so may require changes in detector design or in PMT technology [78, 79]; these considerations may shape the design of IceCube-Gen2 [50], KM3NeT [80, 81], and Baikal-GVD [82]. With multiple nearby PMTs [50, 81, 83], it may be possible to reconstruct individual events, dramatically improving background rejection. The final word on the observability of the echoes will come from detailed studies by the experimental collaborations.

Conclusions.— The rich phenomenology contained in the flavor composition of high-energy astrophysical neutrinos cannot be fully explored due to the difficulty of distinguishing showers initiated by ν_e versus ν_τ in neutrino telescopes. To break this degeneracy, we have introduced two new observables of showers: the delayed, collective light, or “echoes,” from muon decays and neutron captures. This light reflects the size of the hadronic component of a shower, and it is stronger in ν_τ -initiated than ν_e -initiated showers.

Figure 1 shows the promise of our method. With 100 showers with well-detected echoes, this would improve the separation of ν_e and ν_τ by a factor of ~ 9 over present measurements [5]. That is comparable to the estimated sensitivity attainable with present techniques after more than 15 years of exposure of the next-generation detector IceCube-Gen2 [50, 51]. The observation of other flavor-specific event signatures — Glashow resonance, double bangs, double pulses, and lollipops — will further constrain the flavor composition.

The applications of tagging hadronic showers via muon

and neutron echoes extend beyond flavor discrimination. The technique could improve shower energy reconstruction, by folding in the probability of a shower being electromagnetic or hadronic. And, at the considered energies, the echoes are shifted forward along the shower direction by ~ 5 m from the shower peak. If this shift can be detected, it would improve the poor angular resolution [70] of showers.

High-energy neutrino astronomy has just begun. We are still learning the best ways to detect and analyze astrophysical neutrinos. We should pursue all potentially detectable signatures, moving closer to finding the origins and properties of these ghostly messengers.

Note added: Preliminary results from a search for echoes in IceCube data are very encouraging [84].

ACKNOWLEDGMENTS

We thank Carlos Argüelles, Amy Connolly, Eric Huff, Tim Linden, Kenny Ng, David Nygren, Eric Oberla, Annika Peter, Benedikt Riedel, Jakob van Santen, Shigeru Yoshida, Guanying Zhu, and especially Lutz Köpke and Anna Steuer for useful discussions and comments. We also thank Markus Ahlers, Kfir Blum, Claudio Kopper, Carsten Rott, and others for helpful feedback beginning at JFB's presentation of these ideas at the CCAPP Cosmic Messages in Ghostly Bottles workshop in 2014. SWL, MB, and JFB are supported by NSF Grant PHY-1404311.

-
- [1] M. G. Aartsen *et al.* (IceCube Collaboration), “First observation of PeV-energy neutrinos with IceCube,” *Phys. Rev. Lett.* **111**, 021103 (2013), arXiv:1304.5356 [astro-ph.HE].
- [2] M. G. Aartsen *et al.* (IceCube), “Evidence for High-Energy Extraterrestrial Neutrinos at the IceCube Detector,” *Science* **342**, 1242856 (2013), arXiv:1311.5238 [astro-ph.HE].
- [3] M. G. Aartsen *et al.* (IceCube), “Search for a diffuse flux of astrophysical muon neutrinos with the IceCube 59-string configuration,” *Phys. Rev. D* **89**, 062007 (2014), arXiv:1311.7048 [astro-ph.HE].
- [4] M. G. Aartsen *et al.* (IceCube), “Observation of High-Energy Astrophysical Neutrinos in Three Years of IceCube Data,” *Phys. Rev. Lett.* **113**, 101101 (2014), arXiv:1405.5303 [astro-ph.HE].
- [5] M. G. Aartsen *et al.* (IceCube), “A combined maximum-likelihood analysis of the high-energy astrophysical neutrino flux measured with IceCube,” *Astrophys. J.* **809**, 98 (2015), arXiv:1507.03991 [astro-ph.HE].
- [6] M. G. Aartsen *et al.* (IceCube), “Evidence for Astrophysical Muon Neutrinos from the Northern Sky with IceCube,” *Phys. Rev. Lett.* **115**, 081102 (2015), arXiv:1507.04005 [astro-ph.HE].
- [7] J. G. Learned and S. Pakvasa, “Detecting tau-neutrino oscillations at PeV energies,” *Astropart. Phys.* **3**, 267–274 (1995), arXiv:hep-ph/9405296.
- [8] H. Athar, M. Jezabek, and O. Yasuda, “Effects of neutrino mixing on high-energy cosmic neutrino flux,” *Phys. Rev. D* **62**, 103007 (2000), arXiv:hep-ph/0005104 [hep-ph].
- [9] M. Bustamante, J. F. Beacom, and W. Winter, “Theoretically palatable flavor combinations of astrophysical neutrinos,” *Phys. Rev. Lett.* **115**, 161302 (2015), arXiv:1506.02645 [astro-ph.HE].
- [10] J. F. Beacom, N. F. Bell, D. Hooper, S. Pakvasa, and T. J. Weiler, “Decay of high-energy astrophysical neutrinos,” *Phys. Rev. Lett.* **90**, 181301 (2003), arXiv:hep-ph/0211305 [hep-ph].
- [11] G. Barenboim and C. Quigg, “Neutrino observatories can characterize cosmic sources and neutrino properties,” *Phys. Rev. D* **67**, 073024 (2003), arXiv:hep-ph/0301220 [hep-ph].
- [12] J. F. Beacom, N. F. Bell, D. Hooper, S. Pakvasa, and T. J. Weiler, “Measuring flavor ratios of high-energy astrophysical neutrinos,” *Phys. Rev. D* **68**, 093005 (2003), [Erratum: *Phys. Rev. D* **72**, 019901 (2005)], arXiv:hep-ph/0307025 [hep-ph].
- [13] J. F. Beacom, N. F. Bell, D. Hooper, S. Pakvasa, and T. J. Weiler, “Sensitivity to θ_{13} and δ in the decaying astrophysical neutrino scenario,” *Phys. Rev. D* **69**, 017303 (2004), arXiv:hep-ph/0309267 [hep-ph].
- [14] J. F. Beacom, N. F. Bell, D. Hooper, J. G. Learned, S. Pakvasa, and T. J. Weiler, “Pseudo-Dirac neutrinos, a challenge for neutrino telescopes,” *Phys. Rev. Lett.* **92**, 011101 (2004), arXiv:hep-ph/0307151.
- [15] Z. Z. Xing and S. Zhou, “Towards determination of the initial flavor composition of ultrahigh-energy neutrino fluxes with neutrino telescopes,” *Phys. Rev. D* **74**, 013010 (2006), arXiv:astro-ph/0603781 [astro-ph].
- [16] P. Lipari, M. Lusignoli, and D. Meloni, “Flavor Composition and Energy Spectrum of Astrophysical Neutrinos,” *Phys. Rev. D* **75**, 123005 (2007), arXiv:0704.0718 [astro-ph].
- [17] S. Pakvasa, W. Rodejohann, and T. J. Weiler, “Flavor Ratios of Astrophysical Neutrinos: Implications for Precision Measurements,” *JHEP* **02**, 005 (2008), arXiv:0711.4517 [hep-ph].
- [18] A. Esmaili and Y. Farzan, “An Analysis of Cosmic Neutrinos: Flavor Composition at Source and Neutrino Mixing Parameters,” *Nucl. Phys. B* **821**, 197–214 (2009), arXiv:0905.0259 [hep-ph].
- [19] K. C. Lai, G. L. Lin, and T. C. Liu, “Determination of the Neutrino Flavor Ratio at the Astrophysical Source,” *Phys. Rev. D* **80**, 103005 (2009), arXiv:0905.4003 [hep-ph].
- [20] J. L. Bazo, M. Bustamante, A. M. Gago, and O. G. Miranda, “High energy astrophysical neutrino flux and modified dispersion relations,” *Int. J. Mod. Phys. A* **24**, 5819 (2009), arXiv:0907.1979 [hep-ph].
- [21] S. Choubey and W. Rodejohann, “Flavor Composition of UHE Neutrinos at Source and at Neutrino Telescopes,” *Phys. Rev. D* **80**, 113006 (2009), arXiv:0909.1219 [hep-ph].

- [22] M. Bustamante, A. M. Gago, and C. Pena-Garay, “Energy-independent new physics in the flavour ratios of high-energy astrophysical neutrinos,” *JHEP* **1004**, 066 (2010), arXiv:1001.4878 [hep-ph].
- [23] M. Bustamante, A. M. Gago, and J. Jones-Perez, “SUSY Renormalization Group Effects in Ultra High Energy Neutrinos,” *JHEP* **1105**, 133 (2011), arXiv:1012.2728 [hep-ph].
- [24] P. Baerwald, M. Bustamante, and W. Winter, “Neutrino Decays over Cosmological Distances and the Implications for Neutrino Telescopes,” *JCAP* **1210**, 020 (2012), arXiv:1208.4600 [astro-ph.CO].
- [25] R. Laha, J. F. Beacom, B. Dasgupta, S. Horiuchi, and K. Murase, “Demystifying the PeV Cascades in IceCube: Less (Energy) is More (Events),” *Phys. Rev. D* **88**, 043009 (2013), arXiv:1306.2309 [astro-ph.HE].
- [26] C. Y. Chen, P. S. Bhupal Dev, and A. Soni, “Standard model explanation of the ultrahigh energy neutrino events at IceCube,” *Phys. Rev. D* **89**, 033012 (2014), arXiv:1309.1764 [hep-ph].
- [27] O. Mena, S. Palomares-Ruiz, and A. C. Vincent, “Flavor Composition of the High-Energy Neutrino Events in IceCube,” *Phys. Rev. Lett.* **113**, 091103 (2014), arXiv:1404.0017 [astro-ph.HE].
- [28] K. C. Y. Ng and J. F. Beacom, “Cosmic neutrino cascades from secret neutrino interactions,” *Phys. Rev. D* **90**, 065035 (2014), arXiv:1404.2288 [astro-ph.HE].
- [29] X. Xu, H. He, and W. Rodejohann, “Constraining Astrophysical Neutrino Flavor Composition from Leptonic Unitarity,” *JCAP* **1412**, 039 (2014), arXiv:1407.3736 [hep-ph].
- [30] L. Fu, C. M. Ho, and T. J. Weiler, “Aspects of the Flavor Triangle for Cosmic Neutrino Propagation,” *Phys. Rev. D* **91**, 053001 (2015), arXiv:1411.1174 [hep-ph].
- [31] C. Y. Chen, P. S. Bhupal Dev, and A. Soni, “Two-component flux explanation for the high energy neutrino events at IceCube,” *Phys. Rev. D* **92**, 073001 (2015), arXiv:1411.5658 [hep-ph].
- [32] S. Palomares-Ruiz, A. C. Vincent, and O. Mena, “Spectral analysis of the high-energy IceCube neutrinos,” *Phys. Rev. D* **91**, 103008 (2015), arXiv:1502.02649 [astro-ph.HE].
- [33] A. Palladino, G. Pagliaroli, F. L. Villante, and F. Vissani, “What is the Flavor of the Cosmic Neutrinos Seen by IceCube?” *Phys. Rev. Lett.* **114**, 171101 (2015), arXiv:1502.02923 [astro-ph.HE].
- [34] M. G. Aartsen *et al.* (IceCube), “Flavor Ratio of Astrophysical Neutrinos above 35 TeV in IceCube,” *Phys. Rev. Lett.* **114**, 171102 (2015), arXiv:1502.03376 [astro-ph.HE].
- [35] A. Palladino and F. Vissani, “The natural parameterization of cosmic neutrino oscillations,” *Eur. Phys. J. C* **75**, 433 (2015), arXiv:1504.05238 [hep-ph].
- [36] C. A. Argüelles, T. Katori, and J. Salvado, “New Physics in Astrophysical Neutrino Flavor,” *Phys. Rev. Lett.* **115**, 161303 (2015), arXiv:1506.02043 [hep-ph].
- [37] A. Palladino, G. Pagliaroli, F. L. Villante, and F. Vissani, “Double pulses and cascades above 2 PeV in IceCube,” *Eur. Phys. J. C* **76**, 52 (2016), arXiv:1510.05921 [astro-ph.HE].
- [38] I. M. Shoemaker and K. Murase, “Probing BSM Neutrino Physics with Flavor and Spectral Distortions: Prospects for Future High-Energy Neutrino Telescopes,” *Phys. Rev. D* **93**, 085004 (2016), arXiv:1512.07228 [astro-ph.HE].
- [39] Aaron C. Vincent, Sergio Palomares-Ruiz, and Olga Mena, “Analysis of the 4-year IceCube high-energy starting events,” *Phys. Rev. D* **94**, 023009 (2016), arXiv:1605.01556 [astro-ph.HE].
- [40] S. L. Glashow, “Resonant Scattering of Antineutrinos,” *Phys. Rev.* **118**, 316 (1960).
- [41] L. A. Anchordoqui, H. Goldberg, F. Halzen, and T. J. Weiler, “Neutrinos as a diagnostic of high energy astrophysical processes,” *Phys. Lett. B* **621**, 18–21 (2005), arXiv:hep-ph/0410003 [hep-ph].
- [42] A. Bhattacharya, R. Gandhi, W. Rodejohann, and A. Watanabe, “The Glashow resonance at IceCube: signatures, event rates and pp vs. $p\gamma$ interactions,” *JCAP* **1110**, 017 (2011), arXiv:1108.3163 [astro-ph.HE].
- [43] M. G. Aartsen *et al.* (IceCube), “Search for Astrophysical Tau Neutrinos in Three Years of IceCube Data,” *Phys. Rev. D* **93**, 022001 (2016), arXiv:1509.06212 [astro-ph.HE].
- [44] T. DeYoung, S. Razzaque, and D. F. Cowen, “Astrophysical tau neutrino detection in kilometer-scale Cherenkov detectors via muonic tau decay,” *Astropart. Phys.* **27**, 238–243 (2007), arXiv:astro-ph/0608486 [astro-ph].
- [45] R. Gandhi, C. Quigg, M. H. Reno, and I. Sarcevic, “Ultrahigh-energy neutrino interactions,” *Astropart. Phys.* **5**, 81–110 (1996), arXiv:hep-ph/9512364 [hep-ph].
- [46] R. Gandhi, C. Quigg, M. H. Reno, and I. Sarcevic, “Neutrino interactions at ultrahigh-energies,” *Phys. Rev. D* **58**, 093009 (1998), arXiv:hep-ph/9807264 [hep-ph].
- [47] A. Connolly, R. S. Thorne, and D. Waters, “Calculation of High Energy Neutrino-Nucleon Cross Sections and Uncertainties Using the MSTW Parton Distribution Functions and Implications for Future Experiments,” *Phys. Rev. D* **83**, 113009 (2011), arXiv:1102.0691 [hep-ph].
- [48] K. A. Olive *et al.* (Particle Data Group), “Review of Particle Physics,” *Chin. Phys. C* **38**, 090001 (2014).
- [49] J. F. Beacom and J. Candia, “Shower power: Isolating the prompt atmospheric neutrino flux using electron neutrinos,” *JCAP* **0411**, 009 (2004), arXiv:hep-ph/0409046 [hep-ph].
- [50] M. G. Aartsen *et al.* (IceCube), “IceCube-Gen2: A Vision for the Future of Neutrino Astronomy in Antarctica,” (2014), arXiv:1412.5106 [astro-ph.HE].
- [51] M. Kowalski, “Next-Generation IceCube,” Presented at Neutrino Oscillation Workshop (NOW) 2018, September 12, 2018, Ostuni, Italy.
- [52] W. Heitler, *The Quantum Theory of Radiation*, 3rd ed. (Dover Publications, 1954).
- [53] J. Matthews, “A Heitler model of extensive air showers,” *Astropart. Phys.* **22**, 387–397 (2005).
- [54] P. Lipari, “The Concepts of ‘Age’ and ‘Universality’ in Cosmic Ray Showers,” *Phys. Rev. D* **79**, 063001 (2009), arXiv:0809.0190 [astro-ph].
- [55] P. Lipari, “Universality of cosmic ray shower development,” *Nucl. Phys. Proc. Suppl.* **196**, 309–318 (2009).
- [56] C. Rott, J. M. Siegal-Gaskins, and J. F. Beacom, “New Sensitivity to Solar WIMP Annihilation using Low-Energy Neutrinos,” *Phys. Rev. D* **88**, 055005 (2013), arXiv:1208.0827 [astro-ph.HE].
- [57] S. W. Li and J. F. Beacom, “First calculation of cosmic-ray muon spallation backgrounds for MeV astrophysical neutrino signals in Super-Kamiokande,” *Phys. Rev. C* **89**, 045801 (2014), arXiv:1402.4687 [hep-ph].

- [58] S. W. Li and J. F. Beacom, “Spallation Backgrounds in Super-Kamiokande Are Made in Muon-Induced Showers,” *Phys. Rev. D* **91**, 105005 (2015), arXiv:1503.04823 [hep-ph].
- [59] S. W. Li and J. F. Beacom, “Tagging Spallation Backgrounds with Showers in Water-Cherenkov Detectors,” *Phys. Rev. D* **92**, 105033 (2015), arXiv:1508.05389 [physics.ins-det].
- [60] J. K. Tuli, “Evaluated Nuclear Structure Data File (ENSDF),” <http://www.nndc.bnl.gov/ensdf/>, [Online; accessed 2016-05-20].
- [61] A. Ferrari, P. R. Sala, A. Fasso, and J. Ranft, “FLUKA: A multi-particle transport code (Program version 2005),” (2005).
- [62] G. Battistoni *et al.*, “The FLUKA code: Description and benchmarking,” *AIP Conf. Proc.* **896**, 31–49 (2007).
- [63] S. Abe *et al.* (KamLAND), “Production of Radioactive Isotopes through Cosmic Muon Spallation in KamLAND,” *Phys. Rev. C* **81**, 025807 (2010), arXiv:0907.0066 [hep-ex].
- [64] G. Bellini *et al.* (Borexino), “Cosmogenic Backgrounds in Borexino at 3800 m water-equivalent depth,” *JCAP* **1308**, 049 (2013), arXiv:1304.7381 [physics.ins-det].
- [65] Y. Zhang *et al.* (Super-Kamiokande), “First measurement of radioactive isotope production through cosmic-ray muon spallation in Super-Kamiokande IV,” *Phys. Rev. D* **93**, 012004 (2016), arXiv:1509.08168 [hep-ex].
- [66] J. F. Beacom and M. R. Vagins, “GADZOOKS! Antineutrino spectroscopy with large water Cherenkov detectors,” *Phys. Rev. Lett.* **93**, 171101 (2004), arXiv:hep-ph/0309300 [hep-ph].
- [67] M. Askani *et al.* (WATCHMAN), “The Physics and Nuclear Nonproliferation Goals of WATCHMAN: A WATER Cherenkov Monitor for ANTineutrinos,” (2015), arXiv:1502.01132 [physics.ins-det].
- [68] I. Anghel *et al.* (ANNIE), “Letter of Intent: The Accelerator Neutrino Neutron Interaction Experiment (ANNIE),” (2015), arXiv:1504.01480 [physics.ins-det].
- [69] P. Fernandez (Super-Kamiokande), “Status of GADZOOKS!: Neutron Tagging in Super-Kamiokande,” *Proceedings, 37th International Conference on High Energy Physics (ICHEP 2014)*, *Nucl. Part. Phys. Proc.* **273-275**, 353–360 (2016).
- [70] M. G. Aartsen *et al.* (IceCube), “Energy Reconstruction Methods in the IceCube Neutrino Telescope,” *JINST* **9**, P03009 (2014), arXiv:1311.4767 [physics.ins-det].
- [71] C. Kopper, “Recent IceCube Results on High Energy Neutrinos,” http://macros2013.in2p3.fr/_shared/doc/talks/kopper.pdf, MACROS 2013.
- [72] M. G. Aartsen *et al.* (IceCube), “The IceCube Neutrino Observatory Part V: Neutrino Oscillations and Supernova Searches,” (2013), arXiv:1309.7008 [astro-ph.HE].
- [73] M. G. Aartsen *et al.* (IceCube), “The IceCube Neutrino Observatory: Instrumentation and Online Systems,” *JINST* **12**, P03012 (2017), arXiv:1612.05093 [astro-ph.IM].
- [74] M. G. Aartsen *et al.* (IceCube), “Measurement of the Atmospheric ν_e Spectrum with IceCube,” *Phys. Rev. D* **91**, 122004 (2015), arXiv:1504.03753 [astro-ph.HE].
- [75] L. Köpke and A. Steuer, (2016–2017), private communications.
- [76] R. Abbasi *et al.* (IceCube), “Calibration and Characterization of the IceCube Photomultiplier Tube,” *Nucl. Instrum. Meth. A* **618**, 139–152 (2010), arXiv:1002.2442 [astro-ph.IM].
- [77] M. J. Larson, *Simulation and Identification of Non-Poissonian Noise Triggers in the IceCube Neutrino Detector*, Master’s thesis, The University of Alabama (2013).
- [78] J. R. Incandela *et al.*, “The Performance of Photomultipliers Exposed to Helium,” *Nucl. Instrum. Meth. A* **269**, 237–245 (1988).
- [79] M. P. Bristow, “Suppression of afterpulsing in photomultipliers by gating the photocathode,” *Appl. Opt.* **41**, 4975–4987 (2002).
- [80] U. F. Katz, “KM3NeT: Towards a km³ Mediterranean Neutrino Telescope,” *Nucl. Instrum. Meth. A* **567**, 457–461 (2006), arXiv:astro-ph/0606068 [astro-ph].
- [81] S. Adrian-Martinez *et al.* (KM3Net), “Letter of intent for KM3NeT 2.0,” *J. Phys. G* **43**, 084001 (2016), arXiv:1601.07459 [astro-ph.IM].
- [82] A. V. Avrorin *et al.*, “Current status of the BAIKAL-GVD project,” *Nucl. Instrum. Meth. A* **725**, 23–26 (2013).
- [83] E. Resconi, “The stepping stones to proton decay: IceCube, PINGU, MICA,” https://indico.cern.ch/event/224351/contributions/466197/attachments/368604/513012/Aspen_2013_Resconi.pdf, Aspen Center for Physics, New Directions in Neutrino Physics 2013.
- [84] A. Steuer and L. Köpke, “Delayed light emission to distinguish astrophysical neutrino flavors in IceCube,” *Contributions to the 35th International Cosmic Ray Conference (ICRC 2017)*, *PoS ICRC 2017*, 1008 (2018).