



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Majorana Multipole Response of Topological Superconductors

Shingo Kobayashi, Ai Yamakage, Yukio Tanaka, and Masatoshi Sato

Phys. Rev. Lett. **123**, 097002 — Published 27 August 2019

DOI: [10.1103/PhysRevLett.123.097002](https://doi.org/10.1103/PhysRevLett.123.097002)

Majorana Multipole Response of Topological Superconductors

Shingo Kobayashi,^{1,2} Ai Yamakage,³ Yukio Tanaka,² and Masatoshi Sato⁴

¹*Institute for Advanced Research, Nagoya University, Nagoya 464-8601, Japan*

²*Department of Applied Physics, Nagoya University, Nagoya 464-8603, Japan*

³*Department of Physics, Nagoya University, Nagoya 464-8602, Japan*

⁴*Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan*

(Dated: July 25, 2019)

In contrast to elementary Majorana particles, emergent Majorana fermions (MFs) in condensed-matter systems may have electromagnetic multipoles. We developed a general theory of magnetic multipoles for helical MFs on time-reversal-invariant superconductors. The results show that the multipole response is governed by crystal symmetry, and that a one-to-one correspondence exists between the symmetry of Cooper pairs and the representation of magnetic multipoles under crystal symmetry. The latter property provides a way to identify unconventional pairing symmetry via the magnetic response of helical MFs. We also find that most helical MFs exhibit a magnetic-dipole response, but those on superconductors with spin-3/2 electrons may display a magnetic-octupole response in leading order, which uniquely characterizes high-spin superconductors. Detection of such an octupole response provides direct evidence of high-spin superconductivity, such as in half-Heusler superconductors.

Introduction. The emergence of Majorana fermions (MFs) in electron systems has led to intense interest in searching for such exotic new excitations in condensed-matter physics. Particularly, recent developments have shown that emergent MFs appear as gapless Andreev bound states in topological superconductors (TSCs) [1–14], which provide a potential candidate for fault-tolerant qubits for topological quantum computation [15]. The increased interest in topological materials has led to a proposal of versatile three-dimensional (3D) time-reversal-invariant (TR-invariant) TSCs, such as superconducting doped topological insulators (TIs) [16–23] and Dirac semimetals [24–30], which commonly host helical MFs forming Kramers pairs on their surfaces.

Emergent MFs share some properties with elementary Majorana particles [31, 32]: Both of them obey Dirac equations with charge-conjugation symmetry. Furthermore, a pair of MF zero modes are required to define the fermionic creation and annihilation operators, from which zero modes exhibit non-Abelian anyon statistics. However, compared with elementary Majorana particles, emergent MFs respond very differently to electric and magnetic fields. On the one hand, neither electric nor magnetic multipoles are possible for elementary MFs [33–35]: CPT invariance, where \mathcal{C} is charge conjugation, \mathcal{P} is space inversion, and \mathcal{T} is time reversal, is a fundamental symmetry that any relativistic elementary particles is expected to respect. This symmetry forbids intrinsic electric and magnetic multipoles for elementary Majorana particles because they are their own antiparticles under CPT . On the other hand, in superconductors, fundamental symmetry is just charge-conjugation (namely, particle-hole (PH) symmetry), and the emergent MFs are self-conjugate under \mathcal{C} . Therefore, MFs in condensed-matter physics are not subject to such a strong constraint, and no systematic study on their elec-

tromagnetic multipoles has yet been attempted.

In this Letter, we develop a theory describing the electric and magnetic response of MFs in superconductors. For clarity, we focus here on surface helical MFs on 3D TR-invariant TSCs. A key ingredient specific to emergent MFs is crystalline symmetry. In analogy with CPT invariance for elementary MFs, crystal symmetry provides additional symmetry constraints on electromagnetic structures of emergent MFs. Considering the constraints, we establish a response theory for helical MFs in a low-energy limit, in which the problem reduces to the selection rule for crystal-symmetry groups. Applying our theory to possible crystal-symmetry groups, we find that helical MFs can host magnetic-multipole structures of dipole or octupole orders as the leading contribution. Additionally, the results predict a one-to-one correspondence between irreducible representations (IRs) of Cooper pairs and magnetic multipoles, which helps to determine the pairing symmetry experimentally through the magnetic response of MFs.

Particularly, the proposed theory provides a unique way to identify topological superconductivity of spin-3/2 electrons. Although research interest has recently focused on high-spin topological superconductivity [29, 36–51], little is known about distinguishing TSCs of spin-3/2 electrons from those of spin-1/2 electrons. We clarify here that magnetic responses of helical MFs can unambiguously distinguish between these two types of SCs because the magnetic-octupole response is unique to high-spin TSCs. To illustrate this, we apply the proposed theory to superconducting TIs of ordinary spin-1/2 electrons [17] and parity-mixed half-Heusler superconductors of spin-3/2 electrons [41, 43]. The results of both numerical and analytical analyses show that only the latter exhibits the octupole response under the same crystalline symmetry.




FIG. 1. MFs with gapless point and line in the surface Brillouin zone (BZ) that are protected by two-fold rotation and mirror reflection, respectively.

Majorana multipole. Helical MFs are a superconducting analogue of surface Dirac fermions of TIs and can be realized in 3D TR-invariant TSCs. From the bulk-boundary correspondence, the existence of helical MFs is ensured by the so-called 3D winding number [4, 5, 52, 53]. Whereas the 3D winding number is defined only for fully gapped TSCs, its parity is well defined even for nodal superconductors [18]. Provided TR symmetry is maintained, these invariants are well defined and protect surface helical MFs for both nodal and nodeless superconductors.

We consider the quantum response of helical MFs when exposed to external electric or magnetic fields. First, we notice that electric fields only give moderate responses from helical MFs: Since electric fields keep TR symmetry, helical MFs remain gapless so they cannot respond so much. Conversely, magnetic fields may substantially affect them. Magnetic fields break TR symmetry, so the 3D winding number and its parity become invalid. However, this does not mean that helical MFs are not immune to any magnetic fields because actual TSCs have their own crystalline symmetry. Depending on the direction of the applied magnetic field, TR symmetry may be partially preserved by combining it with crystalline symmetry. Such magnetic crystalline symmetry determines the stability of helical MFs under magnetic fields [54].

As relevant point group operations, we consider rotations and mirror reflections that are compatible with the surface. The rotation axis and the mirror plane should be normal to the surface (see Fig. 1). We consider any surface-preserving point group G formed by them: $G = C_2, C_3, C_4, C_6, C_s, C_{2v}, C_{3v}, C_{4v}, C_{6v}$, in addition to TR symmetry \mathcal{T} . (G contains only the unbroken part of crystalline symmetry, if Cooper pairs spontaneously break a part of it.) Under a magnetic field, we retain magnetic two-fold rotation or magnetic mirror reflection. Note that the retained magnetic symmetry is selected by the direction of an applied magnetic field. Only for a magnetic field normal (parallel) to the rotation axis (mirror plane), magnetic two-fold rotation (magnetic mirror

reflection) is preserved: The above magnetic field is easily seen to flip under TR, but it points back to the original when we simultaneously do a two-fold rotation (mirror reflection).

The retained magnetic symmetry enables TSCs to host an additional topological number that is valid even when the TSC is exposed to a magnetic field: Let $g_0 \in G$ be two-fold rotation or mirror reflection that defines the magnetic symmetry. Combining the magnetic symmetry with charge conjugation \mathcal{C} , one can introduce the magnetic chiral operator $\Gamma_M \propto \tilde{g}_0 \mathcal{T} \mathcal{C}$, which involves the magnetic one-dimensional (1D) winding number w_{M1D} [55–59]. If w_{M1D} for magnetic two-fold rotation (magnetic mirror reflection) is nonzero in the absence of magnetic fields, then helical MFs remain gapless even under a magnetic field normal (parallel) to the rotation axis (mirror plane), provided the system maintains the bulk gap. On the other hand, helical MFs do not necessarily remain gapless under other magnetic fields. This direction dependence results in an anisotropic magnetic response of helical MFs. Note that w_{M1D} for magnetic two-fold rotation (magnetic mirror reflection) is defined on the symmetric axis (plane), so it protects the gapless point (line) of helical MFs at the symmetry axis (plane) in the surface Brillouin zone (see Fig. 1).

The gapless points or lines are obtained as zero modes $|u_0^{(a)}\rangle$ of the Bogoliubov-de Gennes (BdG) equation. ($a = 1, 2$ labels the Kramers degeneracy.) The index theorem for w_{M1D} [60] implies that the stable zero modes $|u_0^{(a)}\rangle$ have a common eigenvalue of Γ_M , say

$$\Gamma_M |u_0^{(a)}\rangle = |u_0^{(a)}\rangle. \quad (1)$$

since zero modes with opposite eigenvalues are easily gapped in pairs. (This property is rigorously proven for generic lattice systems [59].) Moreover, from crystalline symmetry, the zero modes $|u_0^{(a)}\rangle$ should transform as a (double-valued) representation under the action of G .

To systematically study the magnetic response of MFs, we examine possible contributions of MFs to a local operator $\hat{O}(x) = \hat{c}_\sigma^\dagger(x) O_{\sigma,\sigma'} \hat{c}_{\sigma'}(x)$ of electrons, where $\hat{c}_\sigma^\dagger(x)$ and $\hat{c}_\sigma(x)$ are the electron operators with internal degrees of freedom σ such as spin and orbital, and $O_{\sigma,\sigma'}$ is a Hermitian matrix. The MFs have a nonzero response to external fields through such a local operator. For instance, if MFs make a nonzero contribution to the electron-spin operator $\hat{S}_i(x) = \hat{c}_\sigma^\dagger(x) [s_i/2]_{\sigma,\sigma'} \hat{c}_{\sigma'}(x)$ with the Pauli matrix s_i , then the MFs show a nonzero magnetic response through the Zeeman term of electrons.

The contribution of MFs is evaluated as follows: In the Nambu space with $\hat{\Psi}^\dagger(x) = (\hat{c}_\sigma^\dagger(x), \hat{c}_\sigma(x))$, $\hat{O}(x)$ is recast into $\hat{O}(x) = (1/2) \hat{\Psi}^\dagger(x) \mathcal{O} \hat{\Psi}(x)$, with $\mathcal{O} = \text{diag}(O, -O^T) = \text{diag}(O, -O^*)$, where we have used the Hermiticity of O . Then, by performing the mode expansion of the quantum field $\hat{\Psi}(x) = \sum_{a=1,2} \hat{\gamma}^{(a)} |u_0^{(a)}\rangle +$ (nonzero modes) and using the PH symmetry $\mathcal{C} \hat{\Psi} = \hat{\Psi}$,

TABLE I. Magnetic multipole for Kramers pair of MFs. From left to right, each column shows two-dimensional point groups G , IRs of Δ with $w_{\text{M1D}} \neq 0$, the basis of Δ , \tilde{g}_0 associated with Γ_{M} , IRs of \mathcal{O} , and the basis of \mathcal{O} . Here, J_i are the spin matrices, “–” means the absence of IRs, and \mathcal{O} is the leading order of the magnetic multipole. We omit $e^{-i\pi J_y}$ in the basis of Δ .

G	IR of Δ	basis of Δ	\tilde{g}_0	IR of \mathcal{O}	basis of \mathcal{O}
C_2, C_4, C_6	A	$\mathbf{k} \cdot \mathbf{J}$	C_2	A	J_z
C_3	–	–	–	–	–
C_s	A	$k_x J_z, k_x J_y, k_y J_x, k_z J_x$	$\sigma_v(yz)$	A	J_x
C_{2v}	A_2	$k_z J_z$	C_2	A_2	J_z
	B_1	$k_x J_z, k_z J_x$	$\sigma_v(yz)$	B_1	J_x
	B_2	$k_y J_z, k_z J_y$	$\sigma_v(xz)$	B_2	J_y
C_{3v}	A_1	$k_z(J_x^3 - J_x J_y J_y - J_y J_x J_y - J_y J_y J_x)$	$\sigma_v(yz)$	A_1	$J_x^3 - J_x J_y J_y - J_y J_x J_y - J_y J_y J_x$
C_{4v}	A_2	$k_z J_z$	C_2	A_2	J_z
C_{6v}	A_2	$k_z J_z$	C_2	A_2	J_z
	B_1	$k_z(J_x^3 - J_x J_y J_y - J_y J_x J_y - J_y J_y J_x)$	$\sigma_v(yz)$	B_1	$J_x^3 - J_x J_y J_y - J_y J_x J_y - J_y J_y J_x$
	B_2	$k_z(J_y^3 - J_y J_x J_x - J_x J_y J_x - J_x J_x J_y)$	$\sigma_d(xz)$	B_2	$J_y^3 - J_y J_x J_x - J_x J_y J_x - J_x J_x J_y$

we obtain the coupling between $\hat{O}(x)$ and the MFs $\hat{\gamma}^{(a)}$ in the low-energy limit,

$$\begin{aligned} \hat{O}_{\text{MF}} &= \frac{1}{2} \sum_{a,b=1,2} \hat{\gamma}^{(b)} \hat{\gamma}^{(a)} \langle \mathcal{C} u_0^{(b)} | \mathcal{O} | u_0^{(a)} \rangle \\ &= \frac{1}{2} \hat{\gamma}^{(2)} \hat{\gamma}^{(1)} \text{tr} [\mathcal{O} \rho^{(12)}], \end{aligned} \quad (2)$$

where $\rho^{(ab)} \equiv |u_0^{(a)}\rangle \langle \mathcal{C} u_0^{(b)}| - |u_0^{(b)}\rangle \langle \mathcal{C} u_0^{(a)}|$ and an irrelevant constant term has been omitted in the last line. A non-zero \hat{O}_{MF} requires the following two conditions: (i) There should be zero modes $|u_0^{(a)}\rangle$ that satisfy Eq. (1). (ii) \mathcal{O} should share the same IR with $\rho^{(12)}$ under the action of G . As discuss below, the former condition determines possible pairing symmetry of Cooper pairs, and the latter decides possible magnetic responses.

We outline here how the condition (i) specifies the pairing symmetry of Cooper pairs. First, we note that the symmetry of Cooper pairs determines the commutation relations between charge-conjugation and point group operations. Let us consider the BdG Hamiltonian

$$\mathcal{H}(\mathbf{k}) = \begin{pmatrix} \mathcal{E}(\mathbf{k}) & \Delta(\mathbf{k}) \\ \Delta^*(\mathbf{k}) & -\mathcal{E}^T(-\mathbf{k}) \end{pmatrix}, \quad (3)$$

where $\mathcal{E}(\mathbf{k})$ is the Hamiltonian of the normal state and $\Delta(\mathbf{k})$ is the gap function of the superconducting state. A point group G implies that $g\mathcal{E}(\mathbf{k})g^{-1} = \mathcal{E}(g\mathbf{k})$ with $g \in G$. The BdG Hamiltonian retains G if the Cooper pairs have symmetry $g\Delta(\mathbf{k})g^T = e^{i\theta_g} \Delta(g\mathbf{k})$ ($e^{i\theta_g}$ is a phase factor) as it holds that $\tilde{g}\mathcal{H}(\mathbf{k})\tilde{g}^{-1} = \mathcal{H}(g\mathbf{k})$, where $\tilde{g} \equiv \text{diag}[g, e^{-i\theta_g} g^*]$ is the point group operator in the Nambu space. Then, the charge conjugation $\mathcal{C} \equiv \tau_x K$ and \tilde{g} obeys the relation $\tilde{g}\mathcal{C} = e^{i\theta_g} \mathcal{C}\tilde{g}$, where τ_i is the Pauli matrix in the Nambu space and K is the complex-conjugation operator. Note that \mathcal{T} always commutes with \tilde{g} (and \mathcal{C}), irrespective of the pairing symmetry. For TR-invariant TSCs, $e^{i\theta_g}$ must be real, so $e^{i\theta_g} = \pm 1$.

Thus, the gap function should be either even or odd under $g \in G$, which leads to $[\mathcal{C}, \tilde{g}] = 0$ ($\{\mathcal{C}, \tilde{g}\} = 0$) for g -even (odd) pairing symmetry.

As we discussed before, all zero modes, if exist, have a common eigenvalue of Γ_{M} . Combining this with the result in the above, we can determine the pairing symmetry that is consistent with the presence of helical MFs: Since a zero mode $|u_0^{(1)}\rangle$ and its Kramers partner $|u_0^{(2)}\rangle \equiv \mathcal{T}|u_0^{(1)}\rangle$ have the same eigenvalue of Γ_{M} , we should have $[\mathcal{T}, \Gamma_{\text{M}}]=0$. This relation determines the phase ambiguity of the magnetic operator as $\Gamma_{\text{M}} = \tilde{g}_0 \mathcal{C} \mathcal{T}$. Then, using $\mathcal{C}^2 = 1$ and $\tilde{g}_0^2 = \mathcal{T}^2 = -1$, we obtain $[\tilde{g}_0, \mathcal{C}] = 0$ so as to be consistent with $\Gamma_{\text{M}}^2 = 1$. This yields that the gap function should be even under g_0 . Moreover, any other $\tilde{g} \in G$ should not anti-commute with Γ_{M} for the same reason, so by considering the multiplication law between \tilde{g} and \tilde{g}_0 as well, we can establish the commutation or anticommutation relation between \mathcal{C} and \tilde{g} [61]. The obtained set of commutation or anti-commutation relations specifies the IR of the gap function under G , namely the pairing symmetry of Cooper pairs. We summarize it for given G and \tilde{g}_0 in Table I.

Now we discuss the condition (ii). Since $|u_0^{(a)}\rangle$ and $|\mathcal{C} u_0^{(a)}\rangle$ ($a = 1, 2$) are double-valued representations of G , $\rho^{(12)}$ is their product representation. Thus, using the standard group theory, we can decompose $\rho^{(12)}$ into IRs under G , which determines the representation of \mathcal{O} with a nonzero \hat{O}_{MF} . (See discussions in S2 of Supplemental Material [61].) We find that \mathcal{O} consists of a single IR, which is summarized in Table I. Remarkably, the IR for \mathcal{O} coincides with the paring symmetry of Cooper pairs. In other words, *helical MFs respond to magnetic fields in accord with the IR of Cooper pairs*. This notable property allows us to determine the pairing symmetry through the magnetic response of MFs.

We note an additional constraint $\Gamma_{\text{M}} \mathcal{O} \Gamma_{\text{M}}^\dagger = \mathcal{O}$ as Eq. (1) yields $\Gamma_{\text{M}} \rho^{(12)} \Gamma_{\text{M}}^\dagger = \rho^{(12)}$. This constraint im-




FIG. 2. (a) Surface states of half-Heusler superconductor in (111) plane. The red line and red areas indicate helical MFs with flat dispersion and the line-node-induced Majorana flat bands. $k_1 = \frac{1}{\sqrt{3}}(k_x + k_y + k_z)$, $k_2 = \frac{1}{\sqrt{2}}(k_x - k_y)$, and $k_3 = \frac{1}{\sqrt{6}}(k_x + k_y - 2k_z)$. (b) Energy gap of helical MF at $k_2 = k_3 = 0$ as a function of \mathbf{B} under the Zeeman magnetic field $\mu\mathbf{B} \cdot \mathbf{J}$.

plies $\mathcal{T}\mathcal{O}\mathcal{T}^{-1} = -\mathcal{O}$ as it holds that $\mathcal{C}\mathcal{O}\mathcal{C}^{-1} = -\mathcal{O}$ and $\tilde{g}_0\mathcal{O}\tilde{g}_0^{-1} = \mathcal{O}$. Therefore, \mathcal{O} should be a magnetic operator, as we expected. In Table I, we provide a representative set of basis for the gap function and the magnetic operator \mathcal{O} .

Majorana octupole in spin-3/2 superconductors. Table I reveals that MFs show a magnetic-octupole response $\propto J^3$ when the surface has C_{3v} or C_{6v} symmetry. This unique behavior is intrinsic to high-spin TSCs of spin-3/2 electrons: In fact, the relevant basis vanishes, $J_x^3 - J_x J_y J_y - J_y J_x J_y - J_y J_y J_x = J_y^3 - J_y J_x J_x - J_x J_y J_x - J_x J_x J_y = 0$ for spin-1/2 electrons ($J_i = \sigma_i/2$).

To confirm this result, we calculate the magnetic response of MFs in half-Heusler superconductors. In these compounds [36–41], a strong spin-orbit interaction (SOI) and high crystal symmetry provide a four-fold degenerate band at the Γ point, which is well described by spin-3/2 fermions [43]. Additionally, recent experiments have suggested the existence of parity-mixed superconductivity with line nodes [40, 41]. We show here that the parity-mixed superconductor exhibits a magnetic-octupole response. Consider the low-energy model with T_d symmetry [43]:

$$H_{\text{LK}}(\mathbf{k}) = \alpha \mathbf{k}^2 + \beta \sum_i k_i^2 J_i^2 + \gamma \sum_{i \neq j} k_i k_j J_i J_j + \delta \sum_i k_i (J_{i+1} J_i J_{i+1} - J_{i+2} J_i J_{i+2}), \quad (4)$$

where $i = x, y, z$ and $i + 1 = y$ if $i = x$, etc., and J_i are the 4×4 spin matrices of spin-3/2 fermions. Because inversion symmetry is absent, the Hamiltonian includes the antisymmetric SOI, which is proportional to δ and causes spin splitting at the Fermi surface. In their superconducting states, Cooper pairs form between spin-3/2 electrons, which allows quintet and septet pairings in addition to the conventional singlet and triplet pairings [43, 62, 63]. Furthermore, the antisymmetric SOI generally mixes the

parity of the gap function, so the even- and odd-parity components coexist in the gap function [64–68] and the odd-parity component is aligned with the antisymmetric SOI [65], providing the spin-septet pairing. Based on this insight, the gap function is given by the mixture of spin-singlet and spin-septet components, $\Delta(\mathbf{k}) = \Delta/\sqrt{1 + \eta^2}[\eta 1_4 + \sum_i k_i (J_{i+1} J_i J_{i+1} - J_{i+2} J_i J_{i+2})]e^{-i\pi J_y}$, even when we choose the conventional A_1 state of T_d , where η parametrizes the mixing between the spin-singlet and spin-septet components and 1_n is the $n \times n$ identity matrix.

The superconducting state hosts six line nodes encircling the k_x , k_y , and k_z axis, in analogy with other parity-mixed superconductors [60, 69–74]. Here, we focus on the (111) surface because the magnetic-octupole response requires C_{3v} symmetry. To verify the existence of helical MFs, we numerically diagonalize the BdG Hamiltonian with the surface normal to the [111] direction and find a helical MF with three flat dispersions (see Fig. S2 [61]), as schematically depicted in Fig. 2(a). Each flat dispersion lies on the mirror planes with mirror-reflection symmetries, σ , $C_3^\dagger \sigma C_3$, and $(C_3^\dagger)^2 \sigma (C_3)^2$, where σ is mirror-reflection with respect to the (110) plane and C_3 is three-fold rotation around the [111] direction. Using these mirror reflections, we obtain three Γ_M and the associated three w_{MID} , which protects zero modes on each flat dispersion. In particular, the three flat dispersions meet at a C_3 invariant point, at which the zero modes become simultaneous eigenstates of Γ_M and C_3 . To demonstrate magnetic response, we add a Zeeman magnetic term $\mu\mathbf{B} \cdot \mathbf{J}$ in Eq. (4), which leads to an anisotropic response with C_{3v} symmetry in Fig. 2(b). The Zeeman magnetic term contributes to the energy gap of the MFs on the order of $3\sqrt{2}\mu^3 B^3/32E_F^2$ (See Fig. S3 [61]), where μ is a coefficient of Zeeman term and E_F the Fermi energy, implying a magnetic-octupole response.

Several remarks are in order. (i) We also examine the magnetic response of a spin-1/2 TSC with C_{3v} symmetry. In sharp contrast to the spin-3/2 case, the helical MFs only show the dipole response (see the magnetic response of superconducting TIs in Fig. S1 [61]). (ii) The octupole response also appears in orbital magnetic effects since the orbital magnetic terms [75–80] also should be the same IR. (iii) Another high-spin superconductor of spin-3/2 electrons was recently proposed for antiperovskite materials with O_h group [28, 29]. We obtain a similar magnetic-octupole response of MFs on the (111) surface when its pairing symmetry is A_{2u} of O_h .

Conclusions. In this paper, we develop a theory of Majorana multipoles for 3D TR-invariant TSCs, which provide novel experimental means to identify bulk pairing symmetry and high-spin superconductivity. The Majorana multipoles may be observed through spin-sensitive measurements such as spatially resolved NMR measurements [81] or the surface tunneling spectroscopy under magnetic fields [75–79, 82].

This work was supported by the Grants-in-Aid for Scientific Research on Innovative Areas “Topological Material Science” (Grant Nos. JP15H05855, JP15H05851, JP15H05853, JP15K21717, and JP18H04224) from JSPS of Japan, and by JSPS KAKENHI Grant Nos. JP17H02922 and JP18H01176. S.K. was supported by the CREST project (JPMJCR16F2) from Japan Science and Technology Agency (JST), and the Building of Consortia for the Development of Human Resources in Science and Technology.

-
- [1] C.-R. Hu, Phys. Rev. Lett. **72**, 1526 (1994).
 [2] S. Kashiwaya and Y. Tanaka, Rep. Prog. Phys. **63**, 1641 (2000).
 [3] G. E. Volovik, *The Universe in a Helium Droplet* (Oxford University Press, Oxford, 2003).
 [4] A. P. Schnyder, S. Ryu, A. Furusaki, and A. W. W. Ludwig, Phys. Rev. B **78**, 195125 (2008).
 [5] M. Sato, Phys. Rev. B **79**, 214526 (2009).
 [6] F. Wilczek, Nature Phys. **5**, 614 (2009).
 [7] M. Z. Hasan and C. L. Kane, Rev. Mod. Phys. **82**, 3045 (2010).
 [8] X.-L. Qi and S.-C. Zhang, Rev. Mod. Phys. **83**, 1057 (2011).
 [9] Y. Tanaka, M. Sato, and N. Nagaosa, Journal of the Physical Society of Japan **81**, 011013 (2012).
 [10] J. Alicea, Rep. Prog. Phys. **75**, 076501 (2012).
 [11] M. Sato and S. Fujimoto, Journal of the Physical Society of Japan **85**, 072001 (2016).
 [12] T. Mizushima, Y. Tsutsumi, T. Kawakami, M. Sato, M. Ichioka, and K. Machida, Journal of the Physical Society of Japan **85**, 022001 (2016).
 [13] C.-K. Chiu, J. C. Y. Teo, A. P. Schnyder, and S. Ryu, Rev. Mod. Phys. **88**, 035005 (2016).
 [14] M. Sato and Y. Ando, Rep. Prog. Phys. **80**, 076501 (2017).
 [15] C. Nayak, S. H. Simon, A. Stern, M. Freedman, and S. Das Sarma, Rev. Mod. Phys. **80**, 1083 (2008).
 [16] Y. S. Hor, A. J. Williams, J. G. Checkelsky, P. Roushan, J. Seo, Q. Xu, H. W. Zandbergen, A. Yazdani, N. P. Ong, and R. J. Cava, Phys. Rev. Lett. **104**, 057001 (2010).
 [17] L. Fu and E. Berg, Phys. Rev. Lett. **105**, 097001 (2010).
 [18] S. Sasaki, M. Kriener, K. Segawa, K. Yada, Y. Tanaka, M. Sato, and Y. Ando, Phys. Rev. Lett. **107**, 217001 (2011).
 [19] S. Sasaki, Z. Ren, A. A. Taskin, K. Segawa, L. Fu, and Y. Ando, Phys. Rev. Lett. **109**, 217004 (2012).
 [20] T. Hashimoto, K. Yada, M. Sato, and Y. Tanaka, Phys. Rev. B **92**, 174527 (2015).
 [21] L. Fu, Phys. Rev. B **90**, 100509(R) (2014).
 [22] K. Matano, M. Kriener, K. Segawa, Y. Ando, and G.-q. Zheng, Nature Physics **12**, 852 (2016).
 [23] S. Yonezawa, K. Tajiri, S. Nakata, Y. Nagai, Z. Wang, K. Segawa, Y. Ando, and Y. Maeno, Nature Physics **13**, 123 (2016).
 [24] L. Aggarwal, A. Gaurav, G. S. Thakur, Z. Haque, A. K. Ganguli, and G. Sheet, Nature Materials **16**, 3237 (2015).
 [25] H. Wang, H. Wang, H. Liu, H. Lu, W. Yang, S. Jia, X.-J. Liu, X. C. Xie, J. Wei, and J. Wang, Nature Materials **15**, 3842 (2015).
 [26] S. Kobayashi and M. Sato, Phys. Rev. Lett. **115**, 187001 (2015).
 [27] T. Hashimoto, S. Kobayashi, Y. Tanaka, and M. Sato, Phys. Rev. B **94**, 014510 (2016).
 [28] M. Oudah, A. Ikeda, J. N. Hausmann, S. Yonezawa, T. Fukumoto, S. Kobayashi, M. Sato, and Y. Maeno, Nature Communications **7**, 13617 (2016).
 [29] T. Kawakami, T. Okamura, S. Kobayashi, and M. Sato, Phys. Rev. X **8**, 041026 (2018).
 [30] P. Zhang *et al.*, Nat. Phys. **15**, 41 (2019).
 [31] E. Majorana, Nuovo Cimento **14**, 171 (1937).
 [32] F. T. Avignone, S. R. Elliott, and J. Engel, Rev. Mod. Phys. **80**, 481 (2008).
 [33] B. Kayser and A. S. Goldhaber, Phys. Rev. D **28**, 2341 (1983).
 [34] E. E. Radescu, Phys. Rev. D **32**, 1266 (1985).
 [35] F. Boudjema, C. Hamzaoui, V. Rahal, and H. C. Ren, Phys. Rev. Lett. **62**, 852 (1989).
 [36] G. Goll, M. Marz, A. Hamann, T. Tomanic, K. Grube, T. Yoshino, and T. Takabatake, Physica B: Condensed Matter **403**, 1065 (2008).
 [37] N. P. Butch, P. Syers, K. Kirshenbaum, A. P. Hope, and J. Paglione, Phys. Rev. B **84**, 220504(R) (2011).
 [38] F. F. Tafti, T. Fujii, A. Juneau-Fecteau, S. René de Cotret, N. Doiron-Leyraud, A. Asamitsu, and L. Taillefer, Phys. Rev. B **87**, 184504 (2013).
 [39] G. Xu, W. Wang, X. Zhang, Y. Du, E. Liu, S. Wang, G. Wu, Z. Liu, and X. X. Zhang, Scientific Reports **4**, 5709 (2014).
 [40] T. V. Bay, T. Naka, Y. K. Huang, and A. de Visser, Phys. Rev. B **86**, 064515 (2012).
 [41] H. Kim, K. Wang, Y. Nakajima, R. Hu, S. Ziemak, P. Syers, L. Wang, H. Hodovanets, J. D. Denlinger, P. M. R. Brydon, D. F. Agterberg, M. A. Tanatar, R. Prozorov, and J. Paglione, Science Advances **4**, eaao4513 (2018).
 [42] I. Boettcher and I. F. Herbut, Phys. Rev. B **93**, 205138 (2016).
 [43] P. M. R. Brydon, L. Wang, M. Weinert, and D. F. Agterberg, Phys. Rev. Lett. **116**, 177001 (2016).
 [44] S. A. A. Ghorashi, S. Davis, and M. S. Foster, Phys. Rev. B **95**, 144503 (2017).
 [45] L. Savary, J. Ruhman, J. W. F. Venderbos, L. Fu, and P. A. Lee, Phys. Rev. B **96**, 214514 (2017).
 [46] C. Timm, A. P. Schnyder, D. F. Agterberg, and P. M. R. Brydon, Phys. Rev. B **96**, 094526 (2017).
 [47] J. W. F. Venderbos, L. Savary, J. Ruhman, P. A. Lee, and L. Fu, Phys. Rev. X **8**, 011029 (2018).
 [48] I. Boettcher and I. F. Herbut, Phys. Rev. Lett. **120**, 057002 (2018).
 [49] J. Yu and C.-X. Liu, Phys. Rev. B **98**, 104514 (2018).
 [50] I. Kuzmenko, T. Kuzmenko, Y. Avishai, and M. Sato, Phys. Rev. B **98**, 165139 (2018).
 [51] B. Roy, S. A. A. Ghorashi, M. S. Foster, and A. H. Nevidomskyy, Phys. Rev. B **99**, 054505 (2019).
 [52] P. G. Grinevich and G. E. Volovik, Journal of Low Temperature Physics **72**, 371 (1988).
 [53] M. Sato, Phys. Rev. B **81**, 220504(R) (2010).
 [54] In principal, a non-magnetic point group symmetry that is unbroken under a magnetic field may also provide an additional stability, but it occurs only for C_s . This case will be discussed elsewhere.

- [55] T. Mizushima, M. Sato, and K. Machida, *Phys. Rev. Lett.* **109**, 165301 (2012).
- [56] K. Shiozaki and M. Sato, *Phys. Rev. B* **90**, 165114 (2014).
- [57] E. Dumitrescu, J. D. Sau, and S. Tewari, *Phys. Rev. B* **90**, 245438 (2014).
- [58] B. Lu, K. Yada, M. Sato, and Y. Tanaka, *Phys. Rev. Lett.* **114**, 096804 (2015).
- [59] Y. Xiong, A. Yamakage, S. Kobayashi, M. Sato, and Y. Tanaka, *Crystals* **7**, 58 (2017).
- [60] M. Sato, Y. Tanaka, K. Yada, and T. Yokoyama, *Phys. Rev. B* **83**, 224511 (2011).
- [61] See the Supplemental Material at [URL will be inserted by publisher] for sections (S1) IRs of gap functions, (S2) IRs of $\rho^{(ab)}$, (S3) magnetic response of superconducting TIs, (S4) two-orbital model with magnetic octupole response, (S5) topological surface states of the A_1 state in half-Heusler superconductors, (S6) energy gap of magnetic octupole response, which includes Refs. [17, 21–23, 43, 60, 68, 83].
- [62] T.-L. Ho and S. Yip, *Phys. Rev. Lett.* **82**, 247 (1999).
- [63] W. Yang, Y. Li, and C. Wu, *Phys. Rev. Lett.* **117**, 075301 (2016).
- [64] L. P. Gor'kov and E. I. Rashba, *Phys. Rev. Lett.* **87**, 037004 (2001).
- [65] P. A. Frigeri, D. F. Agterberg, A. Koga, and M. Sigrist, *Phys. Rev. Lett.* **92**, 097001 (2004).
- [66] S. Fujimoto, *Journal of the Physical Society of Japan* **76**, 034712 (2007).
- [67] S. Fujimoto, *Journal of the Physical Society of Japan* **76**, 051008 (2007).
- [68] E. Bauer, M. Sigrist, and editors, *Non-Centrosymmetric Superconductors: Introduction and Overview* (Springer, Heidelberg, 2012).
- [69] Y. Tanaka, Y. Mizuno, T. Yokoyama, K. Yada, and M. Sato, *Phys. Rev. Lett.* **105**, 097002 (2010).
- [70] K. Yada, M. Sato, Y. Tanaka, and T. Yokoyama, *Phys. Rev. B* **83**, 064505 (2011).
- [71] P. M. R. Brydon, A. P. Schnyder, and C. Timm, *Phys. Rev. B* **84**, 020501(R) (2011).
- [72] A. P. Schnyder, P. M. R. Brydon, and C. Timm, *Phys. Rev. B* **85**, 024522 (2012).
- [73] S. Matsuura, P.-Y. Chang, A. P. Schnyder, and S. Ryu, *New Journal of Physics* **15**, 065001 (2013).
- [74] A. P. Schnyder and P. M. R. Brydon, *Journal of Physics: Condensed Matter* **27**, 243201 (2015).
- [75] Y. Tanaka and S. Kashiwaya, *Phys. Rev. Lett.* **74**, 3451 (1995).
- [76] Y. Tanaka, Y. Tanuma, K. Kuroki, and S. Kashiwaya, *Journal of the Physical Society of Japan* **71**, 2102 (2002).
- [77] Y. Tanuma, K. Kuroki, Y. Tanaka, R. Arita, S. Kashiwaya, and H. Aoki, *Phys. Rev. B* **66**, 094507 (2002).
- [78] Y. Tanaka, T. Yokoyama, A. V. Balatsky, and N. Nagosa, *Phys. Rev. B* **79**, 060505(R) (2009).
- [79] S. Tamura, S. Kobayashi, L. Bo, and Y. Tanaka, *Phys. Rev. B* **95**, 104511 (2017).
- [80] L. Chirulli and F. Guinea, *Phys. Rev. B* **98**, 094515 (2018).
- [81] S. B. Chung and S.-C. Zhang, *Phys. Rev. Lett.* **103**, 235301 (2009).
- [82] M. Fogelström, D. Rainer, and J. A. Sauls, *Phys. Rev. Lett.* **79**, 281 (1997).
- [83] S. Sumita and Y. Yanase, *Phys. Rev. B* **97**, 134512 (2018).