

This is the accepted manuscript made available via CHORUS. The article has been published as:

Macroscopic Thermodynamic Reversibility in Quantum Many-Body Systems

Philippe Faist, Takahiro Sagawa, Kohtaro Kato, Hiroshi Nagaoka, and Fernando G. S. L. Brandão

Phys. Rev. Lett. **123**, 250601 — Published 17 December 2019

DOI: [10.1103/PhysRevLett.123.250601](https://doi.org/10.1103/PhysRevLett.123.250601)

Macroscopic Thermodynamic Reversibility in Quantum Many-Body Systems

Philippe Faist,^{1,2} Takahiro Sagawa,³ Kohtaro Kato,¹ Hiroshi Nagaoka,⁴ and Fernando G. S. L. Brandão^{1,5}

¹*Institute for Quantum Information and Matter,*

California Institute of Technology, Pasadena, CA 91125, USA

²*Institute for Theoretical Physics, ETH Zurich, 8093 Switzerland*

³*Department of Applied Physics, The University of Tokyo, Tokyo 113-8656, Japan*

⁴*The University of Electro-Communications, Tokyo, 182-8585, Japan*

⁵*Google Inc., Venice, CA 90291, USA*

(Dated: October 14, 2019)

The resource theory of thermal operations, an established model for small-scale thermodynamics, provides an extension of equilibrium thermodynamics to nonequilibrium situations. On a lattice of any dimension with any translation-invariant local Hamiltonian, we identify a large set of translation-invariant states that can be reversibly converted to and from the thermal state with thermal operations and a small amount of coherence. These are the spatially ergodic states, i.e., states that have sharp statistics for any translation-invariant observable, and mixtures of such states with the same thermodynamic potential. As an intermediate result, we show for a general state that if the gap between the min- and the max-relative entropies to the thermal state is small, then the state can be approximately reversibly converted to and from the thermal state with thermal operations and a small source of coherence. Our proof provides a quantum version of the Shannon-McMillan-Breiman theorem for the relative entropy and a quantum Stein’s lemma for ergodic states and local Gibbs states. Our results provide a strong link between the abstract resource theory of thermodynamics and more realistic physical systems, as we achieve a robust and operational characterization of the emergence of a thermodynamic potential in translation-invariant lattice systems.

Introduction.—The quantum information approach to thermodynamics has allowed thermodynamic concepts, such as work, to be successfully extended into regimes of small-scale systems that store and process quantum information [1]. Notably, formulating thermodynamics as a resource theory [2–5] allows for a precise characterization of the resources that are required in single-instance state transformations, for instance thermodynamic work [6–8] and quantum coherence [9–12]. This is done by establishing a set of natural rules such as energy conservation, characterizing which possible evolutions a quantum state can undergo under these rules, and studying which external resources allows the system to undergo otherwise forbidden state transformations. A simple such framework is the resource theory of *thermal operations*, where one allows any energy-conserving unitary interaction with a heat bath at a fixed background temperature [4, 8, 13], and can be extended to more general types of reservoirs [14–17]. This approach has strong connections with information-theoretic entropy measures and quantum Shannon theory [18, 19]. More generally, information-theoretic approaches have provided new descriptions of nonequilibrium states and dynamics in statistical mechanics and thermodynamics, both in the classical and quantum regimes [20–24]. The resource theory connects to standard macroscopic thermodynamics in several ways. This approach is equivalent [25–27] to an established abstract and axiomatic formulation of thermodynamics by Lieb and Yngvason [28–31]. Second, one recovers the usual laws of thermodynamics in regimes of many identically and independently distributed (i.i.d.) copies of a state, such as for an ideal gas, or if the states

considered are quantum statistical ensembles [4, 8, 32–34].

The resource theory of thermodynamics extends equilibrium thermodynamics to non-equilibrium situations. In standard macroscopic thermodynamics, a system is defined to be in thermodynamic equilibrium if it no longer presents macroscopic changes or currents, and if it has lost memory of its initial, possibly non-equilibrium state [35]. The purpose of this definition is to ensure that the thermodynamic behavior of the system is entirely specified by a *thermodynamic potential*: The optimal work required to transform one equilibrium state into another by a reversible thermodynamic process is given by the difference of the potentials for the initial and final states, and does not depend on any further details of the process. In the resource theory, this can be verified directly: Is the amount of work required to transform a state A into a state B equal to the amount of work that can be extracted in the reverse process? If so, the resource theory is said to be *reversible*. Crucially, reversibility of a resource theory—i.e., the emergence of a thermodynamic potential—can happen for states that are not necessarily in thermodynamic equilibrium, as we show in this paper.

A natural question is whether the notion of resource-theoretic reversibility can be leveraged to show the emergence of a thermodynamic potential for new classes of states that are physically relevant (cf. also [36–41]), such as interacting particles on a lattice, which go beyond idealized macroscopic settings such as i.i.d. states.

Here, we show that on a translation-invariant lattice of any spatial dimension with a local Hamiltonian, all

ergodic states—i.e., states for which macroscopic quantities have sharply peaked statistics—can be reversibly converted to and from the thermal state. Furthermore, mixtures of ergodic states with the same thermodynamic potential also have this property. This ensures the emergence of a thermodynamic potential for this class of states even if some of these states are far out of equilibrium.

In the following, we first introduce the resource theory of thermodynamics and show that for general states, an equipartition property implies the emergence of a thermodynamic potential. We then consider translation-invariant lattices and explain our main result illustrated with an example of a 1-D Ising spin chain, before concluding with a discussion.

Resource theory of thermal operations.—In this resource theory, one is allowed to (i) bring in any ancilla systems in their thermal state, (ii) to carry out any energy-conserving unitaries, and (iii) to trace out any systems. We may then quantify the amount of work required to transform ρ into another state ρ' by including an explicit *battery system*, initialized in a pure energy eigenstate $|E\rangle$ and which we require to transition into another energy eigenstate $|E'\rangle$ at the end of the process. That is, if the transformation $\rho \otimes |E\rangle\langle E| \rightarrow \rho' \otimes |E'\rangle\langle E'|$ is possible with the operations (i)–(iii), then we define this process as consuming $E - E'$ work [8, 34, 42] (negative work consumption corresponds to work extraction).

We refer to the class of states which are block-diagonal in the energy eigenspaces as *semiclassical states*. For these states, transformations under thermal operations are fully characterized by *thermo-majorization* [8], a generalized notion of matrix majorization [43–45]. Let's consider two natural tasks associated with a semiclassical state ρ : *state formation* and *work distillation* (Fig. 1a). State formation consists in preparing the state ρ starting from the thermal state of the system, $\gamma = e^{-\beta H} / \text{tr}(e^{-\beta H})$. The optimal amount of work that needs to be invested, if we allow an inaccuracy $\epsilon > 0$ in the final state and if ρ is semiclassical, is [7, 8]

$$W_{\text{formation}}(\rho) = \beta^{-1} S_{\text{max}}^{\epsilon}(\rho \| \gamma), \quad (1)$$

with the *max-relative entropy* defined as $S_{\text{max}}^{\epsilon}(\rho \| \sigma) = \min_{\tilde{\rho} \approx_{\epsilon} \rho} \ln \|\sigma^{-1/2} \tilde{\rho} \sigma^{-1/2}\|_{\infty}$ with the optimization ranging over all states $\tilde{\rho}$ that are ϵ -close to ρ in trace distance [46]. On the other hand, *work distillation* consists in extracting as much work as possible from a given state ρ , resulting in the thermal state γ on the system. The optimal amount of work that can be extracted from a semiclassical state ρ is [7, 8]

$$W_{\text{distillable}}(\rho) = \beta^{-1} S_{\text{min}}^{\epsilon}(\rho \| \gamma), \quad (2)$$

with the *min-relative entropy* defined as $S_{\text{min}}^{\epsilon}(\rho \| \sigma) = \max_{\tilde{\rho} \approx_{\epsilon} \rho} \{-\ln \text{tr}(\Pi^{\tilde{\rho}} \sigma)\}$ where $\Pi^{\tilde{\rho}}$ is the projector onto the support of $\tilde{\rho}$ [46]. The min- and max-relative entropies are special cases of the Rényi relative entropies [47–49].




FIG. 1. A thermodynamic potential emerges when the underlying resource theory is reversible. **a.** For a state $\bar{\rho}$ that is block-diagonal in energy, the work that can be extracted is given by the min-relative entropy $W_{\text{dist.}} = \beta^{-1} S_{\text{min}}^{\epsilon}(\bar{\rho} \| \gamma)$, leaving the system in the thermal state $\gamma = e^{-\beta H} / \text{tr}(e^{-\beta H})$. Conversely, the work required to prepare $\bar{\rho}$ from the thermal state is $W_{\text{form.}} = \beta^{-1} S_{\text{max}}^{\epsilon}(\bar{\rho} \| \gamma)$. **b.** Suppose a state ρ (respectively ρ') can be reversibly converted to and from the thermal state with work $F(\rho) - F(\gamma)$ (respectively $F(\rho') - F(\gamma)$). Then ρ and ρ' can be reversibly interconverted. In this case the resource theory is said to be *reversible*, and the thermodynamic potential $F(\rho)$ fully characterizes the work required for state transformations. **c.** As an intermediate result, we show that if the min- and the max-relative entropies of any arbitrary quantum state ρ coincide approximately, then coherences in the state are suppressed, making it nearly block-diagonal in energy. The state is then approximately reversibly convertible to and from the thermal state with thermal operations and a small source of coherence.

There are no known necessary and sufficient conditions for transformations of arbitrary states under thermal operations. The reason is that thermal operations cannot generate any coherent superposition of energy levels, underscoring the role of time asymmetry in thermodynamics [9–12, 50–54]. It is thus necessary to account for coherence as a separate resource that enable operations that cannot be performed with thermal operations alone [55–61].

We resort to a very rudimentary way of accounting for coherence. We allow a system C with a bounded range of energy, which can be prepared in any pure state of our choosing and which we must dispose of in any state that is close to a pure state. This energy range is what we refer to as *amount of coherence* when such a system is used in a thermodynamic process. This crude approach is sufficient for our purposes, since our protocols only require such a system with an energy range that is negligibly small compared to the overall work cost of the transformation, thus forbidding any noticeable embezzling of work [13].

Emergence of a thermodynamic potential.—A resource theory is *reversible* for a class of states if the optimal work cost of any transition between two such states is equal to the optimal work extracted in the corresponding reverse process. This class of states then has a total or-

der, and we can assign a “thermodynamic value” to each state—this is the thermodynamic potential. A sufficient condition for reversibility is to check whether the work required for state formation can fully be recovered in the reverse task of work distillation [8]; any transformation between two such states is then reversible (Fig. 1b). In well-behaved cases, such as in the i.i.d. regime [4] or for statistical ensembles [27], the thermodynamic potential is given by the Kullback-Leibler divergence or Umegaki relative entropy $S(\rho \parallel \gamma)$, defined as

$$S(\rho \parallel \sigma) = \text{tr}(\rho(\ln \rho - \ln \sigma)) . \quad (3)$$

Equipartition implies reversibility with thermal operations.—We first present an intermediate result: If the min- and max-relative entropies coincide approximately, a condition which can be interpreted as a form of equipartition, then the state can approximately be reversibly converted to and from the thermal state (Fig. 1c). Our physical explanations are complemented by a fully rigorous proof that will be published elsewhere [62]. We refer to the Supplemental Material for a precise technical formulation of our main results [63].

Theorem I. *For any ρ and for $\epsilon > 0$, suppose that*

$$S_{\min}^{\epsilon}(\rho \parallel \gamma) \geq S - \Delta ; \quad S_{\max}^{\epsilon}(\rho \parallel \gamma) \leq S + \Delta ,$$

for some $S \in \mathbb{R}$, $\Delta > 0$. Then ρ can be approximately converted to and from the thermal state at a work cost (resp. work yield) of approximately $\beta^{-1}[S + O(\Delta)]$ (resp. $\beta^{-1}[S - O(\Delta)]$), with an amount of coherence of approximately $O(\Delta)$, and with arbitrarily good precision as $\epsilon \rightarrow 0$.

For a system of n particles, if we have $\Delta/n \rightarrow 0$ as $n \rightarrow \infty$, then the extractable work per system and the work of formation per system both converge to $s_{\infty}(\rho) := \lim_{n \rightarrow \infty} S/n$, and the amount of coherence used per copy goes to zero. In this case $s_{\infty}(\rho)$ becomes the thermodynamic potential in the thermodynamic limit $n \rightarrow \infty$.

To prove Theorem I, we first show that a state ρ for which the min-entropy and the max-entropy differ by at most $O(\Delta)$ have off-diagonal elements $\langle E_k | \rho | E_{k'} \rangle$ that are exponentially suppressed in $\beta|E_k - E_{k'}|$ if $\beta|E_k - E_{k'}| \gtrsim O(\Delta)$. In this sense, such a state may not harbor a large amount of coherence. Theorem I is then proven by exhibiting protocols for work distillation and state formation with the claimed properties. For both protocols, we first replace the Hamiltonian by one where the energy levels are integer multiples of some elementary spacing $O(\Delta)$, which can be done by investing an amount of coherence of order $O(\Delta)$. The work distillation protocol is then executed as follows. One dephases ρ in the new energy basis. Then we apply the known protocol for work extraction of semiclassical states. Because ρ has little coherence, the work that was wasted by the dephasing is small and the min-entropy does not




FIG. 2. Ergodic state on a lattice. An ergodic state is one that is translation-invariant and that produces sharp statistics for any translation-invariant observable. Our main result is to show that any two ergodic states can be reversibly inter-converted with thermal operations and a sublinear amount of coherence, with the reversible work cost deriving from a thermodynamic potential given by the Kullback-Leibler divergence. Furthermore, a translation-invariant state has a thermodynamic potential if and only if it is a mixture of ergodic states of equal potential, providing a robust and operational understanding of the emergence of a thermodynamic potential in lattice systems.

change by much, so we can still recover $S - O(\Delta)$ work. For the second protocol, we use the notion of an *internal reference frame*: The state ρ is equivalently described by a completely incoherent state $\tilde{\rho} = \mathcal{D}[\rho \otimes \eta]$, where η is a special state called a *reference frame*, and where $\mathcal{D}[\cdot]$ is the joint dephasing operation on the system and the reference frame [64, 65]. Because ρ has only little coherence, a small reference frame η suffices to achieve an accurate description of ρ . Our protocol consists in first preparing the incoherent state $\tilde{\rho}$ using the known protocol for semiclassical states, and then “shifting” the coherence from η to ρ , a process known as “externalizing” the reference frame [65].

Ergodic states on a lattice.—We now consider a d -dimensional square lattice with a local Hamiltonian that is translation-invariant:

$$H = \sum_{\mathbf{z} \in \mathbb{Z}^d} h_{\mathbf{z}} , \quad (4)$$

where each term $h_{\mathbf{z}}$ is a lattice-translated version of a term $h_{\mathbf{0}}$ that acts on a constant number of sites neighboring the origin. Each site is a quantum system of some finite dimension. Our calculations will be performed for finite lattice sizes, where the total number of sites is denoted by n . For finite n , the Hamiltonian is truncated at the boundary by ignoring any terms that have support outside of the finite region considered.

In statistical mechanics, thermodynamic behavior is often captured in the notion of *ergodicity* (Fig. 2). Ergodic states are defined on the infinite lattice in two equivalent ways [66–70]. First, they are exactly those states that self-average over space translations. I.e., an ergodic state ρ satisfies the following property: For any local observable $a_{\mathbf{0}}$, we have $\text{Var}_{\rho}(\frac{1}{n} \sum a_{\mathbf{z}}) \rightarrow 0$ as $n \rightarrow \infty$. Equivalently, ergodic states are the extremal points of the set of states that are translation-invariant on the infinite lattice. Consequently, any translation-invariant state can be written as a mixture of ergodic states.

Ergodic states are the natural quantum analogue of

classically ergodic probability distributions [71, 72] for spatial translations instead of time evolution. Examples of ergodic states include Gibbs states of a local Hamiltonian at sufficiently high temperature, where the correlation functions of local observables decay exponentially in space (see for example Ref. [73] and references therein). Also, any i.i.d. state is ergodic, being the Gibbs state of a noninteracting Hamiltonian. In contrast, a mixed state of macroscopically different sectors (e.g., different magnetization sectors in a symmetry-broken phase) is not ergodic, as spatial fluctuations do not vanish.

Ergodicity and reversibility under thermal operations.—Our main contribution is to prove that on a lattice of any dimension with a translation-invariant local Hamiltonian, all ergodic states fall into the setting of [Theorem I](#) and are thus reversibly interconvertible:

Theorem II. *In the thermodynamic limit $n \rightarrow \infty$, any two ergodic states can be reversibly converted into one another using thermal operations and a sublinear amount of coherence, and the corresponding reversible work cost rate is given by the thermodynamic potential*

$$s(\rho) = \lim_{n \rightarrow \infty} \frac{1}{n} S(\rho_n \parallel \gamma_n), \quad (5)$$

where ρ_n is the reduced state of ρ on a finite sublattice of size n and $\gamma_n = e^{-\beta H_n} / \text{tr}(e^{-\beta H_n})$ is the Gibbs state with the truncated Hamiltonian H_n on the sublattice.

[Theorem II](#) is a quantum version of the Shannon-McMillan-Breiman theorem for the relative entropy [71]. The proof of [Theorem II](#) proceeds via the *hypothesis testing relative entropy* [74–79], which interpolates between the min- and max-relative entropies [79] and can be formulated as a semidefinite program [80]. Inspired by the proof techniques of [69, 70, 81–84], we construct a quantum relative typical projector for an ergodic state relative to a Gibbs state associated with a truncated local Hamiltonian. This allows us to prove a generalized version of Stein’s lemma for hypothesis testing [69, 81, 85–87] from which it follows that the min- and max-relative entropies must coincide up to sublinear terms in n , and where the limiting value converges to $s(\rho)$. We are then in the setting of [Theorem I](#): Any ergodic state can be reversibly converted to and from the thermal state with the reversible work deriving from the thermodynamic potential $s(\rho)$. We refer to the Supplemental Material [63] for a precise statement of the theorem and how it connects with [Theorem I](#); rigorous proofs will be published elsewhere [62].

Translation-invariant states and reversibility.—We can further ask, is there a larger class of translation-invariant states on a lattice that can be reversibly converted to and from the thermal state? We provide an answer to this question as follows:

Theorem III. *A translation-invariant state ρ that is a mixture of a finite number of ergodic states is reversibly*

convertible to and from the thermal state if and only if all ergodic states in the mixture are of equal potential, i.e., $\rho = \sum p_k \rho^{(k)}$ with $s(\rho^{(1)}) = s(\rho^{(2)}) = \dots$.

By definition of ergodic states, any translation-invariant state is a possibly infinite mixture of ergodic states. For states that are finite mixtures, such as finitely correlated states [88], [Theorem III](#) provides a complete characterization of when a thermodynamic potential emerges.

To prove the above theorem, we note the following property of the min- and max-relative entropy for a mixture $\rho = \sum p_k \rho^{(k)}$:

$$S_{\min}^{\epsilon}(\rho_n \parallel \gamma_n) \approx \min_k S_{\min}^{\epsilon'}(\rho_n^{(k)} \parallel \gamma_n); \quad (6a)$$

$$S_{\max}^{\epsilon}(\rho_n \parallel \gamma_n) \approx \max_k S_{\max}^{\epsilon'}(\rho_n^{(k)} \parallel \gamma_n), \quad (6b)$$

with the approximation holding up to terms that do not scale with n and up to an adjustment of the smoothing parameter ϵ that does not depend on n . If all the $\rho^{(k)}$ in the decomposition have the same potential, $s(\rho^{(1)}) = s(\rho^{(2)}) = \dots$, then $S_{\min}^{\epsilon}(\rho_n \parallel \gamma_n) \approx S_{\max}^{\epsilon}(\rho_n \parallel \gamma_n)$ with equality in the thermodynamic limit, and we can apply [Theorem I](#). Conversely, if the $\rho^{(k)}$ do not all have the same potential, then the min- and max-relative entropies differ even in the thermodynamic limit. This implies that ρ cannot be reversibly convertible to and from the thermal state, because the min- and max-relative entropies are monotones under thermal operations.

Example: 1D Ising spin chain.—This toy example illustrates how a thermodynamic potential can emerge for states that are not in thermodynamic equilibrium. Consider a 1D chain of spin-1/2 particles with an Ising nearest-neighbor (n.n.) coupling and an external field h :

$$H = -J \sum_{i,j \text{ n.n.}} \sigma_z^i \sigma_z^j + h \sum_i \sigma_z^i, \quad (7)$$

where $\sigma_z = |\uparrow\rangle\langle\uparrow| - |\downarrow\rangle\langle\downarrow|$. Since i.i.d. states are ergodic, our results imply that two pure states of the form $|\psi\rangle^{\otimes n}, |\psi'\rangle^{\otimes n}$ can be converted into one another with thermal operations and an asymptotically negligible source of coherence at a reversible work cost of $F_{\psi} - F_{\psi'}$ per copy, where the thermodynamic potential is $F_{\psi} = \beta^{-1} \lim_{n \rightarrow \infty} S(\psi^{\otimes n} \parallel \gamma_n)/n$, which is the free energy per site up to an unimportant additive constant. The thermodynamic potential is well defined on an operational level even for states $\psi^{\otimes n}$ that are not in macroscopic equilibrium. Consider for instance the state $|\psi\rangle = |\uparrow\rangle = [|\uparrow\rangle + |\downarrow\rangle]/\sqrt{2}$. For $h \gg J$, the state $\psi^{\otimes n}$ presents macroscopic changes in the total spin along the X axis under time evolution according to H , but this does not prevent it from being reversibly convertible to and from another state $|\psi'\rangle^{\otimes n}$.

Discussion.—Our results provide a direct link between the abstract theory of thermodynamics at the small scale

formulated in terms of a resource theory, and realistic many-body systems that are commonly studied in statistical mechanics. In statistical mechanics, an ergodic state physically corresponds to a definite macroscopic state; it describes a pure thermodynamic phase without phase coexistence [66]. We endow these ergodic states with a stronger notion of thermodynamic behavior: The notion of reversibility associated with the resource theory—which extends the concept in equilibrium thermodynamics to nonequilibrium situations—is tightly related to the notion of ergodicity. Furthermore, our analysis underscores how reversibility in the resource theory does not imply equilibrium. Indeed, spatially ergodic states, as considered here, can evolve nontrivially in time as illustrated in the toy example above.

Our rigorous proof [62] makes use of advanced information-theoretic techniques, including the information spectrum [89–95], hypothesis testing and quantum Stein’s lemma [78, 79, 85, 86], as well as quantum typical projectors [69, 70, 81, 96]. Our results are an extension of the ergodic theorems of Refs. [69, 70]. We also use Ref. [97] to show that if we consider the reduced state of the infinite-dimensional Gibbs state instead of truncating the Hamiltonian for finite sublattices, then our results persist for sufficiently high temperatures where there is a unique KMS state.

Curiously, it is possible to construct toy situations in which the thermodynamic potential is not given by the Kullback-Leibler divergence [62]. While this does not happen in the setting considered in the present paper, it shows that the Kullback-Leibler divergence is not universally the correct expression of the emergent thermodynamic potential as defined via [Theorem I](#) when the min- and max-relative entropies converge to the same value. Whether this observation is relevant in physically interesting systems is an open question.

It seems plausible that our results could be robust to slight violations of translation invariance. For example, slight spatial inhomogeneity in a hydrodynamic mode could be allowed. Also, the results of [60, 61] would provide a more accurate analysis of the coherence requirements of the state transformations in [Theorem I](#). Furthermore, ergodic states exhibit some similarities with states obeying the eigenstate thermalization hypothesis [98–100], such as exponential decay of off-diagonal entries of the density matrix [62], suggesting that it might be possible to extend our techniques to such settings. We also note that a characterization of infinite or continuous mixtures of ergodic states is lacking, as opposed to the finite mixture considered in [Theorem III](#). Finally, one might hope that our methods can be extended to models exhibiting disorder, where a gap between the min- and max-relative entropies would characterize the irreversibility of conversions between many-body-localized states.

The authors are grateful to Matteo Lostaglio, Keiji Matsumoto, Yoshiko Ogata, David Sutter, and Hiroyasu Tajima for valuable discussions. TS is supported

by JSPS KAKENHI Grant Number JP16H02211 and JP19H05796. PhF is supported by the Institute for Quantum Information and Matter (IQIM) at Caltech which is a National Science Foundation (NSF) Physics Frontiers Center (NSF Grant PHY-1733907), by the Department of Energy Award DE-SC0018407, and by the Swiss National Science Foundation (SNSF) via the NCCR QSIT as well as project No. 200020_165843. KK acknowledge funding provided by the Institute for Quantum Information and Matter, an NSF Physics Frontiers Center (NSF Grant PHY-1733907). FB is supported by the NSF.

-
- [1] J. Goold, M. Huber, A. Riera, L. del Rio, and P. Skrzypczyk, “The role of quantum information in thermodynamics—a topical review,” *Journal of Physics A: Mathematical and Theoretical* **49**, 143001 (2016), [arXiv:1505.07835](#).
 - [2] D. Janzing, P. Wocjan, R. Zeier, R. Geiss, and T. Beth, “Thermodynamic cost of reliability and low temperatures: Tightening Landauer’s principle and the second law,” *International Journal of Theoretical Physics* **39**, 2717–2753 (2000), [arXiv:quant-ph/0002048](#).
 - [3] M. Horodecki, P. Horodecki, and J. Oppenheim, “Reversible transformations from pure to mixed states and the unique measure of information,” *Physical Review A* **67**, 062104 (2003), [arXiv:quant-ph/0212019](#).
 - [4] F. G. S. L. Brando, M. Horodecki, J. Oppenheim, J. M. Renes, and R. W. Spekkens, “Resource theory of quantum states out of thermal equilibrium,” *Physical Review Letters* **111**, 250404 (2013), [arXiv:1111.3882](#).
 - [5] E. Chitambar and G. Gour, “Quantum resource theories,” *Reviews of Modern Physics* **91**, 025001 (2019), [arXiv:1806.06107](#).
 - [6] O. C. O. Dahlsten, R. Renner, E. Rieper, and V. Vedral, “Inadequacy of von Neumann entropy for characterizing extractable work,” *New Journal of Physics* **13**, 53015 (2011), [arXiv:0908.0424](#).
 - [7] J. Berg, “Truly work-like work extraction via a single-shot analysis,” *Nature Communications* **4**, 1925 (2013), [arXiv:1110.6121](#).
 - [8] M. Horodecki and J. Oppenheim, “Fundamental limitations for quantum and nanoscale thermodynamics,” *Nature Communications* **4**, 2059 (2013), [arXiv:1111.3834](#).
 - [9] M. Lostaglio, D. Jennings, and T. Rudolph, “Description of quantum coherence in thermodynamic processes requires constraints beyond free energy,” *Nature communications* **6**, 6383 (2015), [arXiv:1405.2188](#).
 - [10] K. Korzekwa, M. Lostaglio, J. Oppenheim, and D. Jennings, “The extraction of work from quantum coherence,” *New Journal of Physics* **18**, 023045 (2016), [arXiv:1506.07875](#).
 - [11] G. Gour, D. Jennings, F. Buscemi, R. Duan, and I. Marvian, “Quantum majorization and a complete set of entropic conditions for quantum thermodynamics,” *Nature Communications* **9**, 5352 (2018), [arXiv:1708.04302](#).
 - [12] I. Marvian, “Coherence distillation machines are impossible in quantum thermodynamics,” (2018),

- arXiv:1805.01989.
- [13] F. Brando, M. Horodecki, N. Ng, J. Oppenheim, and S. Wehner, “The second laws of quantum thermodynamics,” *Proceedings of the National Academy of Sciences* **112**, 3275–3279 (2015), arXiv:1305.5278.
 - [14] N. Yunger Halpern and J. M. Renes, “Beyond heat baths: Generalized resource theories for small-scale thermodynamics,” *Physical Review E* **93**, 022126 (2016), arXiv:1409.3998.
 - [15] Y. Guryanova, S. Popescu, A. J. Short, R. Silva, and P. Skrzypczyk, “Thermodynamics of quantum systems with multiple conserved quantities,” *Nature Communications* **7**, 12049 (2016), arXiv:1512.01190.
 - [16] N. Yunger Halpern, P. Faist, J. Oppenheim, and A. Winter, “Microcanonical and resource-theoretic derivations of the thermal state of a quantum system with noncommuting charges,” *Nature Communications* **7**, 12051 (2016), arXiv:1512.01189.
 - [17] E. Hinds Mingo, Y. Guryanova, P. Faist, and D. Jennings, “Quantum thermodynamics with multiple conserved quantities,” in *Thermodynamics in the Quantum Regime*, edited by Felix Binder, Luis A. Correa, Christian Gogolin, Janet Anders, and Gerardo Adesso (Springer, Cham, Heidelberg, 2018) pp. 751–771, arXiv:1806.08325.
 - [18] P. Faist, F. Dupuis, J. Oppenheim, and R. Renner, “The minimal work cost of information processing,” *Nature Communications* **6**, 7669 (2015), arXiv:1211.1037.
 - [19] C. T. Chubb, M. Tomamichel, and K. Korzekwa, “Beyond the thermodynamic limit: finite-size corrections to state interconversion rates,” *Quantum* **2**, 108 (2018), arXiv:1711.01193.
 - [20] M. Esposito, U. Harbola, and S. Mukamel, “Nonequilibrium fluctuations, fluctuation theorems, and counting statistics in quantum systems,” *Reviews of Modern Physics* **81**, 1665–1702 (2009), arXiv:0811.3717.
 - [21] T. Sagawa, “Second law-like inequalities with quantum relative entropy: An introduction,” in *Lectures on Quantum Computing, Thermodynamics and Statistical Physics*, edited by Mikio Nakahara and Shu Tanaka (World Scientific, 2012) pp. 125–190, arXiv:1202.0983.
 - [22] U. Seifert, “Stochastic thermodynamics, fluctuation theorems and molecular machines,” *Reports on Progress in Physics* **75**, 126001 (2012), arXiv:1205.4176.
 - [23] J. M. R. Parrondo, J. M. Horowitz, and T. Sagawa, “Thermodynamics of information,” *Nature Physics* **11**, 131–139 (2015), arXiv:0903.2792.
 - [24] H. Tajima, E. Wakakuwa, and T. Ogawa, “Large deviation implies first and second laws of thermodynamics,” (2016), arXiv:1611.06614.
 - [25] M. Weilenmann, L. Kraemer, P. Faist, and R. Renner, “Axiomatic relation between thermodynamic and information-theoretic entropies,” *Physical Review Letters* **117**, 260601 (2016), arXiv:1501.06920.
 - [26] M. Weilenmann, *Quantum Causal Structure and Quantum Thermodynamics*, Ph.D. thesis, University of York (2017), arXiv:1807.06345.
 - [27] M. Weilenmann, L. Krmer, and R. Renner, “Smooth entropy in axiomatic thermodynamics,” in *Thermodynamics in the Quantum Regime*, edited by Felix Binder, Luis A. Correa, Christian Gogolin, Janet Anders, and Gerardo Adesso (Springer, Cham, Heidelberg, 2018) pp. 773–798, arXiv:1807.07583.
 - [28] E. H. Lieb and J. Yngvason, “The physics and mathematics of the second law of thermodynamics,” *Physics Reports* **310**, 1–96 (1999), arXiv:cond-mat/9708200.
 - [29] E. H. Lieb and J. Yngvason, “A guide to entropy and the second law of thermodynamics,” in *Statistical Mechanics*, edited by Bruno Nachtergaele, Jan Philip Solovej, and Jakob Yngvason (Springer Berlin Heidelberg, 2004) pp. 353–363, arXiv:math-ph/9805005.
 - [30] E. H. Lieb and J. Yngvason, “The entropy concept for non-equilibrium states,” *Proceedings of the Royal Society A* **469**, 20130408 (2013), arXiv:1305.3912.
 - [31] E. H. Lieb and J. Yngvason, “Entropy meters and the entropy of non-extensive systems,” *Proceedings of the Royal Society A* **470**, 20140192 (2014), arXiv:1403.0902.
 - [32] K. Matsumoto, “Reverse test and characterization of quantum relative entropy,” (2010), arXiv:1010.1030.
 - [33] Y. Jiao, E. Wakakuwa, and T. Ogawa, “Asymptotic convertibility of entanglement: An information-spectrum approach to entanglement concentration and dilution,” *Journal of Mathematical Physics* **59**, 022201 (2018), arXiv:1701.09050.
 - [34] P. Faist and R. Renner, “Fundamental work cost of quantum processes,” *Physical Review X* **8**, 021011 (2018), arXiv:1709.00506.
 - [35] H. B. Callen, *Thermodynamics and an Introduction to Thermostatistics* (Wiley, 1985).
 - [36] N. Datta, Y. Pautrat, and C. Rouz, “Second-order asymptotics for quantum hypothesis testing in settings beyond i.i.d.—quantum lattice systems and more,” *Journal of Mathematical Physics* **57**, 062207 (2016), arXiv:1510.04682.
 - [37] N. Yunger Halpern, “Toward physical realizations of thermodynamic resource theories,” in *Information and Interaction: Eddington, Wheeler, and the Limits of Knowledge*, The Frontiers Collection, edited by Ian T. Durham and Dean Rickles (Springer International Publishing, Cham, 2017) pp. 135–166, arXiv:1509.03873.
 - [38] . M. Alhambra, M. Lostaglio, and C. Perry, “Heat-bath algorithmic cooling with thermal operations,” (2018), arXiv:1807.07974.
 - [39] N. Yunger Halpern and D. T. Limmer, “Fundamental limitations on photoisomerization from thermodynamic resource theories,” (2018), arXiv:1811.06551.
 - [40] N. Yunger Halpern, “Beyond heat baths II: framework for generalized thermodynamic resource theories,” *Journal of Physics A: Mathematical and Theoretical* **51**, 094001 (2018), arXiv:1409.7845.
 - [41] F. Buscemi, D. Sutter, and M. Tomamichel, “An information-theoretic treatment of quantum dichotomies,” (2019), arXiv:1907.08539.
 - [42] P. Skrzypczyk, A. J. Short, and S. Popescu, “Work extraction and thermodynamics for individual quantum systems,” *Nature communications* **5**, 4185 (2014), arXiv:1307.1558.
 - [43] E. Ruch, R. Schraner, and T. H. Seligman, “The mixing distance,” *The Journal of Chemical Physics* **69**, 386 (1978).
 - [44] R. Bhatia, *Matrix Analysis* (Springer, New York, 1997) series Title: Graduate Texts in Mathematics.
 - [45] A. W. Marshall, I. Olkin, and B. C. Arnold, *Inequalities: Theory of Majorization and Its Applications* (Springer, New York, 2010).
 - [46] N. Datta, “Min- and max-relative entropies and a new entanglement monotone,” *IEEE Transactions on Information Theory* **55**, 2816–2826 (2009), arXiv:0803.2770.

- [47] A. Rnyi, “On measures of entropy and information,” in *Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability, Vol. 1: Contributions to the Theory of Statistics* (1960) pp. 547–561.
- [48] M. Tomamichel, *A Framework for Non-Asymptotic Quantum Information Theory*, Ph.D. thesis, ETH Zurich (2012), arXiv:1203.2142.
- [49] M. Tomamichel, *Quantum Information Processing with Finite Resources*, Vol. 5 (Springer International Publishing, Cham, 2016) series Title: SpringerBriefs in Mathematical Physics, arXiv:1504.00233.
- [50] J. Berg, “Catalytic coherence,” *Physical Review Letters* **113**, 150402 (2014), arXiv:1304.1060.
- [51] I. Marvian and R. W. Spekkens, “Modes of asymmetry: The application of harmonic analysis to symmetric quantum dynamics and quantum reference frames,” *Physical Review A* **90**, 062110 (2014), arXiv:1312.0680.
- [52] I. Marvian and R. W. Spekkens, “Extending noether’s theorem by quantifying the asymmetry of quantum states,” *Nature Communications* **5**, 3821 (2014), arXiv:1404.3236.
- [53] M. Lostaglio and M. P. Müller, “Coherence and asymmetry cannot be broadcast,” *Physical Review Letters* **123**, 020403 (2019), arXiv:1812.08214.
- [54] I. Marvian and R. W. Spekkens, “No-broadcasting theorem for quantum asymmetry and coherence and a trade-off relation for approximate broadcasting,” *Physical Review Letters* **123**, 020404 (2019), arXiv:1812.08766.
- [55] T. Baumgratz, M. Cramer, and M. B. Plenio, “Quantifying coherence,” *Physical Review Letters* **113**, 140401 (2014), arXiv:1311.0275.
- [56] A. Winter and D. Yang, “Operational resource theory of coherence,” *Physical Review Letters* **116**, 120404 (2016), arXiv:1506.07975.
- [57] K. Ben Dana, M. García Daz, M. Mejjaty, and A. Winter, “Resource theory of coherence: Beyond states,” *Physical Review A* **95**, 062327 (2017), arXiv:1704.03710.
- [58] M. G. Daz, K. Fang, X. Wang, M. Rosati, M. Skotiniotis, J. Calsamiglia, and A. Winter, “Using and reusing coherence to realize quantum processes,” *Quantum* **2**, 100 (2018), arXiv:1805.04045.
- [59] S. Popescu, A. B. Sainz, A. J. Short, and A. Winter, “Quantum reference frames and their applications to thermodynamics,” *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* **376**, 20180111 (2018), arXiv:1804.03730.
- [60] H. Tajima, N. Shiraishi, and K. Saito, “Uncertainty relations in implementation of unitary operations,” *Physical Review Letters* **121**, 110403 (2018), arXiv:1709.06920.
- [61] H. Tajima, N. Shiraishi, and K. Saito, “Coherence cost for violating conservation laws,” (2019), arXiv:1906.04076.
- [62] T. Sagawa, P. Faist, K. Kato, K. Matsumoto, H. Nagaoka, and F. G. S. L. Brando, “Asymptotic reversibility of thermal operations for interacting quantum spin systems via generalized quantum Stein’s lemma,” (2019), arXiv:1907.05650.
- [63] See Supplemental Material at [URL] for a precise technical statement of our main results.
- [64] S. D. Bartlett, T. Rudolph, and R. W. Spekkens, “Dialogue concerning two views on quantum coherence: Fac-tist and fictionist,” *International Journal of Quantum Information* **4**, 17–43 (2006), arXiv:quant-ph/0507214.
- [65] S. Bartlett, T. Rudolph, and R. Spekkens, “Reference frames, superselection rules, and quantum information,” *Reviews of Modern Physics* **79**, 555–609 (2007), arXiv:quant-ph/0610030.
- [66] D. Ruelle, *Statistical Mechanics: Rigorous Results* (World Scientific, 1999).
- [67] O. Bratteli and D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics I: C^* - and W^* -Algebras. Symmetry Groups. Decomposition of States* (Springer, Berlin, Heidelberg, 1987).
- [68] O. Bratteli and D. W. Robinson, *Operator Algebras and Quantum Statistical Mechanics II: Equilibrium States. Models in Quantum Statistical Mechanics* (Springer, Berlin, Heidelberg, 1981).
- [69] I. Bjelakovic and R. Siegmund-Schultze, “An ergodic theorem for the quantum relative entropy,” *Communications in Mathematical Physics* **247**, 697–712 (2004), arXiv:quant-ph/0306094.
- [70] I. Bjelakovi, T. Krger, R. Siegmund-Schultze, and A. Szkoła, “The Shannon-McMillan theorem for ergodic quantum lattice systems,” *Inventiones mathematicae* **155**, 203–222 (2004), arXiv:math/0207121.
- [71] T. M. Cover and J. A. Thomas, *Elements of Information Theory*, 2nd ed. (Wiley, 2006).
- [72] R. B. Israel, *Convexity in the theory of lattice gases* (Princeton University Press, 2015).
- [73] H. Tasaki, “On the local equivalence between the canonical and the microcanonical ensembles for quantum spin systems,” *Journal of Statistical Physics* **172**, 905–926 (2018), arXiv:1609.06983.
- [74] F. Buscemi and N. Datta, “The quantum capacity of channels with arbitrarily correlated noise,” *IEEE Transactions on Information Theory* **56**, 1447–1460 (2010), arXiv:0902.0158.
- [75] F. G. S. L. Brando and N. Datta, “One-shot rates for entanglement manipulation under non-entangling maps,” *IEEE Transactions on Information Theory* **57**, 1754–1760 (2011), arXiv:0905.2673.
- [76] L. Wang and R. Renner, “One-shot classical-quantum capacity and hypothesis testing,” *Physical Review Letters* **108**, 200501 (2012), arXiv:1007.5456.
- [77] W. Matthews and S. Wehner, “Finite blocklength converse bounds for quantum channels,” *IEEE Transactions on Information Theory* **60**, 7317–7329 (2014), arXiv:1210.4722.
- [78] M. Tomamichel and M. Hayashi, “A hierarchy of information quantities for finite block length analysis of quantum tasks,” *IEEE Transactions on Information Theory* **59**, 7693–7710 (2013), arXiv:1208.1478.
- [79] F. Dupuis, L. Kraemer, P. Faist, J. M. Renes, and R. Renner, “Generalized entropies,” in *XVIIth International Congress on Mathematical Physics* (2013) pp. 134–153, arXiv:1211.3141.
- [80] J. Watrous, “Semidefinite programs for completely bounded norms,” *Theory of Computing* **5**, 217–238 (2009), arXiv:0901.4709.
- [81] I. Bjelakovic and R. Siegmund-Schultze, “Quantum Stein’s lemma revisited, inequalities for quantum entropies, and a concavity theorem of Lieb,” (2003), arXiv:quant-ph/0307170.
- [82] I. Bjelakovi and A. Szkoła, “The data compression theorem for ergodic quantum information sources,” *Quantum Information Processing* **4**, 49–63 (2005),

- arXiv:quant-ph/0301043.
- [83] I. Bjelakovic, T. Krueger, R. Siegmund-Schultze, and A. Szkola, “Chained typical subspaces - a quantum version of Breiman’s theorem,” (2003), arXiv:quant-ph/0301177.
 - [84] Y. Ogata, “The Shannon-McMillan theorem for AF C^* -systems,” *Letters in Mathematical Physics* **103**, 1367–1376 (2013), arXiv:1303.2288.
 - [85] F. Hiai and D. Petz, “The proper formula for relative entropy and its asymptotics in quantum probability,” *Communications in Mathematical Physics* **143**, 99–114 (1991).
 - [86] H. Nagaoka and T. Ogawa, “Strong converse and Stein’s lemma in quantum hypothesis testing,” *IEEE Transactions on Information Theory* **46**, 2428–2433 (2000), arXiv:quant-ph/9906090.
 - [87] F. G. S. L. Brando and M. B. Plenio, “A generalization of quantum Stein’s lemma,” *Communications in Mathematical Physics* **295**, 791–828 (2010), arXiv:0904.0281.
 - [88] M. Fannes, B. Nachtergaele, and R. F. Werner, “Finitely correlated states on quantum spin chains,” *Communications in Mathematical Physics* **144**, 443–490 (1992).
 - [89] T. S. Han, *Information-Spectrum Methods in Information Theory*, Vol. 50 (Springer, Berlin, Heidelberg, 2003) series Title: Stochastic Modelling and Applied Probability.
 - [90] T. S. Han, “Hypothesis testing with the general source,” *IEEE Transactions on Information Theory* **46**, 2415–2427 (2000), arXiv:math/0004121.
 - [91] H. Nagaoka and M. Hayashi, “An information-spectrum approach to classical and quantum hypothesis testing for simple hypotheses,” *IEEE Transactions on Information Theory* **53**, 534–549 (2007), arXiv:quant-ph/0206185.
 - [92] N. Datta and R. Renner, “Smooth entropies and the quantum information spectrum,” *IEEE Transactions on Information Theory* **55**, 2807–2815 (2009), arXiv:0801.0282.
 - [93] G. Bowen and N. Datta, “Beyond i.i.d. in quantum information theory,” in *IEEE International Symposium on Information Theory (ISIT)* (IEEE, 2006) pp. 451–455, arXiv:quant-ph/0604013.
 - [94] G. Bowen and N. Datta, “Quantum coding theorems for arbitrary sources, channels and entanglement resources,” (2006), arXiv:quant-ph/0610003.
 - [95] B. Schoenmakers, J. Tjoelker, P. Tuijls, and E. Verbitskiy, “Smooth Rényi entropy of ergodic quantum information sources,” in *IEEE International Symposium on Information Theory (ISIT)* (IEEE, 2007) pp. 256–260, arXiv:0704.3504.
 - [96] M. M. Wilde, *Quantum Information Theory* (Cambridge University Press, 2013) arXiv:1106.1445.
 - [97] M. Lenci and L. Rey-Bellet, “Large deviations in quantum lattice systems: One-phase region,” *Journal of Statistical Physics* **119**, 715–746 (2005), arXiv:math-ph/0406065.
 - [98] M. Srednicki, “Chaos and quantum thermalization,” *Physical Review E* **50**, 888–901 (1994), arXiv:cond-mat/9403051.
 - [99] M. Rigol, V. Dunjko, and M. Olshanii, “Thermalization and its mechanism for generic isolated quantum systems,” *Nature* **452**, 854–858 (2008), arXiv:0708.1324.
 - [100] L. D’Alessio, Y. Kafri, A. Polkovnikov, and M. Rigol, “From quantum chaos and eigenstate thermalization to statistical mechanics and thermodynamics,” *Advances in Physics* **65**, 239–362 (2016).