



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Multiparticle Interactions for Ultracold Atoms in Optical Tweezers: Cyclic Ring-Exchange Terms

Annabelle Bohrdt, Ahmed Omran, Eugene Demler, Snir Gazit, and Fabian Grusdt

Phys. Rev. Lett. **124**, 073601 — Published 19 February 2020

DOI: [10.1103/PhysRevLett.124.073601](https://doi.org/10.1103/PhysRevLett.124.073601)

Multiparticle interactions for ultracold atoms in optical tweezers: Cyclic ring-exchange terms

Annabelle Bohrdt,^{1,2,*} Ahmed Omran,^{3,*} Eugene Demler,³ Snir Gazit,⁴ and Fabian Grusdt^{5,2,1,†}

¹*Department of Physics and Institute for Advanced Study,
Technical University of Munich, 85748 Garching, Germany*

²*Munich Center for Quantum Science and Technology (MCQST), Schellingstr. 4, D-80799 München, Germany*

³*Department of Physics, Harvard University, Cambridge, Massachusetts 02138, USA*

⁴*Racah Institute of Physics and The Fritz Haber Research Center for
Molecular Dynamics, The Hebrew University, Jerusalem 91904, Israel*

⁵*Department of Physics and Arnold Sommerfeld Center for Theoretical Physics (ASC),
Ludwig-Maximilians-Universität München, Theresienstr. 37, München D-80333, Germany*

(Dated: January 21, 2020)

Dominant multi-particle interactions can give rise to exotic physical phases with anyonic excitations and phase transitions without local order parameters. In spin systems with a global $SU(N)$ symmetry, cyclic ring-exchange couplings constitute the first higher-order interaction in this class. In this letter we propose a protocol how $SU(N)$ invariant multi-body interactions can be implemented in optical tweezer arrays. We utilize the flexibility to re-arrange the tweezer configuration on time scales short compared to the typical lifetimes, in combination with strong non-local Rydberg interactions. As a specific example we demonstrate how a chiral cyclic ring-exchange Hamiltonian can be implemented in a two-leg ladder geometry. We study its phase diagram using DMRG simulations and identify phases with dominant vector chirality, a ferromagnet, and an emergent spin-1 Haldane phase. We also discuss how the proposed protocol can be utilized to implement the strongly frustrated $J - Q$ model, a candidate for hosting a deconfined quantum critical point.

Introduction.— Ultracold atoms in optical lattices have become a versatile platform for performing analogue quantum simulations, with widely tunable interactions [1] and the ability to control the single-particle band structure [2–8]. Using atoms with permanent electric or magnetic dipole moments [9] or in Rydberg states [10] allows to study systems with long-range dipole-dipole or van-der Waals interactions, which can mimic the long-range Coulomb repulsion between electrons in a solid. These ingredients can be combined to study exotic phenomena in strongly correlated many-body systems, related for example to quantum magnetism [11–17] or the fractional quantum Hall effect [18–20]. Leveraging the capabilities of ultracold atoms, such experiments promise new insights for example to directly measure topological invariants [21–25] or image the quantum mechanical wavefunction with single-site resolution [26–31].

In this letter, we go beyond the two-body interactions realized so far and propose a general protocol to implement highly symmetric multiparticle interactions with ultracold atoms in optical tweezer arrays. Multiparticle interactions can lead to exotic ground states with intrinsic topological order [32, 33], with applications for quantum computation [34, 35], and they are an important ingredient for realizing lattice gauge theories [36–39] central to the quantum simulation of high-energy phenomena or deconfined quantum criticality [40, 41]. If these higher-order couplings possess additional symmetries, e.g. $SU(N)$ invariance in spin systems, models with

strong frustration can be realized whose ground states are strongly correlated quantum liquids.

In condensed matter systems multi-spin interactions of this type emerge from higher-order virtual processes [42], leading to corrections to the pairwise Heisenberg couplings of $SU(2)$ spins in a half-filled Hubbard model. These cyclic ring-exchange terms play a role in frustrated quantum magnets like solid ^3He [43] and possibly also for the phase diagram of high- T_c cuprate superconductors




FIG. 1. Proposed setup: $SU(2)$ -invariant chiral cyclic ring-exchange interactions can be realized by combining state-dependent lattices generated by optical tweezer arrays and strong Rydberg interactions with a central Rydberg-dressed control qubit (C). The auxiliary states $|\tau = 1\rangle|\sigma\rangle$ with $\sigma = \uparrow, \downarrow$ (orange) of the atoms on the sites of the plaquette are subject to a state-dependent tweezer potential which allows to permute them coherently around the center. Our protocol makes use of stroboscopic π pulses between the physical states $\tau = 0$ (green) and the auxiliary states $\tau = 1$, which only take place collectively on all sites and conditioned on the absence of a Rydberg excitation in the control atom.

* These authors contributed equally.

† Corresponding author email: fabian.grusdt@lmu.de

[44, 45]. In this letter we demonstrate how such multi-spin interactions can be realized and independently tuned in ultracold atom systems without resorting to high-order virtual processes.

A promising route to implementing multiparticle processes is to use strong interactions between atoms in different Rydberg states representing spin degrees of freedom. This allows to build a versatile quantum simulator which can be used to realize ring-exchange interactions in spin systems by representing them as sums of products of Pauli matrices [46], or to implement local constraints giving rise to emergent dynamical gauge fields [47, 48].

Here we follow a similar strategy but propose to combine strong Rydberg interactions with the capabilities to quickly change the spatial configuration of atoms trapped in optical tweezer arrays [49–53]. We consider general lattice models with one N -component particle per lattice site (fermionic or bosonic) and show, as an explicit example, how a general class of $SU(N)$ -invariant chiral cyclic ring-exchange (CCR) interactions can be realized. They are described by a Hamiltonian ($\hbar = 1$)

$$\hat{H}_{\text{CCR}}(\phi) = K \sum_p (e^{i\phi} \hat{P}_p + e^{-i\phi} \hat{P}_p^\dagger). \quad (1)$$

where the sum is over all plaquettes p of the underlying lattice, the operator \hat{P}_p^\dagger (\hat{P}_p) cyclically permutes the spin configuration on plaquette p in clockwise (counterclockwise) direction and ϕ is a tunable complex phase. A generalization to finite hole doping, with zero or one particle per lattice site, is straightforward.

Non-chiral cyclic ring-exchange interactions, realized by Eq. (1) for $\phi = 0$, are believed to play a role in the high- T_c cuprate compounds. These materials can be described by the 2D Fermi-Hubbard model on a square lattice, with weak couplings between multiple layers in z -direction [54]. For the relevant on-site interactions U , which dominate over the nearest-neighbor tunneling $t \ll U$, this model can be simplified by an expansion in powers of t/U . To lowest order, one obtains a t - J model [55] with nearest-neighbor spin-exchange interactions of strength $J = 4t^2/U$. Next to leading order, cyclic ring-exchange terms on the plaquettes of the square lattice contribute with strength $K = 20t^4/U^3$. By comparison of first principle calculations and measurements in the high-temperature regime it was shown that $K \approx 0.13 \times J$ in La_2CuO_4 [56] but its effect on the phase diagram remains debated. In ultracold atoms, similar higher-order processes have been used to realize non-chiral cyclic ring-exchange couplings [57, 58].

We start by explaining the general scheme using the example of CCR interactions. Our method is more versatile however, and we discuss how it can be adapted to implement the J - Q model which has been proposed as a candidate system realizing deconfined quantum criticality [40, 41]. We also analyze the phase diagram of the CCR Hamiltonian (1) in a ladder geometry, with exactly one $SU(2)$ spin per lattice site. We show that the phase diagram contains a gapped Haldane phase with topologi-

cally protected edge states [59–61] at intermediate values of $\pi/4 \lesssim \phi \lesssim 3\pi/4$, a ferromagnetic phase for $\phi \gtrsim 3\pi/4$ and a dominant vector chirality for $\phi \lesssim \pi/4$.

Implementation.— For simplicity we consider a single plaquette, restrict ourselves to $N_p = 4$ sites and assume $SU(2)$ symmetry, see Fig. 1. Generalizations of our scheme to more than one plaquette, $N_p \neq 4$ and $SU(N)$ symmetry are possible, see supplements (SM), which includes Refs. [69, 70].

Each of the four sites, labeled $j = 1, \dots, 4$, consists of a static optical tweezer trapping a single atom, where recently demonstrated rearrangement methods [49–53] allow for populating each site with high fidelity. We assume that the atoms remain in the vibrational ground states of the microtraps throughout the sequence. Every atom has two internal states $\sigma = \uparrow, \downarrow$ which we use to implement an effective spin-1/2 system. As a specific configuration we suggest to use ^{133}Cs atoms and utilize their $F = 3$, $m_F = 2, 3$ hyperfine states to represent the two spins. Optical pumping with site-resolved addressing can then be employed to prepare arbitrary initial spin patterns [58] and study their dynamics under Eq. (1).

The key ingredient for our proposed implementation of CCR interactions is to realize collective permutations of the entire spin configuration in the plaquette. This can be achieved by physically rotating the tweezer array around the center of the plaquette while ensuring that the motional and spin states of the atoms are preserved and coherence is not lost. The effect of clockwise rotations of the microtraps on the spin states is described by the operator \hat{P} ,

$$\hat{P}|\sigma_1, \sigma_2, \sigma_3, \sigma_4\rangle = |\sigma_4, \sigma_1, \sigma_2, \sigma_3\rangle. \quad (2)$$

Optimized trajectories can be chosen to cancel heating effects from the motion [62]. These require a timescale set by the quantum speed limit that scales as the inverse energy gap of each traps $t_{\text{rot}} \sim 1/\Delta\varepsilon$. For deep trapping potentials where $\Delta\varepsilon \approx 150$ kHz, rotation times of $t_{\text{rot}} < 10 \mu\text{s}$ are achievable.

In contrast to Eq. (2), the effective Hamiltonian leads to a superposition of permuted and non-permuted states in every infinitesimal time step Δt , as can be seen from a Taylor expansion: $e^{-i\hat{H}_{\text{CCR}}\Delta t} = 1 - i\hat{H}_{\text{CCR}}\Delta t$. To create such superposition states in our time evolution, we assume that every atom has a second internal degree of freedom labeled by $\tau = 0, 1$. Concretely we propose to realize the new states $|\tau = 1\rangle|\sigma\rangle$ in ^{133}Cs atoms by $F = 4$, $m_F = 3, 4$ hyperfine levels, where $m_F = 3$ ($m_F = 4$) corresponds to $\sigma = \downarrow$ ($\sigma = \uparrow$). These additional levels will be used as auxiliary states, whereas the states $|\tau = 0\rangle|\sigma\rangle$ introduced before – implemented as $F = 3$, $m_F = 2, 3$ levels in ^{133}Cs – realize the physical spin states.

One part of our protocol consists of a permutation of the spins σ , but only in the manifold of auxiliary states. This step requires a total time t_{rot} and can be described by the unitary transformations

$$\hat{U}_+ = \prod_j |1\rangle_j \langle 1| \otimes \hat{P} + \prod_j |0\rangle_j \langle 0| \otimes \hat{1}_\sigma, \quad \hat{U}_- = \hat{U}_+^\dagger \quad (3)$$

To implement this evolution, two sets of optical tweezer arrays can be used, of which only one is rotating. We suggest to realize it by the near-magic wavelength $\lambda_{\text{magic}} \approx 871.6 \text{ nm}$ in ^{133}Cs which strongly confines atoms in the state $\tau = 1$ but almost does not affect atoms in $\tau = 0$. By applying \hat{U}_{\pm} to superposition states with either all atoms in $\tau = 1$ or all atoms in $\tau = 0$, one can realize the desired superpositions of permuted and non-permuted spin configurations. Such states can be realized by collective π -pulses conditioned upon a control qubit trapped in the center of the plaquette [63], as described next.

If the control atom is in the state $|+\rangle_c$ it is transferred to a Rydberg state $|r\rangle_c$ with a resonant π -pulse and Rabi frequency Ω_r , see Fig. 1. If the control atom is in state $|-\rangle_c$, the laser Ω_r is off-resonant and no Rydberg excitation is created. Next a Raman transition by lasers $\Omega_j^{(1)}, \Omega_j^{(2)}$ through an intermediate Rydberg state $|r\rangle_j$ is used to implement a π -pulse transferring the physical states $|0\rangle_j$ to $|1\rangle_j$, without changing their spin state $|\sigma\rangle_j$. In the presence of a coupling field Ω^{EIT} that establishes two-photon resonance to the Rydberg state with each Raman laser, electromagnetically induced transparency (EIT) [64] suppresses the transition $|0\rangle_j \leftrightarrow |1\rangle_j$. However, the EIT condition is lifted by the Rydberg blockade mechanism if the control atom is in the Rydberg state $|r\rangle_c$ [63], enabling the transfer. After the transfer is complete, another π -pulse by Ω_r is applied to the control atom. This ensures that the control atom remains trapped during the protocol, even if the Rydberg excited state is not subject to a trapping potential. In summary, this part is described by the unitary transformation

$$\hat{U}_{\text{sw}} = |+\rangle_c \langle +| \otimes \left(\prod_j |1\rangle_j \langle 0| + \text{h.c.} \right) \otimes \hat{\mathbf{1}}_{\sigma} + |-\rangle_c \langle -| \otimes \hat{\mathbf{1}}_{\tau} \otimes \hat{\mathbf{1}}_{\sigma}. \quad (4)$$

The total time required to implement this switch (sw) is denoted by t_{sw} .

Finally, we need to introduce quantum dynamics between the states of the control atom. This can be realized by a dressing laser Ω_c driving transitions between $|\pm\rangle_c$, at a detuning Δ_c . These dynamics take place over a period of time t_c and are described by the unitary evolution $\hat{U}_c = e^{-i\hat{\mathcal{H}}_c t_c}$ with $\hat{\mathcal{H}}_c = \Delta_c |+\rangle_c \langle +| + \Omega_c (|+\rangle_c \langle -| + \text{h.c.})$. During the remaining steps of the protocol, Eqs. (3) - (4), we assume that $\Omega_c = 0$ is off and the control atom picks up a phase $\pm\varphi_c$ if it is in state $|+\rangle_c$. This phase can be adjusted by the detuning Δ_c and the duration $t_{\text{rx}} = 2t_{\text{sw}} + t_{\text{rot}}$, during which the time evolution of the control is $\hat{U}_{\pm\varphi_c} = |+\rangle_c \langle +| e^{\mp i\varphi_c} + |-\rangle_c \langle -|$.

The complete protocol is summarized in Fig. 2. It consists of a periodic repetition of the individual steps described above. At the discrete time steps nT , where $T = 2(t_c + t_{\text{rx}})$, the unitary evolution is described by an effective Hamiltonian $\hat{\mathcal{H}}_{\text{eff}}$:

$$e^{-inT\hat{\mathcal{H}}_{\text{eff}}} = (\hat{U}_T)^n = (\hat{U}_{\text{rx},+} \hat{U}_c \hat{U}_{\text{rx},-} \hat{U}_c)^n, \quad (5)$$




FIG. 2. Proposed protocol: The sequence in (a) is repeated periodically with period $T = 2(t_c + t_{\text{rx}})$. When $t_c \ll 2\pi/\Delta_c$, $1/\Omega_c$ it implements a trotterized time evolution of the effective Hamiltonian Eq. (8), which realizes CCR couplings when $\Delta_c \gg \Omega_c$. The individual time steps are illustrated in (b).

where we defined $\hat{U}_{\text{rx},\pm} = \hat{U}_{\text{sw}} (\hat{U}_{\pm\varphi_c} \otimes \hat{U}_{\pm}) \hat{U}_{\text{sw}}$. As will be shown below, $\hat{\mathcal{H}}_{\text{eff}}$ realizes CCR interactions with a tunable phase $\phi = -\varphi_c$ and amplitude

$$K = -\frac{1}{2T} (t_c \Delta_c) \left(\frac{\Omega_c}{\Delta_c} \right)^2 \quad (6)$$

provided that

$$t_c \ll 2\pi/\Delta_c, \quad \Omega_c \ll \Delta_c. \quad (7)$$

Now we estimate the strength $|K|$ of the CCR interactions that can be achieved with the proposed setup. To satisfy Eq. (7) we assume $\Omega_c = 0.2\Delta_c$ and $t_c \Delta_c = 0.4$. For a rotation time $t_{\text{rot}} = 10\mu\text{s}$ and assuming $t_{\text{sw}}, t_c \ll t_{\text{rot}}$ a reasonable strength of $K/\hbar = 50\text{Hz} \times 2\pi$ can be achieved. This requires $\Omega_c/2\pi \gg 1.3\text{kHz}$, which can be easily realized; the condition $t_{\text{sw}} \ll 10\mu\text{s}$ can also be met, as the Rydberg π -pulses on the control atom can be executed in $\sim 100 \text{ ns}$ each and the Raman transfer between the states $|0\rangle_j$ and $|1\rangle_j$ can be driven with coupling strengths above 1 MHz.

Effective Hamiltonian.— Next we show that our protocol realizes the Hamiltonian in Eq. (1). When $t_c/2\pi \ll 1/\Delta_c, 1/\Omega_c$, we can write $\hat{U}_c = 1 - i\hat{\mathcal{H}}_c t_c$ and calculate $\exp[-i\hat{\mathcal{H}}_{\text{eff}} T]$ to leading order in t_c . Eqs. (3) - (4) yield

$$\hat{\mathcal{H}}_{\text{eff}} = \frac{t_c}{T} \left\{ 2\Delta_c |+\rangle_c \langle +| + \Omega_c \left[|-\rangle_c \langle +| \left(1 + e^{i\varphi_c} \hat{P}^\dagger \right) + \text{h.c.} \right] \right\}. \quad (8)$$

When $\Omega_c \ll \Delta_c$ we can eliminate the state $|+\rangle_c$ which is only virtually excited. This further simplifies the effective Hamiltonian and we obtain

$$\hat{\mathcal{H}}_{\text{eff}} = K \left(2 + e^{-i\varphi_c} \hat{P} + e^{i\varphi_c} \hat{P}^\dagger \right). \quad (9)$$

Up to the energy shift $2K$ this realizes CCR interactions in an isolated plaquette. The result can be extended to




FIG. 3. Phase diagram of the CCR Hamiltonian on a ladder, obtained from DMRG in a system with 64 sites: different observables are evaluated in the ground state of the Hamiltonian (1) to characterize the phases. Upon varying ϕ , three different phases can be identified: A topological Haldane phase featuring a vanishing gap in the entanglement spectrum (a) and edge states with a non-zero local magnetization for $S_{\text{tot}}^z = 1$ (b); A symmetry-broken phase around $\phi = \pi$ with long-range ferromagnetic correlations (c); And a symmetric phase for small ϕ , where the staggered vector chirality remains non-vanishing over long distances (d).

multiple plaquettes by implementing the trotterized time step \hat{U}_T interchangeably on inequivalent plaquettes.

Two-leg ladder with CCR.— Now we discuss the physics of the SU(2) CCR Hamiltonian in a two-leg ladder. We vary the phase ϕ in the Hamiltonian (1) with $K = 1$ and calculate the ground state phase diagram using the density-matrix renormalization group (DMRG). For $\phi = \pi$, the ground state has ferromagnetic correlations, see Fig. 3 (c). It can be readily seen that the variational energy $\langle \hat{\mathcal{H}}_{\text{CCR}}(\pi) \rangle$ is minimized for ferromagnetic configurations. In the sector $S_{\text{tot}}^z = 0$ used in our DMRG in Fig. 3 (c), we find phase separation with two ferromagnetic domains of opposite magnetization.

At intermediate ϕ we find an emergent Haldane phase, with two-fold degenerate states in the entanglement spectrum, see Fig. 3 (a). For a finite $S_{\text{tot}}^z = 1$ the expectation value $\langle \hat{S}_{L,1}^z \rangle$ at the edge is non-zero, see Fig. 3 (b). The spin gap $\Delta E_S = E_{0,S=1} - E_{0,S=0}$, defined as the difference between the ground state energy with and without finite total magnetization, is zero in this phase, since the additional spin can be placed in the spin-1/2 topological edge states of the system without increasing the total energy. We corroborate this picture further by considering the $K - K'$ model with alternating strengths K, K' of the CCR interactions on adjacent plaquettes. In the SM

we provide an explicitly derivation of a spin-1 model with a gapped Haldane ground state [60, 65] for $\phi = \pi/2$ and $K' \ll K$.

For small values of ϕ , the ground state of the CCR Hamiltonian is a dominant vector chirality phase, as discussed in Ref. [66]. This phase is characterized by correlations of the form $\hat{S}_{x,y} \times \hat{S}_{x',y'}$ in a staggered arrangement around each plaquette. We find that the staggered correlation between different rungs, measured from the center $L/2$ of the chain,

$$(-1)^x \left\langle \left(\hat{S}_{L/2,1} \times \hat{S}_{L/2,2} \right) \cdot \left(\hat{S}_{L/2+x,1} \times \hat{S}_{L/2+x,2} \right) \right\rangle, \quad (10)$$

decays slowly as a function of the distance x and retains significant non-zero values over the considered system sizes, see Fig. 3 (d). The transition between the dominant vector chirality and Haldane phases is a symmetry-protected topological (SPT) phase transition.

Using the global SU(2) symmetry, the staggered vector chirality becomes $6 \left\langle \hat{S}_{L/2,1}^x \hat{S}_{L/2,2}^y \left(\hat{S}_{L/2+x,1}^x \hat{S}_{L/2+x,2}^y - \hat{S}_{L/2+x,1}^y \hat{S}_{L/2+x,2}^x \right) \right\rangle (-1)^x$. Measuring it requires access to two four-point functions of the form $\langle \hat{S}_i^\mu \hat{S}_j^\nu \hat{S}_k^\lambda \hat{S}_l^\rho \rangle$ which can be detected by making use of local addressing techniques, see e.g. [67]. To detect the Haldane phase experimentally, we propose to study weakly magnetized systems and image the topological edge states. Alternatively, one could work in the plaquette basis (see SM) and measure the Haldane string order parameter. An interesting future extension would be to use machine learning techniques to retrieve non-local order parameters from a series of quantum projective measurements.

Summary and Outlook.— In summary, we propose a general method for implementing multi-body interactions in ultracold atom experiments using optical tweezer arrays. The approach is particularly useful in the presence of additional, e.g. global SU(N) spin, symmetries. Specifically, we consider a four-body cyclic ring exchange term, which can be realized with a combination of multi-qubit gates based on Rydberg states and movable optical tweezers. We numerically study the ground state of the cyclic ring exchange Hamiltonian and find different dominant correlation functions as the complex phase of the ring exchange term is varied.

Our work paves the way for future studies of the interplay between ring-exchange and pair-exchange terms, as discussed in Ref. [68] for the non-chiral case $\phi = 0$. In the experimental realization proposed here, it is conceptually straightforward to introduce mobile holes into the system, leading to a finite doping. The interplay between spin and charge degrees of freedom could be further studied by adding direct tunneling terms, which lead to rich Hamiltonians in the spirit of $t - J$ like models. The physics of this type of model is completely unknown and provides an exciting prospect for future theoretical and experimental research. The proposed protocol is versatile enough to implement larger classes of models with

multi-spin interactions, such as the $J - Q$ model [41] (as we discuss in the SM). In two dimensions, this model features a phase transition between an antiferromagnet and a valence-bond solid, which has been proposed as a candidate for a deconfined quantum critical point [41]. Moreover, the experimental protocol can be varied to study different types of problems, such as discrete time-evolutions of complex models or impurity models, which can be realized by an inclusion of the control qubits into the models.

Acknowledgements. – The authors would like to thank T. Calarco, M. Endres, M. Greiner, A. Kaufman, M. Knap and M. Lukin for fruitful discussions. A.B. and F.G. acknowledge support from the Technical University of Munich - Institute for Advanced Study, funded by the German Excellence Initiative and the European

Union FP7 under grant agreement 291763, the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy–EXC-2111–390814868, DFG grant No. KN1254/1-1, DFG TRR80 (Project F8). A.B. acknowledges support by the Studienstiftung des deutschen Volkes. A.O. acknowledges support by a research fellowship from the German Research Foundation (DFG). E.D. acknowledges support by Harvard-MIT CUA, AFOSR-MURI Quantum Phases of Matter (grant FA9550-14-1-0035), AFOSR-MURI: Photonic Quantum Matter (award FA95501610323), DARPA DRINQS program (award D18AC00014). S.G. acknowledges support from the Israel Science Foundation, Grant No. 1686/18. F.G. acknowledges support by the Gordon and Betty Moore foundation through the EPiQS program.

-
- [1] Immanuel Bloch, Jean Dalibard, and Wilhelm Zwerger. Many-body physics with ultracold gases. *Reviews of Modern Physics*, 80(3):885–964, 2008.
- [2] Leticia Tarruell, Daniel Greif, Thomas Uehlinger, Gregor Jotzu, and Tilman Esslinger. Creating, moving and merging dirac points with a fermi gas in a tunable honeycomb lattice. *Nature*, 483(7389):302–U91, March 2012.
- [3] Y.-. J. Lin, K. Jimenez-Garcia, and I. B. Spielman. Spin-orbit-coupled bose-einstein condensates. *Nature*, 471(7336):83–U99, March 2011.
- [4] M. Aidelsburger, M. Atala, S. Nascimbene, S. Trotzky, Y.-. A. Chen, and I. Bloch. Experimental realization of strong effective magnetic fields in an optical lattice. *Physical Review Letters*, 107(25):255301, December 2011.
- [5] Lawrence W. Cheuk, Ariel T. Sommer, Zoran Hadzibabic, Tarik Yefsah, Waseem S. Bakr, and Martin W. Zwierlein. Spin-injection spectroscopy of a spin-orbit coupled fermi gas. *PRL*, 109:095302, 2012.
- [6] J. Struck, C. Ölschläger, M. Weinberg, P. Hauke, J. Simonet, A. Eckardt, M. Lewenstein, K. Sengstock, and P. Windpassinger. Tunable gauge potential for neutral and spinless particles in driven optical lattices. *Phys. Rev. Lett.*, 108:225304, May 2012.
- [7] Junru Li, Wujie Huang, Boris Shteynas, Sean Burchesky, Furkan Cagri Top, Edward Su, Jeongwon Lee, Alan O. Jamison, and Wolfgang Ketterle. Spin-orbit coupling and spin textures in optical superlattices. *Phys. Rev. Lett.*, 117:185301, Oct 2016.
- [8] Zhan Wu, Long Zhang, Wei Sun, Xiao-Tian Xu, Bao-Zong Wang, Si-Cong Ji, Youjin Deng, Shuai Chen, Xiong-Jun Liu, and Jian-Wei Pan. Realization of two-dimensional spin-orbit coupling for bose-einstein condensates. *Science*, 354(6308):83–88, 2016.
- [9] T. Lahaye, C. Menotti, Santos L., M. Lewenstein, and T. Pfau. The physics of dipolar bosonic quantum gases. *Reports On Progress In Physics*, 72(12):126401, 2009.
- [10] M. Saffman, T. G. Walker, and K. Molmer. Quantum information with rydberg atoms. *Reviews of Modern Physics*, 82(3):2313–2363, 2010.
- [11] Jonathan Simon, Waseem S. Bakr, Ruichao Ma, M. Eric Tai, Philipp M. Preiss, and Markus Greiner. Quantum simulation of antiferromagnetic spin chains in an optical lattice. *Nature*, 472(7343):307–312, April 2011.
- [12] Daniel Greif, Thomas Uehlinger, Gregor Jotzu, Leticia Tarruell, and Tilman Esslinger. Short-range quantum magnetism of ultracold fermions in an optical lattice. *Science*, 340(6138):1307–1310, 2013.
- [13] Sebastian Hild, Takeshi Fukuhara, Peter Schauß, Johannes Zeiher, Michael Knap, Eugene Demler, Immanuel Bloch, and Christian Gross. Far-from-equilibrium spin transport in heisenberg quantum magnets. *Phys. Rev. Lett.*, 113:147205, Oct 2014.
- [14] Russell A. Hart, Pedro M. Duarte, Tsung-Lin Yang, Xinxing Liu, Thereza Paiva, Ehsan Khatami, Richard T. Scalettar, Nandini Trivedi, David A. Huse, and Randall G. Hulet. Observation of antiferromagnetic correlations in the hubbard model with ultracold atoms. *Nature*, 519(7542):211–214, March 2015.
- [15] Timon A. Hilker, Guillaume Salomon, Fabian Grusdt, Ahmed Omran, Martin Boll, Eugene Demler, Immanuel Bloch, and Christian Gross. Revealing hidden antiferromagnetic correlations in doped hubbard chains via string correlators. *Science*, 357(6350):484–487, 2017.
- [16] Anton Mazurenko, Christie S. Chiu, Geoffrey Ji, Maxwell F. Parsons, Marton Kanasz-Nagy, Richard Schmidt, Fabian Grusdt, Eugene Demler, Daniel Greif, and Markus Greiner. A cold-atom fermi-hubbard antiferromagnet. *Nature*, 545(7655):462–466, May 2017.
- [17] Peter T. Brown, Debayan Mitra, Elmer Guardado-Sanchez, Peter Schauß, Stanimir S. Kondov, Ehsan Khatami, Thereza Paiva, Nandini Trivedi, David A. Huse, and Waseem S. Bakr. Spin-imbalance in a 2d fermi-hubbard system. *Science*, 357(6358):1385–, September 2017.
- [18] Nathan Gemelke, Edina Sarajlic, and Steven Chu. Rotating few-body atomic systems in the fractional quantum hall regime. *arXiv:1007.2677v1*, 2010.
- [19] M. Eric Tai, Alexander Lukin, Matthew Rispoli, Robert Schittko, Tim Menke, Dan Borgnia, Philipp M. Preiss, Fabian Grusdt, Adam M. Kaufman, and Markus Greiner. Microscopy of the interacting harper-hofstadter model in the two-body limit. *Nature*, 546(7659):519–523, June 2017.
- [20] Logan W. Clark, Nathan Schine, Claire Baum, Ningyuan

- Jia, and Jonathan Simon. Observation of Laughlin states made of light. *arXiv:1907.05872*, 2019.
- [21] Marcos Atala, Monika Aidelsburger, Julio T. Barreiro, Dmitry Abanin, Takuya Kitagawa, Eugene Demler, and Immanuel Bloch. Direct measurement of the Zak phase in topological Bloch bands. *Nature Physics*, 9:795, 2013.
- [22] F. Grusdt, N. Y. Yao, D. Abanin, M. Fleischhauer, and E. Demler. Interferometric measurements of many-body topological invariants using mobile impurities. *Nat Commun*, 7:11994, June 2016.
- [23] N. Fläschner, B. S. Rem, M. Tarnowski, D. Vogel, D.-S. Lühmann, K. Sengstock, and C. Weitenberg. Experimental reconstruction of the Berry curvature in a Floquet Bloch band. *Science*, 352(6289):1091–1094, 2016.
- [24] Tracy Li, Lucia Duca, Martin Reitter, Fabian Grusdt, Eugene Demler, Manuel Endres, Monika Schleier-Smith, Immanuel Bloch, and Ulrich Schneider. Bloch state tomography using Wilson lines. *Science*, 352(6289):1094–1097, 2016.
- [25] Luca Asteria, Duc Thanh Tran, Tomoki Ozawa, Matthias Tarnowski, Benno S. Rem, Nick Fläschner, Klaus Sengstock, Nathan Goldman, and Christof Weitenberg. Measuring quantized circular dichroism in ultracold topological matter. *Nature Physics*, 15(5):449–454, 2019.
- [26] Tatjana Gericke, Peter Wuertz, Daniel Reitz, Tim Langen, and Herwig Ott. High-resolution scanning electron microscopy of an ultracold quantum gas. *Nature Physics*, 4(12):949–953, December 2008.
- [27] Waseem S. Bakr, Jonathon I. Gillen, Amy Peng, Simon Foelling, and Markus Greiner. A quantum gas microscope for detecting single atoms in a Hubbard-regime optical lattice. *Nature*, 462(7269):74–U80, November 2009.
- [28] Jacob F. Sherson, Christof Weitenberg, Manuel Endres, Marc Cheneau, Immanuel Bloch, and Stefan Kuhr. Single-atom-resolved fluorescence imaging of an atomic Mott insulator. *Nature*, 467(7311):68–U97, September 2010.
- [29] Maxwell F. Parsons, Anton Mazurenko, Christie S. Chiu, Geoffrey Ji, Daniel Greif, and Markus Greiner. Site-resolved measurement of the spin-correlation function in the Fermi-Hubbard model. *Science*, 353(6305):1253–1256, 2016.
- [30] Martin Boll, Timon A. Hilker, Guillaume Salomon, Ahmed Omran, Jacopo Nespolo, Lode Pollet, Immanuel Bloch, and Christian Gross. Spin- and density-resolved microscopy of antiferromagnetic correlations in Fermi-Hubbard chains. *Science*, 353(6305):1257–1260, 2016.
- [31] Lawrence W. Cheuk, Matthew A. Nichols, Katherine R. Lawrence, Melih Okan, Hao Zhang, Ehsan Khatami, Nandini Trivedi, Thereza Paiva, Marcos Rigol, and Martin W. Zwierlein. Observation of spatial charge and spin correlations in the 2d Fermi-Hubbard model. *Science*, 353(6305):1260–1264, 2016.
- [32] N. Read and E. Rezayi. Beyond paired quantum Hall states: Parafermions and incompressible states in the first excited Landau level. *Physical Review B*, 59(12):8084–8092, 1999.
- [33] A. Kitaev. Anyons in an exactly solved model and beyond. *Annals of Physics*, 321(1):2–111, January 2006.
- [34] A. Y. Kitaev. Fault-tolerant quantum computation by anyons. *Annals of Physics*, 303(1):P11 S0003–4916(02)00018–0, 2003.
- [35] Chetan Nayak, Steven H. Simon, Ady Stern, Michael Freedman, and Sankar Das Sarma. Non-abelian anyons and topological quantum computation. *Reviews of Modern Physics*, 80(3):1083–1159, 2008.
- [36] H. P. Büchler, M. Hermele, S. D. Huber, Matthew P. A. Fisher, and P. Zoller. Atomic quantum simulator for lattice gauge theories and ring exchange models. *Phys. Rev. Lett.*, 95:040402, Jul 2005.
- [37] L. Tagliacozzo, A. Celi, A. Zamora, and M. Lewenstein. Optical abelian lattice gauge theories. *Annals of Physics*, 330:160–191, March 2013.
- [38] U.-J. Wiese. Ultracold quantum gases and lattice systems: quantum simulation of lattice gauge theories. *ANNALEN DER PHYSIK*, 525(10-11):777–796, July 2013.
- [39] Erez Zohar, J Ignacio Cirac, and Benni Reznik. Quantum simulations of lattice gauge theories using ultracold atoms in optical lattices. *Reports on Progress in Physics*, 79(1):014401–, 2016.
- [40] T. Senthil, Ashvin Vishwanath, Leon Balents, Subir Sachdev, and Matthew P. A. Fisher. Deconfined quantum critical points. *Science*, 303(5663):1490–1494, 2004.
- [41] Anders W. Sandvik. Evidence for deconfined quantum criticality in a two-dimensional Heisenberg model with four-spin interactions. *Phys. Rev. Lett.*, 98:227202, Jun 2007.
- [42] P. A. M. Dirac. Quantum mechanics of many-electron systems. *Proc R Soc Lond A Math Phys Sci*, 123(792):714, April 1929.
- [43] M. Roger, J. H. Hetherington, and J. M. Delrieu. Magnetism in solid ^3He . *Rev. Mod. Phys.*, 55:1–64, Jan 1983.
- [44] M. Roger and J. M. Delrieu. Cyclic four-spin exchange on a two-dimensional square lattice: Possible applications in high- T_c superconductors. *Phys. Rev. B*, 39:2299–2303, Feb 1989.
- [45] Michel Roger. Ring exchange and correlated fermions. *Journal of Physics and Chemistry of Solids*, 66(8):1412–1416, 2005.
- [46] Hendrik Weimer, Markus Müller, Igor Lesanovsky, Peter Zoller, and Hans Peter Büchler. A Rydberg quantum simulator. *Nat Phys*, 6(5):382–388, May 2010.
- [47] A. W. Glaetzle, M. Dalmonte, R. Nath, I. Rousochatzakis, R. Moessner, and P. Zoller. Quantum spin-ice and dimer models with Rydberg atoms. *Phys. Rev. X*, 4:041037, Nov 2014.
- [48] Federica M. Surace, Paolo P. Mazza, Giuliano Giudici, Alessio Lerose, Andrea Gambassi, and Marcello Dalmonte. Lattice gauge theories and string dynamics in Rydberg atom quantum simulators. *arXiv:1902.09551*, 2019.
- [49] Daniel Barredo, Sylvain de Leseleuc, Vincent Lienhard, Thierry Lahaye, and Antoine Browaeys. An atom-by-atom assembler of defect-free arbitrary two-dimensional atomic arrays. *Science*, 354(6315):1021–, November 2016.
- [50] Manuel Endres, Hannes Bernien, Alexander Keesling, Harry Levine, Eric R. Anschuetz, Alexandre Krajenbrink, Crystal Senko, Vladan Vuletic, Markus Greiner, and Mikhail D. Lukin. Atom-by-atom assembly of defect-free one-dimensional cold atom arrays. *Science*, 354(6315):1024–1027, 2016.
- [51] Hyosub Kim, Woojun Lee, Han-gyeol Lee, Hanlae Jo, Yunheung Song, and Jaewook Ahn. In situ single-atom array synthesis using dynamic holographic optical tweezers. *Nature Comm.*, 7:13317, 2016.
- [52] C. Robens, J. Zopes, W. Alt, S. Brakhane, D. Meschede, A. Alberti. Low-Entropy States of Neutral Atoms in Polarization-Synthesized Optical Lattices.

- Phys. Rev. Lett.*, 118, 065302, 2017.
- [53] D. Ohl de Mello, D. Schäffner, J. Werkmann, T. Preuschoff, L. Kohfahl, M. Schlosser, G. Birkl. Defect-Free Assembly of 2D Clusters of More Than 100 Single-Atom Quantum Systems. *Phys. Rev. Lett.*, 122, 203601, 2019.
- [54] V. J. Emery. Theory of high- t_c superconductivity in oxides. *Phys. Rev. Lett.*, 58:2794–2797, Jun 1987.
- [55] Assa Auerbach. *Interacting Electrons and Quantum Magnetism*. Springer, Berlin, 1998.
- [56] A. M. Toader, J. P. Goff, M. Roger, N. Shannon, J. R. Stewart, and M. Enderle. Spin correlations in the paramagnetic phase and ring exchange in La_2CuO_4 . *Phys. Rev. Lett.*, 94:197202, May 2005.
- [57] Belén Paredes and Immanuel Bloch. Minimum instances of topological matter in an optical plaquette. *Phys. Rev. A*, 77:023603, Feb 2008.
- [58] Han-Ning Dai, Bing Yang, Andreas Reingruber, Hui Sun, Xiao-Fan Xu, Yu-Ao Chen, Zhen-Sheng Yuan, and Jian-Wei Pan. Four-body ring-exchange interactions and anyonic statistics within a minimal toric-code hamiltonian. *Nature Physics*, 13:1195–, August 2017.
- [59] F. D. M. Haldane. Continuum dynamics of the 1-d heisenberg anti-ferromagnet - identification with the $o(3)$ non-linear sigma-model. *Physics Letters A*, 93(9):464–468, 1983.
- [60] T Kennedy. Exact diagonalisations of open spin-1 chains. *Journal of Physics: Condensed Matter*, 2(26):5737, 1990.
- [61] Tom Kennedy and Hal Tasaki. Hidden symmetry breaking and the haldane phase in $s=1$ quantum spin chains. *Commun. Math. Phys.*, 147:431–484, 1992.
- [62] M. Murphy, L. Jiang, N. Khaneja, and T. Calarco. High-fidelity fast quantum transport with imperfect controls. *Phys. Rev. A*, 79:020301(R), 2009.
- [63] M. Müller, I. Lesanovsky, H. Weimer, H. P. Büchler, and P. Zoller. Mesoscopic Rydberg gate based on electromagnetically induced transparency. *Phys. Rev. Lett.*, 102:170502, 2009.
- [64] M. Fleischhauer, A. Imamoglu, and J. P. Marangos. Electromagnetically induced transparency: Optics in coherent media. *Reviews of Modern Physics*, 77(2):633–673, 2005.
- [65] F. D. M. Haldane. Non-linear field-theory of large-spin heisenberg anti-ferromagnets - semi-classically quantized solitons of the one-dimensional easy-axis neel state. *Physical Review Letters*, 50(15):1153–1156, 1983.
- [66] A. Läuchli, G. Schmid, and M. Troyer. Phase diagram of a spin ladder with cyclic four-spin exchange. *Phys. Rev. B*, 67:100409, Mar 2003.
- [67] Christof Weitenberg, Manuel Endres, Jacob F. Sherson, Marc Cheneau, Peter Schauss, Takeshi Fukuhara, Immanuel Bloch, and Stefan Kuhr. Single-spin addressing in an atomic mott insulator. *Nature*, 471(7338):319–+, March 2011.
- [68] Alexandros Metavitsiadis and Sebastian Eggert. Competing phases in spin ladders with ring exchange and frustration. *Phys. Rev. B*, 95:144415, Apr 2017.
- [69] Ian Affleck, Tom Kennedy, Elliott H. Lieb, and Hal Tasaki. Rigorous results on valence-bond ground states in antiferromagnets. *Phys. Rev. Lett.*, 59:799–802, Aug 1987.
- [70] Frank Pollmann, Erez Berg, Ari M. Turner, and Masaki Oshikawa. Symmetry protection of topological phases in one-dimensional quantum spin systems. *Phys. Rev. B*, 85:075125, Feb 2012.