



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Strong Coupling Thermodynamics of Open Quantum Systems

Ángel Rivas

Phys. Rev. Lett. **124**, 160601 — Published 21 April 2020

DOI: [10.1103/PhysRevLett.124.160601](https://doi.org/10.1103/PhysRevLett.124.160601)

Strong Coupling Thermodynamics of Open Quantum Systems

Ángel Rivas

*Departamento de Física Teórica, Facultad de Ciencias Físicas, Universidad Complutense, 28040 Madrid, Spain.
CCS - Center for Computational Simulation, Campus de Montegancedo UPM, 28660 Boadilla del Monte, Madrid, Spain.*
(Dated: March 17, 2020)

A general thermodynamic framework is presented for open quantum systems in fixed contact with a thermal reservoir. The first and second law are obtained for arbitrary system-reservoir coupling strengths, and including both factorized and correlated initial conditions. The thermodynamic properties are adapted to the generally strong coupling regime, approaching the ones defined for equilibrium, and their standard weak-coupling counterparts as appropriate limits. Moreover, they can be inferred from measurements involving only system observables. Finally, a thermodynamic signature of non-Markovianity is formulated in the form of a negative entropy production rate.

Introduction.— The statement of thermodynamics laws at the quantum level is an open and fundamental task. However, given its practical implications in areas such as quantum transport [1–3], quantum information [4–6], or AMO physics [7–16], the motivation is far from being only fundamental. For these reasons, the field of quantum thermodynamics has emerged lately attracting an extensive attention [17, 18].

A particularly interesting problem concerns the formulation of a universally valid nonequilibrium thermodynamic framework for open quantum systems in contact with thermal reservoirs [19]. By “reservoir” we understand a quantum system with an infinitely large, continuous, number of degrees of freedom, and a “thermal reservoir” is the one which initially remains in its canonical Gibbs state $\rho_R(0) = \rho_{R,\beta} = \exp(-\beta H_R)/Z_R$. Here, $\beta = 1/(k_B T)$ is the inverse temperature (with T temperature and k_B Boltzmann constant), H_R is the reservoir Hamiltonian and $Z_R = \text{Tr}[\exp(-\beta H_R)]$ is its partition function. Strictly speaking, since Z_R becomes infinity for an infinitely large, continuous, system, the density matrix $\rho_{R,\beta}$ is ill-defined in such a case. The rigorous definition of this state is given as a functional in the algebraic formulation of quantum mechanics [20]. However, we shall write $\rho_{R,\beta}$ with a formal meaning. If the interaction Hamiltonian between system and reservoir is denoted by V , the total Hamiltonian reads

$$H(t) = H_S(t) + H_R + V, \quad (1)$$

where $H_S(t)$ is a generally time-dependent system Hamiltonian (unless otherwise stated we shall adopt the Schrödinger picture throughout the text). Considering the system and thermal reservoir initially in the product state

$$\rho_{SR}(0) = \rho_S(0) \otimes \rho_{R,\beta}, \quad (2)$$

after some time interval t , the state changes to

$$\rho_{SR}(t) = U(t,0)\rho_S(0) \otimes \rho_{R,\beta}U^\dagger(t,0), \quad (3)$$

with the evolution family $U(t,0) := \mathcal{T} \exp[-\frac{i}{\hbar} \int_0^t H(s)ds]$. This dynamics induces a time-evolution

in the open system S given by a dynamical map Λ_t , i.e. a family of completely positive and trace-preserving (CPTP) maps [21–23], $\rho_S(0) \rightarrow \rho_S(t) = \Lambda_t \rho_S(0) := \text{Tr}_R[\rho_{SR}(t)]$. We shall address the derivation of the thermodynamics laws for the open quantum system S in this situation.

Weak coupling considerations.— The first step is the identification of system thermodynamic variables. Since the global SR system is isolated, any energy change (which only occurs for time-dependent Hamiltonians) must be identified with work W . Thus, the power is given by

$$\dot{W}(t) := \frac{d\langle H(t) \rangle}{dt} = \text{Tr}[\dot{H}_S(t)\rho_{SR}(t)] = \text{Tr}[\dot{H}_S(t)\rho_S(t)], \quad (4)$$

where the overdot notation indicates time-derivatives. This work is assumed to be performed by/applied to the system as only depends on system variables.

Internal energy and heat are magnitudes more difficult to be properly defined. However, this task can be successfully accomplished in the limit of small interaction V . In such a case, the expectation value of the total Hamiltonian becomes $\langle H(t) \rangle \simeq \langle H_S(t) \rangle + \langle H_R \rangle$, and so $\langle H_S(t) \rangle$ can be unequivocally identified with the system internal energy E_U at time t [24]. Then, taking time-derivative one obtains the first law in the form of

$$\frac{dE_U^{(w)}(t)}{dt} = \frac{d\langle H_S(t) \rangle}{dt} = \dot{W}(t) + \dot{Q}^{(w)}(t), \quad (5)$$

with $\dot{Q}^{(w)}(t) := \text{Tr}[H_S(t)\dot{\rho}_S(t)]$ the heat flow in the weak coupling approximation.

The second law can also be obtained in the weak coupling limit. For a slow time-varying $H_S(t)$ compared to the relaxation time of the reservoir [25], the dynamical map can be rigorously approximated by $\Lambda_t = \mathcal{T} \exp[\int_0^t \mathcal{L}_D(s)ds]$ where $\mathcal{L}_D(t)$ is the time-dependent “Davies generator” [26] with the (time-dependent) GKLS form [27, 28], so that Λ_t is CPTP [21–23]. Thanks to this, and applying a series of results [19, 29] based on the monotonicity of the quantum relative entropy $D(\rho_1||\rho_2) := \text{Tr}(\rho_1 \log \rho_1) - \text{Tr}(\rho_1 \log \rho_2)$ under a CPTP

map Λ [30, 31],

$$D[\Lambda(\rho_1)\|\Lambda(\rho_2)] \leq D(\rho_1\|\rho_2), \quad (6)$$

it is possible to obtain the second law in the differential form $\frac{d}{dt}S_{\text{vN}}(t) - \beta\dot{Q}^{(\text{w})}(t) \geq 0$, where $S_{\text{vN}}(t) := -k_B \text{Tr}[\rho_{\text{S}}(t) \log \rho_{\text{S}}(t)]$ is the (thermodynamic) von Neumann entropy. This can be extended to arbitrarily fast periodic drivings [19, 32, 33]. Recently, other drivings have also been analyzed [34].

Out of the weak coupling regime, several attempts have been performed to formulate a thermodynamic framework, e.g. [35–46]. A possible approach [35, 39, 40] defines $\dot{Q}^{(\text{e})}(t) := -\text{Tr}[H_{\text{R}}(t)\dot{\rho}_{\text{R}}(t)]$ and $E_{\text{U}}^{(\text{e})}(t) := \text{Tr}\{[H_{\text{S}}(t) + V]\rho_{\text{SR}}(t)\}$ as heat and system internal energy, respectively. Here, the superscript “e” stands for “external” as, in this approach, those variables are actually external properties, defined in terms of reservoir mean values. This is undesirable from the open system theory, and implies the experimental difficulty of controlling the state of the reservoir in order to make thermodynamic measurements. Nevertheless, it is possible to obtain the first and the second law in the integrated form $\Delta S_{\text{vN}}(t) - \beta Q^{(\text{e})}(t) \geq 0$ for a “finite size” reservoir [35, 39, 40]. Although they are also expected to hold for true (infinite) reservoirs as an appropriate limit, one should be careful at this point in the continuous limit because quantities such as the reservoir von Neumann entropy are ill-defined. On the other hand, as we shall see in a moment, these definitions do not fit with the expected situation once the system reaches thermal equilibrium. In order to overcome these difficulties an alternative approach is needed.

Equilibrium considerations.— Let us assume for a moment that H_{S} is time-independent in (1). There is a vast literature showing that system and reservoir thermalize after some transient time interval [37, 47–60]. In particular, under certain regularity conditions on reservoirs and couplings, it can be rigorously proven [47–54] that

$$\rho_{\text{SR}}(t) = e^{-iHt/\hbar} \rho_{\text{S}}(0) \otimes \rho_{\text{R},\beta} e^{iHt/\hbar} \xrightarrow{t \rightarrow \infty} \frac{e^{-\beta H}}{Z_{\text{SR}}}. \quad (7)$$

This convergence requires R to be an infinitely large continuum, and this implies $Z_{\text{SR}} = \text{Tr}[\exp(-\beta H)]$ to be singular, as commented. Thus, the limit in (7) must be understood in functional sense [61]. Thermalization suggests that, any suitable choice of nonequilibrium system internal energy must fit the system internal energy obtained from the global canonical state $\rho_{\text{SR},\beta} = Z_{\text{SR}}^{-1} \exp(-\beta H)$, once equilibration is reached (perhaps asymptotically). That equilibrium thermal internal energy has been studied both in the classical [62–67] and in the quantum [44, 63, 68, 69] realm. The method is based on the definition of the “Hamiltonian of mean force” H_{S}^*

by the equation

$$H_{\text{S}}^* := -\beta^{-1} \log \left[\left(\frac{Z_{\text{SR}}}{Z_{\text{R}}} \right) \text{Tr}_{\text{R}}(\rho_{\text{SR},\beta}) \right], \quad (8)$$

so that the reduced system state at thermal equilibrium is given formally by a Gibbs state for H_{S}^* ,

$$\rho_{\text{S,eq}} := \text{Tr}_{\text{R}}(\rho_{\text{SR},\beta}) = \frac{e^{-\beta H_{\text{S}}^*}}{Z_{\text{S}}^*}, \quad (9)$$

with $Z_{\text{S}}^* = Z_{\text{SR}}/Z_{\text{R}}$. Now, one requires the fulfillment of standard equilibrium relations such as $F = -\beta^{-1} \log Z_{\text{S}}$, $E_{\text{U}} = F + \partial_{\beta} F$, $S = \beta^2 \partial_{\beta} F$, and $F = E_{\text{U}} - TS$, for Z_{S}^* playing the role of $Z_{\text{S}} = \text{Tr}[\exp(-\beta H_{\text{S}})]$ (Z_{S}^* approaches Z_{S} for vanishing coupling). Since the Hamiltonian of mean force (8) is a function of β , $H_{\text{S}}^*(\beta)$, this leads to the following redefinitions of internal and free energy and thermodynamic entropy at equilibrium (units of $k_B = 1$):

$$E_{\text{U}}^* := \text{Tr}\{\rho_{\text{S,eq}}[H_{\text{S}}^*(\beta) + \beta \partial_{\beta} H_{\text{S}}^*(\beta)]\}, \quad (10)$$

$$F^* := \text{Tr}\{\rho_{\text{S,eq}}[H_{\text{S}}^*(\beta) + \beta^{-1} \log \rho_{\text{S,eq}}]\}, \quad (11)$$

$$S^* := \text{Tr}\{\rho_{\text{S,eq}}[-\log \rho_{\text{S,eq}} + \beta^2 \partial_{\beta} H_{\text{S}}^*(\beta)]\}. \quad (12)$$

A possible generalization of Eqs. (10)–(12) for nonequilibrium is given by the straightforward substitution $\rho_{\text{S,eq}} \rightarrow \rho_{\text{S}}(t)$ [44, 65, 67]. This choice satisfies thermodynamic laws for some restricted class of initial states [44], but fails for the general initial condition (2) [70]. Therefore, we shall take a different route.

Time-independent system Hamiltonians.— Let us consider first a time-independent H_{S} . Typically, $\text{Tr}_{\text{R}}(V \rho_{\text{R},\beta}) = 0$ (otherwise this can always be achieved by a convenient redefinition of system and interaction Hamiltonians, see e.g. [23]), and so under the initial condition (2) we have

$$\langle H(0) \rangle = \text{Tr}[H \rho_{\text{S}}(0) \otimes \rho_{\text{R},\beta}] = \text{Tr}[H_{\text{S}} \rho_{\text{S}}(0)] + \text{Tr}[H_{\text{R}} \rho_{\text{R},\beta}]. \quad (13)$$

The first term of the right hand side of (13) can be unambiguously identified with the system internal energy at $t = 0$. In addition, because of (7), the internal energy must become equivalent to (10) for asymptotic times. Strictly speaking, since only relative differences appear in the thermodynamic laws, this equivalence must hold up to some time-independent additive constant (which observes $E_{\text{U}} = F + \partial_{\beta} F$, $S = \beta^2 \partial_{\beta} F$, $F = E_{\text{U}} - TS$ at thermal equilibrium). The third property we desire for the choice of internal energy is that it should be given just in terms of the reduced system dynamics. In similar spirit to [45], all these properties are satisfied by defining

$$H_{\text{S}}^{\otimes}(t, \beta) := -\beta^{-1} \log[\Lambda_t e^{-\beta H_{\text{S}}}], \quad (14)$$

such that $\Lambda_t e^{-\beta H_{\text{S}}} = e^{-\beta H_{\text{S}}^{\otimes}(t, \beta)}$ and, in parallelism with (10)–(12),

$$E_{\text{U}}(t) := \text{Tr}\{\rho_{\text{S}}(t)[H_{\text{S}}^{\otimes}(t, \beta) + \beta \partial_{\beta} H_{\text{S}}^{\otimes}(t, \beta)]\}, \quad (15)$$

$$F(t) := \text{Tr}\{\rho_{\text{S}}(t)[H_{\text{S}}^{\otimes}(t, \beta) + \beta^{-1} \log \rho_{\text{S}}(t)]\}, \quad (16)$$

$$S(t) := \text{Tr}\{\rho_{\text{S}}(t)[- \log \rho_{\text{S}}(t) + \beta^2 \partial_{\beta} H_{\text{S}}^{\otimes}(t, \beta)]\}. \quad (17)$$

One should note that

$$H_S^\otimes(0, \beta) = H_S, \quad (18)$$

$$H_S^\otimes(\infty, \beta) = H_S^* + \beta^{-1} \log[Z_{SR}/(Z_S Z_R)], \quad (19)$$

where the last additive constant is due to the fact that Λ_t is trace-preserving [71]. These relations ensure the correct initial and long time limits of $E_U(t)$. Moreover, for small coupling V , the dynamics is given by the Davies semigroup [21–23] which has $\exp(-\beta H_S)$ as a fixed point, and so $E_U(t) \rightarrow E_U^{(w)}(t)$ and the thermodynamic entropy $S(t) \rightarrow -\text{Tr}[\rho_S(t) \log \rho_S(t)]$ approaches the usual von Neumann expression.

By construction, the definitions (15)-(17) satisfy asymptotically, at thermal equilibrium, the ‘standard’ relations $F(\infty) = -\beta^{-1} \log Z_S^\otimes(\infty)$, $E_U(\infty) = F(\infty) + \partial_\beta F(\infty)$, and $S(\infty) = \beta^2 \partial_\beta F(\infty)$. In addition, since Λ_t is trace preserving, $Z_S^\otimes(\infty) := \text{Tr}\{\exp[-\beta H_S^\otimes(\infty, \beta)]\} = \text{Tr}\{\exp(-\beta H_S)\} = Z_S$. This implies that the thermodynamic variables at equilibrium take the same value regardless of the strength of the coupling V [72], so they can be obtained by e.g. their weak coupling expressions. Namely, for the internal energy we have $E_U(\infty) = -\partial_\beta \log Z_S^\otimes(\infty) = -\partial_\beta \log Z_S = E_U^{(w)}(\infty) = \text{Tr}(\rho_{S,\beta} H_S)$. Similarly, $S(\infty) = -\text{Tr}(\rho_{S,\beta} \log \rho_{S,\beta})$ and so the entropy (17) at equilibrium approaches 0 for vanishing temperature (in absence of degeneracy), as expected for a ‘thermodynamic’ entropy. This behavior has also been found for S^* [69], but it is not fulfilled for the von Neumann entropy as entanglement may preclude the reduced state to be pure for non-vanishing V .

On the other hand, since for a time-independent Hamiltonian H there is no work, the *first law* defines heat as

$$\begin{aligned} \dot{Q}(t) &= \frac{dE_U(t)}{dt} \Rightarrow Q(t) = E_U(t) - E_U(0) \\ &= E_U(t) - \langle H_S(0) \rangle. \end{aligned} \quad (20)$$

In regard to the *second law*, it can be derived in the integrated form. From Eq. (6),

$$D\left\{\Lambda_t[\rho_S(0)] \middle| \middle| \Lambda_t(\rho_{S,\beta})\right\} \leq D\left[\rho_S(0) \middle| \middle| \rho_{S,\beta}\right], \quad (21)$$

which can be straightforwardly recast in the form

$$\begin{aligned} &-\text{Tr}[\rho_S(t) \log \rho_S(t)] - S(0) \\ &-\beta\{\text{Tr}[\rho_S(t) H_S^\otimes(t, \beta)] - \langle H_S(0) \rangle\} \geq 0. \end{aligned} \quad (22)$$

By adding and subtracting $\beta^2 \text{Tr}[\rho_S(t) \partial_\beta H_S^\otimes(t, \beta)]$, and using (20), we finally obtain

$$\Delta S(t) - \beta Q(t) \geq 0. \quad (23)$$

One may notice that this equation for the entropy production reaches the zero value if the system is initially in




FIG. 1: Internal energy (left) and entropy production (right) of a qubit interacting with a composite spin-boson reservoir. The qubit is initially in its ground state. The internal energy E_U turns out to be a modest correction to the expectation value of the system Hamiltonian $\langle H_S \rangle$ (lighter color, dotted line). The nonmonotonic entropy production (23) shows the non-Markovian character of the dynamics, see Eq. (40). We have taken $\omega_0 = \omega_1$, $\kappa = 0.9\omega_0$, and $10^{-3}\omega_0$ for the spin-boson decay rates [70].

the Gibbs state $\rho_S(0) = \rho_{S,\beta}$, as in that case the equality in (21) is trivially obtained. Actually, in such a situation, no thermodynamic magnitude in (15)-(17) changes on time. This might seem surprising but it can be understood because, formally,

$$U(t, 0) \rho_{S,\beta} \otimes \rho_{R,\beta} U^\dagger(t, 0) = \frac{e^{-\beta U(t, 0)(H_S + H_R) U^\dagger(t, 0)}}{Z_S Z_R}. \quad (24)$$

Since at $t = 0$ (13) holds, system and reservoir starts effectively and remains canonical throughout the process, at instantaneous ‘thermal equilibrium’ in the Gibbs state of the ‘Hamiltonian’ $U(t, 0)(H_S + H_R)U^\dagger(t, 0)$. Given that, by hypothesis, there is no applied work, the internal energy and the rest of the thermodynamic properties remain constant. This ‘thermodynamic’ reversibility should be considered, in the strong coupling, as a different concept from ‘informational’ reversibility. The density matrix indeed changes on time despite the thermodynamic variables and so the thermodynamic ‘state’ remain constant. However, this does not contradict the standard reversibility notion in weak-coupling thermodynamics. That is a particular case of this more general formalism where both informational variables (e.g. the von Neumann entropy) and thermodynamic variables (15)-(17) coincide.

Time-dependent system Hamiltonians.— For a general time-dependent $H_S(t)$, we still demand the same initial condition for the internal energy $E_U(0) = \langle H_S(0) \rangle = \text{Tr}[H_S(0) \rho_S(0)]$ as before, and the recovery of the previous results if $H_S(t)$ becomes time-independent. Moreover, we retain the requirement for a formulation just in terms of system observables. This can be done by keeping the definitions (15)-(17) and redefining the operator $H_S^\otimes(t, \beta)$ by

$$H_S^\otimes(t, \beta) := -\beta^{-1} \log \left[\Lambda_t \left\{ e^{-\beta H_S(0) - \beta \int_0^t \Lambda_s^* [H_S(s)] ds} \right\} \right] \quad (25)$$

where Λ^* denotes the Heisenberg adjoint of Λ , $\text{Tr}[\Lambda(A)B] = \text{Tr}[A\Lambda^*(B)]$ [45]. Note that $H_S^{\otimes}(0, \beta) = H_S(0)$ and for time-independent H_S the equation (14) is recovered, as required.

The *first law* in this case reads

$$\frac{dE_U(t)}{dt} = \dot{Q}(t) + \dot{W}(t), \quad (26)$$

and that is used to define heat,

$$Q(t) := E_U(t) - \langle H_S(0) \rangle - \int_0^t \text{Tr}[\rho_S(s)\dot{H}_S(s)]ds. \quad (27)$$

Note that, if we consider a quasistatic and small coupling regime where $\Lambda_t \left(\exp \left\{ -\beta H_S(0) - \beta \int_0^t \Lambda_s^* [\dot{H}_S(s)] ds \right\} \right) \approx \exp[-\beta H_S(t)]$ such that $H_S^{\otimes}(t, \beta) \approx H_S(t)$, the weak-coupling first law (5) is obtained. This reinforces the definition (25).

In order to derive the second law, we define the auxiliary object

$$\Omega(t, r) := -\beta^{-1} \log \left[\Lambda_t \left\{ e^{-\beta H_S(0) - \beta \int_0^r \Lambda_s^* [\dot{H}_S(s)] ds} \right\} \right], \quad (28)$$

which satisfies $\Omega(t, t) = H_S^{\otimes}(t, \beta)$. A straightforward computation in (27) gives

$$Q(t) = \text{Tr}\{\rho_S(t)[\Omega(t, t) + \beta \partial_\beta H_S^{\otimes}(t, \beta)]\} - \text{Tr}\{\rho_S(0)\Omega(0, t)\}. \quad (29)$$

For the state

$$\rho_0(\beta, r) := Z_S^{-1}(r) e^{-\beta H_S(0) - \beta \int_0^r \Lambda_s^* [\dot{H}_S(s)] ds}, \quad (30)$$

with $Z_S(r) = \text{Tr} \left(\exp \left\{ -\beta H_S(0) - \beta \int_0^r \Lambda_s^* [\dot{H}_S(s)] ds \right\} \right)$, monotonicity of the relative entropy (6) yields $D \left\{ \Lambda_t[\rho_S(0)] \middle| \middle| \Lambda_t[\rho_0(\beta, r)] \right\} \leq D \left[\rho_S(0) \middle| \middle| \rho_0(\beta, r) \right]$, which can be recast in the form

$$-\text{Tr}[\rho_S(t) \log \rho_S(t)] - S(0) - \beta \{ \text{Tr}[\rho_S(t)\Omega(t, r)] - \text{Tr}[\rho_S(0)\Omega(0, r)] \} \geq 0. \quad (31)$$

Since this is fulfilled for all r , and particularly for $r = t$, again by adding and subtracting $\beta^2 \text{Tr}[\rho_S(t)\partial_\beta H_S^{\otimes}(t, \beta)]$ and using (29), we obtain the *second law*

$$\Delta S(t) - \beta Q(t) \geq 0. \quad (32)$$

This completes the thermodynamic formulation for general open quantum systems in contact with a thermal reservoir.

As a simple example of this approach we may consider a qubit with Hamiltonian $H_S = \frac{\omega_0}{2} \sigma_S^z$, in dipolar contact $V = \kappa(\sigma_S^+ \sigma_R^- + \sigma_S^- \sigma_R^+)$ with a composite spin-boson reservoir [73] with Hamiltonian $H_R = H_{\text{spin}} + H_{\text{boson}} + V_{\text{spin-boson}}$. Here, $H_{\text{spin}} = \frac{\omega_1}{2} \sigma_R^z$, $H_{\text{boson}} = \int dk \omega(k) a^\dagger(k) a(k)$, $V_{\text{spin-boson}} = \alpha \int dk g(k) \sigma_R^x [a(k) + a^\dagger(k)]$, $a(k)$ denotes bosonic annihilation operators, and

$\sigma_S^{z,\pm}$, $\sigma_R^{z,x,\pm}$ stand for Pauli matrices for system and spin part of the reservoir, respectively. Note that the bosonic part of the reservoir is not directly coupled to the system qubit. The coupling between system and reservoir is mediated by κ , whereas α is supposed small enough such that a weak coupling treatment of the spin-boson degrees of freedom is justified [70]. Figure 1 shows the comparison between the internal energy E_U , Eq. (15), in the strong coupling and the mean value of the system Hamiltonian $\langle H_S \rangle$ for several temperatures. One can notice that, for this model, the latter turns out to be a modest correction to $\langle H_S \rangle$ sharing a very similar time-dependency.

Initially correlated states.— It is worth to examine whether the previous approach can be extended to different initial system-reservoir states. We do not expect that for any initial state, but for those sufficiently close to the thermodynamic paradigm of a system coupled to a thermal reservoir. Namely, we should consider just those initial system-reservoir states where the reservoir can be well-described via the temperature parameter β . This condition can be rigorously formulated in the framework of operator algebras [47, 51–53], but, for our purposes, there are another two natural classes of states which can be considered in addition to (2). They correspond to the displacement from the global equilibrium $\rho_{\text{SR},\beta}$ either by system driving $H_S(t)$ or by system quantum measurements, respectively [44].

For the first case $\rho_{\text{SR}}(0) = \rho_{\text{SR},\beta}$, and we can assume, formally, a former product ‘initial’ condition $\rho_{\text{S},\beta} \otimes \rho_{\text{R},\beta}$ at $t_0 = -\infty$ and $\dot{H}_S(t) = \dot{H}_S(t)\theta(t)$, with $\theta(t)$ the step function. Following the same steps as before we conclude

$$S(t) - S(-\infty) - \beta Q_{-\infty}(t) \geq 0, \quad (33)$$

where $Q_{-\infty}(t) := \int_{-\infty}^t \dot{Q}(s) ds$ is the heat in the interval $(-\infty, t)$. Then, by splitting this integral into positive and negative time values, and adding and subtracting $S(0)$, which in this case is the entropy (17) of the reduced equilibrium state (9), we have

$$S(t) - S(0) + S(0) - S(-\infty) - \beta \left[\int_0^t \dot{Q}(s) ds + \int_{-\infty}^0 \dot{Q}(s) ds \right] \geq 0. \quad (34)$$

Since we have taken $\dot{H}_S(t) = 0$ for $t < 0$, and the entropy production for the canonical system state $\rho_{\text{S},\beta}$ reaches the zero value for time-independent system Hamiltonians, $S(0) - S(-\infty) - \beta \int_{-\infty}^0 \dot{Q}(s) ds = 0$. Hence, we obtain the desired result (32) for $\rho_{\text{SR}}(0) = \rho_{\text{SR},\beta}$.

For the second case, the joint initial state (after a generally nonselective, projective measurement) is written as

$$\rho_{\text{SR}}(0) = \sum_k p_k \Pi_k \otimes \rho_{\text{R}|k} \quad (35)$$

with $\Pi_k = |k\rangle\langle k|$ a complete set of orthonormal projectors and

$$p_k = \text{Tr}(\Pi_k \otimes \mathbb{I}_{\rho_{\text{SR},\beta}}), \quad \rho_{\text{R}|k} = \frac{\text{Tr}_{\text{S}}(\Pi_k \otimes \mathbb{I}_{\rho_{\text{SR},\beta}})}{p_k}. \quad (36)$$

For this kind of states it is possible to write the reduced system dynamics as $\tilde{\Lambda}_t \rho_{\text{S}}(0) = \rho_{\text{S}}(t)$ for $\rho_{\text{S}}(0) = \sum_k p_k \Pi_k$, with $\tilde{\Lambda}_t$ a CPTP map [74, 75].

On the other hand, $\rho_{\text{SR},\beta}$ remains static before the measurement, with system internal energy (15) given by

$$E_{\text{U}}(\text{eq}) := \text{Tr}\{\rho_{\text{S},\text{eq}}[H_{\text{S}}^{\otimes}(\text{eq}, \beta) + \beta \partial_{\beta} H_{\text{S}}^{\otimes}(\text{eq}, \beta)]\}, \quad (37)$$

where $H_{\text{S}}^{\otimes}(\text{eq}, \beta) = H_{\text{S}}^* + \beta^{-1} \log[Z_{\text{SR}}/(Z_{\text{S}}Z_{\text{R}})]$ according to (19). Therefore, it seems reasonable to take the internal energy after the measurement

$$E_{\text{U}}(0) = \text{Tr}\{\rho_{\text{S}}(0)[H_{\text{S}}^{\otimes}(\text{eq}, \beta) + \beta \partial_{\beta} H_{\text{S}}^{\otimes}(\text{eq}, \beta)]\}, \quad (38)$$

with $\rho_{\text{S}}(0) = \sum_k p_k \Pi_k$. A finer choice could be possible with a microscopic model for the measurement interaction where the measurement change was not ‘instantaneous’. Then by redefining $H_{\text{S}}^{\otimes}(t, \beta)$ as in (25) with $\tilde{\Lambda}_t$ and $H_{\text{S}}^{\otimes}(\text{eq}, \beta)$ in the roles of Λ_t and $H_{\text{S}}(0)$, respectively, the derivation of (32) follows from the same argument as in previous sections.

Non-Markovianity. — Finally, we show that, within this approach, it is possible to establish a thermodynamic signature of non-Markovianity (see also [44, 46, 76, 77]). Suppose the dynamical map Λ_t to be CP divisible [78–80] (actually P-divisible is enough [81]); namely, it can be decomposed as $\Lambda_t = \Lambda_{t,s} \Lambda_s$ for any pair $t > s$ with $\Lambda_{t,s}$ CPTP. Then, monotonicity of the relative entropy (6) implies

$$S\left\{\Lambda_{t+\epsilon}[\rho_{\text{S}}(0)] \middle| \middle| \Lambda_{t+\epsilon}[\rho_0(\beta, r)]\right\} \leq S\left\{\Lambda_t[\rho_{\text{S}}(0)] \middle| \middle| \Lambda_t[\rho_0(\beta, r)]\right\}, \quad (39)$$

for $\epsilon > 0$. From here, following similar steps as for (32) and dividing by ϵ in the limit $\epsilon \rightarrow 0$, we obtain a positive entropy production rate

$$\frac{dS(t)}{dt} - \beta \dot{Q}(t) \geq 0. \quad (40)$$

Hence, a negative production rate for some t is a rigorous indicator of the non-Markovian character of the dynamics. It is clear the presence of intervals with a strong negative production rate in Fig. 1.

Conclusions. — We have presented a general thermodynamic framework for open quantum systems in contact with a thermal reservoir. This was done by identifying the nonequilibrium internal energy imposing suitable initial and asymptotic conditions, and the recovery of the standard weak-coupling result as an appropriate limit.

The factorized initial condition was analyzed in detail and generalized to two natural extensions of correlated initial states. Furthermore, we have found that Markovian dynamics imply monotonically increasing entropy production. This provides quantum non-Markovianity with a thermodynamic meaning, and allows for the introduction of new physical quantifiers of non-Markovianity. Notably, all quantities in this approach can be inferred from measurements involving only system observables. At most, several preparations might be needed to determine $H_{\text{S}}^{\otimes}(t, \beta)$ and its derivatives, but no controlled reservoirs are required. This greatly simplifies the approach and opens the possibility to measure these strong coupling thermodynamic variables in the lab.

The author is grateful to M. Merkli and P. Strasberg for enlightening discussions. Financial support from Spanish MINECO grants FIS2017-91460-EXP, PGC2018-099169-B-I00, the CAM research consortium QUITEMAD S2018/TCS-4342, and US Army Research Office through grant W911NF-14-1-0103 is acknowledged.

-
- [1] J. P. Pekola, *Towards quantum thermodynamics in electronic circuits*, Nat. Phys. **11**, 118 (2015).
 - [2] S. Jezouin et al., *Quantum limit of heat flow across a single electronic channel*, Science **342**, 601 (2013).
 - [3] A. Bermudez, M. Bruderer, and M. B. Plenio, *Controlling and Measuring Quantum Transport of Heat in Trapped-Ion Crystals*, Phys. Rev. Lett. **111**, 040601 (2013).
 - [4] J. Goold, M. Huber, A. Riera, L. del Rio, and P. Skrzypczyk, *The role of quantum information in thermodynamics? a topical review*, J. Phys. A: Math. Theor. **49**, 143001 (2016).
 - [5] S. Deffner and S. Campbell, *Quantum Thermodynamics: An Introduction to the Thermodynamics of Quantum Information* (Morgan & Claypool, San Rafael, CA, 2019)
 - [6] A. Bérut et al., *Experimental verification of Landauer’s principle linking information and thermodynamics*, Nature **483**, 187 (2012).
 - [7] M. Esposito, U. Harbola and S. Mukamel, *Nonequilibrium fluctuations, fluctuation theorems, and counting statistics in quantum systems*, Rev. Mod. Phys. **81**, 1665 (2009).
 - [8] U. Seifert, *Stochastic thermodynamics, fluctuation theorems and molecular machines*, Rep. Prog. Phys. **75**, 126001 (2012)
 - [9] K. Zhang, F. Bariani, and P. Meystre, *Quantum Optomechanical Heat Engine*, Phys. Rev. Lett. **112**, 150602 (2014).
 - [10] C. Bergenfeldt, P. Samuelsson, B. Sothmann, C. Flindt, and M. Büttiker, *Hybrid Microwave-Cavity Heat Engine*, Phys. Rev. Lett. **112**, 076803 (2014).
 - [11] T. Langen et al., *Experimental observation of a generalized Gibbs ensemble*, Science **348**, 207 (2015).
 - [12] S. An et al., *Experimental test of the quantum Jarzynski equality with a trapped-ion system*, Nat. Phys. **11**, 193 (2015).

- [13] J. Jaramillo, M. Beau, and A. del Campo, *Quantum supremacy of many-particle thermal machines*, New J. Phys. **18**, 075019 (2016).
- [14] J. Roßnagel et al., *A single-atom heat engine*, Science **352**, 325 (2016).
- [15] A. Ronzani et al., *Tunable photonic heat transport in a quantum heat valve*, Nat. Phys. **14**, 991 (2018).
- [16] R. M. Fereidani and D. Segal, *Phononic heat transport in molecular junctions: Quantum effects and vibrational mismatch*, J. Chem. Phys. **150**, 024105 (2019).
- [17] J. Gemmer, M. Michel, and G. Mahler, *Quantum Thermodynamics: Emergence of Thermodynamic Behavior within Composite Quantum Systems* (Springer, Berlin, 2004).
- [18] F. Binder, L. A. Correa, C. Gogolin, J. Anders, G. Adesso, (Eds.) *Thermodynamics in the Quantum Regime* (Springer, Cham, Switzerland, 2018).
- [19] R. Kosloff, *Quantum Thermodynamics: A Dynamical Viewpoint*, Entropy **15**, 2100(2013).
- [20] O. Bratteli and D. Robinson, *Operator Algebras and Quantum Statistical Mechanics I, II*, (Springer-Verlag, Berlin, 2002).
- [21] R. Alicki and K. Lendi *Quantum Dynamical Semigroups and Applications* (Springer, Berlin, 1987).
- [22] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, Oxford, 2002).
- [23] A. Rivas and S. F. Huelga, *Open Quantum Systems. An Introduction* (Springer, Heidelberg, 2011).
- [24] We write E_U instead the common notation U to avoid any confusion with the unitary evolution operator.
- [25] R. Alicki, *Quantum open systems as a model of a heat engine*, J. Phys. A: Math. Gen. **12**, L103 (1979).
- [26] E. B. Davies and H. Spohn, *Open quantum systems with time-dependent Hamiltonians and their linear response*, J. Stat. Phys. **19**, 511 (1978).
- [27] V. Gorini, A. Kossakowski, E.C.G. Sudarshan, *Completely positive dynamical semigroups of N -level systems*, J. Math. Phys. **17**, 821 (1976).
- [28] G. Lindblad, *On the generators of quantum dynamical semigroups*, Commun. Math. Phys. **48**, 119 (1976).
- [29] H. Spohn, *Entropy production for quantum dynamical semigroups*, J. Math. Phys. **19**, 1227 (1978).
- [30] G. Lindblad, *Completely positive maps and entropy inequalities*, Commun. Math. Phys. **40**, 147 (1975).
- [31] A. Uhlmann, *Relative entropy and the Wigner-Yanase-Dyson-Lieb concavity in an interpolation theory*, Commun. Math. Phys. **54**, 21 (1977).
- [32] R. Alicki, D. A. Lidar and P. Zanardi, *Internal consistency of fault-tolerant quantum error correction in light of rigorous derivations of the quantum Markovian limit*, Phys. Rev. A **73**, 052311 (2006).
- [33] R. Alicki, D. Gelbwaser-Klimovsky, G. Kurizki, *Periodically driven quantum open systems: Tutorial*, e-print: arXiv:1205.4552.
- [34] R. Dann, A. Levy, and R. Kosloff, *Time-dependent Markovian quantum master equation*, Phys. Rev. A **98**, 052129 (2018).
- [35] M. Esposito, K. Lindberg, and C. van den Broek, *Entropy production as correlation between system and reservoir*, New J. Phys. **12**, 013013 (2010).
- [36] F. C. Binder, S. Vinjanampathy, K. Modi, and J. Goold, *Quantum thermodynamics of general quantum processes*, Phys. Rev. E **91**, 032119 (2015).
- [37] P. Strasberg, G. Schaller, N. Lambert, and T. Brandes, *Nonequilibrium thermodynamics in the strong coupling and non-Markovian regime based on a reaction coordinate mapping*, New J. Phys. **18**, 073007 (2016).
- [38] S. Alipour, F. Benatti, F. Bakhshinezhad, M. Afsary, S. Marcantoni, and A. T. Rezakhani, *Correlations in quantum thermodynamics*, Sci. Rep. **6**, 35568 (2016).
- [39] A. Kato and Y. Tanimura, *Quantum heat current under non-perturbative and non-Markovian conditions: Applications to heat machines*, J. Chem. Phys. **145**, 224105 (2016).
- [40] P. Strasberg, G. Schaller, T. Brandes, and M. Esposito, *Quantum and Information Thermodynamics: A Unifying Framework Based on Repeated Interactions*, Phys. Rev. X **7**, 021003 (2017).
- [41] G. Thomas, N. Siddharth, S. Banerjee, and S. Ghosh, *Thermodynamics of non-Markovian reservoirs and heat engines*, Phys. Rev. E **97**, 062108 (2018).
- [42] M. Perarnau-Llobet, H. Wilming, A. Riera, R. Gallego, and J. Eisert, *Strong coupling corrections in quantum thermodynamics*, Phys. Rev. Lett. **120**, 120602 (2018).
- [43] J.-T. Hsiang, C. H. Chou, Y. Subasi, and B. L. Hu, *Quantum thermodynamics from the nonequilibrium dynamics of open systems: Energy, heat capacity, and the third law*, Phys. Rev. E **97**, 012135 (2018).
- [44] P. Strasberg and M. Esposito, *Non-Markovianity and negative entropy production rates*, Phys. Rev. E **99**, 012120 (2019).
- [45] A. Rivas, *Quantum Thermodynamics in the Refined Weak Coupling Limit*, Entropy **21**, 725 (2019).
- [46] P. Strasberg, *Repeated Interactions and Quantum Stochastic Thermodynamics at Strong Coupling*, Phys. Rev. Lett. **123**, 180604 (2019).
- [47] V. Bach, J. Fröhlich, and I. M. Sigal, *Return to equilibrium*, J. Math. Phys. **41**, 3985 (2000).
- [48] J. Dereziński and V. Jakšić, *Spectral theory of Pauli-Fierz operators*, J. Funct. Anal. **180**, 243 (2001).
- [49] M. Merkli, *Positive commutators in non-equilibrium statistical mechanics*, Commun. Math. Phys. **223**, 327(2001).
- [50] J. Dereziński and V. Jakšić, *Return to equilibrium for Pauli-Fierz systems*, Ann. Henri Poincaré **4**, 739 (2003).
- [51] J. Fröhlich and M. Merkli, *Another return of “return to equilibrium”*, Commun. Math. Phys. **251**, 235 (2004).
- [52] M. Merkli, I. M. Sigal, and P. Berman, *Resonance theory of decoherence and thermalization*, Ann. Phys. (N.Y.) **323**, 373 (2008).
- [53] M. Könenberg and M. Merkli, *On the irreversible dynamics emerging from quantum resonances*, J. Math. Phys. **57**, 033302 (2016).
- [54] M. Merkli, *Quantum Markovian master equations: Resonance theory shows validity for all time scales*, Ann. Phys. (N.Y.) **412**, 167996 (2020).
- [55] N. Linden, S. Popescu, A. J. Short, and A. Winter, *Quantum mechanical evolution towards thermal equilibrium*, Phys. Rev. E **79**, 061103 (2009).
- [56] P. Reimann, *Canonical thermalization*, New J. Phys. **12**, 055027 (2010).
- [57] A. J. Short and T. C. Farrelly, *Quantum equilibration in finite time*, New J. Phys. **14**, 013063 (2012).
- [58] C. Gogolin and J. Eisert, *Equilibration, thermalisation, and the emergence of statistical mechanics in closed quantum systems*, Rep. Prog. Phys. **79**, 056001 (2016).
- [59] Y. Subasi, C. H. Fleming, J. M. Taylor, and B. L. Hu,

- Equilibrium states of open quantum systems in the strong coupling regime*, Phys. Rev. E. **86**, 061132 (2012).
- [60] J. Iles-Smith, N. Lambert, and A. Nazir, *Environmental dynamics, correlations, and the emergence of noncanonical equilibrium states in open quantum systems*, Phys. Rev. A **90**, 032114 (2014).
- [61] The limit holds in the weak operator topology in the algebra of observables \mathfrak{A} of the joint system: $\lim_{t \rightarrow \infty} \rho_{\text{SR}}[A(t)] = \rho_{\text{SR},\beta}(A)$ for $A \in \mathfrak{A}$. See details in [47–54].
- [62] J. G. Kirkwood, *Statistical Mechanics of Fluid Mixtures*, J. Chem. Phys. **3**, 300 (1935).
- [63] M. F. Gelin and M. Thoss, *Thermodynamics of a subensemble of a canonical ensemble*, Phys. Rev. E **79**, 051121 (2009).
- [64] C. Jarzynski, *Stochastic and Macroscopic Thermodynamics of Strongly Coupled Systems*, Phys. Rev. X **7**, 011008 (2017).
- [65] U. Seifert, *First and Second Law of Thermodynamics at Strong Coupling*, Phys. Rev. Lett. **116**, 020601 (2016).
- [66] H. J. D. Miller and J. Anders, *Entropy production and time asymmetry in the presence of strong interactions*, Phys. Rev. E **95**, 062123 (2017).
- [67] P. Strasberg and M. Esposito, *Stochastic thermodynamics in the strong coupling regime: An unambiguous approach based on coarse graining*, Phys. Rev. E **95**, 062101 (2017).
- [68] M. Campisi, P. Talkner, and P. Hänggi, *Fluctuation Theorem for Arbitrary Open Quantum Systems*, Phys. Rev. Lett. **102**, 210401 (2009).
- [69] J.-T. Hsiang and B. L. Hu, *Quantum thermodynamics at strong coupling: Operator thermodynamic functions and relations*, Entropy **20**, 423 (2018).
- [70] See Supplemental Material for further details.
- [71] Note that the change $H_S \rightarrow H_S^*$ in the definition (14) leads to the ‘symmetric’ result $H_S^{\otimes}(0, \beta) = H_S^{\otimes}(\infty, \beta) = H_S^*$, and allows for a similar derivation of thermodynamic laws. However, it breaks the nice condition $E_U(0) = \langle H_S(0) \rangle$ for the product state (2) and requires the knowledge of $Z_{\text{SR}}/Z_{\text{R}}$.
- [72] In this approach, at equilibrium, we may write $Z_{\text{SR}} = Z_S Z_{\text{R}} Z_{\text{int}}$, with $Z_{\text{int}} = Z_{\text{SR}}/(Z_S Z_{\text{R}})$, so that $\mathcal{X}_{\text{SR}} = \mathcal{X}_S + \mathcal{X}_{\text{R}} + \mathcal{X}_{\text{int}}$ for $\mathcal{X} = E_U, F, S$, where \mathcal{X}_j is the thermodynamic variable \mathcal{X} computed from Z_j . Therefore, the strength of the coupling V affects to \mathcal{X}_{SR} and \mathcal{X}_{int} , but not to \mathcal{X}_S and \mathcal{X}_{R} .
- [73] N. P. Oxtoby, A. Rivas, S. F. Huelga, and R. Fazio, *Probing a composite spin-boson environment*, New J. Phys. **11**, 063028 (2009).
- [74] C. A. Rodríguez-Rosario, K. Modi, A. Kuah, A. Shaji, and E. C. G. Sudarshan, *Completely positive maps and classical correlations*, J. Phys. A: Math. Theor. **41**, 205301 (2008).
- [75] The CPTP map is formally given by the Kraus operators $K_{m,n} = \sum_k \sqrt{\lambda_{m|k}} \langle \psi_{\text{R},m|k} | U(t, 0) | \psi_{\text{R},n|k} \rangle \Pi_k$, where $\rho_{\text{R}|k} = \sum_m \lambda_{m|k} |\psi_{\text{R},m|k}\rangle \langle \psi_{\text{R},m|k}|$ is the spectral decomposition of $\rho_{\text{R}|k}$ [74].
- [76] S. Bhattacharya, A. Misra, C. Mukhopadhyay, and A. K. Pati, *Exact master equation for a spin interacting with a spin bath: Non-Markovianity and negative entropy production rate*, Phys. Rev. A **95**, 012122 (2017).
- [77] S. Marcantoni, S. Alipour, F. Benatti, R. Floreanini, A. T. Rezakhani, *Entropy production and non-Markovian dynamical maps*, Sci. Rep. **7**, 12447 (2017).
- [78] A. Rivas, S. F. Huelga, and M. B. Plenio, *Quantum non-Markovianity: Characterization, quantification and detection*, Rep. Prog. Phys. **77**, 094001 (2014).
- [79] H.-P. Breuer, E.-M. Laine, J. Piilo, and B. Vacchini, *Colloquium: Non-Markovian dynamics in open quantum systems*, Rev. Mod. Phys. **88**, 021002 (2016).
- [80] L. Li, M. J. W. Hall, and H. M. Wiseman, *Concepts of quantum non-Markovianity: A hierarchy*, Phys. Rep. **759**, 1 (2018).
- [81] A. Müller-Hermes and D. Reeb, *Monotonicity of the Quantum Relative Entropy Under Positive Maps*, Ann. Henri Poincaré **18**, 1777 (2017).