

This is the accepted manuscript made available via CHORUS. The article has been published as:

# Nonadiabatic Phase Transition with Broken Chiral Symmetry

Bin Yan, Vladimir Y. Chernyak, Wojciech H. Zurek, and Nikolai A. Sinitsyn

Phys. Rev. Lett. **126**, 070602 — Published 19 February 2021

DOI: [10.1103/PhysRevLett.126.070602](https://doi.org/10.1103/PhysRevLett.126.070602)

# Nonadiabatic Phase Transition with Broken Chiral Symmetry

Bin Yan,<sup>1,2</sup> Vladimir Y. Chernyak,<sup>3</sup> Wojciech H. Zurek,<sup>2</sup> and Nikolai A. Sinitsyn<sup>2</sup>

<sup>1</sup>*Center for Nonlinear Studies, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA*

<sup>2</sup>*Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA*

<sup>3</sup>*Department of Chemistry and Department of Mathematics, Wayne State University, Detroit, USA*

We explore nonadiabatic quantum phase transitions in an Ising spin chain with a linearly time-dependent transverse field and two different spins per unit cell. Such a spin system passes through critical points with gapless excitations, which support nonadiabatic transitions. Nevertheless, we find that the excitations on one of the chain sub-lattices are suppressed in the nearly-adiabatic regime exponentially. Thus, we reveal a coherent mechanism to induce exponentially large density separation for different quasiparticles.

The separation of two slightly different types of particles is often encountered both in applied and fundamental research. One example is the apparent asymmetry between matter and antimatter in our Universe. Observations prove that the symmetry between matter and antimatter was broken, presumably early on in the history of the Universe. However, it is still unclear how the subtle CP symmetry violation could lead to the large observed differences at the cosmological scale [1]; although it is known that when the characteristic asymmetry energy scale is very small, the large density difference has to be produced during a nonequilibrium process [2].

The goal of this article is to introduce a mechanism for controlling different quasiparticles separately using exponential sensitivity of quantum nonadiabatic transitions to symmetry-breaking interactions. Namely, when parameters of a quantum system change with time adiabatically, the system remains in the instantaneous eigenstate, e.g., the ground state. However, the Kibble-Zurek mechanism [3–11] predicts that the number of nonadiabatic excitations is not suppressed exponentially if a macroscopic system passes through a quantum critical point at which the energy gap to excitations closes. Without an asymmetry of interactions, different particle species would pass through the same phase transition simultaneously. However, even a small asymmetry generally opens the gap to certain excitation types, even though the critical point is not destroyed. Thus, we can harvest the excitations of one type and suppress the others.

A broadly known quantum example that confirms the Kibble-Zurek mechanism is the model of a uniform Ising spin chain in a transverse magnetic field [12–14], with the Hamiltonian

$$\hat{H}_u = \sum_{n=1}^N [B\hat{\sigma}_n^z + J\hat{\sigma}_n^x\hat{\sigma}_{n+1}^x], \quad B = -\beta t, \quad (1)$$

where  $\hat{\sigma}_n^{x,z}$  are Pauli operators for the  $n$ -th spin,  $J$  is the spin-spin coupling, and  $B$  is the transverse time-dependent field that changes with rate  $\beta > 0$ . This model has two Dirac points at  $B = \pm J$ , at which its spectral gaps close, and which mark boundaries between three phases. The phase with strong quantum correlations (Fig. 1 top) contains the point  $B = 0$  with two




FIG. 1. Top: Phase diagram of Ising chain (1) in a transverse magnetic field. Dashed horizontal lines mark the critical points with the spectral gap closings. At  $B = \pm\infty$ , the ground state is fully spin-polarized along  $z$ -axis, whereas at  $B = 0$  the ground state is a superposition of  $x$ -polarized spin states. Bottom: Spin chain model (5) with broken chiral symmetry. We consider the case of  $b_1 > b_2$ . In the adiabatic limit, the initial spin polarized ground state (top) at  $t = -\infty$  transfers to the spin polarized state at  $t = +\infty$  with all spins flipped. The nonadiabatic excitations (non-flipped spins) on the odd-site sublattice (red) are suppressed as a power law (31), respecting the Kibble-Zurek mechanism. In contrast, nonadiabatic excitations on the even-site sublattice (blue) are suppressed exponentially (25).

degenerate ground states:

$$|\rightarrow \rightarrow \dots \rightarrow \rightarrow\rangle \quad \text{and} \quad |\leftarrow \leftarrow \dots \leftarrow \leftarrow\rangle. \quad (2)$$

For  $B(t) = -\beta t$ , the model is exactly solvable [12]. If the system starts from the fully polarized ground state

$$|\downarrow \downarrow \downarrow \downarrow \dots \downarrow \downarrow\rangle \quad (3)$$

as  $t \rightarrow -\infty$ , then in the thermodynamic limit of this

solution, after the passage through a critical point the number of non-adiabatic excitations follows a power law:

$$\rho \sim J^{-1} \beta^{1/2}. \quad (4)$$

Let us now explore robustness of the Ising model predictions by considering a spin chain whose unit cell contains two different spins, with the Hamiltonian

$$\hat{H} = \sum_{n=1}^N \left[ -\frac{b_1 t}{2} \hat{\sigma}_{2n}^z - \frac{b_2 t}{2} \hat{\sigma}_{2n+1}^z \right] + \sum_{n=1}^N [J_1 \hat{\sigma}_{2n}^x \hat{\sigma}_{2n+1}^x + J_2 \hat{\sigma}_{2n+1}^x \hat{\sigma}_{2n+2}^x]. \quad (5)$$

The difference between  $b_1$  and  $b_2$  is due to different spin  $g$ -factors, and  $J_1 \neq J_2$  reflects the lack of inversion symmetry at zero field. Periodic boundary condition is imposed. Figure 1 (bottom) illustrates the structure of this spin chain.

Without loss of generality, here and in what follows, we assume  $b_1 > b_2$ ; namely, the spins on the even-site sublattice have a larger  $g$ -factor than those on the odd-sites. We note that for weak symmetry breaking,  $|J_1 - J_2| \ll |J_1 + J_2|$  and  $|b_1 - b_2| \ll |b_1 + b_2|$ , the ground state polarizations at  $B = 0, \pm\infty$  are the same for spins with odd and even indices, and the spin excitations on different sublattices then resemble particles of two slightly different types of matter.

Via the Jordan-Wigner transformation, Hamiltonian (5) reduces to a quadratic form of fermionic operators,  $\hat{c}$  and  $\hat{d}$ , on the even and odd sites, respectively. The details are presented in Appendix A of the Supplemental Material (SM). Thus,

$$\hat{\sigma}_{2n}^z = 1 - 2\hat{c}_n^\dagger \hat{c}_n, \quad \hat{\sigma}_{2n+1}^z = 1 - 2\hat{d}_n^\dagger \hat{d}_n. \quad (6)$$

It is convenient to work with the Fourier transformed operators

$$\hat{c}_p = \frac{1}{\sqrt{N}} \sum_{n=1}^N \hat{c}_n e^{-ipn}, \quad \hat{c}_p^\dagger = \frac{1}{\sqrt{N}} \sum_{n=1}^N \hat{c}_n^\dagger e^{ipn}, \quad (7)$$

and similarly defined  $\hat{d}_p, \hat{d}_p^\dagger$ . We will assume that  $N$  is even, so the momentum takes discrete values  $p = \pm(2k-1)\pi/N$ ,  $k = 1, \dots, N/2$ . Hamiltonian (5) then is

$$\hat{H} = \sum_{p>0} \hat{\mathbf{a}}_p^\dagger H_p \hat{\mathbf{a}}_p, \quad (8)$$

where  $\hat{\mathbf{a}}_p$  and  $H_p$  are given by

$$\hat{\mathbf{a}}_p \equiv \begin{pmatrix} \hat{c}_p \\ \hat{c}_{-p}^\dagger \\ \hat{d}_p \\ \hat{d}_{-p}^\dagger \end{pmatrix}, \quad H_p \equiv \begin{pmatrix} b_1 t & 0 & g & \gamma \\ 0 & -b_1 t & -\gamma & -g \\ g^* & -\gamma^* & b_2 t & 0 \\ \gamma^* & -g^* & 0 & -b_2 t \end{pmatrix}, \quad (9)$$

with the couplings

$$g \equiv J_1 + J_2 e^{-ip}, \quad \gamma \equiv J_1 - J_2 e^{-ip}. \quad (10)$$




FIG. 2. The time-dependent spectrum of the Hamiltonian  $H_p(t)$  from (9). (a) Two exact level crossing points appear at  $p = 0$ . (b) At finite  $p$ , which is here  $p = 0.3$ , small gaps open up near places of the former crossing points. Other parameters are  $b_1 = 2$ ,  $b_2 = 1$ ,  $J_1 = 3$ ,  $J_2 = 1$  in both figures. The numbers 1-4 in (a) mark the diagonal elements of  $H_p$ , with which the corresponding eigenenergies merge as  $t \rightarrow \pm\infty$ , e.g., 1 is for  $(H_p)_{11}$ . The dashed blue and green arrows in (b) show two semiclassical trajectories that contribute to the amplitude  $S_{33}$ . Each trajectory jumps through one of the gaps but avoids the other gap.

In the Heisenberg's picture, the evolution of  $\hat{\mathbf{a}}_p$  is described by a Schrödinger-like equation:

$$i\hat{\mathbf{a}}_p(t) = H_p(t)\hat{\mathbf{a}}_p(t), \quad (11)$$

and the initial ground state (3) corresponds to the initially fully filled Fermi sea, for all fermions as  $t \rightarrow -\infty$ .

In Fig. 2, we show the time-dependent eigenvalues of the matrix  $H_p$  for a broken chiral symmetry:  $J_1 \neq J_2$  and  $b_1 > b_2$ . Figure 2(a) shows that at  $p = 0$  two of the four energy levels experience exact level crossings, whereas Fig. 2(b) illustrates that this degeneracy is lifted for nonzero  $p$ . This means that the chiral asymmetry does not destroy the Dirac points in the spectrum, which must produce a power law density of excitations according to the Kibble-Zurek mechanism.

We will consider the evolution with the Hamiltonian (5) during the time interval  $t \in (-\infty, +\infty)$ . In terms of the phase diagram in Fig. 1, this means that the system passes through two phase transition points. In the adiabatic limit, the initial ground state (3) transfers, as  $t \rightarrow +\infty$ , into the ground state with all spins up

$$|\uparrow\uparrow\uparrow\uparrow \dots \uparrow\uparrow\rangle. \quad (12)$$

This is the fermionic vacuum, and the free fermions are the elementary excitations. The numbers of excitations (non-flipped spins) on the even and odd sublattices at the end of the protocol are given by, respectively,

$$N_{ex}^e \equiv \sum_{n=1}^N \langle \hat{c}_n^\dagger \hat{c}_n \rangle = \sum_p^N \langle \hat{c}_p^\dagger \hat{c}_p \rangle, \quad (13)$$

$$N_{ex}^o \equiv \sum_{n=1}^N \langle \hat{d}_n^\dagger \hat{d}_n \rangle = \sum_p^N \langle \hat{d}_p^\dagger \hat{d}_p \rangle, \quad (14)$$

where, in the Schrödinger's picture, the averaging is taken over the state at  $t = +\infty$ .

The Hamiltonian (9) has the form of a multistate Landau-Zener (MLZ) model, i.e., can be written as  $H(t) = A + Bt$  with hermitian matrices  $A$  and  $B$ . The MLZ models have been extensively studied previously [15–17]. Our case with  $J_1 \neq J_2$  is generally not solvable [18] but the MLZ theory provides exact formulas for some of the evolution matrix elements [19], which we will utilize.

Let us define the  $S$ -matrix

$$S = S(p) \equiv U^p(T, -T)_{T \rightarrow \infty}, \quad (15)$$

where  $U^p(T, -T)$  is the evolution matrix over the time interval  $t \in (-T, T)$  with the Hamiltonian  $H_p$ . We can say that the operators at  $t = \pm\infty$  are related by

$$\hat{a}_p(+\infty) = S\hat{a}_p, \quad (16)$$

where  $\hat{a}_p \equiv \hat{a}_p(-\infty)$ . Hence,

$$\begin{aligned} \hat{c}_p(+\infty) &= S_{11}\hat{c}_p + S_{12}\hat{c}_{-p}^\dagger + S_{13}\hat{d}_p + S_{14}\hat{d}_{-p}^\dagger, \\ \hat{c}_p^\dagger(+\infty) &= S_{11}^*\hat{c}_p^\dagger + S_{12}^*\hat{c}_{-p} + S_{13}^*\hat{d}_p^\dagger + S_{14}^*\hat{d}_{-p}. \end{aligned} \quad (17)$$

The number of excitations (13) can be evaluated in the Heisenberg's picture, i.e., taking the average of the operators at  $t = +\infty$  with respect to the initial state (3):

$$N_{ex}^e(+\infty) = \sum_p (|S_{11}|^2 + |S_{13}|^2). \quad (18)$$

The survival amplitudes for states with the highest energy level slopes are known exactly for any MLZ model [15]:

$$S_{11} = S_{22} = e^{-\frac{\pi|g|^2}{b_1 - b_2} - \frac{\pi|\gamma|^2}{b_1 + b_2}}. \quad (19)$$

Another exact relation of MLZ theory is for the transition amplitudes between the levels with the two highest slopes [19]:

$$S_{11}S_{33} + |S_{13}|^2 = e^{-\frac{2\pi|\gamma|^2}{b_1 + b_2}}. \quad (20)$$

This does not fix  $S_{13}$  because  $S_{33}$  is not known. Fortunately, model (9) has discrete symmetries leading to

$$\begin{aligned} S_{11} + S_{22} &= S_{33} + S_{44}, \\ S_{33}(-p) &= S_{44}(p), \end{aligned} \quad (21)$$

the derivations of which are given in Appendix B of the SM. Equations (19)-(21) then predict

$$\sum_p |S_{13}|^2 = \sum_p \left[ e^{-\frac{2\pi|\gamma|^2}{b_1 + b_2}} - |S_{11}|^2 \right]. \quad (22)$$

This formula is exact. Hence, without approximations

$$N_{ex}^e(+\infty) = \sum_p e^{-\frac{2\pi|\gamma|^2}{b_1 + b_2}}. \quad (23)$$




FIG. 3. (a): Density of excitations of the even-site sublattice as a function of the quenching rate,  $b_1 = 2\beta$ ,  $b_2 = \beta$ . Black solid curve and blue dashed curve correspond to, respectively, the exact prediction (24) at  $J_1 = J_2 = 1.5$  and perturbed coupling  $J_1 = 2$ ,  $J_2 = 1$ . (b): The numerically calculated transition probability  $\sum_p |S_{33}|^2$  at  $J_1 = 2$ ,  $J_2 = 1$ ,  $b_1 = 2\beta$  and  $b_2 = \beta$ . Triangles are the numerical data. The black dashed line is the power-law scaling (31) prediction. The insets resolve the fast Stueckelberg oscillations around the mean power-law tail. The filled triangles are the same points in the main figure and the insets.

As  $N \rightarrow \infty$ , we replace  $\sum_p/N \rightarrow \int_{-\pi}^{\pi} dp/(2\pi)$ , and find the density of excitations (per unit cell) in the thermodynamic limit

$$\rho_{ex}^e(+\infty) = e^{-\frac{2\pi(J_1^2 + J_2^2)}{|b_1 + b_2|}} I_0 \left( \frac{4\pi J_1 J_2}{b_1 + b_2} \right), \quad (24)$$

where  $I_0(x)$  is the modified Bessel function of the first kind. In the leading order of small  $b_1 + b_2$

$$\rho_{ex}^e(+\infty) \approx e^{-\frac{2\pi(J_1 - J_2)^2}{|b_1 + b_2|}} \frac{\sqrt{b_1 + b_2}}{2\pi\sqrt{2J_1 J_2}}. \quad (25)$$

This is the central result of our article. For any  $J_1 \neq J_2$  the excitations on the even-sublattice are suppressed exponentially rather than by a power-law. The latter is found only for a symmetric case with  $J_1 = J_2$  but  $b_1 > b_1$ . Complete exact solution for this special case is presented in Appendix D of the SM. We also note that the above solutions do not smoothly carry over  $b_1 = b_2$  point, to which our exact MLZ analysis does not apply because of the degeneracy of the time-dependent Zeeman coupling.

Equation (25) does not contradict the Kibble-Zurek mechanism because the power-law for excitation density does appear on the odd-site sublattice. Consider

$$N_{ex}^o(+\infty) = \sum_p (|S_{33}|^2 + |S_{31}|^2). \quad (26)$$

Here,  $|S_{31}|^2$  can be obtained exactly because  $|S_{31}| = |S_{13}|$ . According to Eq. (22), it produces an exponentially suppressed contribution. We also prove in Appendix C of the SM that  $\sum_p |S_{33}|^2 \geq \sum_p |S_{11}|^2$ , so,  $N_{ex}^o \geq N_{ex}^e$ . Hence, only  $S_{33}$  contains the information about the power-law scaling.

The exact crossings of the 2nd and 3rd adiabatic levels of  $H_p$  happen at  $p = 0$  and time moments

$$t_{1,2} = \mp 2\sqrt{J_1 J_2 / (b_1 b_2)}. \quad (27)$$

We interpret  $t_{1,2}$  as the moments of the phase transitions. By setting  $t = t_{1,2} + \delta t$  and  $p = \delta p$ , with small  $\delta p$  and  $\delta t$ , we find an effective Hamiltonian for the interactions within the subspace of  $H_p$  spanned by the two eigenstates, whose energies become nearly degenerate near the critical points:

$$\hat{H}_{\text{eff}} \approx \frac{2\sqrt{J_1 J_2 b_1 b_2}(\sqrt{J_1 J_2} \delta p \hat{\tau}_x + \sqrt{b_1 b_2} \delta t \hat{\tau}_z)}{\sqrt{(b_2 J_1 + b_1 J_2)(b_1 J_1 + b_2 J_2)}}, \quad (28)$$

where  $\hat{\tau}_x$  and  $\hat{\tau}_z$  are the Pauli operators that act in subspace of two states with closest to one Dirac point energies. The Landau-Zener formula applied to Hamiltonian (28) returns the probability of a non-adiabatic excitation after a transition through one Dirac point:

$$P_{ex} = \exp\left(-\frac{2\pi(\delta p)^2(J_1 J_2)^{3/2}}{\sqrt{(b_2 J_1 + b_1 J_2)(b_1 J_1 + b_2 J_2)}}\right). \quad (29)$$

Our system passes through two Dirac points. An excitation created near the first point should remain excited after the second point. The system may also not produce an excitation during the first crossing but produce it at the second one. Generally, there is interference between such evolution paths (Stueckelberg's oscillations). If we disregard this interference, the probabilities of the two possibilities simply sum, i.e.,

$$P_{2ex} = 2P_{ex}(1 - P_{ex}). \quad (30)$$

Integrating  $P_{2ex}$  over  $\delta p \in (-\infty, +\infty)$ , we find that in the adiabatic limit:

$$\sum_p \frac{|S_{33}|^2}{N} \approx \frac{2 - \sqrt{2}}{4\pi(J_1 J_2)^{3/4}} [(b_2 J_1 + b_1 J_2)(b_1 J_1 + b_2 J_2)]^{1/4}. \quad (31)$$

This is the estimate of the excitation density on the odd-site sublattice in the nearly-adiabatic limit. In Fig. 3, the numerical simulations confirm the power-law scaling trend for the parameter dependence in (31), which is modulated by fast Stueckelberg oscillations. We also note that (31) reproduces correctly the scaling prediction of the uniform chain model (4) if we set  $b_1 = b_2 = 2\beta$  and  $J_1 = J_2$ .

Hence, as  $t \rightarrow +\infty$ , the remaining excitations on the odd-site sublattice are suppressed according to a power-law, in contrast to the exponential suppression on the even-site sublattice. At the intermediate moment,  $t = 0$ ,

excitations take the form of superpositions of kinks. The asymmetry appears then too, i.e., some of the excitations are exponentially suppressed in comparison to the others but this effect does not have a simple interpretation in terms of different kink types. We leave such details to Appendix E of the SM.

Our results illustrate how the Kibble-Zurek mechanism works when a part of a system is not observable. The latter may happen in atomic gases, in which optically observable spins interact via spins of other atomic species [20]. In the asymmetric Ising chain, if the spin excitations on the odd sites are not observable, the even-site spins still create a visible ferromagnetic state at zero external field and polarize when the external field is strong. Hence, looking only at the even spins, one can expect that this system goes through the same phase transitions as the uniform Ising chain and thus produces excitations with a power-law scaling. Our solution shows, however, that all observable spin excitations are suppressed in this case exponentially. A power-law tail of the excitations is hidden then in the non-observable spins.

Summarizing, we demonstrated that small differences between interacting spins in a simple spin chain can lead to an exponentially large effect when passing slowly through a phase transition. The underlying mechanism arguably processes universalities: strong but symmetric interactions compete with each other near a critical point, where subtle interaction differences play a decisive role. As long as a symmetry breaking opens the gap for certain types of quasi-particles while not destroying the critical point, the exponentially large density separation of different quasi-particles should happen after the passage through the phase transition independently of the model's microscopic details.

The Kibble-Zurek mechanism for Ising-type quantum phase transition has been recently studied experimentally in ultracold atoms [9]. In such systems an asymmetry can be introduced by placing atoms in a periodic potential without the inversion symmetry. If the time-linear field ramp is induced by changing the AC-frequency in the rotating-wave approximation, the sweeping of the detuning frequency across an optical resonance effectively mimics the field changes in range  $B_z \in (+\infty, -\infty)$  [21, 22]. Hence, the demonstration of our effect requires only simple modification of the control protocols in already accessible for studies dynamic phase transitions.

Let us also speculate about possible applications. Different isotopes have different spin interactions in an ultracold atomic mixture [23]. Hence, an adiabatic passage through a quantum critical point can induce spin excitations in this mixture, overwhelmingly, for only one of the isotopes. One can then separate such excited atoms using the magnetic deflection approach from [24], and thus develop a technology for isotope separation. The asymmetry of matter and antimatter in our Universe may also be viewed now as a result of a hypothetical transition through a quantum critical point during cosmic inflation, when the matter could be excited from the vacuum via



quantum nonadiabatic processes [25, 26].

The dynamic phase transitions are found broadly, from cosmological models to the experiments with superconductors and ultracold atoms. They are likely responsible for the scaling of mistakes that limit the quantum annealing computation techniques [27]. Fortunately, different systems show universal behavior, which is driven by relatively simple effects near the critical point. The quasiparticle separation is one of such effects that remain relevant near the critical point and thus can be used for control of complex quantum systems.

## ACKNOWLEDGMENTS

This work was carried out under the support of the U.S. Department of Energy, Office of Science, Basic Energy Sciences, Materials Sciences and Engineering Division, Condensed Matter Theory Program. B.Y. also acknowledges partial support from the Center for Nonlinear Studies.

- 
- [1] L. Canetti, M. Drewes, and M. Shaposhnikov, “Matter and antimatter in the universe,” *New Journal of Physics* **14**, 095012 (2012).
  - [2] A. D. Sakharov, “Violation of CP invariance, C asymmetry, and baryon asymmetry of the universe,” *Soviet Physics Uspekhi* **34**, 392 (1991).
  - [3] T. W. B. Kibble, “Some implications of a cosmological phase transition,” *Physics Reports* **67**, 183–199 (1980).
  - [4] T. W. B. Kibble, “Topology of cosmic domains and strings,” *Journal of Physics A: Mathematical and General* **9**, 1387 (2001).
  - [5] T. W. B. Kibble, “Phase-transition dynamics in the lab and the universe,” *Physics Today* **60**, 47–52 (2007).
  - [6] W. H. Zurek, “Cosmological experiments in superfluid helium?” *Nature* **317**, 505–508 (1985).
  - [7] W. H. Zurek, “Cosmological experiments in condensed matter systems,” *Physics Reports* **276**, 177–221 (1996).
  - [8] Adolfo del Campo and Wojciech H Zurek, “Universality of phase transition dynamics: Topological defects from symmetry breaking,” *International Journal of Modern Physics. A, Particles and Fields, Gravitation, Cosmology* **29**, 1430018 (2014).
  - [9] A. Keesling, A. Omran, H. Levine, H. Bernien, H. Pichler, S. Choi, R. Samajdar, S. Schwartz, P. Silvi, S. Sachdev, P. Zoller, M. Endres, M. Greiner, V. Vuletić, and M. D. Lukin, “Quantum Kibble-Zurek mechanism and critical dynamics on a programmable rydberg simulator,” *Nature* **568**, 207–211 (2019).
  - [10] M. M. Rams, J. Dziarmaga, and W. H. Zurek, “Symmetry breaking bias and the dynamics of a quantum phase transition,” *Physical Review Letters* **123**, 130603 (2019).
  - [11] L.-Y. Qiu, Y.-B. Liang, H.-Y. and Yang, H.-X. Yang, T. Tian, Y. Xu, and L.-M. Duan, “Observation of generalized Kibble-Zurek mechanism across a first-order quantum phase transition in a spinor condensate,” *Science Advances* **6**, 7292 (2020).
  - [12] J. Dziarmaga, “Dynamics of a quantum phase transition: exact solution of the quantum ising model,” *Physical Review Letters* **95**, 245701 (2005).
  - [13] W. H. Zurek, U. Dorner, and P. Zoller, “Dynamics of a quantum phase transition,” *Physical Review Letters* **95**, 105701 (2005).
  - [14] M. Heyl, A. Polkovnikov, and S. Kehrein, “Dynamical quantum phase transitions in the transverse-field ising model,” *Physical Review Letters* **110**, 135704 (2013).
  - [15] S Brundobler and Veit Elser, “S-matrix for generalized Landau-Zener problem,” *Journal of Physics A: Mathematical and General* **26**, 1211 (1993).
  - [16] Nikolai A Sinitsyn, Emil A Yuzbashyan, Vladimir Y Chernyak, Aniket Patra, and Chen Sun, “Integrable Time-Dependent quantum hamiltonians,” *Physical Review Letters* **120**, 190402 (2018).
  - [17] N. A. Sinitsyn, “Solvable four-state Landau-Zener model of two interacting qubits with path interference,” *Physical Review B* **92**, 205431 (2015).
  - [18] R. K. Malla and M. E. Raikh, “Loss of adiabaticity with increasing tunneling gap in nonintegrable multistate Landau-Zener models,” *Physical Review B* **96**, 115437 (2017).
  - [19] F. Li, C. Sun, V. Y. Chernyak, and N. A Sinitsyn, “Multistate Landau-Zener models with all levels crossing at one point,” *Physical Review A* **96**, 022107 (2017).
  - [20] D. Roy, L. Yang, S. A. Crooker, and N. A. Sinitsyn, “Cross-correlation spin noise spectroscopy of heterogeneous interacting spin systems,” *Scientific Reports* **5**, 9573 (2015).
  - [21] C. Sun and N. A. Sinitsyn, “Exact transition probabilities for a linear sweep through a Kramers-Kronig resonance,” *J. Phys. A: Math. Theor.* **48**, 505202 (2015).
  - [22] R. T. Brierley, C. Creatore, P. B. Littlewood, and P. R. Eastham, “Adiabatic state preparation of interacting two-level systems,” *Phys. Rev. Lett.* **109**, 043002 (2012).
  - [23] M. Tomza, “Energetics and control of ultracold Isotope-Exchange reactions between heteronuclear dimers in external fields,” *Physical Review Letters* **115**, 063201 (2015).
  - [24] Thomas R Mazur, Bruce Klappauf, and Mark G Raizen, “Demonstration of magnetically activated and guided isotope separation,” *Nature Physics* **10**, 601–605 (2014).
  - [25] Michael A Fedderke, Edward W Kolb, and Mark Wyman, “Irruption of massive particle species during inflation,” *Physical Review D* **91**, 063505 (2015).
  - [26] Seishi Enomoto and Tomohiro Matsuda, “The exact WKB for cosmological particle production,” (2020), [arXiv:2010.14835 \[hep-ph\]](https://arxiv.org/abs/2010.14835).
  - [27] A. Pearson, A. Mishra, Hen I., and D. A. Lidar, “Analog errors in quantum annealing: doom and hope,” *NPJ Quantum Information* **5** (2019).