



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Observing Dynamical Quantum Phase Transitions through Quasilocal String Operators

Souvik Bandyopadhyay, Anatoli Polkovnikov, and Amit Dutta

Phys. Rev. Lett. **126**, 200602 — Published 17 May 2021

DOI: [10.1103/PhysRevLett.126.200602](https://doi.org/10.1103/PhysRevLett.126.200602)

Observing dynamical quantum phase transitions through quasi-local string operators

Souvik Bandyopadhyay¹, Anatoli Polkovnikov² and Amit Dutta¹

¹ *Department of Physics, Indian Institute of Technology Kanpur, Kanpur 208016, India and*

² *Department of Physics, Boston University, Boston, Massachusetts, USA*

We analyze signatures of the dynamical quantum phase transitions (DQPTs) in physical observables. In particular, we show that both the expectation value and various out of time order correlation functions of the finite length product or string operators develop cusp singularities following quench protocols, which become sharper and sharper as the string length increases. We illustrated our ideas analyzing both integrable and non-integrable one-dimensional Ising model showing that these transitions are robust both to the details of the model and to the choice of the initial state.

Understanding out-of-equilibrium dynamics of quantum many body systems is an exciting field of recent research both from theoretical and experimental viewpoints [1–6]. In this regard, dynamical and equilibrium quantum phase transitions (DQPTs), manifested as real time singularities in time-evolving integrable and non-integrable quantum systems, are indeed an emerging and intriguing phenomena [7–11]. To probe DQPTs, a quantum many body system is prepared initially in the ground state $|\psi(0)\rangle$ of some Hamiltonian. At time $t = 0$ a parameter λ of the Hamiltonian is suddenly changed to say $\lambda = \lambda_f$ and the subsequent temporal evolution of the system generated by the time-independent final Hamiltonian $H(\lambda_f)$ is tracked. DQPTs occur at those instants of time t when the evolved state $|\psi(t)\rangle = \exp(-iH(\lambda_f)t)|\psi(0)\rangle$, becomes orthogonal to the initial state $|\psi(0)\rangle$, i.e., the so-called Loschmidt overlap (LO), $\mathcal{L}(t) = |\langle\psi(0)|\psi(t)\rangle|^2$ vanishes. At those critical instants, the so-called dynamical free energy density (or the rate function of the return probability) defined as $\mathcal{F} = -(1/N) \log |\mathcal{L}(t)|$, N being the system size, develop nonanalytic singularities (cusps in 1D systems) in the thermodynamic limit [7].

Following the initial proposal [7], there have been a plethora of studies investigating intricacies of DQPTs in several integrable and non-integrable, one dimensional (as well as two dimensional) quantum systems occurring subsequent to both sudden [12–45], and smooth [46–49] ramping protocols. The notion of DQPT has also been generalised for *mixed* initial states [50–53] and finally also in open quantum systems [54]. Analogous to equilibrium phase transitions, it has been established that one expects universal scaling of the dynamical free energy density near the critical instants with identifiable critical exponents. (For reviews on various aspects of DQPTs, we refer to [55–57].) Remarkably, these non-analyticities have been detected experimentally subsequent to a rapid quench from a topologically trivial system into a Haldane-like system [58].

Recently, DQPTs were also experimentally [59] detected in trapped-ion setups simulating a long-range interacting transverse field Ising model (TFIM). Starting from a degenerate ground state manifold, it was established that following a quench in an interacting chain of

$^{40}\text{Ca}^+$ ions, the dynamical free energy density develops cusp like singularities at critical instants signalling DQPTs. However, a thorough understanding of the phenomena in harmony with the now well-understood notion of equilibrium quantum phase transitions is far from being complete. [Although DQPTs may be characterised by a topological dynamical order parameter \[20, 30\] indicating the emergence of momentum space vortices at critical instants, there is an ongoing search for spatially local observables which are able to capture these non-equilibrium quantum phase transitions \[10, 11, 60\].](#) There has been many attempts aiming at finding real-time observable effects of the dynamical transitions on many-body observables such as work distributions and the growth of entanglement in quenched systems [43, 61]. A very interesting perspective on DQPTs was put forward in another recent trapped ion quantum simulator [62], where the authors experimentally studied singularities in the domain wall statistics following a sudden quench of the transverse magnetic field in the system with long range Ising type interactions.

One obvious drawback of DQPTs is that they are manifested in the overlap of the wave functions, which is difficult to observe experimentally. However, the experiment of J. Zhang et. al. (Ref. [62]) showed that DQPTs can be also manifested in the behavior of non-local, string like observables. But the precise mathematical connection between DQPTs and this experiment remains unclear. Currently, the broad questions which remain under scrutiny are: (i) Can the singular transitions at DQPTs be captured in the real time behavior of strong observable quantities? (ii) What can one infer about the spatio-temporal locality of the observables required to detect the DQPTs? (iii) Is it possible to obtain the critical exponents associated with the dynamical transitions through a measurement of time-evolving observables [38]? In this work, we approach these issues by defining finite length string operators. As the length of these operators increases they effectively play the role of the projection operators to polarized spin states. Using an exact diagonalisation scheme [63, 64], we show that these observables are able

to capture the critical singularities in their time evolving expectations and temporal correlators. We observe that early time rate function of the out-of-time order correlator (OTOC), which is an important quantity to study scrambling of information in chaotic systems [65] and quantum phase transitions [42, 66], quickly becomes non-analytic at the critical instants with the operator length and show universal critical scaling near DQPTs.

To exemplify, we consider a ferromagnetic transverse field Ising model (TFIM) with nearest and next nearest neighbour interactions ($J > 0$ and $J_2 > 0$) and a non-commuting external field h , having N spins,

$$H = -J \sum_{i=1}^N \sigma_z^i \sigma_z^{i+1} - J_2 \sum_{i=1}^N \sigma_z^i \sigma_z^{i+2} - h \sum_{i=1}^N \sigma_x^i \quad (1)$$

under periodic boundary conditions where σ 's are the Pauli Matrices satisfying standard $\text{su}(2)$ commutation relations. The presence of both the nearest neighbour and the next nearest neighbour interactions renders the model non-integrable with an integrable point at $J_2 = 0$.

For concreteness we start from a fully spin-polarised state $|\uparrow\uparrow\uparrow \dots\rangle$, which is a ground state corresponding to zero transverse field. However, this assumption can be lifted without affecting the results of this work as we will observe similar non-analyticities in the infinite temperature OTOCs (also see Supplemental Material Ref. 67 for a discussion on generic initial states). As an observable we consider the translationally invariant Pauli string operator,

$$P_n(0) = \frac{1}{N} \sum_{i=1}^N \frac{1}{2^n} \prod_i^{i+n} (\mathbb{I}_i + \sigma_i^z), \quad (2)$$

having a finite string size n of a system of size N and probe its time evolution following a sudden quench of the transverse field h at $t = 0$. When the strings in Eq. (2) span the whole system (i.e., when $n = N$), the quantity $P_n(0)$ simply reduces to the projector over the complete initial state $|\psi(0)\rangle$.

Originally DQPTs are defined through emergent non-analyticities of the rate function [7, 12] defined as

$$f(t) = - \lim_{N \rightarrow \infty} \frac{1}{N} \log \text{LO}(t), \quad \text{where } \mathcal{L}(t) = |\langle \psi(0) | \psi(t) \rangle|^2. \quad (3)$$

These non-analyticities develop at critical instants of time ($t = t_c$). It is easy to see that formally the rate function can be understood through the time-dependent expectation value of the ground state projection operator $P = |\psi(0)\rangle\langle\psi(0)|$ in the Heisenberg representation: $P(t) = \exp[iHt]P(0)\exp[-iH(t)]$, such that,

$$f(t) = - \lim_{N \rightarrow \infty} \frac{1}{N} \log \langle P(t) \rangle, \quad (4)$$

where, $\langle \dots \rangle = \langle \psi(0) | \dots | \psi(0) \rangle$. In the one-dimensional Ising models, close to the critical point $|f(t) - f(t_c)| \sim |t - t_c|^\alpha$ with the universal critical exponent $\alpha = 1$ [7, 12, 19]. For the initially polarized state $|\uparrow\uparrow\uparrow \dots\rangle$, clearly $P \equiv P_N(0)$. The idea of this work is to look into the expectation value of the operator $P_n(t)$ ($n \leq N$) instead of the full projector through the observable

$$\mathcal{O}_n(t) = -\frac{1}{n} \log \langle P_n(t) \rangle, \quad (5)$$

We find (see Fig. 1) that the observable $\mathcal{O}_n(t)$ develops emergent cusp singularities at the critical instants in both integrable and integrability-broken systems, thus establishing that DQPTs can be detected using normal physical observables. As it is evident from the plot the singularities quickly develop with increasing n , becoming very sharp for $n \gtrsim 6$ for the parameters used to generate this plot. The singularities were also seen to develop with increasing string length in the thermodynamic limit ($N \rightarrow \infty$) following an exact calculation for the integrable situation as the expectation $\langle P_n(t_c) \rangle$ vanish at the critical instants exponentially fast in n [67].




Figure 1. (Color online) Emergent cusp singularities in the observable $\mathcal{O}_n(t)$ (see Eq. (5)) for finite string lengths n following an integrable sudden quench in the transverse field (with $4J = 1.0, 2J_2 = 0.0$), from the completely polarised ferromagnetic ground state ($2h = 0$) to a paramagnetic phase ($2h = 4.0$). The emergence of the singularities are shown for various string lengths comparing the cusps with increasing string length n . (Inset) The same observable following a quench in the non-integrable Ising chain ($4J = 1.0, 4J_2 = 0.5$). The simulations have been performed in a chain containing $N = 16$ spins using exact diagonalisation.

These results can be generalized for quenches starting from an arbitrary initial state [67], like a ground state corresponding to a finite transverse field within the paramagnetic phase or a mixed initial density matrix, for example, corresponding to a finite temperature ensemble. To see how it works, let us observe that for any initial density matrix $\rho(0)$ we have,




Figure 2. (Color online) (a) The universal scaling of the OTOCs for an integrable chain near the critical point. The deviation $\log|C_n(t) - C_n(t_0)|$ vs $\log|t - t_0|$ (in the region shown within the vertical dashed lines in inset) exhibits a scaling collapse for different string lengths n (see also Ref. [68]); here we have chosen an instant $t_0 = 1.3$ to be a point near the critical point such that it captures the universal linear collapse region. (Inset) Corresponding emergent cusp singularities in the rate function of the OTOC C_n (see Eq. (8)) for finite string lengths following a sudden quench in the transverse field (with $4J = 1.5, 4J_2 = 0$ for $N = 16$) from the completely polarised ferromagnetic ground state ($2h = 0$) to a paramagnetic phase ($2h = 2.5$). (b) The linear scaling of C_n (in the region shown within the vertical dashed lines in inset) and the corresponding cusps (Inset) at critical times in a non-integrable chain (with $4J = 1.0, 4J_2 = 1.0$ for $N = 16$) subsequent to a quench from $2h = 0.0$ to $2h = 4.5$ and choosing a $t_0 = 2.1$ near the critical point. Both the integrable and the non-integrable situations exhibit a universal critical exponent of $\alpha \sim 1$ as seen by the linear fits depicted by black solid lines. The linear scaling in both (a) and (b) have been shown for $t < t_c$ (left of the critical instant). Similar scaling has also been checked to hold for $t > t_c$.

$$C(t) \equiv \text{Tr}[\rho(t)P(0)P(-t)P(0)] = p(t)\mathcal{L}(t), \quad (6)$$

where, $p(t) = \langle \psi(0)|\rho(t)|\psi(0) \rangle$. It is easy to check that,

$$\text{Tr}[\rho(t)P(0)P(-t)P(0)] = \text{Tr}[\rho(0)P(t)P(0)P(t)],$$

such that,

$$C(t) = p(t)\mathcal{L}(t) = \text{Tr}[\rho(0)P(t)P(0)P(t)]. \quad (7)$$

The function $p(t)$ is a projection of the time dependent density matrix to the ground state $|\psi(0)\rangle$. In some situations, like when the initial state is the ground state of the initial Hamiltonian, $p(t) = \mathcal{L}(t)$ such that $C(t) \propto \mathcal{L}^2(t)$. If the initial state, on the other hand, is stationary with respect to the final Hamiltonian $H(\lambda_f)$ then $p(t) = \text{const}(t)$ and $C(t) \propto \mathcal{L}(t)$. In both cases that $-\frac{1}{N} \log[C(t)]$ is expected to show the exact same singularities as the rate function and scale similar to $f(t)$ with the same exponent α near the critical point.

The function $C(t)$ is nothing but the OTOC of the projection operator, which can be measured through, for example, quantum echo protocols [22, 69–72].

Like before instead of the full $C(t)$ we define the quasi-local truncated OTOC

$$C_n(t) = -\frac{1}{n} \log \langle P_n(t)P_n(0)P_n(t) \rangle, \quad (8)$$

where the average now is over the initial density matrix $\rho(0)$, which we first take to be the same as before $\rho(0) = |\psi(0)\rangle\langle\psi(0)|$ and later show that the results qualitatively do not change if we start from general mixed ensembles.

Similar to the expectation $\mathcal{O}_n(t)$ the post-quench rate function $C_n(t)$ of the OTOC develops non-analytic cusp singularities (see Fig. 2(a),2(b)) even for finite length string operators $P_n(0)$. We find that the observable $C_n(t)$ apart from being singular at critical times, also show a collapse to an universal scaling for sufficiently long strings near the critical point $t = t_c$, having critical exponent $\alpha \sim 1$ for quenches in both the integrable and non-integrable chains. Consequently, the growth rate of the OTOC in its early time dynamics, shows singular discontinuous jumps at the critical instants of DQPTs. In Fig. 3 we show that the jump singularities in the early-time OTOC growth rate at critical instants,

$$\mathcal{G}_n(t) = \frac{dC_n(t)}{dt}, \quad (9)$$

for finite string operators $P_n(0)$ emerge with increasing sharpness for increasing string lengths.

Although we have demonstrated the development of emergent singularities through a quenched TFIM, the phenomena is not explicitly model dependent as has been




Figure 3. (Color online) Emergent jump discontinuities in the OTOC growth rate \mathcal{G}_n (see Eq. (9)) following an integrable sudden quench in the transverse field with the quench parameters being same as in Fig. 2(a). The quantity $\mathcal{G}_n(t)$ vs t , is shown for various n comparing the sharp jumps at the critical instant $t = t_c(n)$ with increasing string length. The critical instants $t_c(n)$ corresponding to the peaks in the rate function of the OTOC, for a string length n can be determined by the emergent jump singularities in the OTOC growth rate (shown by vertical dashed lines in the figure where the rate crosses zero, with $t_c(14)$ explicitly marked). Similar jump discontinuities were also observed in the non-integrable situations.

previously established in literature. We also stress that the information about the complete initial ground state is not a necessary requirement to study the non-equilibrium phase transitions which can also be detected in temporal correlators of the strings when summed over *arbitrary* complete basis states. To elaborate, consider the infinite temperature auto-correlator and OTOCs,

$$\begin{aligned}\tilde{\mathcal{O}}_n(t) &= -\frac{1}{n} \log \text{Tr} [P_n(t)P_n(0)], \\ \tilde{\mathcal{C}}_n(t) &= -\frac{1}{n} \log \text{Tr} [P_n(t)P_n(0)P_n(t)],\end{aligned}\quad (10)$$

respectively. Remarkably, we find that though the traced correlations are independent of the full initial state $|\psi(0)\rangle$, they develop cusp like singularities at the critical instants for finite but sufficiently long strings and are therefore sufficient to observe the non-equilibrium transitions (see Fig. 4). This also establishes that the emergent critical behaviour stays robust in the observables even when the dynamics starts from *arbitrary* excited initial states.

In conclusion, we demonstrated that DQPTs can be observed experimentally as post-quench singularities developing in time for string like observables. These singularities become sharper with increasing string length. We showed that for the initial ground state these singularities emerge in the expectation values of string operators, while for generic, even infinite temperature, initial states they appear in the OTOC of such operators




Figure 4. (Color online) (Solid lines) The logarithm of the infinite temperature OTOC $\tilde{\mathcal{C}}_n(t)$ (see Eq. (10)) at the critical instants show sharper singularities with increasing peak heights as the string size n increases. (Dotted lines) The corresponding post-quench auto-correlation functions $\tilde{\mathcal{O}}_n(t)$. The system is initially chosen to be in a completely polarised ground state of an integrable TFIM having a transverse field $h = 0.0$, $J = 0.1$ and suddenly quenched to a final field in the paramagnetic phase $h = 2.5$ at $t = 0$. The simulations have been performed for a chain of $N = 8$ spins while considering finite strings of lengths n . The singularities were seen to become sharper for higher string sizes.

and can be detected through echo-type protocols. It is interesting that similar signatures of post-quench singularities in two-spin observables were found in Ref. [10, 60]. The precise relation of the results of that work to the present one regarding DQPTs are yet to be understood.

One can also check (see Ref. [67]) that the coefficient of variation of the observable $P_n(t_c)$ at a critical point remains finite and non-zero for local strings (having a finite length n) even in the thermodynamic limit, unlike that of the full projector $P_L(t_c)$ which diverges exponentially in system size. This suggests a further experimental advantage of the proposed observables as unlike the full Loschmidt overlap, it takes only a finite number of measurements to accurately determine the expectations of finite projectors even for infinitely large system sizes.

Note added:

During preparation of the manuscript we came across a similar study [68] which also exhibit the efficacy of local operators in capturing DQPTs.

S.B. acknowledges support from a PMRF fellowship, MHRD, India. A.D. and A.P. acknowledge financial support from SPARC program, MHRD, India. A.P. was supported by NSF DMR-1813499 and AFOSR FA9550-16-1-0334. We acknowledge Sourav Bhattacharjee and

Somnath Maity for comments. We acknowledge QuSpin and HPC-2010, IIT Kanpur for computational facilities.

-
- [1] A. Polkovnikov, K. Sengupta, A. Silva, & M. Vengalattore, Colloquium: Nonequilibrium dynamics of closed interacting quantum systems, *Rev. Mod. Phys.* **83**, 863 (2011).
- [2] P. Calabrese, F. H. L. Essler, and M. Fagotti, "Quantum Quench in the Transverse-Field Ising Chain", *Phys. Rev. Lett.* **106**, 227203 (2011).
- [3] A. Dutta, G. Aeppli, B. K. Chakrabarti, U. Divakaran, T. Rosenbaum, T. & D. Sen, *Quantum Phase Transitions in Transverse Field Spin Models: From Statistical Physics to Quantum Information* (Cambridge University Press, Cambridge, 2015).
- [4] J. Eisert, M. Friesdorf, & C. Gogolin, Quantum many-body systems out of equilibrium, *Nat. Phys.* **11**, 124 (2015).
- [5] L. D'Alessio, Y. Kafri, A. Polkovnikov, & M. Rigol, From Quantum Chaos and Eigenstate Thermalization to Statistical Mechanics and Thermodynamics, *Adv. Phys.* **65**, 239 (2016).
- [6] P. Calabrese, F. H. L. Essler, & G. , J. Mussardo, *Stat. Mech.: Theo. and Expt, special issue Quantum Integrability in Out of Equilibrium Systems* **2016** .
- [7] M. Heyl, A. Polkovnikov, & S. Kehrein, Dynamical Quantum Phase Transitions in the Transverse-Field Ising Model, *Phys. Rev. Lett.* , **110**, 135704 (2013).
- [8] M. Białończyk and B. Damski, Locating quantum critical points with Kibble-Zurek quenches, *Phys. Rev. B* **102**, 134302 (2020).
- [9] P. Titum, J. T. Iosue, J. R. Garrison, A. V. Gorshkov, and Z-X Gong, Probing Ground-State Phase Transitions through Quench Dynamics, *Phys. Rev. Lett.* **123**, 115701 (2019).
- [10] S. Bhattacharyya, S. Dasgupta, A. Das, "Signature of a Continuous Quantum Phase Transition in Nonequilibrium Energy Absorption: Footprints of Criticality on Highly Excited States", *Scientific Reports* **5**, 16490 (2015).
- [11] S. Roy, R. Moessner, and A. Das, Locating topological phase transitions using nonequilibrium signatures in local bulk observables, *Phys. Rev. B* **95**, 041105(R) (2017).
- [12] C. Karrasch, & D. Schuricht, Dynamical phase transitions after quenches in nonintegrable models, *Phys. Rev. B*, **87**, 195104 (2013).
- [13] N. Kriel, C. Karrasch, & S. Kehrein, Dynamical quantum phase transitions in the axial next-nearest-neighbor Ising chain, *Phys. Rev. B* **90**, 125106 (2014).
- [14] F. Andraschko, & J. Sirker, Dynamical quantum phase transitions and the Loschmidt echo: A transfer matrix approach, *Phys. Rev. B* **89**, 125120 (2014).
- [15] E. Canovi, P. Werner, & M. Eckstein, First-Order Dynamical Phase Transitions, *Phys. Rev. Lett.* **113**, 265702 (2014).
- [16] M. Heyl, Dynamical Quantum Phase Transitions in Systems with Broken-Symmetry Phases, *Phys. Rev. Lett.*, **113**, 205701 (2014).
- [17] S. Vajna, & B. Dora, Disentangling dynamical phase transitions from equilibrium phase transitions, *Phys. Rev. B* **89**, 161105(R) (2014).
- [18] S. Vajna, & B. Dora, Topological classification of dynamical phase transitions, *Phys. Rev. B* **91**, 155127 (2015).
- [19] M. Heyl, Scaling and Universality at Dynamical Quantum Phase Transitions, *Phys. Rev. Lett.*, **115**, 140602 (2015) .
- [20] J. C. Budich, & M. Heyl, Dynamical topological order parameters far from equilibrium, *Phys. Rev. B* **93**, 085416 (2016).
- [21] T. Palmai, Edge exponents in work statistics out of equilibrium and dynamical phase transitions from scattering theory in one-dimensional gapped systems, *Phys. Rev. B* **92**, 235433 (2015).
- [22] M. Schmitt, & S. Kehrein, Dynamical Quantum Phase Transitions in the Kitaev Honeycomb Model, *Phys. Rev. B* **92**, 075114 (2015).
- [23] U. Divakaran, S. Sharma, & A. Dutta, Tuning the presence of dynamical phase transitions in a generalized XY spin chain, *Phys. Rev. E* **93**, 052133 (2016).
- [24] Z. Huang, & A. V. Balatsky, Dynamical Quantum Phase Transitions: Role of Topological Nodes in Wave Function Overlaps, *Phys. Rev. Lett.* **117**, 086802 (2016).
- [25] T. Puskarov, & , D. Schuricht, Time evolution during and after finite-time quantum quenches in the transverse-field Ising chain, *SciPost Phys.* **1**, 003 (2016).
- [26] J. M. Zhang, & H.-T. Yang, Sudden jumps and plateaus in the quench dynamics of a Bloch state, *EPL*. **116**, 10008 (2016).
- [27] M. Heyl, Quenching a quantum critical state by the order parameter: Dynamical quantum phase transitions and quantum speed limits *Phys. Rev. B* **95**, 060504 (2017).
- [28] B. Zunkovic, A. Silva, & M. Fabrizio, Dynamical phase transitions and Loschmidt echo in the infinite-range XY model, *Philos Trans A Math Phys Eng Sci.* **374**(2069), (2016).
- [29] T. Obuchi, S. Suzuki, & K. Takahashi, Complex semiclassical analysis of the Loschmidt amplitude and dynamical quantum phase transitions, *Phys. Rev. B* **95**, 174305 (2017).
- [30] U. Bhattacharya. & A. Dutta, Emergent topology and dynamical quantum phase transitions in two-dimensional closed quantum systems, *Phys. Rev. B* **96**, 014302 (2017).
- [31] U. Bhattacharya . & A. Dutta, Interconnections between equilibrium topology and dynamical quantum phase transitions in a linearly ramped Haldane model, *Phys. Rev. B* **95**, 184307 (2017).
- [32] T. Fogarty, A. Usui, T. Busch, A. Silva, & J. Gool, Dynamical phase transitions, temporal orthogonality and the dynamics of observables in one dimensional ultra-cold quantum gases: from the continuum to the lattice, *New J. Phys.* **19** 113018 (2017).
- [33] J. C. Halimeh, & V. Zauner-Stauber, Dynamical phase diagram of quantum spin chains with long-range interactions, *Phys. Rev. B* **96**, 134427 (2017).
- [34] I. Homrighausen, N. O. Abeling, V. Zauner-Stauber, & J. C. Halimeh, Anomalous dynamical phase in quantum spin chains with long-range interactions, *Phys. Rev. B* **96**, 104436 (2017).
- [35] A. Dutta, & A. Dutta, Probing the role of long-range interactions in the dynamics of a long-range Kitaev chain, *Phys. Rev. B* **96**, 125113 (2017).
- [36] B. Mera., C. Vlachou, N. Paunkovic, V. R. Vieira, V & O. Viyuela, Dynamical phase transitions at finite temperature from fidelity and interferometric Loschmidt echo induced metrics, *Phys. Rev. B* **97**, 094110 (2018).
- [37] N. Sedlmay, P. Jger, M. Maiti, & J. Sirker, A bulk-

- boundary correspondence for dynamical phase transitions in one-dimensional topological insulators and superconductors, *Phys. Rev. B* **97**, 064304 (2018)
- [38] D. Trapin, & M. Heyl, Constructing effective free energies for dynamical quantum phase transitions in the transverse-field Ising chain, *Phys. Rev. B* **97**, 174303 (2018).
- [39] S. Bhattacharjee, & A. Dutta, Dynamical Quantum Phase Transitions in Extended Transverse Ising Models, *Phys. Rev. B* **97**, 134306 (2018).
- [40] D. M. Kennes, D. Schuricht, & C. Karrasch, Controlling Dynamical Quantum Phase Transitions, *Phys. Rev. B* **97**, 184302 (2018).
- [41] L. Piroli, B. Pozsgay, & E. Vernier, , Non-analytic behavior of the Loschmidt echo in XXZ spin chains: exact results, *Nucl. Phys. B* **933**, 454 (2018).
- [42] M. Heyl, F. Pollmann, and B. Dóra, Detecting Equilibrium and Dynamical Quantum Phase Transitions in Ising Chains via Out-of-Time-Ordered Correlators, *Phys. Rev. Lett.* **121**, 016801 (2018).
- [43] S. D. Nicola, A. A. Michailidis, M. Serbyn, *Entanglement view of dynamical quantum phase transitions*, arXiv:2008.04894 (2020).
- [44] S. Zamani, R. Jafari, and A. Langari, "Floquet dynamical quantum phase transition in the extended XY model: Nonadiabatic to adiabatic topological transition", *Phys. Rev. B* **102**, 144306 (2020).
- [45] J.C. Halimeh, N. Yegovtsev, V. Gurarie, Dynamical quantum phase transitions in many-body localized systems, arXiv:1903.03109 (2019).
- [46] F. Pollmann, S. Mukerjee, ., A. G. Green, & J. E. Moore, Dynamics after a sweep through a quantum critical point, *Phys. Rev. E* **81**, 020101(R) (2010).
- [47] S. Sharma, S. Suzuki, & A. Dutta, Quenches and dynamical phase transitions in a nonintegrable quantum Ising model, *Phys. Rev. B* **92**, 104306 (2015).
- [48] S. Sharma, U. Divakaran, A. Polkovnikov, & A. Dutta, Slow quenches in a quantum Ising chain; dynamical phase transitions and topology, *Phys. Rev. B* **93**, 144306 (2016).
- [49] S. Porta, F. Cavaliere, M. Sasseti & N. Traverso Ziani, "Topological classification of dynamical quantum phase transitions in the xy chain", *Scientific Reports* **10**, 12766 (2020).
- [50] U. Bhattacharya, S. Bandyopadhyay, & A. Dutta, Mixed state dynamical quantum phase transitions, *Phys. Rev. B* **96**, 180303(R) (2017).
- [51] M. Heyl, & J. C. Budich, Dynamical topological quantum phase transitions for mixed states, *Phys. Rev. B* **96**, 180304(R) (2017) .
- [52] Nils O. Abeling, & S. Kehrein, Quantum quench dynamics in the transverse field Ising model at nonzero temperatures, *Phys. Rev. B* **93**, 104302 (2016).
- [53] N. Sedlmayr, M. Fleischhauer, M., & J. Sirker, The fate of dynamical phase transitions at finite temperatures and in open systems, *Phys. Rev. B* **97**, 045147 (2018)(2018).
- [54] S. Bandyopadhyay, S. Laha, U. Bhattacharya and A. Dutta, Exploring the possibilities of dynamical quantum phase transitions in the presence of a Markovian bath, *Scientific Reports*, **8** 1 (2018).
- [55] A.A. Zvyagin, Dynamical quantum phase transitions, *Low Temp. Phys.* **42**, 971 (2016).
- [56] V. Gurarie, , Quantum Phase Transitions Go Dynamical10.1103/Physics.10.95 (2017).
- [57] M. Heyl, Dynamical quantum phase transitions: a review, *Reports on Progress in Physics*, **81**, 054001 (2018).
- [58] D. Fläschner, M. Vogel, B. Tarnowski,, S. Rem, D. S. Luhmann, M. Heyl, J. Budich, L. Mathey, K. Sengstock, & C. Weitenberg, Observation of dynamical vortices after quenches in a system with topology, *Nat. Phys.* **14**, 265–268 (2018).
- [59] P. Jurcevic, H. Shen, P. Hauke, C. Maier, T. Brydges, C. Hempel, B.P. Lanyon, M. Heyl, R. Blatt, & C.F. Roos, Direct Observation of Dynamical Quantum Phase Transitions in an Interacting Many-Body System , *Phys. Rev. Lett.* **119**, 080501 (2017).
- [60] A. Haldar, K. Mallayya, M. Heyl, F. Pollmann, M. Rigol, A. Das, Signatures of quantum phase transitions after quenches in quantum chaotic one-dimensional systems, arXiv:2004.02905 (2020).
- [61] R. Jafari, A. Akbari, "Floquet dynamical phase transition and entanglement spectrum", arXiv:2009.09484 (2020).
- [62] J. Zhang, G. Pagano, P. W. Hess, A. Kyprianidis, P. Becker, H. Kaplan, A. V. Gorshkov, Z.-X. Gong & C. Monroe, Observation of a many-body dynamical phase transition with a 53-qubit quantum simulator, *Nature* **551**, 601–604 (2017).
- [63] P. Weinberg, M. Bukov, QuSpin: a Python package for dynamics and exact diagonalisation of quantum many body systems part I: spin chains, *SciPost Phys.* **2**, 003 (2017).
- [64] P. Weinberg, M. Bukov, QuSpin: a Python package for dynamics and exact diagonalisation of quantum many body systems. Part II: bosons, fermions and higher spins, *SciPost Phys.* **7**, 020 (2019).
- [65] S. Xu and B. Swingle, "Locality, Quantum Fluctuations, and Scrambling", *Phys. Rev. X* **9**, 031048 (2019).
- [66] Ceren B. Dağ, Kai Sun, and L.-M. Duan, "Detection of Quantum Phases via Out-of-Time-Order Correlators", *Phys. Rev. Lett.* **123**, 140602 (2019).
- [67] Supplemental Material.
- [68] J.C. Halimeh, D. Trapin, M. V. Damme, M. Heyl, Local measures of dynamical quantum phase transitions, arXiv:2010.07307 (2020).
- [69] R. J. L-Swan, A. S-Naini, J. J. Bollinger, A. M. Rey, Unifying scrambling, thermalization and entanglement through measurement of fidelity out-of-time-order correlators in the Dicke model, *Nature Communications* **10**, 1581 (2019).
- [70] S. Pappalardi, A. Polkovnikov, A. Silva, Quantum echo dynamics in the Sherrington-Kirkpatrick model, *SciPost Phys.* **9**, 021 (2020).
- [71] B. V. Fine, T. A. Elsayed, C. M. Kropf, and A. S. de Wijn *Phys. Rev. E* **89**, 012923 (2014).
- [72] T. A. Elsayed and B. V. Fine, Sensitivity to small perturbations in systems of large quantum spins, *Physica Scripta* **2015** (T165), 014011 (2015).