



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Efficient Entanglement of Spin Qubits Mediated by a Hot Mechanical Oscillator

Emma Rosenfeld, Ralf Riedinger, Jan Gieseler, Martin Schuetz, and Mikhail D. Lukin

Phys. Rev. Lett. **126**, 250505 — Published 24 June 2021

DOI: [10.1103/PhysRevLett.126.250505](https://doi.org/10.1103/PhysRevLett.126.250505)

Efficient entanglement of spin qubits mediated by a hot mechanical oscillator

Emma Rosenfeld,¹ Ralf Riedinger,¹ Jan Gieseler,^{1,*} Martin Schuetz,^{2,3,†} and Mikhail D. Lukin¹

¹Physics Department, Harvard University, Cambridge, MA 02318, USA.‡

²Amazon Quantum Solutions Lab, Seattle, Washington 98170, USA.

³AWS Center for Quantum Computing, Pasadena, California 91125, USA.

(Dated: May 5, 2021)

Localized electronic and nuclear spin qubits in the solid state constitute a promising platform for storage and manipulation of quantum information, even at room temperature. However, the development of scalable systems requires the ability to entangle distant spins, which remains a challenge today. We propose and analyze an efficient, heralded scheme that employs a parity measurement in a decoherence free subspace to enable fast and robust entanglement generation between distant spin qubits mediated by a hot mechanical oscillator. We find that high-fidelity entanglement at cryogenic and even ambient temperatures is feasible with realistic parameters, and show that the entangled pair can be subsequently leveraged for deterministic controlled-NOT operations between nuclear spins. Our results open the door for novel quantum processing architectures for a wide variety of solid-state spin qubits.

Introduction. Electronic and nuclear spin qubits in the solid state are encouraging candidates for the realization of quantum information systems. Over the past decade, long-lived quantum memories and few-qubit registers have been demonstrated in several different platforms, including under ambient conditions. The key, outstanding challenge is engineering fast, programmable interactions between spin qubits separated by micrometer-scale distances. For example, color centers such as the nitrogen vacancy (NV) center in diamond are promising contenders as robust qubits, owing to their long coherence times at room temperature [1], well-developed microwave control, and optical initialization and readout. However, generating entanglement on-demand between spins remains a challenge: the short-range dipole-dipole interaction limits connectivity [2], while optical entanglement schemes are inefficient [3–5], require cryogenic temperatures, and decohere nuclear memories [6, 7].

In a complementary approach, it was suggested [8] to transduce interactions via magnetically functionalized oscillators, leveraging recent advances in the control of micro-mechanical resonators [9, 10], which enables quantum control of solid-state electron spins [11–15]. Robustness against thermal noise is desirable for such applications at elevated temperatures, to avoid phonon-induced gate errors [16]. Previous approaches for such ‘hot’ gates require large qubit-resonator cooperativities $C \gg 1$ for low error rates [17], with error scaling as $\mathcal{E} \propto 1/\sqrt{C}$ ($C \equiv \lambda^2/\Gamma\kappa n_{th}$ compares the coherent coupling rate λ to the dissipation rates of the spin and resonator, Γ and κn_{th} , respectively). These regimes are experimentally challenging to achieve, such that a demonstration of mechanically mediated entanglement remains elusive.

In this Letter, we propose and analyze a fast and robust




FIG. 1. Entanglement protocol. **(a)** Two qubits are coupled with equal strength λ to a high Q resonator which is independently measured and has thermal occupation $n_{th} \gg 1$. **(b)** Spin-dependent resonator potential. Spin states $|01\rangle$, $|10\rangle$ are decoupled, whereas $|00\rangle$, $|11\rangle$ shift the potential, such that toggling between them every half period drives the resonator. **(c)** The spins are initialized in $|++\rangle$, and the resonator states before (M_1) and after (M_2) applying the spin pulse sequence (black) for duration t_I are compared. Absence of displacement indicates spins are in the entangled anti-parallel states. **(d)** Proposed implementation. A diamond with NV centers is placed near a microresonator (grey) functionalized with nanomagnets (black), which is measured interferometrically. The entangled NV spin state is used to teleport gates between proximal ^{13}C or ^{15}N nuclear spins.

entanglement protocol for two spins (with eigenstates $|0\rangle$ and $|1\rangle$), linearly coupled to a common mode of a high-temperature mechanical resonator (Fig 1(a)), via parity measurements in a decoherence-free subspace (DFS) [18–21]. The Bell states $|\Psi^\pm\rangle \propto |01\rangle \pm |10\rangle$ decouple from the resonator, forming a DFS insensitive to its thermal fluctuations. In contrast, the aligned states $|00\rangle$ and $|11\rangle$ exert a force on the oscillator, resulting in a large, ob-

* Current address: IAV GmbH DigitalLab

† Previous address: Physics Department, Harvard University. This work was done prior to joining AWS.

‡ rosenfeld@g.harvard.edu

servable displacement (Fig 1(b)). We can thus herald the entangled $|\Psi^\pm\rangle$ state by monitoring the absence of an excess force on the mechanical oscillator, constituting a (half) parity measurement in a measurement-free subspace [19, 20, 22] and in analogy to previous proposals considering atoms in optical cavities [18]. By design, this protocol is robust to thermal noise, and neither requires strong coupling, nor cooling to the mechanical ground state. We show that entanglement can be generated at high success rates with relaxed cooperativity requirements, $C \gtrsim 1$, and with error scaling approaching $\mathcal{E} \propto \ln(C)/C$ at large cooperativities. We specifically analyze an experimental realization involving NV centers in diamond, coupled to magnetically functionalized mechanical nano-beam resonators [12, 13, 15, 23], but note that it can equally be applied to other qubit species coupled to bosonic modes at high temperatures [24–34], even when high-fidelity readout is not available by other means [11, 35–37].

The entangled pair of electronic spins can be subsequently leveraged to herald two-qubit gates between nearby, coherently-coupled nuclear spin memories in the solid state (Fig 1(d)). Assuming state-of-the-art quality (Q) factors, spin-mechanics coupling strength, and spin coherence times [9, 38–41], we expect that our entangling gate can achieve error rates below 1% at cryogenic temperatures. With modest improvements in the coupling strength, similarly low error rates can be achieved at room temperature.

Entanglement protocol. The key idea of our approach can be understood by considering two spin qubits, characterized by the Pauli operators $\sigma_{x,y,z}^{(i)}$ ($i = 1, 2$), that are linearly coupled with equal strength λ to a micromechanical oscillator (Fig 1(a), see [42], which contains refs. [43–53], for inhomogenous λ). If the qubit frequencies $\omega_s^{(i)}$ ($i = 1, 2$) strongly exceed that of the resonator $\omega_r \ll \omega_s^{(i)}$, the transverse coupling terms can be ignored, and the system Hamiltonian is:

$$\mathcal{H}/\hbar = \frac{\omega_s^{(1)}}{2}\sigma_z^{(1)} + \frac{\omega_s^{(2)}}{2}\sigma_z^{(2)} + \omega_r a^\dagger a + \lambda S_z (a + a^\dagger), \quad (1)$$

where $S_z = \sigma_z^{(1)} + \sigma_z^{(2)}$ and a (a^\dagger) are the bosonic annihilation (creation) operators of the resonator mode. For the two states $|01\rangle$ and $|10\rangle$, the qubits are decoupled from the resonator: the $S_z = 0$ states comprise a DFS, i.e. their phase is independent of the mechanical state. The other two states exert a force $\sim \pm 2\hbar\lambda/z_p$ on the resonator, where z_p is the mechanical zero point fluctuation.

In our entanglement protocol (Fig 1(c)), (i), the state of the resonator is first measured, while the two spins are initialized in the separable state $|+\rangle \otimes |+\rangle \propto \sqrt{2}|\Psi^+\rangle + |11\rangle + |00\rangle$, with $|+\rangle = (|1\rangle + |0\rangle)/\sqrt{2}$. Then, (ii) the spins interact with the resonator for a time t_I , while being subjected to a special resonant decoupling sequence, such that the spin states $|11\rangle$ and $|00\rangle$ displace the resonator state. Finally (iii), the resonator displacement is measured. If it is below a threshold, the spins are

projected into the Bell state $|\Psi^+\rangle$, indicating successful entanglement generation. The protocol can be made deterministic by repeating steps (i-iii) until success (~ 2 -3 repetitions in the regime of interest).

To estimate its practical performance, we assume the mechanical system can be described by a master equation, is weakly coupled ($Q \gg 1$) to a hot thermal bath at rate $\kappa = \omega_r/Q$ and temperature $T \gg \hbar\omega_r/k_B$, and each qubit is dispersively coupled to an independent reservoir with dephasing rate Γ . The Gaussian state of the oscillator can be estimated interferometrically and independently of the spins. In practice, the effects of the measurement backaction and duration are negligible in near term realizations (see SI). To simplify the derivation, in the following we assume short backaction evading measurements of the mechanical quadrature [54], neglecting the measurement duration and avoiding a lower limit on the measurement uncertainty.

Figure 2 illustrates the key ingredients of the scheme. In step (i) we perform an initial linear measurement M_1 on the momentum quadrature $p \equiv i(a^\dagger - a)/\sqrt{2}$, with measurement uncertainty Δm (Fig 2(a)). In step (ii), the spins are resonantly coupled to the oscillator by a series of π pulses (here assumed to be ideal), with a pulse separation $2\tau = \pi/\omega_r$. This simultaneously maximizes the conditional mechanical displacement and the spin coherence by dynamically decoupling from their bath (Fig 1(b) and 1(c)). Throughout this pulse sequence, the force acting on the resonator is a square wave with amplitude $\sim -\hbar\lambda S_z/z_p$ (Fig 1(c)) and frequency ω_r . In the high Q limit, higher harmonics of the force can be neglected, resulting in the effective interaction Hamiltonian in the rotating and toggling frame, under a rotating wave approximation [42]

$$\mathcal{H}_{int}/\hbar = \frac{2\lambda}{\pi} S_z (a^\dagger + a), \quad (2)$$

leading to a momentum shift of the resonator $\mu(S_z, t_I) = -4\sqrt{2}\lambda S_z(1 - e^{-\kappa t_I/2})/\pi \kappa \approx -2\sqrt{2}\lambda S_z t_I/\pi$ in natural units, for $\kappa t_I \ll 1$. As the conditional equations of motion are linear, the motional states after step (ii) remain Gaussian with an uncertainty $\Delta d(t_I) \approx \sqrt{\kappa n_{th} t_I + \Delta m^2}$ (dashed circles in Fig. 2(b)) and a spin-dependent expectation value of the momentum quadrature of $M_1 e^{-\kappa t_I/2} + \mu(S_z, t_I)$. Then, (iii) a second measurement M_2 localizes the resonator with uncertainty Δm , projecting the spin population to $\langle S_z \rangle \in \{0, +2, -2\}$ if the distributions are separable. If this is achieved within the coherence time $1/\Gamma$ of the spins, an $\langle S_z \rangle = 0$ measurement projects the two spins into the entangled state $|\Psi^+\rangle$.

A simple estimate shows the minimum requirements for this protocol. For negligible measurement uncertainties $\Delta m^2 \ll \kappa n_{th} t_I$, and an interaction time comparable to the spin coherence time $t_I \sim 1/\Gamma$, yet shorter than the mechanical lifetime, $t_I \ll 1/\kappa$, the distributions become separable if the displacement $|\mu(2, 1/\Gamma)|/2$ exceeds the uncertainty $\Delta d(1/\Gamma)$, i.e. $\lambda^2/\Gamma \kappa n_{th} = C \gtrsim 1$.

To obtain an estimate of the fidelity, we compute the




FIG. 2. Mechanical phase space picture. (a) The thermal resonator (ρ_{th}) is localized by measurement M_1 (uncertainty Δm , purple) at time $t = 0$ [55–58]. (b) After interaction time t_I , the spin-conditional resonator states (dashed circles, $S_z = 2$ pink, $S_z = 0$ blue, $S_z = -2$ green) separate, and the resonator is measured again (M_2 , purple). (c) The conditional distribution of measured displacement (dashed circles) and (d) its projection onto the momentum basis used to define a threshold $\alpha|\mu|/2$ (black dashed line for $\alpha \approx 0.6$). If a specific measurement (purple dot) lies within this threshold, $\langle S_z \rangle = 0$ is assigned and entanglement is heralded.

(Gaussian) probability density function $\mathcal{P}_{\langle S_z \rangle}(\Delta M)$ of the momentum difference $\Delta M \equiv M_2 - e^{-\kappa t_I/2} M_1$, conditional on the spin state S_z , which has expectation value $\mu(S_z, t_I)$ and variance

$$\sigma(t_I)^2 = \Delta m^2(1 + e^{-\kappa t_I}) + n_{th}(1 - e^{-\kappa t_I}), \quad (3)$$

corresponding to the contributions by the measurement uncertainties and diffusion during the interaction. The state is assumed to have $S_z = 0$ if

$$|\Delta M| < \alpha\mu(2, t_I)/2, \quad (4)$$

where the threshold $\alpha \in (0, 1]$ can be tuned to trade between a high acceptance rate ($\alpha \rightarrow 1$) and low false positive acceptance ($\alpha \rightarrow 0$). The probability of an accepted event being a true positive is $S(\alpha, g(t_I)) \approx 1/(1 + e^{-2g(t_I)^2}) - \mathcal{O}(\alpha^2)$, given by the integrals of $\mathcal{P}_{0,\pm 1}$ within the thresholds, and weighted by the initial spin populations [42], where we define the normalized displacement $g(t_I) \equiv \mu(2, t_I)/2\sigma(t_I)$. The error in entanglement fidelity $\mathcal{F} = \langle \Psi^\pm | \rho | \Psi^\pm \rangle$ is the result of two independent error sources, namely spin dephasing and false positive $\langle S_z \rangle = 0$ assignments, yielding

$$\mathcal{F} = \frac{1 + e^{-2\Gamma t_I}}{2} S(\alpha, g(t_I)) \approx \frac{1}{2} \frac{1 + e^{-2\Gamma t_I}}{1 + e^{-2g(t_I)^2}} - \mathcal{O}(\alpha^2). \quad (5)$$

It follows directly that entanglement can be generated ($\mathcal{F} > 1/2$) for $g(t_I)^2 > \Gamma t_I$. We note that this simple




FIG. 3. Performance. (a) Fidelity \mathcal{F} as a function of Γt_I under (6). Thresholds α are color-coded and cooperativities C have the associated linestyles. Analytic form in eq. (5), corresponding to $\alpha \ll 1$, blue curves. (b) Optimal pulse sequence duration (solid lines), and its analytic approximation eq. (7) (dashed line). (c) Infidelity \mathcal{E} of the Bell state preparation (solid lines) and analytic approximation (5) for $\alpha \ll 1$ (dashed line). The black dotted line represents the optimal infidelity of a deterministic hot gate [17], and the dashed-dotted black line is the asymptotic $\pi^2 \ln C / 16C$ scaling. (d) Normalized rate of true positive entanglement heralding events r_p / Γ . The fast repetition rate allows multiple protocol attempts within the spin coherence time for large C .

estimate for $\alpha \rightarrow 0$ is a good approximation for general \mathcal{F} [42].

Analysis of the Bell state preparation. In the following, we consider the experimentally relevant regime $\Delta m^2 \ll \kappa n_{th} t_I$, i.e. the linearized diffusion term dominates the variance of ΔM , and that $\kappa t_I \ll 1$. In these limits

$$g(t_I)^2 \approx (8/\pi^2) C \Gamma t_I, \quad (6)$$

such that \mathcal{F} can be described only in terms of α , C and Γt_I (Fig 3a). The fidelity exceeds 1/2 for $C > \pi^2/8 \sim 1.2$, and exceeds 96% for $C \sim 100$, demonstrating that our protocol can be applied with relatively low C . The optimal interaction time Γt^* is determined numerically for each C (Fig 3(b), dashed lines) and can be analytically approximated as

$$\Gamma t^* \approx \frac{\pi^2}{16C} \ln(16C/\pi^2 - 1) \quad (7)$$

for $C > \pi^2/8$ and Δm , $\alpha \rightarrow 0$ [42], showing that the entanglement protocol is fast compared to the spin coherence time. We note that in the regime of interest ($C \gtrsim 8$, $t_I \ll 1/\Gamma$) decreasing the threshold α e.g. from 1 to 0.5 reduces the optimal interaction time (Fig. 3(b)). At high C , this can compensate the reduced acceptance rate for small α and increase the absolute rate

of true positive entanglement heralding events, given by $r_p = \int_{-\theta}^{+\theta} \mathcal{P}_0(p) dp / 2t^*$ for threshold $\theta = \alpha\mu(+2, t^*)/2$ (Fig. 3d) [42]. Inserting equation (7) into (5), we find a lower bound to the fidelity

$$\mathcal{F} \geq \frac{1}{2} \frac{1 + (16C/\pi^2 - 1)^{-\frac{\pi^2}{8C}}}{1 + (16C/\pi^2 - 1)^{-1}} \quad (8)$$

again for $C > \pi^2/8$ and $\Delta m, \alpha \rightarrow 0$. The error $\mathcal{E} = 1 - \mathcal{F}$ is shown in Fig 3(c). Remarkably, the cooperativity required to achieve an error $\mathcal{E} < 10^{-3}$ is more than two orders of magnitude lower than for previous mechanically mediated gates [8, 17]. For large C , $\mathcal{E} \sim (\pi^2/16) \ln(C)/C$.

Potential applications. The entanglement protocol presented here is inherently probabilistic, approaching a heralding probability of 1/2 for $\alpha \rightarrow 1$. However, it can be extended to yield deterministic controlled-NOT gates, as required for quantum information processing, between associated qubit registers by feedback, assuming a simple repeat-until-success scheme [4, 18]. In the following, we consider two electronic spins, such as NV centers (using $|m_s\rangle = |\pm 1\rangle$ as qubit states, for maximal displacements), interacting with the mechanical resonator and coupled to nearby ^{13}C (or ^{15}N) nuclear spins in the diamond host (Fig. 1d). The entangled NV spin state is used to teleport a gate between the nuclear spins [59]. Contributions to the gate error \mathcal{E}_T include infidelities related to the ideal entanglement protocol \mathcal{E} , control (\mathcal{E}_C), initialization (\mathcal{E}_{init}) and readout (\mathcal{E}_{RO}) of the NV spins, and the electron-nuclear CNOT gate (\mathcal{E}_{CNOT}). Nuclear qubit errors arise from coupling to a bath (\mathcal{E}_{nuc}) at rate $\Gamma_{\gamma_N}/\gamma_e$, where γ_N (γ_e) is the nuclear (electron) gyromagnetic ratio, as well as dephasing due to electron spin control errors in failed entanglement attempts. As the latter depends on the heralding probability and the hyperfine coupling, we attribute it to \mathcal{E}_C with a factor η , which is below 1 in the regime of interest [42]. Combining state-of-the-art spin control [60, 61] with robust decoupling sequences [62], \mathcal{E}_C can be neglected, however, if left unaddressed without optimal spin control, \mathcal{E}_C can limit the fidelity [7]. As the repetition rate is high (Fig. 3(b, d)), we further neglect the small probability of failure after a large number of repetitions in a synchronous circuit [4]. In this case, the total error of the deterministic nuclear gate is $\mathcal{E}_T = \mathcal{E} + 2((1+\eta)\mathcal{E}_C + \mathcal{E}_{init} + \mathcal{E}_{RO} + \mathcal{E}_{CNOT} + \mathcal{E}_{nuc})$.

Experimental Implementation. In NV-based operations on nearby nuclear spins, optical excitation of the NV induces decoherence on the nuclear spins through the hyperfine coupling [6, 7, 65, 66], which can limit \mathcal{E}_T due to repeated spin initialization and readout, unless compensated by a DFS of nuclear spins [67]. In our system, the mechanical oscillator can also be used for single-shot readout and initialization [11], eliminating the need for optical illumination of the color center [68].

While state-of-the-art continuous interferometric measurement schemes are sufficient for initial experiments




FIG. 4. Error \mathcal{E}_T for teleported CNOT gate between two ^{13}C (or ^{15}N) nuclear spins, as a function of C , for thresholds α and under (6). A mechanical NV spin readout using an interaction time of $20t^*$ is assumed. The CNOT gate error between the NV and nuclear spin spin is set to 10^{-4} (dashed grey line) [63, 64] and is also included in the representation of state-of-the-art hot gates [17]. Gate errors are numerically calculated (solid lines) and analytically approximated as \mathcal{E} (eq. (8)), neglecting nuclear spin decoherence, dashed line). For details see text.

[38, 40], their performance could be optimized using a Kalman filter (see e.g. Fig. 2) [55–58, 69], to account for mechanical spectator modes and other technical noise sources [42]. For further improvement, the estimation of the spin-induced displacement can be achieved with a multiple model adaptive estimation [70, 71], while performing steps (ii) and (iii) simultaneously, as well as feedback on the spin state with additional global spin rotations during the pulse sequence, further increasing the entanglement fidelity and rate [21]. Finally, errors arising from small inhomogeneities in the coupling strength can be suppressed with additional electron spin control [42].

Fig. 4 shows the controlled-NOT gate error \mathcal{E}_T as a function of C . For realistic parameters described in [42], at very high C , the total error \mathcal{E}_T is limited by the electron-nuclear two qubit gate fidelity, while at more modest cooperativity, the error scales favorably compared to the existing state-of-the-art [17]. Note that an experimental demonstration of our protocol (with $\mathcal{E}_T \sim 10^{-1}$) may be possible at room temperature and $C \sim 8$, corresponding to state-of-the-art spin-mechanical systems ($1/\Gamma \sim 10\text{ms}$ [72]; $Q \sim 10^9$ [9, 73]; $\lambda/2\pi \sim 0.9\text{ kHz}$ [41]; $\Delta m^2 \sim 27$ [40]). These parameters are within reach for a soft-clamped, silicon nitride nano-beam resonator, functionalized with a nano-magnet at the antinode of motion, and placed adjacent to diamond hosting NV centers (Fig. 1(d)) [72, 73]. A modest further improvement in the spin-mechanical coupling strength ($\lambda \sim 2\text{ kHz}$) enables high fidelity gates ($\mathcal{E}_T < 10^{-2}$). Conversely, at cryogenic temperatures ($T \sim 4\text{ K}$, coherence time $1/\Gamma \sim 1\text{ s}$ [72, 74]) the same system can achieve this fidelity already with $\lambda/2\pi \sim 100\text{ Hz}$ [42]. Such parameters yield high probability of success (approaching 50% per run) and average gate duration approaching 10ms, faster or comparable to deterministic

protocols with the reported coupling strengths [17].

Conclusion. We proposed and analyzed a half-parity measurement protocol in a decoherence free subspace for entangling two qubits through a hot resonator, with error scaling that nears $\mathcal{E} \propto \ln(C)/C$. Our protocol is fast, robust to thermalization errors, and does not require ground state cooling. A teleported controlled-NOT gate employing the generated Bell pair is feasible with magnetically functionalized resonators and solid state electronic spins featuring long coherence times [72, 73]. While we analyzed an implementation involving NV centers, as readout and initialization can be realized mechanically [11], the protocol can also be applied to other promising paramagnetic defects, such as spins in silicon [75]. Further directions for analysis include leveraging continuous feedback to increase the entanglement rate [21, 22, 76], and application of our protocol to generate multi-partite entangled states. Lastly, for further improvements, nanobeam resonators can be electrostatically coupled [8], using hybridized mechanical modes to selectively couple spins adjacent to distinct resonators, enabling multi-qubit connectivity far beyond reach of

the magnetic dipole-dipole interactions. While our work leverages decades of development of micromechanical devices and solid state qubits, it simultaneously eliminates the need for high-fidelity single qubit optical or electronic addressing. With substantial technical improvement beyond the current state-of-the-art, in the long term, this approach could pave the way for realization of solid-state, room-temperature quantum information systems.

Acknowledgements We would like to thank S. Hofer for insightful discussions and Samantha Dakoulas for assistance during this project. The authors thank C. Maxwell for design assistance with Figs. 1 and 2. This work was supported by NSF, CUA, ARO MURI and V. Bush Faculty Fellowship. R.R. was supported by the Alexander von Humboldt Foundation. This material is based upon work supported by the National Science Foundation Graduate Research Fellowship under Grants No. DGE1144152 and No. DGE1745303. Any opinion, findings, and conclusions or recommendations expressed in this material are those of the authors(s) and do not necessarily reflect the views of the National Science Foundation.

-
- [1] E. D. Herbschleb, H. Kato, Y. Maruyama, T. Danjo, T. Makino, S. Yamasaki, I. Ohki, K. Hayashi, H. Morishita, M. Fujiwara, and N. Mizuochi. Ultra-long coherence times amongst room-temperature solid-state spins. *Nature Communications*, 10(1):1–6, August 2019.
- [2] F. Dolde, I. Jakobi, B. Naydenov, N. Zhao, S. Pezzagna, C. Trautmann, J. Meijer, P. Neumann, F. Jelezko, and J. Wrachtrup. Room-temperature entanglement between single defect spins in diamond. *Nature Physics*, 9(3):139–143, March 2013.
- [3] H. Bernien, B. Hensen, W. Pfaff, G. Koolstra, M. S. Blok, L. Robledo, T. H. Taminiau, M. Markham, D. J. Twitchen, L. Childress, and R. Hanson. Heralded entanglement between solid-state qubits separated by three metres. *Nature*, 497(7447):86–90, May 2013.
- [4] Peter C. Humphreys, Norbert Kalb, Jaco P. J. Morits, Raymond N. Schouten, Raymond F. L. Vermeulen, Daniel J. Twitchen, Matthew Markham, and Ronald Hanson. Deterministic delivery of remote entanglement on a quantum network. *Nature*, 558(7709):268–273, June 2018.
- [5] J. Borregaard, P. Kómár, E. M. Kessler, A. S. Sørensen, and M. D. Lukin. Heralded Quantum Gates with Integrated Error Detection in Optical Cavities. *Physical Review Letters*, 114(11):110502, March 2015.
- [6] L. Jiang, M. V. Gurudev Dutt, E. Togan, L. Childress, P. Cappellaro, J. M. Taylor, and M. D. Lukin. Coherence of an Optically Illuminated Single Nuclear Spin Qubit. *Physical Review Letters*, 100(7):073001, February 2008.
- [7] N. Kalb, P. C. Humphreys, J. J. Slim, and R. Hanson. Dephasing mechanisms of diamond-based nuclear-spin memories for quantum networks. *Physical Review A*, 97(6):062330, June 2018.
- [8] P. Rabl, S. J. Kolkowitz, F. H. L. Koppens, J. G. E. Harris, P. Zoller, and M. D. Lukin. A quantum spin transducer based on nanoelectromechanical resonator arrays. *Nature Physics*, 6(8):602–608, August 2010.
- [9] Y. Tsaturyan, A. Barg, E. S. Polzik, and A. Schliesser. Ultracoherent nanomechanical resonators via soft clamping and dissipation dilution. *Nature Nanotechnology*, 12(8):776–783, August 2017.
- [10] Gregory S. MacCabe, Hengjiang Ren, Jie Luo, Justin D. Cohen, Hengyun Zhou, Alp Sipahigil, Mohammad Mirhosseini, and Oskar Painter. Nano-acoustic resonator with ultralong phonon lifetime. *Science*, 370(6518):840–843, 2020.
- [11] D. Rugar, R. Budakian, H. J. Mamin, and B. W. Chui. Single spin detection by magnetic resonance force microscopy. *Nature*, 430(6997):329–332, July 2004.
- [12] Sungkun Hong, Michael S. Grinolds, Patrick Maletinsky, Ronald L. Walsworth, Mikhail D. Lukin, and Amir Yacoby. Coherent, Mechanical Control of a Single Electronic Spin. *Nano Letters*, 12(8):3920–3924, August 2012.
- [13] T. Delord, P. Huillery, L. Nicolas, and G. Hétet. Spin-cooling of the motion of a trapped diamond. *Nature*, 580(7801):56–59, April 2020.
- [14] Donghun Lee, Kenneth W. Lee, Jeffrey V. Cady, Preeti Ovarthaiyapong, and Ania C. Bleszynski Jayich. Topical review: spins and mechanics in diamond. *Journal of Optics*, 19(3):033001, February 2017.
- [15] D. D. Bhaktavatsala Rao, S. Ali Momenzadeh, and Jörg Wrachtrup. Heralded Control of Mechanical Motion by Single Spins. *Physical Review Letters*, 117(7):077203, August 2016.
- [16] Anders Sørensen and Klaus Mølmer. Quantum Computation with Ions in Thermal Motion. *Physical Review Letters*, 82(9):1971–1974, March 1999.

- [17] M. J. A. Schuetz, G. Giedke, L. M. K. Vandersypen, and J. I. Cirac. High-fidelity hot gates for generic spin-resonator systems. *Physical Review A*, 95(5):052335, May 2017.
- [18] Anders S. Sørensen and Klaus Mølmer. Measurement Induced Entanglement and Quantum Computation with Atoms in Optical Cavities. *Physical Review Letters*, 91(9):097905, August 2003.
- [19] Charles Hill and Jason Ralph. Weak measurement and control of entanglement generation. *Physical Review A*, 77(1):014305, January 2008.
- [20] N. Roch, M.E. Schwartz, F. Motzoi, C. Macklin, R. Vijay, A.W. Eddins, A.N. Korotkov, K.B. Whaley, M. Sarovar, and I. Siddiqi. Observation of Measurement-Induced Entanglement and Quantum Trajectories of Remote Superconducting Qubits. *Physical Review Letters*, 112(17):170501, April 2014.
- [21] Leigh Martin, Felix Motzoi, Hanhan Li, Mohan Sarovar, and Birgitta Whaley. Deterministic generation of remote entanglement with active quantum feedback. *Physical Review A*, 92(6):062321, December 2015.
- [22] Leigh Martin, Mahrud Sayrafi, and K Birgitta Whaley. What is the optimal way to prepare a Bell state using measurement and feedback? *Quantum Science and Technology*, 2(4):044006, December 2017.
- [23] S. Kolkowitz, A. C. Bleszynski Jayich, Q. P. Unterreithmeier, S. D. Bennett, P. Rabl, J. G. E. Harris, and M. D. Lukin. Coherent Sensing of a Mechanical Resonator with a Single-Spin Qubit. *Science*, 335(6076):1603–1606, March 2012.
- [24] David J. Christle, Abram L. Falk, Paolo Andrich, Paul V. Klimov, Jawad Ul Hassan, Nguyen T. Son, Erik Janzén, Takeshi Ohshima, and David D. Awschalom. Isolated electron spins in silicon carbide with millisecond coherence times. *Nature Materials*, 14(2):160–163, February 2015.
- [25] S. Ali Momenzadeh, Felipe Fávoro de Oliveira, Philipp Neumann, D.D. Bhaktavatsala Rao, Andrej Denisenko, Morteza Amjadi, Zhiqin Chu, Sen Yang, Neil B. Manson, Marcus W. Doherty, and Jörg Wrachtrup. Thin Circular Diamond Membrane with Embedded Nitrogen-Vacancy Centers for Hybrid Spin-Mechanical Quantum Systems. *Physical Review Applied*, 6(2):024026, August 2016.
- [26] A. Barfuss, J. Teissier, E. Neu, A. Nunnenkamp, and P. Maletinsky. Strong mechanical driving of a single electron spin. *Nature Physics*, 11(10):820–824, October 2015.
- [27] Michael S. J. Barson, Phani Peddibhotla, Preeti Ovarthaiyapong, Kumaravelu Ganesan, Richard L. Taylor, Matthew Gebert, Zoe Mielens, Berndt Koslowski, David A. Simpson, Liam P. McGuinness, Jeffrey McCallum, Steven Prawer, Shinobu Onoda, Takeshi Ohshima, Ania C. Bleszynski Jayich, Fedor Jelezko, Neil B. Manson, and Marcus W. Doherty. Nanomechanical Sensing Using Spins in Diamond. *Nano Letters*, 17(3):1496–1503, March 2017.
- [28] D. Andrew Golter, Thein Oo, Mayra Amezcua, Kevin A. Stewart, and Hailin Wang. Optomechanical Quantum Control of a Nitrogen-Vacancy Center in Diamond. *Physical Review Letters*, 116(14):143602, April 2016.
- [29] Behzad Khanaliloo, Harishankar Jayakumar, Aaron C. Hryciw, David P. Lake, Hamidreza Kaviani, and Paul E. Barclay. Single-Crystal Diamond Nanobeam Waveguide Optomechanics. *Physical Review X*, 5(4):041051, December 2015.
- [30] E. R. MacQuarrie, T. A. Gosavi, N. R. Jungwirth, S. A. Bhave, and G. D. Fuchs. Mechanical Spin Control of Nitrogen-Vacancy Centers in Diamond. *Physical Review Letters*, 111(22):227602, November 2013.
- [31] Srujan Meesala, Young-Ik Sohn, Haig A. Atikian, Samuel Kim, Michael J. Burek, Jennifer T. Choy, and Marko Lončar. Enhanced Strain Coupling of Nitrogen-Vacancy Spins to Nanoscale Diamond Cantilevers. *Physical Review Applied*, 5(3):034010, March 2016.
- [32] Preeti Ovarthaiyapong, Kenneth W. Lee, Bryan A. Myers, and Ania C. Bleszynski Jayich. Dynamic strain-mediated coupling of a single diamond spin to a mechanical resonator. *Nature Communications*, 5(1):4429, July 2014.
- [33] J. Teissier, A. Barfuss, P. Appel, E. Neu, and P. Maletinsky. Strain Coupling of a Nitrogen-Vacancy Center Spin to a Diamond Mechanical Oscillator. *Physical Review Letters*, 113(2):020503, July 2014.
- [34] J. Gieseler, A. Kabcenell, E. Rosenfeld, J.D. Schaefer, A. Safira, M. J. A. Schuetz, C. Gonzalez-Ballester, C. C. Rusconi, O. Romero-Isart, and M.D. Lukin. Single-Spin Magnetomechanics with Levitated Micromagnets. *Physical Review Letters*, 124(16):163604, April 2020.
- [35] Jonilyn G. Longenecker, H. J. Mamin, Alexander W. Senko, Lei Chen, Charles T. Rettner, Daniel Rugar, and John A. Marohn. High-Gradient Nanomagnets on Cantilevers for Sensitive Detection of Nuclear Magnetic Resonance. *ACS nano*, 6(11):9637–9645, November 2012.
- [36] John M. Nichol, Eric R. Hemesath, Lincoln J. Lauhon, and Raffi Budakian. Nanomechanical detection of nuclear magnetic resonance using a silicon nanowire oscillator. *Physical Review B*, 85(5):054414, February 2012.
- [37] R Fischer, D P McNally, C Reetz, G G T Assumpção, T Knief, Y Lin, and C A Regal. Spin detection with a micromechanical trampoline: towards magnetic resonance microscopy harnessing cavity optomechanics. *New Journal of Physics*, 21(4):043049, April 2019.
- [38] Massimiliano Rossi, David Mason, Junxin Chen, Yeghishe Tsaturyan, and Albert Schliesser. Measurement-based quantum control of mechanical motion. *Nature*, 563(7729):53–58, November 2018.
- [39] M. H. Abobeih, J. Cramer, M. A. Bakker, N. Kalb, M. Markham, D. J. Twitchen, and T. H. Taminau. One-second coherence for a single electron spin coupled to a multi-qubit nuclear-spin environment. *Nature Communications*, 9(1):2552, December 2018.
- [40] Jingkun Guo, Richard Norte, and Simon Gröblacher. Feedback Cooling of a Room Temperature Mechanical Oscillator close to its Motional Ground State. *Physical Review Letters*, 123(22):223602, November 2019.
- [41] O. Arcizet, V. Jacques, A. Siria, P. Poncharal, P. Vincent, and S. Seidelin. A single nitrogen-vacancy defect coupled to a nanomechanical oscillator. *Nature Physics*, 7(11):879–883, November 2011.
- [42] Supplementary information.
- [43] C.E. Bradley, J. Randall, M.H. Abobeih, R.C. Berrevoets, M.J. Degen, M.A. Bakker, M. Markham, D. J. Twitchen, and T.H. Taminau. A Ten-Qubit Solid-State Spin Register with Quantum Memory up to One Minute. *Physical Review X*, 9(3):031045, September 2019.
- [44] C. T. Nguyen, D. D. Sukachev, M. K. Bhaskar, B. Machielse, D. S. Levonian, E. N. Knall, P. Stroganov, C. Chia, M. J. Burek, R. Riedinger, H. Park, M. Lončar,

- and M. D. Lukin. An integrated nanophotonic quantum register based on silicon-vacancy spins in diamond. *Physical Review B*, 100(16):165428, October 2019.
- [45] G. Waldherr, Y. Wang, S. Zaiser, M. Jamali, T. Schulte-Herbrüggen, H. Abe, T. Ohshima, J. Isoya, J. F. Du, P. Neumann, and J. Wrachtrup. Quantum error correction in a solid-state hybrid spin register. *Nature*, 506(7487):204–207, February 2014.
- [46] T. van der Sar, Z. H. Wang, M. S. Blok, H. Bernien, T. H. Taminiau, D. M. Toyli, D. A. Lidar, D. D. Awschalom, R. Hanson, and V. V. Dobrovitski. Decoherence-protected quantum gates for a hybrid solid-state spin register. *Nature*, 484(7392):82–86, April 2012.
- [47] L. Jiang, J. S. Hodges, J. R. Maze, P. Maurer, J. M. Taylor, D. G. Cory, P. R. Hemmer, R. L. Walsworth, A. Yacoby, A. S. Zibrov, and M. D. Lukin. Repetitive Readout of a Single Electronic Spin via Quantum Logic with Nuclear Spin Ancillae. *Science*, 326(5950):267–272, October 2009.
- [48] Genyue Liu, Mo Chen, Yi-Xiang Liu, David Layden, and Paola Cappellaro. Repetitive readout enhanced by machine learning. *Machine Learning: Science and Technology*, 1(1):015003, February 2020.
- [49] Y. Doi, T. Makino, H. Kato, D. Takeuchi, M. Ogura, H. Okushi, H. Morishita, T. Tashima, S. Miwa, S. Yamasaki, P. Neumann, J. Wrachtrup, Y. Suzuki, and N. Mizuochi. Deterministic Electrical Charge-State Initialization of Single Nitrogen-Vacancy Center in Diamond. *Physical Review X*, 4(1):011057, March 2014.
- [50] Yuki Doi, Takahiro Fukui, Hiromitsu Kato, Toshiharu Makino, Satoshi Yamasaki, Toshiyuki Tashima, Hiroki Morishita, Shinji Miwa, Fedor Jelezko, Yoshishige Suzuki, and Norikazu Mizuochi. Pure negatively charged state of the NV center in N^0 -type diamond. *Physical Review B*, 93(8):081203, February 2016.
- [51] A. Dréau, P. Spinicelli, J. R. Maze, J.-F. Roch, and V. Jacques. Single-Shot Readout of Multiple Nuclear Spin Qubits in Diamond under Ambient Conditions. *Physical Review Letters*, 110(6):060502, February 2013.
- [52] M. Chiani, D. Dardari, and M.K. Simon. New exponential bounds and approximations for the computation of error probability in fading channels. *IEEE Transactions on Wireless Communications*, 2(4):840–845, July 2003.
- [53] David A. Hopper, Joseph D. Lauigan, Tzu-Yung Huang, and Lee C. Bassett. Real-Time Charge Initialization of Diamond Nitrogen-Vacancy Centers for Enhanced Spin Readout. *Physical Review Applied*, 13(2):024016, February 2020.
- [54] M. R. Vanner, I. Pikovski, G. D. Cole, M. S. Kim, C. Brukner, K. Hammerer, G. J. Milburn, and M. Aspelmeyer. Pulsed quantum optomechanics. *Proceedings of the National Academy of Sciences*, 108(39):16182–16187, September 2011.
- [55] Sebastian G. Hofer and Klemens Hammerer. Quantum Control of Optomechanical Systems. In *Advances In Atomic, Molecular, and Optical Physics*, volume 66, pages 263–374. Elsevier, 2017.
- [56] Vahid Hassani, A. Pedro Aguiar, António M. Pascoal, and Michael Athans. Further results on plant parameter identification using continuous-time multiple-model adaptive estimators. In *Proceedings of the 48th IEEE Conference on Decision and Control (CDC) held jointly with 2009 28th Chinese Control Conference*, pages 7261–7266, December 2009.
- [57] Robert F. Stengel. *Optimal Control and Estimation*. Dover Publications, New York, reissue edition edition, September 1994.
- [58] Jr Arthur E. Bryson and Yu-Chi Ho. *Applied Optimal Control: Optimization, Estimation and Control*. CRC Press, New York, 1 edition edition, January 1975.
- [59] Kevin S. Chou, Jacob Z. Blumoff, Christopher S. Wang, Philip C. Reinhold, Christopher J. Axline, Yvonne Y. Gao, L. Frunzio, M. H. Devoret, Liang Jiang, and R. J. Schoelkopf. Deterministic teleportation of a quantum gate between two logical qubits. *Nature*, 561(7723):368–373, September 2018.
- [60] Xing Rong, Jianpei Geng, Fazhan Shi, Ying Liu, Kebiao Xu, Wenchao Ma, Fei Kong, Zhen Jiang, Yang Wu, and Jiangfeng Du. Experimental fault-tolerant universal quantum gates with solid-state spins under ambient conditions. *Nature Communications*, 6, November 2015.
- [61] T. P. Harty, D. T. C. Allcock, C. J. Ballance, L. Guidoni, H. A. Janacek, N. M. Linke, D. N. Stacey, and D. M. Lucas. High-Fidelity Preparation, Gates, Memory, and Readout of a Trapped-Ion Quantum Bit. *Physical Review Letters*, 113(22):220501, November 2014.
- [62] Genko T. Genov, Daniel Schraft, Nikolay V. Vitanov, and Thomas Halfmann. Arbitrarily Accurate Pulse Sequences for Robust Dynamical Decoupling. *Physical Review Letters*, 118(13):133202, March 2017.
- [63] Yi Chou, Shang-Yu Huang, and Hsi-Sheng Goan. Optimal control of fast and high-fidelity quantum gates with electron and nuclear spins of a nitrogen-vacancy center in diamond. *Physical Review A*, 91(5):052315, May 2015.
- [64] Wenzheng Dong, F A Calderon-Vargas, and Sophia E Economou. Precise high-fidelity electron–nuclear spin entangling gates in NV centers via hybrid dynamical decoupling sequences. *New Journal of Physics*, 22(7):073059, jul 2020.
- [65] Andreas Reiserer, Norbert Kalb, Machiel S. Blok, Koen J.M. van Bemmelen, Tim H. Taminiau, Ronald Hanson, Daniel J. Twitchen, and Matthew Markham. Robust Quantum-Network Memory Using Decoherence-Protected Subspaces of Nuclear Spins. *Physical Review X*, 6(2):021040, June 2016.
- [66] David A. Hopper, Henry J. Shulevitz, and Lee C. Bassett. Spin Readout Techniques of the Nitrogen-Vacancy Center in Diamond. *Micromachines*, 9(9), August 2018.
- [67] Michael A. Perlin, Zhen-Yu Wang, Jorge Casanova, and Martin B. Plenio. Noise-resilient architecture of a hybrid electron-nuclear quantum register in diamond. *Quantum Sci. Technol.*, 4(1):015007, September 2018. Publisher: IOP Publishing.
- [68] G. P. Berman, D. I. Kamenev, and V. I. Tsifrinovich. Stationary cantilever vibrations in oscillating-cantilever-driven adiabatic reversals: Magnetic-resonance-force-microscopy technique. *Physical Review A*, 66(2):023405, August 2002.
- [69] Witel Wieczorek, Sebastian G. Hofer, Jason Hoelscher-Obermaier, Ralf Riedinger, Klemens Hammerer, and Markus Aspelmeyer. Optimal State Estimation for Cavity Optomechanical Systems. *Physical Review Letters*, 114(22):223601, June 2015.
- [70] A. Pedro Aguiar, Michael Athans, and António M. Pascoal. Convergence properties of a Continuous-Time Multiple-Model Adaptive Estimator. In *2007 European Control Conference (ECC)*, pages 1530–1536, July 2007.

- [71] P.D. Hanlon and P.S. Maybeck. Multiple-model adaptive estimation using a residual correlation Kalman filter bank. *IEEE Transactions on Aerospace and Electronic Systems*, 36(2):393–406, April 2000.
- [72] N. Bar-Gill, L.M. Pham, A. Jarmola, D. Budker, and R.L. Walsworth. Solid-state electronic spin coherence time approaching one second. *Nature Communications*, 4(1):1743, June 2013.
- [73] A. H. Ghadimi, S. A. Fedorov, N. J. Engelsen, M. J. Breyhi, R. Schilling, D. J. Wilson, and T. J. Kippenberg. Elastic strain engineering for ultralow mechanical dissipation. *Science*, 360(6390):764–768, May 2018.
- [74] M. H. Abobeih, J. Randall, C. E. Bradley, H. P. Bartling, M. A. Bakker, M. J. Degen, M. Markham, D. J. Twitchen, and T. H. Taminiau. Atomic-scale imaging of a 27-nuclear-spin cluster using a quantum sensor. *Nature*, 576(7787):411–415, December 2019.
- [75] Jarryd J. Pla, Kuan Y. Tan, Juan P. Dehollain, Wee H. Lim, John J. L. Morton, David N. Jamieson, Andrew S. Dzurak, and Andrea Morello. A single-atom electron spin qubit in silicon. *Nature*, 489(7417):541–545, September 2012.
- [76] Leigh S. Martin and K. Birgitta Whaley. Single-shot deterministic entanglement between non-interacting systems with linear optics. *arXiv:1912.00067 [quant-ph]*, November 2019.