



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

# High Temperature Superconductivity in a Lightly Doped Quantum Spin Liquid

Hong-Chen Jiang and Steven A. Kivelson

Phys. Rev. Lett. **127**, 097002 — Published 27 August 2021

DOI: [10.1103/PhysRevLett.127.097002](https://doi.org/10.1103/PhysRevLett.127.097002)

# High Temperature Superconductivity in a Lightly Doped Quantum Spin Liquid

Hong-Chen Jiang<sup>1,\*</sup> and Steven A. Kivelson<sup>2</sup>

<sup>1</sup>*Stanford Institute for Materials and Energy Sciences,*

*SLAC National Accelerator Laboratory and Stanford University, Menlo Park, California 94025, USA*

<sup>2</sup>*Department of Physics, Stanford University, Stanford, California 94305, USA*

(Dated: July 20, 2021)

We have performed density-matrix renormalization group studies of a square lattice  $t$ - $J$  model with small hole doping,  $\delta \ll 1$ , on long 4 and 6 leg cylinders. We include frustration in the form of a second-neighbor exchange coupling,  $J_2 = J_1/2$ , such that the undoped ( $\delta = 0$ ) “parent” state is a quantum spin liquid. In contrast to the relatively short range superconducting (SC) correlations that have been observed in recent studies of the 6-leg cylinder in the absence of frustration, we find power law SC correlations with a Luttinger exponent,  $K_{sc} \approx 1$ , consistent with a strongly diverging SC susceptibility,  $\chi \sim T^{-(2-K_{sc})}$  as the temperature  $T \rightarrow 0$ . The spin-spin correlations - as in the undoped state - fall exponentially suggesting that the SC “pairing” correlations evolve smoothly from the insulating parent state.

Although the physics of the cuprate high temperature superconductors is surely complex, there are a variety of reasons[1–3] to believe that the “essential”[4] physics is captured by the two-dimensional (2D) Hubbard model or its close relatives. To begin with, as is the case in the cuprates, in an appropriate regime of parameters, the Hubbard model on a square lattice with  $n = 1$  electrons per site exhibits an undoped “parent” state that is a Mott insulating antiferromagnet. However, two key theoretical issues concerning this proposition remain unsettled: 1) Does d-wave superconductivity (SC) “robustly” arise in this model upon light doping, i.e. for  $0 < \delta \equiv (1-n) \ll 1$ . 2) If so, how does it arise (i.e. what is the “mechanism”) and under what circumstances (e.g. does it depend on specific features of the band structure)?

For parametrically small values of the Hubbard  $U \ll W$  (where  $W$  is the bandwidth), it is possible to establish[5] that such a superconducting state arises, but here (except under extremely fine tuned circumstances in which the Fermi surface is perfectly nested) the undoped state at  $n = 1$  is also superconducting, and the superconducting  $T_c$  is exponentially small in units of  $W$ . For intermediate  $U \sim W$ , no controlled analytic approach exists, but calculations based on a variety of physically motivated approximations [6–8] yield results suggestive of values of  $T_c$  as large as  $T_c \sim W$  (where the proportionality is a number of order 1 but may be small, e.g.  $\sim (2\pi)^{-2}$ ). This was further supported by density-matrix normalization group (DMRG) studies of the Hubbard and  $t$ - $J$  models on 4-leg square cylinders.[9–13] However, recent[14] DMRG calculations on 6-leg square cylinders, as well as variational Monte Carlo[15] calculations on 2D models, have called this proposition into question. Specifically, the tendency of a doped antiferromagnet to phase separation[15, 16] or to charge-density wave (CDW) formation[11, 13, 14, 17–21] appear to play a much more dominant role in the physics at small  $\delta$  than accounted for by most approximate approaches.

One attractive notion that was suggested early on

is that high temperature superconductivity could arise naturally[1, 22–27] under circumstances in which the insulating parent state is a quantum spin liquid (QSL) rather than an ordered antiferromagnet. In particular, a QSL with a gap (even a partial gap with nodes), can in some sense be thought of as a state with pre-existing Cooper pairs but with vanishing superfluid stiffness. Then, upon light doping, one might naturally expect SC with a gap scale that is inherited from the QSL (i.e. evolves continuously as  $\delta \rightarrow 0$ ) and with a superfluid stiffness - that rises with  $\delta$ .

In the present paper, we explore the possibility of SC in a doped spin liquid using DMRG to treat the  $t$ - $J$  model (a proxy for the Hubbard model) on cylinders of circumference 4 and 6. A number of studies of the spin-1/2 Heisenberg model on the square lattice with first and second neighbor exchange couplings,  $J_1$  and  $J_2$ , have led to a consensus[28–35] that there is a QSL phase in the range of  $0.46 < J_2/J_1 < 0.52$ . [35] In this range, DMRG on cylinders of circumference up to  $L_y = 10$  show a pronounced spin-gap and exponentially falling spin-spin correlations with a correlation length  $\xi_s$  considerably smaller than  $L_y$ . [30, 32] However, there is still some debate about whether this gap persists in the 2D limit, or if instead the QSL phase has a gapless nodal spinon spectrum.

Here, we study the model with  $J_2/J_1 = 0.5$ , and correspondingly we take the ratio of nearest to next-nearest neighbor hopping matrix elements,  $t_2/t_1 = 0.7 \approx \sqrt{J_2/J_1}$ , and a value of  $J_1/t_1 = 1/3$  corresponding loosely to a value of  $U \approx 4t^2/J = 12t$ . On the cylinders we study, the undoped system is fully gapped, so effectively corresponds to a compactified version of a gapped  $Z_2$  spin liquid of the sort that arises in the quantum dimer model[36, 37] and the toric code model[38]. Upon lightly doping we find a state which still shows exponentially falling spin-spin correlations, with correlation lengths that are longer than but of the same order as in the undoped system. Most importantly, we find that even at the smallest  $\delta$  and on our largest 6-leg cylinders, the

SC correlations are strong and decay with a slow power law,  $\sim |r|^{-K_{sc}}$ , with  $K_{sc} \approx 1$ . This slow decay implies a SC susceptibility that diverges as  $\chi_{sc} \sim T^{-(2-K_{sc})}$  as  $T \rightarrow 0$ . As far as we know, to date, *this is the strongest indication of SC that has been found in any DMRG study of a system on the square lattice of width  $L_y > 4$* . Moreover, the SC correlations dominate over the CDW correlations in the sense that in all cases  $K_c > K_{sc}$ . This is suggestive that SC order is realized in the 2D limit, although given that in all cases the CDW correlations are substantial, this inference remains speculative.

**Model and Method:** We employ DMRG[39] to study the ground state properties of the hole-doped  $t$ - $J$  model on the square lattice, which is defined by the Hamiltonian

$$H = - \sum_{ij\sigma} t_{ij} \left( \hat{c}_{i\sigma}^\dagger \hat{c}_{j\sigma} + h.c. \right) + \sum_{ij} J_{ij} \left( \vec{S}_i \cdot \vec{S}_j - \frac{\hat{n}_i \hat{n}_j}{4} \right).$$

Here  $\hat{c}_{i\sigma}^\dagger$  ( $\hat{c}_{i\sigma}$ ) is the electron creation (annihilation) operator on site  $i = (x_i, y_i)$  with spin polarization  $\sigma$ ,  $\vec{S}_i$  is the spin operator and  $\hat{n}_i = \sum_{\sigma} \hat{c}_{i\sigma}^\dagger \hat{c}_{i\sigma}$  is the electron number operator. The electron hopping amplitude  $t_{ij}$  is equal to  $t_1$  ( $t_2$ ) if  $i$  and  $j$  are NN (NNN) sites.  $J_1$  and  $J_2$  are the spin superexchange interactions between NN and NNN sites, respectively. The Hilbert space is constrained by the no-double occupancy condition,  $n_i \leq 1$ . At half-filling, i.e.,  $n_i = 1$ ,  $H$  reduces to the spin-1/2 antiferromagnetic  $J_1$ - $J_2$  Heisenberg model.

We take the lattice geometry to be cylindrical with periodic and open boundary conditions in the  $\hat{y}$  and  $\hat{x}$  directions, respectively. Here  $\hat{y} = (0, 1)$  and  $\hat{x} = (1, 0)$  are the two basis vectors of the square lattice. Here, we focus on cylinders with width  $L_y$  and length  $L_x$ , where  $L_x$  and  $L_y$  are the number of sites along the  $\hat{x}$  and  $\hat{y}$  directions, respectively. The total number of sites is  $N = L_x \times L_y$ , the number of electrons  $N_e$ , and the doping level of the system is defined as  $\delta = N_h/N$ , where  $N_h = N - N_e$  is the number of doped holes relative to the half-filled insulator with  $N_e = N$ . In the present study, we focus on  $L_y = 4$  cylinders of length up to  $L_x = 128$  and  $L_y = 6$  cylinders of length up to  $L_x = 48$ , and for values of  $\delta = 1/18, 1/16$ , and  $1/12$ . We set  $J_1=1$  as an energy unit and  $J_2 = 0.5$  such that the undoped system is deep in the QSL phase at half-filling.[28, 30, 32, 35] We consider  $t_1 = 3$  and  $t_2 = t_1 \sqrt{J_2/J_1}$  to make a connection to the corresponding Hubbard model. We perform up to 90 sweeps and keep up to  $m = 10000$  states for  $L_y = 4$  cylinders with a typical truncation error  $\epsilon < 10^{-7}$ , and up to  $m = 40000$  states for  $L_y = 6$  cylinders with a typical truncation error  $\epsilon < 10^{-6}$ . Further details of the numerical simulation are provided in the Supplemental Material (SM).[40]

**Superconducting pair-field correlations:** We have calculated the equal-time spin-singlet SC pair-field




FIG. 1. (Color online) Superconducting pair-field correlations  $\Phi_{yy}(r)$  on double-logarithmic scales for (A)  $L_y = 4$  cylinders at  $\delta = 1/12$  and  $\delta = 1/16$ , and (B)  $L_y = 6$  cylinders at  $\delta = 1/12$  and  $\delta = 1/18$ .  $r$  is the distance between two Cooper pairs in the  $\hat{x}$  direction. The dashed lines denote power-law fitting to  $\Phi_{yy}(r) \sim r^{-K_{sc}}$ .

correlation function

$$\Phi_{\alpha\beta}(r) = \frac{1}{L_y} \sum_{y=1}^{L_y} |\langle \Delta_{\alpha}^{\dagger}(x_0, y) \Delta_{\beta}(x_0 + r, y) \rangle|. \quad (1)$$

$\Delta_{\alpha}^{\dagger}(x, y) = \frac{1}{\sqrt{2}} [\hat{c}_{(x,y),\uparrow}^{\dagger} \hat{c}_{(x,y)+\alpha,\downarrow}^{\dagger} + \hat{c}_{(x,y)+\alpha,\uparrow}^{\dagger} \hat{c}_{(x,y),\downarrow}^{\dagger}]$  is the spin-singlet pair creation operator on bond  $\alpha = \hat{x}$  or  $\hat{y}$ , where  $(x_0, y)$  is a reference bond taken as  $x_0 \sim L_x/4$  and  $r$  is the displacement between bonds in the  $\hat{x}$  direction.

Fig.1 shows  $\Phi_{yy}(r)$  for both  $L_y = 4$  and  $L_y = 6$  cylinders at different doping levels. At long distance,  $\Phi(r)$  is characterized by a power-law with the appropriate Luttinger exponent  $K_{sc}$  defined by

$$\Phi(r) \sim r^{-K_{sc}}. \quad (2)$$

The exponent  $K_{sc}$ , which is obtained by fitting the results using Eq.(2), is  $K_{sc} = 1.08(4)$  for  $\delta = 1/12$  and  $K_{sc} = 0.95(2)$  for  $\delta = 1/16$  on  $L_y = 4$  cylinders, and  $K_{sc} = 1.26(7)$  for  $\delta = 1/12$  and  $K_{sc} = 1.14(5)$  for  $\delta = 1/18$  on  $L_y = 6$  cylinders. This establishes that the lightly doped QSL on both  $L_y = 4$  and  $L_y = 6$  cylinders has quasi-long-range SC correlations. In addition to  $\Phi_{yy}(r)$ , we have also calculated components of the tensor  $-\Phi_{xx}(r)$  and  $\Phi_{xy}(r)$  – and find that  $\Phi_{xx}(r) \sim \Phi_{yy}(r) \sim -\Phi_{xy}(r)$ . In short, the SC correlations have a d-wave form.

**CDW correlations:** To measure the charge order, we define the rung density operator  $\hat{n}(x) =$




FIG. 2. (Color online) Charge density profiles  $n(x)$  for (A)  $L_y = 4$  cylinders at  $\delta = 1/12$  and  $\delta = 1/16$ , and (B)  $L_y = 6$  cylinders at  $\delta = 1/12$  and  $\delta = 1/18$ . The exponent  $K_c$  is extracted using Eq. (3), with the data points in grey neglected to minimize boundary effects.

$L_y^{-1} \sum_{y=1}^{L_y} \hat{n}(x, y)$  and its expectation value  $n(x) = \langle \hat{n}(x) \rangle$ . Fig. 2A shows the charge density distribution  $n(x)$  for  $L_y = 4$  cylinders, which is consistent with “half-filled charge stripes” with wavelength  $\lambda = 1/2\delta$ . This corresponds to an ordering wavevector  $Q = 4\pi\delta$  corresponding to half a doped hole per 2D unit cell, i.e. viewing the cylinder as a 1D system, 2 holes per 1D unit cell. The charge density profile  $n(x)$  for  $L_y = 6$  cylinders is shown in Fig. 2B, which has wavelength  $\lambda = 1/3\delta$ , consistent with “third-filled” charge stripes. This corresponds to an ordering wavevector  $Q = 6\pi\delta$  and one third of a doped hole per 2D unit cell - again corresponding to 2 holes per 1D unit cell.

At long distance, the spatial decay of the CDW correlation is dominated by a power-law with the Luttinger exponent  $K_c$ . The exponent  $K_c$  can be obtained by fitting the charge density oscillations (Friedel oscillations) induced by the boundaries of the cylinder[41]

$$n(x) = n_0 + A_Q * \cos(Qx + \phi)x^{-K_c/2}. \quad (3)$$

Here  $A_Q$  is an amplitude,  $\phi$  is a phase shift,  $n_0 = 1 - \delta$  is the mean density, and  $Q = 4\pi\delta$ . Note that a few data points (Fig. 2A and B, light grey color) are excluded to minimize the boundary effect and improve the fitting quality. The extracted exponents for  $L_y = 4$  cylinders are  $K_c = 1.29(3)$  when  $\delta = 1/12$  and  $K_c = 1.37(3)$  when  $\delta = 1/16$ . For  $L_y = 6$  cylinders,  $K_c = 1.42(5)$  when  $\delta = 1/12$  and  $K_c = 1.55(5)$  when  $\delta = 1/18$ . Similarly,  $K_c$  can also be obtained from the charge density-density fluctuation correlation which gives qualitatively consistent results (see SM).




FIG. 3. (Color online) Spin-spin correlations  $F(r)$  for  $L_y = 6$  cylinders at  $\delta = 0$ ,  $\delta = 1/12$  and  $\delta = 1/18$  on the semi-logarithmic scale. Dashed lines denote exponential fit  $F(r) \sim e^{-r/\xi_s}$ , where  $r$  is the distance between two sites in the  $\hat{x}$  direction. Inset: Spin gap  $\Delta_s$  for  $L_y = 6$  cylinders at  $\delta = 0$  and  $\delta = 1/12$ . Solid lines denote second-order polynomial fitting.

**Spin-spin correlations:** To describe the magnetic properties of the ground state, we calculate the spin-spin correlation functions defined as

$$F(r) = \frac{1}{L_y} \sum_{y=1}^{L_y} |\langle \vec{S}_{x_0, y} \cdot \vec{S}_{x_0+r, y} \rangle|. \quad (4)$$

Fig. 3 shows  $F(r)$  for  $L_y = 6$  cylinders at different doping levels, which decays exponentially as  $F(r) \sim e^{-r/\xi_s}$  at long-distances, with a correlation length  $\xi_s = 3.98(1)$  lattice spacings for  $\delta = 1/12$  and  $\xi_s = 3.06(2)$  lattice spacings for  $\delta = 1/18$ . For comparison, the spin-spin correlation  $F(r)$  at half-filling, i.e.,  $\delta = 0$ , is also shown, which decays exponentially with a correlation length  $\xi_s = 1.42(1)$ . Therefore, the spin-spin correlations at finite doping levels are short-ranged and similar to those of the QSL at half-filling. In the inset of Fig. 3, we show the spin gap, defined as  $\Delta_s = E_0(S_z = 1) - E_0(S_z = 0)$ , where  $E_0(S_z)$  is the ground state energy of a system with total spin  $S_z$ . At half-filling, i.e.,  $\delta = 0$ ,  $\Delta_s = 0.40(1)$  which is consistent with previous studies.[30, 32] At  $\delta = 1/12$ ,  $\Delta_s = 0.24(1)$ , which is consistent with the short-range nature of  $F(r)$ .

**Single particle Green function:** We have also calculated the single-particle Green function, defined as

$$G(r) = \frac{1}{L_y} \sum_{y=1}^{L_y} \langle c_{(x_0, y), \sigma}^\dagger c_{(x_0+r, y), \sigma} \rangle. \quad (5)$$

Fig. 4 shows  $G(r)$  for both  $L_y = 4$  and  $L_y = 6$  cylinders at different doping levels, the long distance behavior of  $G$  is consistent with exponential decay  $G(r) \sim e^{-r/\xi_G}$ . The extracted correlation lengths for  $L_y = 4$  cylinders are  $\xi_G = 30(2)$  when  $\delta = 1/12$  and  $\xi_G = 18(1)$  when  $\delta = 1/16$ , while for  $L_y = 6$  cylinders,  $\xi_G = 21(1)$  when  $\delta = 1/12$  and  $\xi_G = 20(2)$  when  $\delta = 1/18$ .




FIG. 4. (Color online) Single-particle Green function  $G(r)$  for (A)  $L_y = 4$  cylinders at  $\delta = 1/12$  and  $\delta = 1/16$ , and (B)  $L_y = 6$  cylinders at  $\delta = 1/12$  and  $\delta = 1/18$  on the semi-logarithmic scale. Dashed line denote exponential fitting  $G(r) \sim e^{-r/\xi_G}$  where  $r$  is the distance between two sites in the  $\hat{x}$  direction.




FIG. 5. (Color online) Hole momentum distribution function  $n^h(k_x, k_y)$  for  $L_y = 4$  cylinders at (A)  $\delta = 1/12$  and (B)  $\delta = 1/16$ , and  $L_y = 6$  cylinders at (C)  $\delta = 1/12$  and (D)  $\delta = 1/18$  at different  $k_y$  as a function of  $k_x$  in unit of  $\pi$ .

We have also measured the hole momentum distribution function defined as

$$n^h(\mathbf{k}) = \frac{1}{2} \left[ 2 - \sum_{\sigma} n_{\sigma}(\mathbf{k}) \right]. \quad (6)$$

Here  $n_{\sigma}(\mathbf{k}) = \frac{1}{N} \sum_{ij} e^{i\mathbf{k} \cdot (\mathbf{r}_i - \mathbf{r}_j)} \langle \hat{c}_{i\sigma}^{\dagger} \hat{c}_{j\sigma} \rangle$  is the electron momentum distribution function for electron with spin- $\sigma$ . Fig.5 shows  $n^h(\mathbf{k})$  for both  $L_y = 4$  and  $L_y = 6$  cylinders at different doping levels. Not surprisingly, there are no clear discontinuities in  $n^h(\mathbf{k})$  of the sort that would be expected at the Fermi momenta of a Fermi liquid.

However, there are sharp drops in  $n^h(\mathbf{k})$  (which can be identified as maxima of  $|dn^h(\mathbf{k})/dk_x|$ ) (See SM for details) that are suggestive of the “near existence” of a Fermi surface. These features are most prominent for  $k_y = 0$ , where they occur at  $k_x \approx \pi \pm k_0$ , but there are slightly broader features of the same general sort for  $k_y = \pi$ , at  $k_x = \pm k_{\pi}$ . For  $L_y = 4$  and  $\delta = 1/16$ ,  $k_0 = 0.075\pi$  and  $k_{\pi} = 0.175\pi$ ; for  $L_y = 4$  and  $\delta = 1/12$ ,  $k_0 = 0.175\pi$  and  $k_{\pi} = 0.192\pi$ ; for  $L_y = 6$  and  $\delta = 1/18$ ,  $k_0 = 0.17\pi$  and  $k_{\pi} = 0.18\pi$ ; for  $L_y = 6$  and  $\delta = 1/12$ ,  $k_0 = 0.25\pi$  and  $k_{\pi} = 0.25\pi$ . Within the numerical uncertainty, there is a direct relation between these quasi-Fermi momenta and the CDW ordering vector:  $Q = 2(k_0 + k_{\pi})$ . Moreover, since  $Q = \pi L_y \delta$ , this corresponds to the expected value of  $2k_F$  that would correspond to the “volume” of the Fermi surface under conditions (not satisfied in the present case) in which Luttinger’s theorem applies.

**Conclusion:** There is necessarily a speculative leap from results on finite cylinders to the 2D limit. However, we feel that the present results - and those of a similar study by one of us on the triangular lattice  $t$ - $J$  model on 4 and 6 leg cylinders[42, 43] - can plausibly be taken as representative of the solution of the corresponding 2D problem. In particular, they support the proposition that SC can emerge upon light doping of a QSL.[44]

Conversely, our earlier observation of an insulating holon crystal in a lightly doped Kagome system[45, 46] and CDW order in a lightly doped honeycomb Kiatev spin liquid[47] imply that SC is not the universal result of doping a QSL. Indeed, for otherwise identical cylinders to those reported above, reversing the sign of  $t_2$  (i.e. taking  $t_2 = -t_1 \sqrt{J_2/J_1}$ ) reduces the long distance SC correlations by many orders of magnitude although whether some weak SC power-law correlations persist is still unsettled. Moreover, we have also found greatly enhanced SC correlations on 4 and 6 leg cylinders with a spatially modulated (“striped”) version of the square-lattice Hubbard model[48]; it thus may be aspects of doping a quantum paramagnet (i.e. a system in which quantum fluctuations are sufficient to destroy magnetic order) rather than specific features of a doped QSL that is responsible for the strong SC tendencies.

It is harder still to make inferences about  $T_c$  itself in the 2D limit. The large values of the spin-gaps,  $\Delta_s \sim J/4$ , are suggestive that pairing is sufficiently strong to persist to very high  $T$ . It is therefore likely that  $T_c$  is determined by the phase ordering scale,[49] in other words that the zero temperature superfluid stiffness and hence  $T_c$  itself rise roughly linearly with  $\delta$  for  $\delta \ll 1$ . [22]

*Note added:* We have become aware of two independent but closely related DMRG studies of the  $t$ - $t'$ - $J$  model by Gong, Zhu, and Sheng[50] (GZS) and Jiang, Scalapino, and White[51] (JSW). Both report results for 6 leg cylinders, while JSW also have results for 8 leg cylinders. Both studies investigated  $t'$  in the range  $0 \leq t' \leq 0.3t$

- i.e. neither include the large value  $t' = t/\sqrt{2}$  studied here. JSW also studied negative  $t'$  - the sign that is thought to be relevant to the hole-doped cuprates - in the range  $-0.3t \leq t \leq 0$ . Overall, the two papers agree that increasing (positive)  $t'$  tends to increase the tendency to d-wave SC order and decrease the tendency to various sorts of competing SDW and CDW orders, which also correlates well with our observations at larger  $t'$ . Conversely, JSW find that negative  $t'$  strengthens stripe order and depresses SC order, consistent with our already mentioned failure to find strong SC tendencies for  $t' = -t/\sqrt{2}$ . One should notice, however, that there are significant differences in other aspects of the inferred phase diagrams reported by GZS and JSW - which likely reflects the delicate nature of the phase competition between multiple phases that occurs in the range of  $t'$  and doping they have explored.

*Acknowledgments:* We would like to thank Erez Berg, Johannes Motruk, Zheng-Yu Weng and Yi-Fan Jiang for helpful discussions. This work was supported by the Department of Energy, Office of Science, Basic Energy Sciences, Materials Sciences and Engineering Division, under Contract DE-AC02-76SF00515.

---

\* [hcjiang@stanford.edu](mailto:hcjiang@stanford.edu)

- [1] P. W. Anderson. The Resonating valence bond state in  $\text{La}_2\text{CuO}_4$  and superconductivity. *Science*, 235(4793):1196–1198, 1987.
- [2] V. J. Emery. Theory of high- $T_c$  superconductivity in oxides. *Phys. Rev. Lett.*, 58:2794–2797, Jun 1987.
- [3] D. J. Scalapino. A common thread: The pairing interaction for unconventional superconductors. *Reviews of Modern Physics*, 84(4):1383–1417, October 2012.
- [4] Sophia Kivelson and Steven Kivelson. Understanding complexity. *Nature Physics*, 14(5):426–427, May 2018.
- [5] S. Raghu, S. A. Kivelson, and D. J. Scalapino. Superconductivity in the repulsive Hubbard model: An asymptotically exact weak-coupling solution. *Phys. Rev. B*, 81:224505, Jun 2010.
- [6] E. Gull and A. J. Millis. Quasiparticle properties of the superconducting state of the two-dimensional hubbard model. *Phys. Rev. B*, 91:085116, Feb 2015.
- [7] P. Staar, M. Jiang, U. R. Hähner, T. C. Schulthess, and T. A. Maier. Interlaced coarse-graining for the dynamic cluster approximation. *Phys. Rev. B*, 93:165144, Apr 2016.
- [8] Yuxuan Wang, Artem Abanov, Boris L. Altshuler, Emil A. Yuzbashyan, and Andrey V. Chubukov. Superconductivity near a quantum-critical point: The special role of the first matsubara frequency. *Phys. Rev. Lett.*, 117:157001, Oct 2016.
- [9] Hong-Chen Jiang, Zheng-Yu Weng, and Steven A. Kivelson. Superconductivity in the doped  $t-J$  model: Results for four-leg cylinders. *Phys. Rev. B*, 98:140505, Oct 2018.
- [10] Hong-Chen Jiang and Thomas P. Devereaux. Superconductivity in the doped hubbard model and its interplay with next-nearest hopping  $t'$ . *Science*, 365(6460):1424–1428, 2019.
- [11] Yi-Fan Jiang, Jan Zaanen, Thomas P. Devereaux, and Hong-Chen Jiang. Ground state phase diagram of the doped hubbard model on the four-leg cylinder. *Phys. Rev. Research*, 2:033073, Jul 2020.
- [12] Hong-Chen Jiang, Shuai Chen, and Zheng-Yu Weng. Critical role of the sign structure in the doped mott insulator: Luther-emery versus fermi-liquid-like state in quasi-one-dimensional ladders. *Phys. Rev. B*, 102:104512, Sep 2020.
- [13] Chia-Min Chung, Mingpu Qin, Shiwei Zhang, Ulrich Schollwöck, and Steven R. White. Plaquette versus ordinary  $d$ -wave pairing in the  $t'$ -hubbard model on a width-4 cylinder. *Phys. Rev. B*, 102:041106, Jul 2020.
- [14] J. P. F. LeBlanc, Andrey E. Antipov, Federico Becca, Ireneusz W. Bulik, Garnet Kin-Lic Chan, Chia-Min Chung, Youjin Deng, Michel Ferrero, Thomas M. Henderson, Carlos A. Jiménez-Hoyos, E. Kozik, Xuan-Wen Liu, Andrew J. Millis, N. V. Prokof'ev, Mingpu Qin, Gustavo E. Scuseria, Hao Shi, B. V. Svistunov, Luca F. Tocchio, I. S. Tupitsyn, Steven R. White, Shiwei Zhang, Bo-Xiao Zheng, Zhenyue Zhu, and Emanuel Gull. Solutions of the two-dimensional Hubbard model: benchmarks and results from a wide range of numerical algorithms. *Phys. Rev. X*, 5:041041, Dec 2015.
- [15] Sandro Sorella. The phase diagram of the Hubbard model by variational auxiliary field quantum Monte Carlo. *arXiv e-print*, page arXiv:2101.07045, Jan 2021.
- [16] V. J. Emery, H. Q. Lin, and S.A. Kivelson. Phase separation in the  $t$ - $J$  model. *Phys. Rev. Lett.*, 64:475–478, Jan 1990.
- [17] Jan Zaanen and Olle Gunnarsson. Charged magnetic domain lines and the magnetism of high- $T_c$  oxides. *Phys. Rev. B*, 40:7391–7394, Oct 1989.
- [18] Schulz, H.J. Domain walls in a doped antiferromagnet. *J. Phys. France*, 50(18):2833–2849, 1989.
- [19] Kazushige Machida. Magnetism in  $\text{La}_2\text{CuO}_4$  based compounds. *Physica C: Superconductivity*, 158(1):192 – 196, 1989.
- [20] S. R. White and D. J. Scalapino. Ground states of the doped four-leg  $t$ - $J$  ladder. *Phys. Rev. B*, 55(22):14701–14704, June 1997.
- [21] Mingpu Qin, Chia-Min Chung, Hao Shi, Ettore Vitali, Claudius Hubig, Ulrich Schollwöck, Steven R. White, and Shiwei Zhang. Absence of superconductivity in the pure two-dimensional Hubbard model. *Phys. Rev. X*, 10:031016, Jul 2020.
- [22] S. A. Kivelson, D. S. Rokhsar, and J. P. Sethna. Topology of the resonating valence-bond state - solitons and High  $T_c$  superconductivity. *Phys. Rev. B*, 35(16):8865–8868, June 1 1987.
- [23] R. B. Laughlin. The relationship between high-temperature superconductivity and the fractional quantum Hall effect. *Science*, 242:525–533, October 1988.
- [24] G. Kotliar. Resonating valence bonds and  $d$ -wave superconductivity. *Phys. Rev. B*, 37(7):3664–3666, MAR 1 1988.
- [25] L. Balents, M. P. A. Fisher, and C. Nayak. Nodal liquid theory of the pseudo-gap phase of high- $T_c$  superconductors. *Int. Jour. Mod. Phys. B*, 12(10):1033–1068, Apr 1998.
- [26] P. W. Anderson, P. A. Lee, M. Randeria, T. M. Rice, N. Trivedi, and F. C. Zhang. The physics behind

- high-temperature superconducting cuprates: the ‘plain vanilla’ version of RVB. *Journal of Physics-Condensed Matter*, 16(24):R755–R769, June 23 2004.
- [27] Xue-Yang Song, Ashvin Vishwanath, and Ya-Hui Zhang. Doping the chiral spin liquid – topological superconductor or chiral metal? [arXiv:2011.10044](https://arxiv.org/abs/2011.10044).
- [28] F. Figueirido, A. Karlhede, S. Kivelson, S. Sondhi, M. Rocek, and D. S. Rokhsar. Exact diagonalization of finite frustrated spin- $\frac{1}{2}$  Heisenberg models. *Phys. Rev. B*, 41:4619–4632, Mar 1990.
- [29] Luca Capriotti, Federico Becca, Alberto Parola, and Sandro Sorella. Resonating valence bond wave functions for strongly frustrated spin systems. *Phys. Rev. Lett.*, 87:097201, Aug 2001.
- [30] Hong-Chen Jiang, Hong Yao, and Leon Balents. Spin liquid ground state of the spin- $\frac{1}{2}$  square  $J_1$ - $J_2$  heisenberg model. *Phys. Rev. B*, 86:024424, Jul 2012.
- [31] Wen-Jun Hu, Federico Becca, Alberto Parola, and Sandro Sorella. Direct evidence for a gapless  $Z_2$  spin liquid by frustrating néel antiferromagnetism. *Phys. Rev. B*, 88:060402, Aug 2013.
- [32] Shou-Shu Gong, Wei Zhu, D. N. Sheng, Olexei I. Motrunich, and Matthew P. A. Fisher. Plaquette ordered phase and quantum phase diagram in the spin- $\frac{1}{2}$   $J_1$ - $J_2$  square heisenberg model. *Phys. Rev. Lett.*, 113:027201, Jul 2014.
- [33] Satoshi Morita, Ryui Kaneko, and Masatoshi Imada. Quantum spin liquid in spin 1/2  $j_1$ - $j_2$  heisenberg model on square lattice: Many-variable variational monte carlo study combined with quantum-number projections. *Journal of the Physical Society of Japan*, 84(2):024720, 2015.
- [34] Ling Wang, Zheng-Cheng Gu, Frank Verstraete, and Xiao-Gang Wen. Tensor-product state approach to spin- $\frac{1}{2}$  square  $J_1$ - $J_2$  antiferromagnetic heisenberg model: Evidence for deconfined quantum criticality. *Phys. Rev. B*, 94:075143, Aug 2016.
- [35] Ling Wang and Anders W. Sandvik. Critical level crossings and gapless spin liquid in the square-lattice spin-1/2  $J_1$  -  $J_2$  heisenberg antiferromagnet. *Phys. Rev. Lett.*, 121:107202, Sep 2018.
- [36] Daniel S. Rokhsar and Steven A. Kivelson. Superconductivity and the Quantum Hard-Core Dimer Gas. *Phys. Rev. Lett.*, 61:2376–2379, Nov 1988.
- [37] R. Moessner and S. L. Sondhi. Resonating valence bond phase in the triangular lattice quantum dimer model. *Phys. Rev. Lett.*, 86:1881–1884, Feb 2001.
- [38] A. Y. Kitaev. Fault-tolerant quantum computation by anyons. *Annals of Physics*, 303(1):2–30, Jan 2003.
- [39] Steven R. White. Density matrix formulation for quantum renormalization groups. *Phys. Rev. Lett.*, 69:2863–2866, Nov 1992.
- [40] See Supplemental Material [url] for more details and results, which includes Refs. [52-53]. [52] P. Calabrese and J. Cardy, *J. Stat. Mech. Theory Exp.*, 2004(6), 2004. [53] M. Fagotti and P. Calabrese., *J. Stat. Mech. Theory Exp.*, 2011(1), Jan 2011.
- Density matrix formulation for quantum renormalization groups. *Phys. Rev. Lett.*, 69:2863–2866, Nov 1992.
- [41] Steven R. White, Ian Affleck, and Douglas J. Scalapino. Friedel oscillations and charge density waves in chains and ladders. *Phys. Rev. B*, 65:165122, Apr 2002.
- [42] Hong-Chen Jiang. Superconductivity in the doped quantum spin liquid on the triangular lattice. [arXiv:1912.06624](https://arxiv.org/abs/1912.06624), December 2019.
- [43] Yi-Fan Jiang and Hong-Chen Jiang. Topological superconductivity in the doped chiral spin liquid on the triangular lattice. *Phys. Rev. Lett.*, 125:157002, Oct 2020.
- [44] It is difficult to determine whether the likely SC state in 2D is fully gapped or has gapless nodal quasi-particles. The large value of  $\xi_G$  we find might be taken as evidence of a small finite-size gap in what would become a gapless spectrum in the 2D limit, but this is far from certain.
- [45] Hong-Chen Jiang, T. Devereaux, and S. A. Kivelson. Holon Wigner crystal in a lightly doped kagome quantum spin liquid. *Phys. Rev. Lett.*, 119:067002, Aug 2017.
- [46] Cheng Peng, Yi-Fan Jiang, Dong-Ning Sheng, and Hong-Chen Jiang. Doping quantum spin liquids on the kagome lattice. *Advanced Quantum Technologies*, 4(3):2000126, 2021.
- [47] Cheng Peng, Yi-Fan Jiang, Thomas P. Devereaux, and Hong-Chen Jiang. Evidence of pair-density wave in doping Kitaev spin liquid on the honeycomb lattice. page [arXiv:2008.03858](https://arxiv.org/abs/2008.03858), Aug 2020.
- [48] Hong-Chen Jiang and Steven A. Kivelson. Stripe order enhanced superconductivity in the hubbard model. [arXiv:2105.07048](https://arxiv.org/abs/2105.07048), 2021.
- [49] V. J. Emery and S. A. Kivelson. Importance of phase fluctuations in superconductors with small superfluid density. *Nature*, 374(6521):434–437, Mar 30 1995.
- [50] Shoushu Gong, W. Zhu, and D. N. Sheng. Robust d-wave superconductivity in the square-lattice  $t$ - $J$  model. *arXiv e-prints*, page [arXiv:2104.03758](https://arxiv.org/abs/2104.03758), April 2021.
- [51] Shengtao Jiang, Douglas J. Scalapino, and Steven R. White. Ground State Phase Diagram of the  $t$ - $t'$ - $J$  model. *arXiv e-prints*, page [arXiv:2104.10149](https://arxiv.org/abs/2104.10149), April 2021.