



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Improved Constraints on Primordial Gravitational Waves using *Planck*, WMAP, and BICEP/Keck Observations through the 2018 Observing Season

P. A. R. Ade et al. (BICEP/Keck Collaboration)

Phys. Rev. Lett. **127**, 151301 — Published 4 October 2021

DOI: [10.1103/PhysRevLett.127.151301](https://doi.org/10.1103/PhysRevLett.127.151301)

BICEP / *Keck* XIII: Improved Constraints on Primordial Gravitational Waves using *Planck*, WMAP, and BICEP/*Keck* Observations through the 2018 Observing Season

BICEP/*Keck* Collaboration: P. A. R. Ade,¹ Z. Ahmed,² M. Amiri,³ D. Barkats,⁴ R. Basu Thakur,⁵ C. A. Bischoff,⁶ D. Beck,^{2,7} J. J. Bock,^{5,8} H. Boenish,⁴ E. Bullock,⁹ V. Buza,¹⁰ J. R. Cheshire IV,⁹ J. Connors,⁴ J. Cornelison,⁴ M. Crumrine,¹¹ A. Cukierman,^{7,2} E. V. Denison,¹² M. Dierickx,⁴ L. Duband,¹³ M. Eiben,⁴ S. Fatigoni,³ J. P. Filippini,^{14,15} S. Fliescher,¹¹ N. Goeckner-Wald,⁷ D. C. Goldfinger,⁴ J. Grayson,⁷ P. Grimes,⁴ G. Hall,¹¹ G. Halal,⁷ M. Halpern,³ E. Hand,⁶ S. Harrison,⁴ S. Henderson,² S. R. Hildebrandt,^{5,8} G. C. Hilton,¹² J. Hubmayr,¹² H. Hui,⁵ K. D. Irwin,^{7,2,12} J. Kang,^{7,5} K. S. Karkare,^{4,10} E. Karpel,⁷ S. Kefeli,⁵ S. A. Kernasovskiy,⁷ J. M. Kovac,^{4,16} C. L. Kuo,^{7,2} K. Lau,¹¹ E. M. Leitch,¹⁰ A. Lennox,¹⁴ K. G. Megerian,⁸ L. Minutolo,⁵ L. Monceli,⁵ Y. Nakato,⁷ T. Namikawa,¹⁷ H. T. Nguyen,⁸ R. O’Brien,^{5,8} R. W. Ogburn IV,^{7,2} S. Palladino,⁶ T. Prouve,¹³ C. Pryke,^{11,9,*} B. Racine,^{4,18} C. D. Reintsema,¹² S. Richter,⁴ A. Schillaci,⁵ R. Schwarz,¹¹ B. L. Schmitt,⁴ C. D. Sheehy,¹⁹ A. Soliman,⁵ T. St. Germaine,^{4,16} B. Steinbach,⁵ R. V. Sudiwala,¹ G. P. Teply,⁵ K. L. Thompson,^{7,2} J. E. Tolan,⁷ C. Tucker,¹ A. D. Turner,⁸ C. Umiltà,^{6,14} C. Vergès,⁴ A. G. Viereggs,^{20,10} A. Wandui,⁵ A. C. Weber,⁸ D. V. Wiebe,³ J. Willmert,¹¹ C. L. Wong,^{4,16} W. L. K. Wu,² H. Yang,⁷ K. W. Yoon,^{7,2} E. Young,^{7,2} C. Yu,⁷ L. Zeng,⁴ C. Zhang,⁵ and S. Zhang⁵

¹*School of Physics and Astronomy, Cardiff University, Cardiff, CF24 3AA, United Kingdom*

²*Kavli Institute for Particle Astrophysics and Cosmology,*

SLAC National Accelerator Laboratory, 2575 Sand Hill Rd, Menlo Park, California 94025, USA

³*Department of Physics and Astronomy, University of British Columbia, Vancouver, British Columbia, V6T 1Z1, Canada*

⁴*Center for Astrophysics, Harvard & Smithsonian, Cambridge, MA 02138, U.S.A*

⁵*Department of Physics, California Institute of Technology, Pasadena, California 91125, USA*

⁶*Department of Physics, University of Cincinnati, Cincinnati, Ohio 45221, USA*

⁷*Department of Physics, Stanford University, Stanford, California 94305, USA*

⁸*Jet Propulsion Laboratory, Pasadena, California 91109, USA*

⁹*Minnesota Institute for Astrophysics, University of Minnesota, Minneapolis, Minnesota 55455, USA*

¹⁰*Kavli Institute for Cosmological Physics, University of Chicago, Chicago, IL 60637, USA*

¹¹*School of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota 55455, USA*

¹²*National Institute of Standards and Technology, Boulder, Colorado 80305, USA*

¹³*Service des Basses Températures, Commissariat à l’Energie Atomique, 38054 Grenoble, France*

¹⁴*Department of Physics, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA*

¹⁵*Department of Astronomy, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA*

¹⁶*Department of Physics, Harvard University, Cambridge, MA 02138, USA*

¹⁷*Kavli Institute for the Physics and Mathematics of the Universe (WPI),*

UTIAS, The University of Tokyo, Kashiwa, Chiba 277-8583, Japan

¹⁸*Aix-Marseille Université, CNRS/IN2P3, CPPM, Marseille, France*

¹⁹*Physics Department, Brookhaven National Laboratory, Upton, NY 11973*

²⁰*Department of Physics, Enrico Fermi Institute, University of Chicago, Chicago, IL 60637, USA*

(Draft August 10, 2021. Submitted to PRL)

We present results from an analysis of all data taken by the BICEP2, *Keck Array* and BICEP3 CMB polarization experiments up to and including the 2018 observing season. We add additional *Keck Array* observations at 220 GHz and BICEP3 observations at 95 GHz to the previous 95/150/220 GHz data set. The Q/U maps now reach depths of 2.8, 2.8 and 8.8 μK_{CMB} arcmin at 95, 150 and 220 GHz respectively over an effective area of ≈ 600 square degrees at 95 GHz and ≈ 400 square degrees at 150 & 220 GHz. The 220 GHz maps now achieve a signal-to-noise on polarized dust emission exceeding that of *Planck* at 353 GHz. We take auto- and cross-spectra between these maps and publicly available WMAP and *Planck* maps at frequencies from 23 to 353 GHz and evaluate the joint likelihood of the spectra versus a multicomponent model of lensed- $\Lambda\text{CDM}+r+\text{dust}+\text{synchrotron}+\text{noise}$. The foreground model has seven parameters, and no longer requires a prior on the frequency spectral index of the dust emission taken from measurements on other regions of the sky. This model is an adequate description of the data at the current noise levels. The likelihood analysis yields the constraint $r_{0.05} < 0.036$ at 95% confidence. Running maximum likelihood search on simulations we obtain unbiased results and find that $\sigma(r) = 0.009$. These are the strongest constraints to date on primordial gravitational waves.

PACS numbers: 98.70.Vc, 04.80.Nn, 95.85.Bh, 98.80.Es

Introduction.—The ΛCDM standard model of cosmology is able to describe the observable universe in a sta-

tistical manner using only six free parameters. Measurements of the cosmic microwave background (CMB) [1] are one of the key pillars of this model and now constrain its parameters with percent-level precision (see most recently Ref. [2]).

The Λ CDM model describes how the universe evolved from an initial high energy state ($T \gg 10^{12}$ K), and the conditions at that time can be inferred from observations: fractionally small, Gaussian, adiabatic perturbations with a slightly red power law spectrum ($n_s \lesssim 1$). Inflationary theories naturally explain such conditions as the outcome of a pre-phase of exponential expansion during which the scale of the proto-universe increased by a factor of $\sim e^{60}$. Inflation makes an additional prediction which has not yet been observed—a background of tensor perturbations, also known as gravitational waves (see Ref. [3] for a review and citations to the original literature). There are many specific inflationary models and classes thereof. If we can detect or set limits on primordial gravitational waves we can set limits on these models [4], and probe physics at energy scales far higher than can ever be accessed in laboratory experiments.

A polarization pattern can be decomposed into E -mode (gradient) and B -mode (curl) components. Under the Λ CDM standard model the CMB polarization pattern is mostly E -mode, with a much smaller B -mode component which arises due to gravitational deflections (lensing) of the CMB photons after their last scattering [5]. Since primordial gravitational waves will produce E -modes and B -modes approximately equally it was realized in the late 1990's that the best way to search for them is to look for an excess B -mode signal [6–8]. Additional non-primordial B -modes are produced by astrophysical foreground emissions, primarily from our own galaxy, but these have different frequency spectra than the CMB, and can be separated from it using multi frequency measurements.

Our BICEP/Keck program first reported detection of an excess over the lensing B -mode expectation at 150 GHz in Ref. [9]. In a joint analysis using multi-frequency data from the *Planck* experiment it was shown that most or all of this is due to polarized emission from dust in our own galaxy [10, hereafter BKP]. In Ref. [11, hereafter BK14] we improved the constraint using *Keck Array* data at 95 GHz taken during the 2014 season, and in Ref. [12, hereafter BK15] we improved again adding *Keck Array* data at 95 GHz and 220 GHz taken during the 2015 season. In this letter [hereafter BK18] we add large amounts of new data taken by *Keck Array* at 220 GHz and BICEP3 at 95 GHz during the 2016, 2017 and 2018 observing seasons. This paper follows BK15 very closely in the methods, structure, and, in places, even the wording, mainly just adding additional experimental data. This improves the constraint on primordial gravitational waves parameterized by the tensor-to-scalar ratio r by more than a factor of two over our previous result to

$r_{0.05} < 0.036$ at 95% confidence, setting important additional limits on inflationary models.

Instrument and observations.—The BICEP2 receiver observed at 150 GHz from 2010–2012 [13]. The *Keck Array* was essentially five copies of BICEP2 running in parallel from 2012–2019, initially at 150 GHz but switching over time to 95 and 220 GHz [14]. BICEP3 is a single similar, but scaled up, receiver which commenced science observations in the 2016 Austral winter season [15]. Whereas the BICEP2 and *Keck* 150 & 220 GHz receivers each contained ≈ 500 bolometric detectors BICEP3 contains ≈ 2500 detectors. The aperture size is also increased from ≈ 0.25 m to ≈ 0.5 m. The *Keck* receivers were mounted on a single telescope mount (movable platform), while BICEP3 occupies a separate mount previously used for BICEP2 on a nearby building. All of these telescopes are located at the South Pole Station in Antarctica. The mounts scan the receivers across the sky, and the cryogenic detectors track the intensity of the incoming microwave radiation. The detectors are arranged as interleaved orthogonally polarized pairs in the focal planes and the pair difference timestreams are thus measures of the polarized emission from the sky [16]. At the South Pole the atmosphere is exceptionally transparent and stable at the observation frequencies [17, Fig. 5].

BICEP2 and *Keck Array* both mapped a region of sky centered at RA 0h, Dec. -57.5° with an effective area of ≈ 400 square degrees. BICEP3 has a larger instantaneous field of view and hence naturally maps a larger sky area with an effective area of ≈ 600 square degrees. We have perturbed the center of the BICEP3 scan region such that most of this additional area falls on the higher declination side of the sky patch in an attempt to stay away from regions where the *Planck* data indicates polarized dust contamination may be higher. The BK15 data set consisted of 4/17/2 receiver-years at 95/150/220 GHz respectively. BICEP3 is equivalent to about eight of the *Keck Array* 95 GHz receivers [15] so the BK18 data set is equivalent to about 28/18/14 *Keck* receiver-years at 95/150/220 GHz respectively.

Maps and Power Spectra—We make maps and power spectra using the same procedures as in our previous series of papers. The timestream data are binned into pixels on the sky using knowledge of the pointing direction of the telescope at each moment in time, together with the relative angles from the telescope boresight to each individual detector pair. By taking data with the receivers rotated at a range of angles, maps of the Stokes parameters Q and U can be constructed.

The maps at each observing frequency are subjected to a matrix purification operation [9, 18] such that they contain only structures sourced by B -modes of the underlying sky pattern. This allows us to measure the B -modes in the presence of the much brighter Λ CDM E -modes. The maps are then inverse noise variance apodized, Fourier transformed and rotated from the

Q/U to the E/B basis. In this paper we use our own maps at 95, 150 and 220 GHz plus the 23 & 33 GHz bands of WMAP [19][20] and the 30, 44, 143, 217 and 353 GHz maps from the NPIPE processing of the *Planck* data [21][22]. For illustration purposes we can inverse Fourier transform to form E/B maps. Fig. 1 shows E - and B -mode maps at 95, 150 and 220 GHz. (See Appendix A for the full set of $T/Q/U$ maps [23].)

We take the variance within annuli of the Fourier plane to estimate the angular power spectra. Fig. 2 shows the EE and BB auto- and cross-spectra for the BICEP/*Keck* bands plus the *Planck* 353 GHz band which remains important for constraining the polarized dust contribution. Comparing this plot to Fig. 2 of BK15 we can see that the uncertainties are dramatically reduced for the auto- and cross-spectra of the 95 and 220 GHz bands. The model plotted is a “baseline” lensed- Λ CDM+dust model from our previous BK15 analysis, which remains a good description of the data. The EE spectra were not used to derive the model but agree well with it under the assumption that $EE/BB = 2$ for dust, as is known to be close to the case [24, 25].

To test for systematic contamination we carry out our usual “jackknife” internal consistency (null) tests on the new 95 GHz and 220 GHz data as described in Appendices B and C. Fig. 3 upper shows the noise spectra for the three main BK18 bands after correction for the filter and beam suppression. In an auto-spectrum the quantity which determines the ability to constrain r is the fluctuation of the noise bandpowers rather than their mean. The lower panel therefore shows the effective sky fraction as inferred from the fractional noise fluctuation. Together, these panels provide a useful synoptic measure of the loss of information due to noise, filtering, and EE/BB separation in the lowest bandpowers (and we are glad to see taken up by others as e.g. Fig. 6 of Ref. [26]).

Likelihood Analysis.—We perform likelihood analysis using the methods introduced in BKP and refined in BK14 & BK15. We use the Hamimeche-Lewis approximation [27] to the joint likelihood of the ensemble of 66 BB auto- and cross-spectra taken between the BICEP/*Keck*, WMAP and *Planck* maps. We compare the observed bandpower values for $20 < \ell < 330$ (9 bandpowers per spectrum) to an eight parameter model of lensed- Λ CDM+ r +dust+synchrotron+noise and explore the parameter space using COSMOMC [28] (which implements a Markov chain Monte Carlo). As in our previous analyses the bandpower covariance matrix is derived from 499 simulations of signal and noise, explicitly setting to zero terms such as the covariance of signal-only bandpowers with noise-only bandpowers or covariance of BICEP/*Keck* noise bandpowers with WMAP/*Planck* noise bandpowers (see Appendix H of BK15 and Appendix B of Ref. [29] for details). We deal with the differing sky coverage of the BICEP3 and BICEP2/*Keck* maps as described in Appendix D. The tensor/scalar power ratio

r is evaluated at a pivot scale of 0.05 Mpc^{-1} , and we fix the tensor spectral index $n_t = 0$. A COSMOMC module containing the data and model is available for download at <http://bicepkeck.org>. The following paragraphs briefly summarize the foreground model.

We include dust with amplitude $A_{d,353}$ evaluated at 353 GHz and $\ell = 80$. The frequency spectral behavior is taken as a modified black body spectrum with $T_d = 19.6 \text{ K}$ and frequency spectral index β_d . In a significant change from the baseline analysis choices of BK15, we remove the prior on the dust frequency spectral index which was previously applied based on *Planck* data in other regions of sky—with the improvement in the *Keck* 220 GHz sensitivity this prior is no longer needed. The spatial power spectrum is taken as a power law $\mathcal{D}_\ell \propto \ell^{\alpha_d}$ marginalizing uniformly over the (generous) range $-1 < \alpha_d < 0$ (where $\mathcal{D}_\ell \equiv \ell(\ell+1)C_\ell/2\pi$). *Planck* analysis consistently finds approximate power law behavior of both the EE and BB dust spectra with exponents ≈ -0.4 [24, 25].

We include synchrotron with amplitude $A_{\text{sync},23}$ evaluated at 23 GHz (the lowest WMAP band) and $\ell = 80$, assuming a simple power law for the frequency spectral behavior $A_{\text{sync}} \propto \nu^{\beta_s}$, and using a Gaussian prior $\beta_s = -3.1 \pm 0.3$ taken from the analysis of WMAP 23 and 33 GHz data in Ref. [30]. We note that analysis of 2.3 GHz data from S-PASS in conjunction with WMAP and *Planck* finds $\beta_s = -3.2$ with no detected trends with galactic latitude or angular scale [31], and that Ref. [32] analyzed the S-PASS and WMAP 23 GHz data and found $\beta_s = -3.22 \pm 0.06$ in the BICEP2 sky patch. The spatial power spectrum is taken as a power law $\mathcal{D}_\ell \propto \ell^{\alpha_s}$ marginalizing over the range $-1 < \alpha_s < 0$ [33]. Ref. [31] finds a value at the bottom end of this range (≈ -1) from the S-PASS data for BB at high galactic latitude.

Finally we include sync/dust correlation parameter ϵ (called ρ in some other papers [25, 31, 34]). As in BK15 we marginalize over the full possible range $-1 < \epsilon < 1$.

We hold the lensing B -mode spectrum fixed at that predicted for the *Planck* 2018 cosmological parameters [2, Table 2]. Results of our baseline analysis are shown in Fig. 4 and yield the following statistics: $r_{0.05} = 0.014_{-0.011}^{+0.010}$ ($r_{0.05} < 0.036$ at 95% confidence), $A_{d,353} = 4.4_{-0.7}^{+0.8} \mu\text{K}^2$, $A_{\text{sync},23} < 1.4 \mu\text{K}^2$ at 95% confidence, and $\beta_d = 1.49_{-0.12}^{+0.13}$. For r , the zero-to-peak likelihood ratio is 0.46. Taking $\frac{1}{2}(1 - f(-2 \log L_0/L_{\text{peak}}))$, where f is the χ^2 CDF (for one degree of freedom), we estimate that the probability to get a likelihood ratio smaller than this is 11% if, in fact, $r = 0$. As compared to the previous BK15 analysis, the likelihood curve for r tightens considerably with the peak position shifting down slightly, and the A_d curve tightens slightly. In addition the A_{sync} curve now peaks at zero—the weak evidence for synchrotron we saw in BK15 is no longer present. (Using the S-PASS data [31] we estimate that the expectation




FIG. 1. E -mode (left column) and B -mode (right column) maps at 95, 150 and 220 GHz in CMB units, and filtered to degree angular scales ($50 < \ell < 120$). Note the differing color ranges left and right. The E maps are dominated by Λ CDM signal, and hence are highly correlated across all three bands. The 95 GHz B map is approximately equal parts lensed- Λ CDM signal and noise. At 150 and 220 GHz the B maps are dominated by polarized dust emission.

is $A_{\text{sync},23} \approx 0.4 \mu\text{K}^2$ in the BICEP/Keck field, which is consistent with our A_{sync} likelihood curve.) In the BK15 analysis the constraint on β_d was prior dominated, but for BK18 we see that the data is able to constrain this parameter almost as well as the prior previously did. Interestingly the peak value selected is very close to the mean value from *Planck* 2018 analysis of larger regions of sky $\beta_d = 1.53$ [25].

The maximum likelihood model has parameters $r_{0.05} = 0.011$, $A_{d,353} = 4.4 \mu\text{K}^2$, $A_{\text{sync},23} = 0.6 \mu\text{K}^2$, $\beta_d = 1.5$, $\beta_s = -3.0$, $\alpha_d = -0.66$, $\alpha_s = 0.00$, and $\epsilon = -0.11$. This model is an acceptable fit to the data with the PTE of the observed value of χ^2 being 0.94. Thus, while the dust spectrum might in general be expected to exhibit fluctu-

ations about power law spatial spectral behavior greater than that expected for a Gaussian random field, for the present the model continues to be an adequate description of the data—see Appendix D for further details.

In Appendix E we explore variation and validation of the likelihood. In Appendix E2 we vary the baseline analysis choices and data selection, finding that these do not significantly alter the results, and that the data do not prefer allowing decorrelation of the dust pattern in the model. We also find that the value of A_d is very similar when evaluated over the larger BICEP3 sky coverage region and the smaller BICEP2/Keck sky region. Freeing the amplitude of the lensing power we obtain $A_L^{\text{BB}} = 1.03^{+0.08}_{-0.09}$, and the r constraint hardly changes.




FIG. 2. EE (green) and BB (blue) auto- and cross-spectra calculated using the BICEP3 95 GHz map, the BICEP2/*Keck* 150 GHz map, the *Keck* 220 GHz map, and the *Planck* 353 GHz map (with the auto-spectra in darker colors). The BICEP/*Keck* maps use all data taken up to and including the 2018 observing season—we refer to these as BK18. The black lines show the model expectation values for lensed- Λ CDM, while the red lines show the expectation values of a baseline lensed- Λ CDM+dust model from our previous BK15 analysis ($r = 0$, $A_{d,353} = 4.7 \mu\text{K}^2$, $\beta_d = 1.6$, $\alpha_d = -0.4$). Note that the model shown was fit to BB only and did not use the BICEP3 95 GHz points shown (which are entirely new). The agreement with the spectra involving 95 GHz and all the EE spectra (under the assumption that $EE/BB = 2$ for dust) is therefore a validation of the model.

In Appendix E3 we verify that the likelihood analysis is unbiased, and in Appendix E4 we explore a suite of alternate foreground models. As part of our standard data reduction we “deproject” leading order temperature to polarization leakage [9, 35]—in Appendix F we quantify possible residual leakage and some other possible systematics.

Fig. 5 shows the constraints in the r vs. n_s plane for the *Planck* 2018 baseline analysis [2] and when adding in BK18 & BAO. The BK18 data shrinks the contours in the vertical (r) direction while the BAO data shrinks

the contours in the horizontal (n_s) direction and shifts the centroid slightly to the right. The $\phi^{2/3}$ model now lies outside the 95% contour as does the band of natural inflation models.

Conclusions.—The BKP analysis yielded a 95% confidence constraint $r_{0.05} < 0.12$, which BK14 improved to $r_{0.05} < 0.09$, and BK15 improved to $r_{0.05} < 0.07$. The BK18 result described in this letter, $r_{0.05} < 0.036$, represents a fractional improvement equivalent to the two previous steps combined. The BK18 simulations have a median 95% upper limit of $r_{0.05} < 0.019$.




FIG. 3. *Upper:* The noise spectra of the BICEP3 95 GHz map (red), the BICEP2/Keck 150 GHz map (green) and the Keck 220 GHz maps (blue). The spectra are shown after correction for the filtering of signal which occurs due to the beam roll-off, timestream filtering, and B -mode purification. (Note that no ℓ^2 scaling is applied.) *Lower:* The effective sky fraction as calculated from the ratio of the mean noise realization bandpowers to their fluctuation $f_{\text{sky}}(\ell) = \frac{1}{2\ell\Delta\ell} \left(\frac{\sqrt{2N_b}}{\sigma(N_b)} \right)^2$, i.e. the observed number of B -mode degrees of freedom divided by the nominal full-sky number. The turn-down at low ℓ is due to mode loss to the timestream filtering and matrix purification.

The distributions of maximum likelihood r values in simulations where the true value of r is zero gave $\sigma(r_{0.05}) = 0.020$ for BK15 which is reduced to $\sigma(r_{0.05}) = 0.009$ for BK18 (see Appendix E 3 for details). Such simulations can also be used to investigate the degree to which the analysis is limited by foregrounds and lensing. Running the baseline BK18 analysis on simulations which contain no lensing B -modes gives $\sigma(r_{0.05}) = 0.004$, while running without foreground parameters on simulations which contain no dust gives $\sigma(r_{0.05}) = 0.007$. Running without foreground parameters on simulations which contain neither lensing or dust gives $\sigma(r_{0.05}) = 0.002$.

Fig. 6 shows the BK18 noise uncertainties in the $\ell \approx 80$ bandpowers as compared to the signal levels. The signal-to-noise on polarized dust emission of our 220 GHz band is now considerably higher than that of the *Planck* 353 GHz band—i.e. the 220×220 noise point is much further below the dust band than the $P353 \times P353$ point. Additional BICEP3 data taken during 2019–2021 will reduce the noise by a factor greater than 2 and $\sqrt{2}$ for 95×95 and $95 \times W23$ respectively, and we have also recorded additional data at 220 and 270 GHz.

Fig. 7 shows the estimated CMB-only component of the BK18 B -mode bandpowers versus measurements from other experiments. See Appendix D for a descrip-

tion of how the CMB-only power spectrum estimate is calculated.

Figure 2 shows that the BK18 data is consistent with Λ CDM plus a remarkably simple dust only foreground model. Nevertheless as we move forward to even higher levels of sensitivity dust decorrelation, and foreground complexity more generally, will remain a serious concern. In addition, we are already in the regime where the sample variance of the lensing component dominates $\sigma(r)$. However, the lensing B -modes can be spatially separated from a primordial component and in this regard we have recently demonstrated a path forward by adding a “lensing template” derived from SPTPOL and *Planck* data to the BK14 analysis, resulting in an improved constraint on r [50].

The *Keck Array* mount has now been replaced by a larger, more capable machine and we are in the process of upgrading to a new system we call BICEP Array [51, 52]. A BICEP3 class receiver is now operating in the 30/40 GHz band and in the coming years additional receivers will be installed at 95, 150 and 220/270 GHz. The system is projected to reach $\sigma(r) \sim 0.003$ within five years with delensing in conjunction with SPT3G.

The BICEP/Keck projects have been made possible through a series of grants from the National Science Foundation including 0742818, 0742592, 1044978, 1110087, 1145172, 1145143, 1145248, 1639040, 1638957, 1638978, & 1638970, and by the Keck Foundation. The development of antenna-coupled detector technology was supported by the JPL Research and Technology Development Fund, and by NASA Grants 06-ARPA206-0040, 10-SAT10-0017, 12-SAT12-0031, 14-SAT14-0009 & 16-SAT16-0002. The development and testing of focal planes were supported by the Gordon and Betty Moore Foundation at Caltech. Readout electronics were supported by a Canada Foundation for Innovation grant to UBC. Support for quasi-optical filtering was provided by UK STFC grant ST/N000706/1. The computations in this paper were run on the Odyssey/Cannon cluster supported by the FAS Science Division Research Computing Group at Harvard University. The analysis effort at Stanford and SLAC is partially supported by the U.S. DOE Office of Science. We thank the staff of the U.S. Antarctic Program and in particular the South Pole Station without whose help this research would not have been possible. Most special thanks go to our heroic winter-overs Robert Schwarz, Steffen Richter, Sam Harrison, Grantland Hall and Hans Boenish. We thank all those who have contributed past efforts to the BICEP/Keck series of experiments, including the BICEP1 team. We also thank the *Planck* and WMAP teams for the use of their data, and are grateful to the *Planck* team for helpful discussions.




FIG. 4. Results of a multicomponent multi-spectral likelihood analysis of BICEP/*Keck*+WMAP/*Planck* data. The red faint curves are the baseline result from the previous BK15 paper (the black curves from Fig. 4 of that paper). The bold black curves are the new baseline BK18 results, adding a large amount of additional data at 95 and 220 GHz taken by BICEP3 and *Keck Array* during the 2016–2018 observing seasons. The upper limit on the tensor-to-scalar ratio tightens to $r_{0.05} < 0.036$ at 95% confidence. The parameters A_d and A_{sync} are the amplitudes of the dust and synchrotron B -mode power spectra, where β and α are the respective frequency and spatial spectral indices. The correlation coefficient between the dust and synchrotron patterns is ϵ . In the β , α and ϵ panels the dashed lines show the priors placed on these parameters (either Gaussian or uniform). Note that the Gaussian prior on β_d has been removed going from BK15 to BK18.




FIG. 5. Constraints in the r vs. n_s plane for the *Planck* 2018 baseline analysis, and when also adding BICEP/*Keck* data through the end of the 2018 season plus BAO data to improve the constraint on n_s . The constraint on r tightens from $r_{0.05} < 0.11$ to $r_{0.05} < 0.035$. This figure is adapted from Fig. 28 of Ref. [2] with the green contours being identical. Some additional inflationary models are added from Fig. 8 of Ref. [36] with the purple region being natural inflation.




FIG. 6. Expectation values and noise uncertainties of the $\ell \sim 80$ *BB* bandpower in the BICEP/*Keck* field. The solid and dashed black lines show the expected signal power of lensed- Λ CDM and $r_{0.05} = 0.01$. Since CMB units are used, the levels corresponding to these are flat with frequency. The blue bands show the 1 and 2σ ranges of dust, and the red shaded region shows the 95% upper limit on synchrotron in the baseline analysis including the uncertainties in the amplitude and frequency spectral index parameters ($A_{\text{sync},23}, \beta_s$ and $A_{d,353}, \beta_d$). The BICEP/*Keck* auto-spectrum noise uncertainties are shown as large blue circles, and the noise uncertainties of the used WMAP/*Planck* single-frequency spectra evaluated in the BICEP/*Keck* field are shown in black. The blue crosses show the noise uncertainty of selected cross-spectra, and are plotted at horizontal positions such that they can be compared vertically with the dust and sync curves.




FIG. 7. Summary of CMB B -mode polarization upper limits [20, 26, 37–46] and detections [47–49]. Theoretical predictions are shown for the lensing B -modes (solid red) which peak at arcminute scales (multipole $\ell \sim 1000$), and for gravitational wave B -modes (dashed red) for two values of r peaking at degree scales ($\ell \sim 80$). The BK18 data are shown after removing Galactic foregrounds.

-
- * pryke@physics.umn.edu
- [1] A. A. Penzias and R. W. Wilson, *Astrophys. J.* **142**, 419 (1965).
 - [2] Planck Collaboration 2018 VI, *Astron. Astrophys.* **641**, A6 (2020), arXiv:1807.06209.
 - [3] M. Kamionkowski and E. D. Kovetz, *Annual Review of Astronomy and Astrophysics* **54**, 227 (2016), arXiv:1510.06042.
 - [4] CMB-S4 Collaboration, *ArXiv e-prints* (2016), arXiv:1610.02743.
 - [5] M. Zaldarriaga and U. Seljak, *Phys. Rev. D* **58**, 023003 (1998), astro-ph/9803150.
 - [6] U. Seljak, *Astrophys. J.* **482**, 6 (1997), astro-ph/9608131.
 - [7] M. Kamionkowski, A. Kosowsky, and A. Stebbins, *Phys. Rev. Lett.* **78**, 2058 (1997), astro-ph/9609132.
 - [8] U. Seljak and M. Zaldarriaga, *Phys. Rev. Lett.* **78**, 2054 (1997), astro-ph/9609169.
 - [9] BICEP2 Collaboration I, *Physical Review Letters* **112**, 241101 (2014), arXiv:1403.3985.
 - [10] BICEP2/Keck and Planck Collaborations, *Physical Review Letters* **114**, 101301 (2015), arXiv:1502.00612.
 - [11] Keck Array and BICEP2 Collaborations VI, *Physical Review Letters* **116**, 031302 (2016), arXiv:1510.09217.
 - [12] Keck Array and BICEP2 Collaborations X, *Phys. Rev. Lett.* **121**, 221301 (2018), arXiv:1810.05216.
 - [13] BICEP2 Collaboration II, *Astrophys. J.* **792**, 62 (2014), arXiv:1403.4302.
 - [14] Keck Array and BICEP2 Collaborations V, *Astrophys. J.* **811**, 126 (2015), arXiv:1502.00643.
 - [15] BICEP/Keck Collaboration XIV, In preparation (2021).
 - [16] BICEP2/Keck and Spider Collaborations, *Astrophys. J.* **812**, 176 (2015), arXiv:1502.00619.
 - [17] C.-L. Kuo, *Astrophys. J.* **848**, 64 (2017), arXiv:1707.08400.
 - [18] Keck Array and BICEP2 Collaborations VII, *Astrophys. J.* **825**, 66 (2016), arXiv:1603.05976.
 - [19] See http://lambda.gsfc.nasa.gov/product/map/dr5/m_products.cfm.
 - [20] C. L. Bennett, D. Larson, J. L. Weiland, N. Jarosik, G. Hinshaw, N. Odegard, K. M. Smith, R. S. Hill, B. Gold, M. Halpern, E. Komatsu, M. R. Nolta, L. Page, D. N. Spergel, E. Wollack, J. Dunkley, A. Kogut, M. Limon, S. S. Meyer, G. S. Tucker, and E. L. Wright, *Astrophys. J. Suppl. Ser.* **208**, 20 (2013), arXiv:1212.5225.
 - [21] See https://irsa.ipac.caltech.edu/data/Planck/release_3/ancillary-data/HFI_Products.html.
 - [22] Planck Collaboration Int. LVII, *Astron. Astrophys.* **643**, A42 (2020), arXiv:2007.04997.
 - [23] See Supplemental Material at <http://xxx.yyy> for appendixes, which include references [53–65].
 - [24] Planck Collaboration Int. XXX, *Astron. Astrophys.* **586**, A133 (2016), arXiv:1409.5738.
 - [25] Planck Collaboration 2018 XI, *ArXiv e-prints* (2018), arXiv:1801.04945.
 - [26] POLARBEAR Collaboration, *Astrophys. J.* **897**, 55 (2020), arXiv:1910.02608.
 - [27] S. Hamimeche and A. Lewis, *Phys. Rev. D* **77**, 103013 (2008), arXiv:0801.0554.
 - [28] A. Lewis and S. Bridle, *Phys. Rev. D* **66**, 103511 (2002), astro-ph/0205436.
 - [29] CMB-S4 Collaboration, submitted to *Astrophys. J.* (2020), arXiv:2008.12619.
 - [30] U. Fuskeland, I. K. Wehus, H. K. Eriksen, and S. K. Næss, *Astrophys. J.* **790**, 104 (2014), arXiv:1404.5323.
 - [31] N. Krachmalnicoff, E. Carretti, C. Baccigalupi, G. Bernardi, S. Brown, B. M. Gaensler, M. Haverkorn, M. Kesteven, F. Perrotta, S. Poppi, and L. Staveley-Smith, *Astron. Astrophys.* **618**, A166 (2018), arXiv:1802.01145.
 - [32] U. Fuskeland, K. J. Andersen, R. Aurlien, R. Banerji, M. Brilenkov, H. K. Eriksen, M. Galloway, E. Gjerløw, S. K. Næss, T. L. Svalheim, and I. K. Wehus, *Astron. Astrophys.* **646**, A69 (2021), arXiv:1909.05923.
 - [33] J. Dunkley, A. Amblard, C. Baccigalupi, M. Betoule, D. Chuss, A. Cooray, J. Delabrouille, C. Dickinson, G. Dobler, J. Dotson, H. K. Eriksen, D. Finkbeiner, D. Fixsen, P. Fosalba, A. Fraisse, C. Hirata, A. Kogut, J. Kristiansen, C. Lawrence, A. M. Magalhães, M. A. Miville-Deschenes, *et al.*, *AIP Conf. Proc.* **1141**, 222 (2009), arXiv:0811.3915.
 - [34] S. K. Choi and L. A. Page, *J. Cosmol. Astropart. Phys.* **12**, 020 (2015), arXiv:1509.05934.
 - [35] BICEP2 Collaboration III, *ArXiv e-prints* (2015), arXiv:1502.00608.
 - [36] Planck Collaboration 2018 X, *Astron. Astrophys.* **641**, A10 (2020), arXiv:1807.06211.
 - [37] E. M. Leitch, J. M. Kovac, N. W. Halverson, J. E. Carlstrom, C. Pryke, and M. W. E. Smith, *Astrophys. J.* **624**, 10 (2005), astro-ph/0409357.
 - [38] T. E. Montroy, P. A. R. Ade, J. J. Bock, J. R. Bond, J. Borrill, A. Boscaleri, P. Cabella, C. R. Contaldi, B. P. Crill, P. de Bernardis, G. De Gasperis, A. de Oliveira-Costa, G. De Troia, G. di Stefano, E. Hivon, A. H. Jaffe, T. S. Kisner, W. C. Jones, A. E. Lange, S. Masi, *et al.*, *Astrophys. J.* **647**, 813 (2006), astro-ph/0507514.
 - [39] J. L. Sievers, C. Achermann, J. R. Bond, L. Bronfman, R. Bustos, C. R. Contaldi, C. Dickinson, P. G. Ferreira, M. E. Jones, A. M. Lewis, B. S. Mason, J. May, S. T. Myers, N. Oyarce, S. Padin, T. J. Pearson, M. Pospieszalski, A. C. S. Readhead, R. Reeves, A. C. Taylor, and S. Torres, *Astrophys. J.* **660**, 976 (2007), astro-ph/0509203.
 - [40] CAPMAP Collaboration, *Astrophys. J.* **684**, 771 (2008), arXiv:0802.0888.
 - [41] QUaD Collaboration, *Astrophys. J.* **705**, 978 (2009), arXiv:0906.1003.
 - [42] QUIET Collaboration, *Astrophys. J.* **760**, 145 (2012), arXiv:1207.5034.
 - [43] BICEP1 Collaboration, *Astrophys. J.* **783**, 67 (2014), arXiv:1310.1422.
 - [44] A. Kusaka, J. Appel, T. Essinger-Hileman, J. A. Beall, L. E. Campusano, H.-M. Cho, S. K. Choi, K. Crowley, J. W. Fowler, P. Gallardo, M. Hasselfield, G. Hilton, S.-P. P. Ho, K. Irwin, N. Jarosik, M. D. Niemack, G. W. Nixon, M. Nolta, L. A. Page, Jr., G. A. Palma, L. Parker, S. Raghunathan, C. D. Reintsema, J. Sievers, S. M. Simon, S. T. Staggs, K. Visnjic, and K.-W. Yoon, *J. Cosmol. Astropart. Phys.* **9**, 005 (2018), arXiv:1801.01218.
 - [45] M. Tristram, A. J. Banday, K. M. Górski, R. Keskitalo, C. R. Lawrence, K. J. Andersen, R. B. Barreiro, J. Borrill, H. K. Eriksen, R. Fernandez-Cobos, T. S. Kisner, E. Martínez-González, B. Partridge, D. Scott, T. L. Svalheim, H. Thommesen, and I. K. Wehus, *Astron. Astrophys.* **647**, A128 (2021), arXiv:2010.01139.
 - [46] SPIDER Collaboration, submitted to *Astrophys. J.*

- (2021), arXiv:2103.13334.
- [47] J. T. Sayre, C. L. Reichardt, J. W. Henning, P. A. R. Ade, *et al.*, Phys. Rev. D **101**, 122003 (2020), arXiv:1910.05748.
- [48] S. K. Choi, M. Hasselfield, S.-P. P. Ho, B. Koopman, M. Lungu, M. H. Abitbol, G. E. Addison, P. A. R. Ade, *et al.*, J. Cosmol. Astropart. Phys. **2020**, 045 (2020), arXiv:2007.07289.
- [49] POLARBEAR Collaboration, Astrophys. J. **848**, 121 (2017), arXiv:1705.02907.
- [50] BICEP2/Keck and SPTPOL Collaborations, Phys. Rev. D **103**, 022004 (2021), arXiv:2011.08163.
- [51] H. Hui and the BICEP/Keck Collaboration, in *SPIE Conference Series*, Vol. 10708 (2018) p. 1070807, arXiv:1808.00568.
- [52] L. Moncelsi and the BICEP/Keck Collaboration, in *SPIE Conference Series*, Vol. 11453 (2020) p. 1145314, arXiv:2012.04047.
- [53] Keck Array and BICEP2 Collaborations XI, Astrophys. J. **884**, 114 (2019), arXiv:1904.01640.
- [54] J. Willmert, *Constraining Inflationary B-modes with the BICEP/Keck Array Telescopes*, Ph.D. thesis, University of Minnesota (2019).
- [55] Planck Collaboration Int. XXII, Astron. Astrophys. **576**, A107 (2015), arXiv:1405.0874.
- [56] V. Pelgrims, S. E. Clark, B. S. Hensley, G. V. Panopoulou, V. Pavlidou, K. Tassis, H. K. Eriksen, and I. K. Wehus, Astron. Astrophys. **647**, A16 (2021), arXiv:2101.09291.
- [57] G. V. Panopoulou and D. Lenz, Astrophys. J. **902**, 120 (2020), arXiv:2004.00647.
- [58] Planck Collaboration Int. L, Astron. Astrophys. **599**, A51 (2017), arXiv:1606.07335.
- [59] B. Thorne, J. Dunkley, D. Alonso, and S. Næss, Mon. Not. R. Astron. Soc. **469**, 2821 (2017), arXiv:1608.02841.
- [60] A. G. Kritsuk, S. D. Ustyugov, and M. L. Norman, New Journal of Physics **19**, 065003 (2017), arXiv:1705.01912.
- [61] A. G. Kritsuk, R. Flauger, and S. D. Ustyugov, Phys. Rev. Lett. **121**, 021104 (2018), arXiv:1711.11108.
- [62] G. Martínez-Solaèche, A. Karakci, and J. Delabrouille, Mon. Not. R. Astron. Soc. **476**, 1310 (2018), arXiv:1706.04162.
- [63] F. Vansyngel, F. Boulanger, T. Ghosh, B. Wandelt, J. Aumont, A. Bracco, F. Levrier, P. G. Martin, and L. Montier, Astron. Astrophys. **603**, A62 (2017), arXiv:1611.02577.
- [64] Keck Array and BICEP2 Collaborations IV, Astrophys. J. **806**, 206 (2015), arXiv:1502.00596.
- [65] R. A. Battye, I. W. A. Browne, M. W. Peel, N. J. Jackson, and C. Dickinson, Mon. Not. R. Astron. Soc. **413**, 132 (2011), arXiv:1003.5846.