

This is the accepted manuscript made available via CHORUS. The article has been published as:

## Neural Network Representation of Tensor Network and Chiral States

Yichen Huang (黄一晨) and Joel E. Moore

Phys. Rev. Lett. **127**, 170601 — Published 18 October 2021

DOI: [10.1103/PhysRevLett.127.170601](https://doi.org/10.1103/PhysRevLett.127.170601)

# Neural network representation of tensor network and chiral states

Yichen Huang (黄溢辰)\*

*Institute for Quantum Information and Matter, California Institute of Technology, Pasadena, California 91125, USA<sup>†</sup>*

Joel E. Moore<sup>‡</sup>

*Department of Physics, University of California, Berkeley, California 94720, USA  
Materials Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA*

(Dated: July 20, 2021)

We study the representational power of Boltzmann machines (a type of neural network) in quantum many-body systems. We prove that any (local) tensor network state has a (local) neural network representation. The construction is almost optimal in the sense that the number of parameters in the neural network representation is almost linear in the number of nonzero parameters in the tensor network representation. Despite the difficulty of representing (gapped) chiral topological states with local tensor networks, we construct a quasi-local neural network representation for a chiral  $p$ -wave superconductor. These results demonstrate the power of Boltzmann machines.

A generic state in quantum many-body systems is classically intractable for the reason that the dimension of the Hilbert space grows exponentially with the system size. However, physically relevant states are often non-generic in the sense of having structures, making use of which we may overcome the curse of dimensionality. Traditionally, tensor networks are used to characterize such structures and efficiently represent states in classical simulations [1]. Recently, Carleo and Troyer [2] proposed neural networks as an (alternative) ansatz for quantum many-body states. Benchmark calculations suggest that this is a promising approach.

Besides numerical experiments, it is also important to explain the working principle of neural network methods. One step in this direction is to characterize the representational power of neural networks. Here we specialize to Boltzmann machines [3], and our main contributions are:

- We prove that any (local) tensor network state can be converted into a (local) neural network without significantly increasing the number of parameters.
- Despite the difficulty of representing (gapped) chiral topological states with local tensor networks [4], we construct a quasi-local neural network representation for a chiral  $p$ -wave superconductor.

The first result states that the representational power of neural networks is at least not weaker than that of tensor networks. The second gives a physically relevant example where neural networks may go beyond tensor networks. In combination, these results provide complementary evidence that neural networks are a promising ansatz.

*Boltzmann machines.*—We provide a minimum background for those people with no prior knowledge of Boltzmann machines. The goal is to motivate the definition of neural network states, rather than a general-purpose introduction from the perspective of machine learning.

Formally, a Boltzmann machine is a type of stochastic recurrent neural network. In the language of physicists, it

is a classical Ising model on a weighted undirected graph. Each vertex (also known as a unit or a neuron) of the graph carries a classical Ising variable  $s_j = \pm 1$  and a local field  $h_j \in \mathbb{R}$ , where  $j$  is the index of the vertex. For a reason that will soon be clear, the set of vertices is divided into the disjoint union of two subsets  $V$  and  $H$  so that  $|V| + |H|$  is the total number of units. Vertices in  $V$  are called visible units, and those in  $H$  are called hidden units. For notational simplicity, we assume that visible units have small indices  $1, 2, \dots, |V|$ , and hidden units have large indices  $|V| + 1, |V| + 2, \dots, |V| + |H|$ . Each edge of the graph carries a weight  $w_{jk} \in \mathbb{R}$  that describes the interaction between  $s_j$  and  $s_k$ . The energy of a configuration is given by

$$E(\{s_j\}) = \sum_j h_j s_j + \sum_{j,k} w_{jk} s_j s_k. \quad (1)$$

A restricted Boltzmann machine [5] is a Boltzmann machine on a bipartite graph, i.e.,  $w_{jk} \neq 0$  only if the edge  $(j, k)$  connects a visible unit with a hidden unit.

At thermal equilibrium, the configurations follow the Boltzmann distribution. Without loss of generality, we fix the temperature  $T = 1$ . Let  $Z = \sum_{\{s_j\} \in \{\pm 1\}^{\times(|V|+|H|)}} e^{-E(\{s_j\})}$  be the partition function. The probability of each configuration  $\{s_j\}$  is given by  $e^{-E(\{s_j\})}/Z$ . Furthermore, the probability of each configuration  $\{s_{j \leq |V|}\}$  of visible units is the marginal probability obtained by summing over hidden units:

$$P\{s_{j \leq |V|}\} = \frac{1}{Z} \sum_{\{s_{j > |V|}\} \in \{\pm 1\}^{\times |H|}} e^{-E(\{s_j\})}. \quad (2)$$

The process of training a Boltzmann machine is specified as follows. The input is a (normalized) probability distribution  $Q$  over the configurations of the visible units. (Here we assume, for simplicity, that  $Q$  is given. In practice,  $Q$  may not be explicitly given but we are allowed to sample from  $Q$ .) The goal is to adjust the weights

$w_{jk}$  such that the probability distribution  $P$  (2) best approximates (as measured by Kullback-Leibler divergence or some other distance function) the desired distribution  $Q$ . This is a variational minimization problem. We minimize a given objective function of  $P$  with respect to the ansatz (2), in which the weights are variational parameters. Note that Monte Carlo techniques are usually used in training a Boltzmann machine.

An interesting question is whether an arbitrary  $Q$  can be exactly represented by a Boltzmann machine. Due to the strict positivity of exponential functions, it is easy to see that the answer is no if the probability of some configuration is zero. Nevertheless,

**Theorem 1** (Le Roux and Bengio [6]). *Any probability distribution  $Q$  can be arbitrarily well approximated by an restricted Boltzmann machine, provided that the number  $|H|$  of hidden units is the number of configurations with nonzero probability. In general,  $|H| \leq 2^{|V|}$ .*

This result can be slightly improved [7]. Indeed, a simple counting argument suggests that an exponential number of hidden units is necessary in general. As a function from  $\{\pm 1\}^{\times |V|}$  to  $[0, 1]$  with one constraint (normalization), the probability distribution  $Q$  has  $2^{|V|} - 1$  degrees of freedom. Therefore, a sufficiently well approximation of  $Q$  requires an exponential number of bits.

*Neural network states.*—Carleo and Troyer [2] developed a minimum extension of Boltzmann machines to the quantum world. In this extension, each vertex still carries a classical spin (bit). This is very different from another extension [8], in which each vertex carries a quantum spin (qubit). It should be clear that the latter extension is more “quantum,” for it is operated by quantum computers. In the former extension, we use classical computers to simulate quantum many-body systems.

Expanded in the computational basis  $\{\pm 1\}^{\times |V|}$ , a quantum state  $|\psi\rangle$  of  $|V|$  qubits can be viewed as a function from  $\{\pm 1\}^{\times |V|}$  to  $\bar{B}(0, 1)$  with one constraint (normalization), where  $\bar{B}(0, 1)$  denotes the closed region  $|z| \leq 1$  in the complex plane. Recall that a probability distribution  $Q$  is characterized by the probability of each configuration. In comparison, a state  $|\psi\rangle$  is characterized by the probability amplitude of each configuration. This analog between classical probability distributions and quantum states motivates the following ansatz dubbed neural network states.

Consider a graph as before. The visible units correspond to physical qubits, and hidden units are auxiliary degrees of freedom (to be summed over). The local field  $h_j \in \mathbb{C}$  at each vertex and the weight  $w_{jk} \in \mathbb{C}$  carried by each edge are promoted to complex numbers because probability amplitudes are generally complex. An (unnormalized) neural network state based on a Boltzmann machine is given by

$$|\psi\rangle = \sum_{\{s_1, s_2, \dots, s_{|V|+|H|}\} \in \{\pm 1\}^{\times (|V|+|H|)}} e^{-\sum_j h_j s_j - \sum_{j,k} w_{jk} s_j s_k} |\{s_1, s_2, \dots, s_{|V|}\}\rangle. \quad (3)$$

Note that summing over hidden units is very different from taking partial trace, for the latter usually results in a mixed state. Similarly, a neural network state based on a restricted Boltzmann machine is given by Eq. (3) on a bipartite graph as described previously.

A very minor modification of the proof of Theorem 1 leads to

**Corollary 1.** *Any  $N$ -qubit quantum state  $|\psi\rangle$  can be arbitrarily well approximated by a neural network state based on a restricted Boltzmann machine with  $w_{jk} \in \mathbb{R}, h_j \in \mathbb{C}$ , provided that the number  $|H|$  of hidden units is the number of configurations with nonzero probability amplitude. In general,  $|H| \leq 2^N$ .*

The formalism of neural network states developed thus far applies to general quantum many-body systems. We now consider the situation that the qubits are arranged on a lattice. The lattice allows us to introduce the notion of locality [9] (with respect to the shortest path metric), which underlies almost all successful methods for simulating quantum lattice systems. Hence, it is desirable to

incorporate locality into the neural network. To do this, we define a position for each unit. Let each site of the lattice carry a visible unit and some hidden units. We require that  $w_{jk} \neq 0$  only if the units  $j$  and  $k$  are close to each other. As an example, Deng et al. [10] showed that the ground state of the toric code in two dimensions can be exactly represented as a local neural network state based on a restricted Boltzmann machine.

Using variational quantum Monte Carlo techniques (“quantum Monte Carlo” is not a quantum algorithm; rather, it is just a classical Monte Carlo algorithm applied to quantum systems), Carleo and Troyer [2] performed practical calculations for quantum lattice systems in one and two spatial dimensions. For the models they studied, they observed that variational approaches based on neural networks are at least as good as those based on tensor networks. It seems worthwhile to devote more study to the practical performance of neural network states.

*Tensor network states.*—We provide a very brief introduction to tensor network states following the presenta-

tion in Ref. [11], Subsection 6.3 or [12], Subsection 2.3. Then, it will be clear almost immediately that any (local) tensor network state has a (local) neural network representation.

An  $l$ -dimensional tensor  $T$  is a multivariable function  $T : \{1, 2, \dots, d_1\} \times \{1, 2, \dots, d_2\} \times \dots \times \{1, 2, \dots, d_l\} \rightarrow \mathbb{C}$ , and  $D = \max_j d_j$  is called the bond dimension. It is easy to see that  $T$  can be reshaped to an  $l'$ -dimensional ( $l' = \sum_j \lceil \log_2 d_j \rceil$  with  $\lceil \cdot \rceil$  the ceiling function) tensor  $T'$  with bond dimension 2 by representing every input variable in binary. Moreover, there is a straightforward way to identify  $T'$  with an  $l'$ -qubit unnormalized state expanded in the computational basis:

$$|\psi\rangle = \sum_{\{s_j\} \in \{\pm 1\}^{l'}} T' \left( \frac{s_1 + 3}{2}, \frac{s_2 + 3}{2}, \dots, \frac{s_n + 3}{2} \right) |\{s_j\}\rangle. \quad (4)$$

Thus, a neural network state with  $|V|$  visible units is a  $|V|$ -dimensional tensor with bond dimension 2. Indeed,

---


$$\sum_{\substack{\{s_1, s_2, \dots, s_{|V|+|H|}\} \in \{\pm 1\}^{\times(|V|+|H|)} \\ \{s'_1, s'_2, \dots, s'_{|V'|+|H'|}\} \in \{\pm 1\}^{\times(|V'|+|H'|)} \\ \text{s. t. } s_1 = s'_1, s_2 = s'_2, \dots, s_c = s'_c}} e^{-\sum_j h_j s_j - \sum_{j,k} w_{jk} s_j s_k - \sum_j h'_j s'_j - \sum_{j,k} w'_{jk} s'_j s'_k} |\{s_{c+1}, s_{c+2}, \dots, s_{|V|}\}\rangle \otimes |\{s'_{c+1}, s'_{c+2}, \dots, s'_{|V'|}\}\rangle. \quad (5)$$


---

Note that only visible units are allowed to be contracted on, and such visible units become hidden after the contraction. Thus, the Boltzmann machine after contraction has  $|V| + |V'| - 2c$  visible and  $|H| + |H'| + c$  hidden units.

A multi-qubit state  $|\psi\rangle$  is a tensor network state if its tensor representation  $T'$  (4) can be obtained by contracting a network of  $C$ -dimensional tensors, where  $C$  is a small absolute constant. The bond dimension  $D$  of a tensor network is defined as the maximum bond dimension of each constituting tensors in the network.

**Theorem 2.** *Any tensor network state  $|\psi\rangle$  can be arbitrarily well approximated by a neural network state based on a Boltzmann machine with  $w_{jk} \in \mathbb{R}, h_j \in \mathbb{C}$ , provided that the number  $|H|$  of hidden units is sufficiently large. Let  $\emptyset(T_j)$  be the number of nonzero elements in a constituting tensor  $T_j$ , and  $\#(T_j)$  be the number of elements. It suffices that*

$$|H| = \sum_{j=1}^n (\emptyset(T_j) + \log_2 \#(T_j) + O(1)), \quad (6)$$

where  $n$  is the number of constituting tensors. Furthermore, the number of parameters in the neural network representation is upper bounded by

$$\sum_j \emptyset(T_j) \log_2 \#(T_j) + \text{less significant terms.} \quad (7)$$

each visible unit corresponds to an input variable to the tensor. As a restatement of Corollary 1,

**Corollary 2.** *Any  $l$ -dimensional tensor  $T$  can be arbitrarily well approximated by a restricted Boltzmann machine with  $w_{jk} \in \mathbb{R}, h_j \in \mathbb{C}$ , provided that the number  $|H|$  of hidden units is the number of nonzero elements in  $T$ . In general,  $|H| \leq \prod_{j=1}^l d_j$ .*

Informally, tensor contraction is defined as follows. Suppose we have two 3-dimensional tensors  $T_1(j_1, j_2, j_3), T_2(j'_1, j'_2, j'_3)$  of shapes  $d_1 \times d_2 \times d_3, d'_1 \times d'_2 \times d'_3$ , respectively. Their contraction on the middle indices is a 4-dimensional tensor  $\sum_{j=1}^{d_2} T_1(j_1, j, j_3) T_2(j'_1, j, j'_3)$ .

Similarly, suppose we have two Boltzmann machines with units  $\{s_j\} = V \cup H, \{s'_j\} = V' \cup H'$ , weights  $w_{jk}, w'_{jk}$ , and local fields  $h_j, h'_j$ , respectively. Their contraction on the first  $c$  visible units is defined as identifying  $s_j$  with  $s'_j$  and then summing over  $s_j$  for  $j = 1, 2, \dots, c$ :

---

*The neural network is local (translationally invariant) if the tensor network is local (translationally invariant).*

*Proof.* We first represent each  $T_j$  with a restricted Boltzmann machine as in Corollary 2, and then contract the restricted Boltzmann machines in the same way as  $T_j$ 's are contracted. Note that the Boltzmann machine after contraction is generically not restricted. The first term on the right-hand side in Eq. (6) is the total number of hidden units in all constituting restricted Boltzmann machines, and the other terms are responsible for the production of hidden units in the contraction process. Equation (7) is the total number of parameters in all constituting restricted Boltzmann machines, and the contraction process does not introduce any new parameters. It is obvious that the construction is locality- and translational-invariance-preserving.  $\square$

This result is almost optimal in the sense that the number of parameters in the neural network representation is at most a logarithmic (in the bond dimension) multiple of the number of nonzero parameters in the tensor network representation. In cases that the tensors are sparse (possibly due to the presence of symmetries), this conversion to neural network states has the desirable property of automatically compressing the representation.

As an example, we specialize Theorem 2 to matrix product states [13].

**Corollary 3.** *To leading order, any  $N$ -qubit matrix product state with bond dimension  $D$  has a neural network representation with  $2ND^2$  hidden units and  $4ND^2 \log_2 D$  parameters.*

*Chiral topological states.*—It seems difficult to obtain a local tensor network representation for gapped chiral topological states. Early attempts did not impose locality [14] or just target at expectation values of local observables rather than the wave function [15]. Recent progress [4, 16] shows that local tensor networks can describe chiral topological states, but the examples there are all gapless. Indeed, Dubail and Read [4] even proved a no-go theorem, which roughly states that for any chiral local free-fermion tensor network state, any local parent Hamiltonian is gapless. Here we construct a Boltzmann machine that approximates the unique ground state of a (gapped) chiral  $p$ -wave superconductor. The error of approximation is an inverse polynomial in the system size, and the neural network is quasi-local in the sense that the maximum distance of connections between units is logarithmic in the system size. This example explicitly demonstrates the power of Boltzmann machines.

From now on, we consider fermionic systems, for which it is necessary and there are multiple ways to account for the exchange statistics of fermions. We take a straightforward approach: Each vertex carries a Grassmann variable  $\xi_j$  rather than an Ising variable  $s_j$ , and the sum over Ising variables in hidden units are replaced by the Grassmann integral. We work in the second quantization formalism. Let  $c_j, c_j^\dagger$  for  $1 \leq j \leq |V|$  be the fermionic annihilation/creation operators, and  $|0\rangle$  be the vacuum state with no fermions. We identify  $c_j^\dagger$  with  $\xi_j$  so that  $\Xi|0\rangle$  represents a fermionic state, where  $\Xi$  is an arbitrary Grassmann variable in the algebra generated by  $\xi_1, \xi_2, \dots, \xi_{|V|}$ . As an analog of Eq. (3),

$$|\psi\rangle = \left( \int e^{-\sum_{j,k} w_{jk} \xi_j \xi_k} \prod_{l=1}^{|H|} d\xi_{|V|+l} \right) |0\rangle \quad (8)$$

is an (unnormalized Gaussian) fermionic neural network state. Note that we have set  $h_j = 0$ . This class of states appeared previously in Ref. [4].

One of the simplest examples of a chiral topological phase is the  $p+ip$  superconductor [17]. For concreteness, we consider the lattice model in Ref. [18]. Let  $c_{\vec{x}}, c_{\vec{x}}^\dagger$  be the fermionic annihilation and creation operators at the site  $\vec{x} \in \mathbb{Z}^2$  on a two-dimensional square lattice. Let  $\vec{i} = (1, 0)$  and  $\vec{j} = (0, 1)$  be the unit vectors in the  $x$  and  $y$  axes, respectively. The Hamiltonian is

$$H = \sum_{\vec{x} \in \mathbb{Z}^2} c_{\vec{x}+\vec{i}}^\dagger c_{\vec{x}} + c_{\vec{x}+\vec{j}}^\dagger c_{\vec{x}} + c_{\vec{x}+\vec{i}}^\dagger c_{\vec{x}}^\dagger + i c_{\vec{x}+\vec{j}}^\dagger c_{\vec{x}}^\dagger + \text{h.c.} \\ - 2\mu \sum_{\vec{x} \in \mathbb{Z}^2} c_{\vec{x}}^\dagger c_{\vec{x}}, \quad \mu \in \mathbb{R}. \quad (9)$$

Let  $\vec{k} = (k_x, k_y)$  be the lattice momentum, and  $\int_{\text{BZ}} d\vec{k}$  be the integral over the Brillouin zone  $(-\pi, \pi]^{\times 2}$ . To slightly simplify the presentation, we will be sloppy about unimportant overall prefactors in the calculations below. The Fourier transform

$$c_{\vec{x}}^\dagger = \int_{\text{BZ}} e^{-i\vec{k} \cdot \vec{x}} c_{\vec{k}}^\dagger d\vec{k}, \quad c_{\vec{x}} = \int_{\text{BZ}} e^{i\vec{k} \cdot \vec{x}} c_{\vec{k}} d\vec{k} \quad (10)$$

leads to

$$H = \int_{\text{BZ}} H_{\vec{k}} d\vec{k}, \quad H_{\vec{k}} = \Delta_{\vec{k}} c_{\vec{k}}^\dagger c_{-\vec{k}}^\dagger + \Delta_{\vec{k}}^* c_{-\vec{k}} c_{\vec{k}} + 2M_{\vec{k}} c_{\vec{k}}^\dagger c_{\vec{k}} \quad (11)$$

in the momentum space, where

$$\Delta_{\vec{k}} = \sin k_x + i \sin k_y, \quad M_{\vec{k}} = \cos k_x + \cos k_y - \mu. \quad (12)$$

The quasi-particle spectrum is given by

$$E_{\vec{k}} = 2\sqrt{|\Delta_{\vec{k}}|^2 + M_{\vec{k}}^2} \\ = 2\sqrt{\sin^2 k_x + \sin^2 k_y + (\cos k_x + \cos k_y - \mu)^2}. \quad (13)$$

Hence,  $H$  is gapless for  $\mu = 0, \pm 2$  and gapped otherwise. The unique ground state of  $H$  is

$$|\psi\rangle \propto e^{\frac{1}{2} \int_{\text{BZ}} \frac{v_{\vec{k}}}{u_{\vec{k}}} c_{\vec{k}}^\dagger c_{-\vec{k}}^\dagger d\vec{k}} |0\rangle, \quad (14)$$

where  $|0\rangle$  is the vacuum state, and

$$v_{\vec{k}} = \Delta_{\vec{k}}, \quad u_{\vec{k}} = \sqrt{|\Delta_{\vec{k}}|^2 + M_{\vec{k}}^2} - M_{\vec{k}} \quad (15)$$

so that  $v_{\vec{k}} = -v_{-\vec{k}}$  and  $u_{\vec{k}} = u_{-\vec{k}}$ . It is not difficult to see that the model (9) is a trivial superconductor for  $|\mu| > 2$ . It is topological superconductors with opposite chirality for  $-2 < \mu < 0$  and  $0 < \mu < 2$ , respectively.

Indeed, a state of the form (14) has an exact local neural network representation (8) [4] if  $u_{\vec{k}}, v_{\vec{k}}$  are constant-degree trigonometric polynomials, i.e., polynomials in  $e^{\pm i k_x}, e^{\pm i k_y}$ . In particular, we have each site  $\vec{x}$  carry a visible and a hidden unit. To simplify the notation, the Grassmann variable in the visible unit is identified with  $c_{\vec{x}}^\dagger$ , and that in the hidden unit is denoted by  $\xi_{\vec{x}}$ . The Fourier transform of Grassmann variables is defined as  $\xi_{\vec{x}} = \int_{\text{BZ}} e^{-i\vec{k} \cdot \vec{x}} \xi_{\vec{k}} d\vec{k}$ . In the momentum space, it is easy to see that the state (14) can be represented as

$$|\psi\rangle \propto \left( \int [d\xi_{\vec{k}}] e^{\int_{\text{BZ}} d\vec{k} (v_{\vec{k}} \xi_{-\vec{k}} c_{\vec{k}}^\dagger - \frac{1}{2} \xi_{-\vec{k}} v_{-\vec{k}} u_{\vec{k}} \xi_{\vec{k}})} \right) |0\rangle, \quad (16)$$

where  $\int [d\xi_{\vec{k}}]$  denotes the integral over all Grassmann variables in the momentum space with a proper measure. Transforming to the real space,  $[d\xi_{\vec{k}}]$  becomes  $\prod_{\vec{x} \in \mathbb{Z}^2} d\xi_{\vec{x}}$ , and the exponent in parentheses is local because  $u_{\vec{k}}, v_{\vec{k}}$  are trigonometric polynomials (the maximum distance of connections between units is proportional to the degree of the trigonometric polynomials). Thus, Eq. (16) reduces

to Eq. (8) with additional properties: (i) the neural network representation is translationally invariant; (ii) there are no connections between visible units.

Generically, exact representations are too much to ask for; hence approximate representations are acceptable. Furthermore, we usually have to work with a finite system size in order to rigorously quantify the error of the approximation and the succinctness of the representation. We now justify these statements with an example. Matrix product states constitute an excellent ansatz for ground states in one-dimensional gapped systems [19]. A generic gapped ground state in a chain of  $N$  spins (i) cannot be exactly written as a matrix product state with bond dimension  $e^{O(N)}$ ; (ii) is not expected to be approximated (in the sense of 99% fidelity) by a matrix product state with bond dimension  $O(1)$  because errors may accumulate while we truncate the Schmidt coefficients across every cut [20]; (iii) can be approximated by a matrix product state with bond dimension  $N^{o(1)}$  [21].

The goal is to construct an as-local-as-possible neural network representation (8) for the ground state of the model (9).

**Proposition 1.** *Let  $|\psi\rangle$  be the ground state of the model (9) on a (finite) square lattice of size  $L \times L$  with periodic boundary conditions. There exists a neural network state  $|\phi\rangle$  such that the fidelity  $|\langle\phi|\psi\rangle| \geq 1 - 1/\text{poly}(L)$  and that the maximum distance of connections between units is  $O(\epsilon^{-1} \log L)$ , where  $\epsilon$  is the energy gap.*

*Proof.* For concreteness, we consider  $0 < \mu \leq 2$  so that the only common zero of  $u_{\vec{k}}$  and  $v_{\vec{k}}$  is at  $\vec{k} = \vec{0}$ . It should be clear the same result applies to other values of  $\mu$  provided that the system is gapped. The main idea is to approximate  $u_{\vec{k}}$  (15) with a trigonometric polynomial. The approximation error is exponentially small in the degree of the polynomial because  $u_{\vec{k}}$  is a real analytic function of  $\vec{k}$ . The locality of the neural network is due to the smallness of the degree.

We now provide the details of the construction. Assume  $L$  is odd. This slightly simplifies the analysis, but it should be clear that a minor modification of the analysis leads to the same result for even  $L$ . We abuse the notation by letting BZ denote the set of viable points  $\{0, \pm 2\pi/L, \dots, \pm(L-1)\pi/L\}^{\times 2}$  for lattice momenta in the Brillouin zone, and  $\text{hBZ} = \{\vec{k} \in \text{BZ} | k_x > 0 \vee (k_x = 0 \wedge k_y > 0)\}$  be the “right half” of BZ. The Fourier transform (10) becomes discrete:

$$c_{\vec{x}}^\dagger = \frac{1}{L} \sum_{\vec{k} \in \text{BZ}} e^{-i\vec{k} \cdot \vec{x}} c_{\vec{k}}^\dagger, \quad c_{\vec{x}} = \frac{1}{L} \sum_{\vec{k} \in \text{BZ}} e^{i\vec{k} \cdot \vec{x}} c_{\vec{k}}, \quad (17)$$

and the Hamiltonian in the momentum space is given by

$$H = 2 \sum_{\vec{k} \in \text{hBZ}} M_{\vec{k}} c_{\vec{k}}^\dagger c_{\vec{k}} + M_{\vec{k}}^* c_{-\vec{k}}^\dagger c_{-\vec{k}} + \Delta_{\vec{k}} c_{\vec{k}}^\dagger c_{-\vec{k}}^\dagger + \Delta_{\vec{k}}^* c_{-\vec{k}} c_{\vec{k}} + 2M_{\vec{k}=\vec{0}} c_{\vec{k}=\vec{0}}^\dagger c_{\vec{k}=\vec{0}}. \quad (18)$$

The normalized ground state is

$$|\psi\rangle = \bigotimes_{\vec{k} \in \text{hBZ}} |\psi_{\vec{k}}\rangle, \quad |\psi_{\vec{k}}\rangle \propto e^{v_{\vec{k}} c_{\vec{k}}^\dagger c_{-\vec{k}}^\dagger / u_{\vec{k}}} |0\rangle. \quad (19)$$

Consider the normalized state

$$|\phi^{(m)}\rangle = \bigotimes_{\vec{k} \in \text{hBZ}} |\phi_{\vec{k}}^{(m)}\rangle, \quad |\phi_{\vec{k}}^{(m)}\rangle \propto e^{v_{\vec{k}} c_{\vec{k}}^\dagger c_{-\vec{k}}^\dagger / u_{\vec{k}}^{(m)}} |0\rangle, \quad (20)$$

where  $u_{\vec{k}}^{(m)}$  is a degree- $m$  trigonometric polynomial obtained by expanding  $u_{\vec{k}}$  in Fourier series and computing the partial sum up to order  $m$ . Similar to Eq. (16),  $|\phi^{(m)}\rangle$  has an exact neural network state representation (8) such that the maximum distance of connections between units is  $O(m)$ .

As  $u_{\vec{k}}$  is a real analytic function of  $\vec{k}$ , its Fourier coefficients of order  $m$  decay exponentially as  $e^{-m/\xi}$  for some constant  $\xi > 0$ . The decay rate  $\xi$  is a function of  $\mu$  and can be solved analytically; see, e.g., Ref. [22], Section 2. In the regime the energy gap  $\epsilon$  is small, we obtain  $\xi = \Theta(1/\epsilon)$ . Therefore,

$$|u_{\vec{k}}^{(m)} - u_{\vec{k}}| = O(e^{-m/\xi}), \quad \forall \vec{k} \in \text{hBZ}. \quad (21)$$

Furthermore, the absolute values of  $u_{\vec{k}}, v_{\vec{k}}$  are bounded away from 0:

$$|u_{\vec{k}}| \geq \Omega(L^{-2}), \quad |v_{\vec{k}}| \geq \Omega(L^{-1}), \quad \forall \vec{k} \in \text{hBZ}. \quad (22)$$

Equations (21), (22) imply

$$f_{\vec{k}} := |\langle\phi_{\vec{k}}^{(m)}|\psi_{\vec{k}}\rangle| \geq 1 - 1/\text{poly}(L), \quad \forall \vec{k} \in \text{hBZ} \quad (23)$$

for  $m = O(\xi \log L)$  with a sufficiently large constant prefactor hidden in the big-O notation. As  $|\psi\rangle, |\phi^{(m)}\rangle$  are product states in the momentum space, the fidelity is given by

$$|\langle\phi^{(m)}|\psi\rangle| = \prod_{\vec{k} \in \text{hBZ}} f_{\vec{k}} \geq (1 - 1/\text{poly}(L))^{O(L^2)} = 1 - \frac{1}{\text{poly}(L)}. \quad (24)$$

We complete the proof by letting  $|\phi\rangle = |\phi^{(m)}\rangle$ .  $\square$

It is not difficult to see that  $|\psi\rangle$  is well approximated by a thermal state at inverse temperature  $O(\log L)$ . Since the thermal state has a projected entangled pair approximation with quasi-polynomial bond dimension  $e^{O(\log^2 L)}$  [23–25], there exists a projected entangled pair state  $|\varphi\rangle$  with bond dimension  $e^{O(\log^2 L)}$  such that  $|\langle\varphi|\psi\rangle| \geq 1 - 1/\text{poly}(L)$ . It is an open problem to improve the bond dimension of  $|\varphi\rangle$  to  $\text{poly}(L)$ .

The authors would like to thank Xie Chen for collaboration in the early stages of this project. Y.H. acknowledges funding provided by the Institute for Quantum Information and Matter, an NSF Physics Frontiers Center (NSF Grant PHY-1733907) with support of the Gordon

and Betty Moore Foundation (GBMF-2644). J.E.M. is supported by NSF DMR-1507141, DMR-1918065 and a Simons Investigatorship.

*Note added.*—Part of this work was presented on November 27, 2014 (Thanksgiving day!) at the Perimeter Institute for Theoretical Physics. Very recently, we became aware of some related papers [26–28], which studied the relationship between neural and tensor network states using different approaches. In particular, Theorem 2 and Corollary 3 are stronger than Theorem 3 in Ref. [28]. After the present work was on arXiv, some other related papers [29–36] appeared.

---

\* yichuang@mit.edu

† Present address: Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

‡ jemooore@berkeley.edu

- [1] F. Verstraete, V. Murg, and J.I. Cirac, “Matrix product states, projected entangled pair states, and variational renormalization group methods for quantum spin systems,” *Advances in Physics* **57**, 143–224 (2008).
- [2] Giuseppe Carleo and Matthias Troyer, “Solving the quantum many-body problem with artificial neural networks,” *Science* **355**, 602–606 (2017).
- [3] David H. Ackley, Geoffrey E. Hinton, and Terrence J. Sejnowski, “A learning algorithm for Boltzmann machines,” *Cognitive Science* **9**, 147–169 (1985).
- [4] J. Dubail and N. Read, “Tensor network trial states for chiral topological phases in two dimensions and a no-go theorem in any dimension,” *Physical Review B* **92**, 205307 (2015).
- [5] P. Smolensky, “Information processing in dynamical systems: Foundations of harmony theory,” in *Parallel Distributed Processing: Explorations in the Microstructure of Cognition, Volume 1: Foundations* (MIT Press, Cambridge, MA, USA, 1986) Chap. 6, pp. 194–281.
- [6] Nicolas Le Roux and Yoshua Bengio, “Representational power of restricted Boltzmann machines and deep belief networks,” *Neural Computation* **20**, 1631–1649 (2008).
- [7] Guido Montufar and Nihat Ay, “Refinements of universal approximation results for deep belief networks and restricted Boltzmann machines,” *Neural Computation* **23**, 1306–1319 (2011).
- [8] Mohammad H. Amin, Evgeny Andriyash, Jason Rolfe, Bohdan Kulchitsky, and Roger Melko, “Quantum Boltzmann machine,” *Physical Review X* **8**, 021050 (2018).
- [9] M. B. Hastings, “Locality in quantum systems,” in *Quantum Theory from Small to Large Scales*, Lecture Notes of the Les Houches Summer School, Vol. 95 (Oxford University Press, Oxford, UK, 2012) Chap. 3, pp. 171–212.
- [10] Dong-Ling Deng, Xiaopeng Li, and S. Das Sarma, “Machine learning topological states,” *Physical Review B* **96**, 195145 (2017).
- [11] Sevag Gharibian, Yichen Huang, Zeph Landau, and Seung Woo Shin, “Quantum Hamiltonian complexity,” *Foundations and Trends in Theoretical Computer Science* **10**, 159–282 (2015).
- [12] Yichen Huang, *Classical simulation of quantum many-body systems*, Ph.D. thesis, University of California, Berkeley (2015).
- [13] David Perez-Garcia, Frank Verstraete, Michael M. Wolf, and J. Ignacio Cirac, “Matrix product state representations,” *Quantum Information and Computation* **7**, 401–430 (2007).
- [14] Zheng-Cheng Gu, Frank Verstraete, and Xiao-Gang Wen, “Grassmann tensor network states and its renormalization for strongly correlated fermionic and bosonic states,” (2010), arXiv:1004.2563.
- [15] B. Béri and N. R. Cooper, “Local tensor network for strongly correlated projective states,” *Physical Review Letters* **106**, 156401 (2011).
- [16] T. B. Wahl, H.-H. Tu, N. Schuch, and J. I. Cirac, “Projected entangled-pair states can describe chiral topological states,” *Physical Review Letters* **111**, 236805 (2013).
- [17] N. Read and Dmitry Green, “Paired states of fermions in two dimensions with breaking of parity and time-reversal symmetries and the fractional quantum Hall effect,” *Physical Review B* **61**, 10267–10297 (2000).
- [18] Andrei Bernevig and Titus Neupert, “Topological superconductors and category theory,” in *Topological Aspects of Condensed Matter Physics*, Lecture Notes of the Les Houches Summer School, Vol. 103 (Oxford University Press, Oxford, UK, 2014) Chap. 2, pp. 63–122.
- [19] M B Hastings, “An area law for one-dimensional quantum systems,” *Journal of Statistical Mechanics: Theory and Experiment* **2007**, P08024 (2007).
- [20] F. Verstraete and J. I. Cirac, “Matrix product states represent ground states faithfully,” *Physical Review B* **73**, 094423 (2006).
- [21] Itai Arad, Alexei Kitaev, Zeph Landau, and Umesh Vazirani, “An area law and sub-exponential algorithm for 1D systems,” (2013), arXiv:1301.1162.
- [22] Lloyd N. Trefethen, “Finite difference and spectral methods for ordinary and partial differential equations,” (1996), unpublished text, available at <http://people.maths.ox.ac.uk/trefethen/pdetext.html>.
- [23] M. B. Hastings, “Solving gapped Hamiltonians locally,” *Physical Review B* **73**, 085115 (2006).
- [24] M. Kliesch, C. Gogolin, M. J. Kastoryano, A. Riera, and J. Eisert, “Locality of temperature,” *Physical Review X* **4**, 031019 (2014).
- [25] Andras Molnar, Norbert Schuch, Frank Verstraete, and J. Ignacio Cirac, “Approximating Gibbs states of local Hamiltonians efficiently with projected entangled pair states,” *Physical Review B* **91**, 045138 (2015).
- [26] Jing Chen, Song Cheng, Haidong Xie, Lei Wang, and Tao Xiang, “Equivalence of restricted Boltzmann machines and tensor network states,” *Physical Review B* **97**, 085104 (2018).
- [27] Dong-Ling Deng, Xiaopeng Li, and S. Das Sarma, “Quantum entanglement in neural network states,” *Physical Review X* **7**, 021021 (2017).
- [28] Xun Gao and Lu-Ming Duan, “Efficient representation of quantum many-body states with deep neural networks,” *Nature Communications* **8**, 662 (2017).
- [29] Yusuke Nomura, Andrew S. Darmawan, Youhei Yamaji, and Masatoshi Imada, “Restricted Boltzmann machine learning for solving strongly correlated quantum systems,” *Physical Review B* **96**, 205152 (2017).
- [30] Elina Robeva and Anna Seigal, “Duality of classical models and tensor networks,” *Information and Inference: A Journal of the IMA* **8**, 273–288 (2018).

- [31] Stephen R. Clark, “Unifying neural-network quantum states and correlator product states via tensor networks,” *Journal of Physics A: Mathematical and Theoretical* **51**, 135301 (2018).
- [32] Ivan Glasser, Nicola Pancotti, Moritz August, Ivan D. Rodriguez, and J. Ignacio Cirac, “Neural-network quantum states, string-bond states, and chiral topological states,” *Physical Review X* **8**, 011006 (2018).
- [33] Raphael Kaubruegger, Lorenzo Pastori, and Jan Carl Budich, “Chiral topological phases from artificial neural networks,” *Physical Review B* **97**, 195136 (2018).
- [34] Giuseppe Carleo, Yusuke Nomura, and Masatoshi Imada, “Constructing exact representations of quantum many-body systems with deep neural networks,” *Nature Communications* **9**, 5322 (2018).
- [35] Sirui Lu, Xun Gao, and L.-M. Duan, “Efficient representation of topologically ordered states with restricted Boltzmann machines,” *Physical Review B* **99**, 155136 (2019).
- [36] Sujie Li, Feng Pan, Pengfei Zhou, and Pan Zhang, “Boltzmann machines as two-dimensional tensor networks,” (2021), arXiv:2105.04130.