

CHCRUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Mechanics of Metric Frustration in Contorted Filament Bundles: From Local Symmetry to Columnar Elasticity

Daria W. Atkinson, Christian D. Santangelo, and Gregory M. Grason Phys. Rev. Lett. **127**, 218002 — Published 18 November 2021 DOI: 10.1103/PhysRevLett.127.218002

Mechanics of metric frustration in contorted filament bundles: From local symmetry to columnar elasticity

Daria W. Atkinson

Department of Physics and Astronomy, University of Pennsylvania^{*}

Christian D. Santangelo Department of Physics, Syracuse University

Gregory M. Grason

Department of Polymer Science and Engineering, University of Massachusetts Amherst

Bundles of filaments are subject to geometric frustration: certain deformations (e.g. bending while twisted) require longitudinal variations in spacing between filaments. While bundles are common—from protein fibers to yarns—the mechanical consequences of longitudinal frustration are unknown. We derive a geometrically-nonlinear formalism for bundle mechanics, using a gauge-like symmetry under *reptations* along filament backbones. We relate force balance to orientational geometry and assess the elastic cost of frustration in twisted toroidal bundles.

Elastic zero modes are a ubiquitous feature of soft materials, from mechanical metamaterials [1, 2] to liquid crystal elastomers [3]. Such systems can undergo large deformations with minimal strain, as geometrically coupled rotations and translations preserve local spacing between microscopic constituents. The smectic and columnar liquid crystalline phases provide paradigmatic examples of zero modes in soft elastic systems, permitting relative "sliding" of 2D layers and 1D columns, respectively. Smectic and columnar phases with zero-cost sliding displacements bear a striking resemblance to a much broader class of laminated and filamentous structures with largely inelastic, disipative forces tangent to their layers or columns and without longitudinal correlations between material points, ranging from multi-layer graphene materials [4] and stacked paper [5] to biopolymer bundles [6, 7], nanotube yarns [8], and wire ropes [9].

While there are well established frameworks which capture the geometric nonlinearities of smectic liquid crystals (i.e., the strain tensor accurately describes arbitrarily large and complex deformations) [10], no such framework exists for columnar and filamentous materials. The orientations of column backbones impose constraints on inter-filament spacing, generating rich modes of geometric frustration without counterpart in smectic liquid crystals. In the simplest non-trivial case of helical bundles, predictions from a minimally non-linear approximation of columnar elasticity [11] and tomographic analysis of elastic filament bundles [12] show that twist in straight bundles gives rise to non-uniform inter-filament stress and spacing in transverse sections (see Fig 1a). Except for the restrictive classes of straight, twisted bundles [13] and twist-free developable domains [14, 15], bundle textures also generate *longitudinal* frustration, requiring local spacings to vary *along* a bundle [16]. Although deformations that introduce longitudinal frustration are the



FIG. 1. In (a), an equilibrium twisted bundle with straight, untwisted reference configuration, $\Omega R = 1$, and 2D Poisson ratio $\nu = 0.8$, colored by the local pressure $P = -S_{\alpha}^{\alpha}/2$. *Reptation* of the gold filament by σ along its contour, draws the filament (in the direction indicated by the gold arrow) above the planar cross-section of the bundle, but leaves the local separation $d\Delta$ unchanged. In (b), a twisted-toroidal bundle found by bending the same bundle to $\kappa_0 = 0.2/R$ and optimizing inter-filament elastic cost.

rule rather than the exception—for example, wire ropes or toroidal biopolymer condensates are both *twisted* and *bent* (e.g. Fig. 1)—existing frameworks of columnar elasticity fail to capture this effect.

In this Letter, we develop a fully geometrically nonlinear Lagrangian elasticity theory of columnar materials, which completely captures the interplay between orientation, and both lateral and longitudinal frustration of inter-filament spacing. We construct this theory by imposing a gauge-like local symmetry under *reptations*, deformations that slide filaments along their contours without changing the inter-filament spacing. The resultant equilibrium equations point to the role geometrical measures of non-equidistance play in bundles' mechanics. Within this framework, we compute the energetic costs of longitudinal frustration in twisted, toroidal bundles, and give evidence that 1) optimal configurations generically incorporate splay and 2) the bending cost that derives from non-uniform compression depends nonmonotonically on pretwist.

To construct the elastic theory, we divide space into points on filament backbones, labeled by two coordinates: \mathbf{v} , a 2D label of filaments; and s, a length coordinate along filaments. Hence, the location of each point in the bundle is described by a function $\mathbf{r}(s, \mathbf{v})$, with $\partial_s \mathbf{r}(s, \mathbf{v})$ parallel to the tangent vector $\mathbf{t} = \partial_s \mathbf{r}/|\partial_s \mathbf{r}|$. Because they lack positional order along their backbone curves, filament bundles and columnar liquid crystals have a family of continuous zero modes, corresponding to *reptations* (Fig. 1),

$$\mathbf{r}'(s, \mathbf{v}) = \mathbf{r}(s + \sigma(s, \mathbf{v}), \mathbf{v}). \tag{1}$$

We assume that changes in local spacing can be described by an energy density function \mathcal{W} , that depends only on the deformation gradient [17].

To account for reptation symmetry, we demand that \mathcal{W} depend on a modified deformation gradient, which transforms as a scalar under $\sigma(s, \mathbf{v})$, depends solely on the deformation, \mathbf{r} , of the material itself, and recovers the well-established 2D elasticity of developable bundles (i.e. parallel arrays) [14, 15]. Specifically, we construct a covariant derivative $D_I \mathbf{r} = \nabla_I \mathbf{r} - \mathbf{A}_I$, where ∇_I is the usual covariant derivative on tensors in the material space and \mathbf{r} determines \mathbf{A}_{I} , such that if two configurations, \mathbf{r} and $\mathbf{r'}$, are related by Eq. (1), then $D_I \mathbf{r} = D'_I \mathbf{r}'$. In order for $D_I \mathbf{r}$ to be reptation invariant, we must have that $-\mathbf{A}'_I + \mathbf{A}_I = \nabla_I \sigma \partial_s \mathbf{r}$. Therefore, $-\mathbf{A}'_{I}+\mathbf{A}_{I}=-(\mathbf{t}\cdot\nabla_{I}\mathbf{r}')\mathbf{t}+(\mathbf{t}\cdot\nabla_{I}\mathbf{r})\mathbf{t}$. To construct an elastic theory of columnar materials, we set $\mathbf{A}_I = (\mathbf{t} \cdot \nabla_I \mathbf{r}) \mathbf{t}$, which is manifestly reptation invariant, and leads to a deformation gradient that only measures deformations *transverse* to the local backbones.

$$D_I \mathbf{r} \equiv \nabla_I \mathbf{r} - (\mathbf{t} \cdot \nabla_I \mathbf{r}) \mathbf{t}.$$
 (2)

Notably, for two nearby filaments at $\mathbf{r}(\mathbf{v})$ and $\mathbf{r}(\mathbf{v} + d\mathbf{v})$ in material coordinates, the covariant derivative gives the local distance of closest approach $d\mathbf{\Delta} = dv^I D_I \mathbf{r}$, for which $d\mathbf{\Delta} \cdot \mathbf{t} = 0$ (see again inset of Fig. 1a). As shown explicitly in the Supplemental Information, this covariant deformation gradient captures the standard 2D deformation gradients of developable domains.

From this deformation gradient, we construct an effective metric $g_{IJ}^{\text{eff}} = D_I \mathbf{r} \cdot D_J \mathbf{r}$, which is naturally invariant under rotations of \mathbf{r} , and which encodes the metric inherited by the local 2D section transverse to the filaments in the bundle [18]. Because $D_s \mathbf{r} = 0$, the effective metric only has components for 2×2 block $I, J \neq s$, which we denote using index notation $\alpha, \beta \in \{1, 2\}$. We can now define the Green - Saint-Venant strain tensor [17, 19, 20] as

$$\epsilon_{\alpha\beta} = \frac{1}{2} \left[D_{\alpha} \mathbf{r} \cdot D_{\beta} \mathbf{r} - g_{\alpha\beta}^{\text{tar}} \right], \tag{3}$$

where $g_{\alpha\beta}^{\text{tar}}$ is the *target metric* corresponding to strainfree state, which for this Letter, we take to be developable with uniform spacing, so $g_{\alpha\beta}^{\text{tar}} = \delta_{\alpha\beta}$. For weak deflections from the uniform parallel state, Eq. (3) reduces to the small-tilt approximation to the non-linear columnar strain [11, 21, 22], which captures the lowest-order dependence of spacing on orientation [23].

Assuming that strains are small though deformations may be large, we can treat the elastic energy as quadratic in the strain, so that

$$\mathcal{E}_s = \int dV \ \mathcal{W}(\epsilon) = \frac{1}{2} \int dV S^{\alpha\beta} \epsilon_{\alpha\beta}, \qquad (4)$$

where $S^{\alpha\beta} = \frac{\partial W}{\partial \epsilon_{\alpha\beta}} = C^{\alpha\beta\gamma\delta}\epsilon_{\gamma\delta}$ is the nominal stress tensor, and $C^{\alpha\beta\gamma\delta}$ is a tensor of elastic constants which depends on both the crystalline symmetries of the underlying columnar order and the target metric, $g_{\alpha\beta}^{\text{tar}}$ [24]. Energetics of columnar materials also include other gaugeinvariant costs, including the Frank-Oseen orientational free energy (particularly filament bending) and the cost of local density changes *along* columns [25]. Here, for clarity, we focus only on the energetics of columnar strain and detail the combined effects of other contributions elsewhere [26]. Given this gauge-invariant formulation of the columnar strain energy, we first illustrate the mechanical effects of orientational geometry on local forces. This follows from the bulk Euler-Lagrange equations of Eq. (4) (see Supplemental Information for a complete derivation),

$$\frac{\delta \mathcal{E}_s}{\delta \mathbf{r}} = -\nabla_\alpha \left(S^{\alpha\beta} D_\beta \mathbf{r} \right) + \nabla_s \left(S^{\alpha\beta} D_\beta \mathbf{r} \frac{\mathbf{t} \cdot \nabla_\alpha \mathbf{r}}{|\nabla_s \mathbf{r}|} \right).$$
(5)

The bulk terms represent body forces generated by the columnar strains, which must be balanced by other internal stresses or external forces. To cast them in a more geometrical light, we consider separately the components tangential and perpendicular to \mathbf{t} , F_{\parallel} and \mathbf{F}_{\perp} , respectively.

Making use of the identity, $\mathbf{t} \cdot \nabla_{\alpha} D_{\beta} \mathbf{r} = -\partial_{\alpha} \mathbf{t} \cdot \partial_{\beta} \mathbf{r}$, the tangential forces can be recast simply as

$$F_{\parallel} = S^{\alpha\beta} h_{\alpha\beta}, \tag{6}$$

where

$$h_{\alpha\beta} = \frac{1}{2} \Big[\partial_{\alpha} \mathbf{t} \cdot \partial_{\beta} \mathbf{r} + \partial_{\beta} \mathbf{t} \cdot \partial_{\alpha} \mathbf{r} \\ - \frac{\mathbf{t} \cdot \partial_{\alpha} \mathbf{r}}{|\partial_{s} \mathbf{r}|} \partial_{s} \mathbf{t} \cdot \partial_{\beta} \mathbf{r} - \frac{\mathbf{t} \cdot \partial_{\beta} \mathbf{r}}{|\partial_{s} \mathbf{r}|} \partial_{s} \mathbf{t} \cdot \partial_{\alpha} \mathbf{r} \Big],$$
(7)

is the convective flow tensor that measures longitudinal variations in inter-filament spacing [16]. Just as the second fundamental form measures gradients of a surface's normal vector [27], $h_{\alpha\beta}$ measure the symmetric gradients of **t** in its normal plane (i.e. its trace is the splay of filament tangents).



FIG. 2. A 2D schematic of local forces in a filament bundle (with a uniform, parallel reference state), illustrating transverse stress S_{\perp}^{\perp} with regions under compression in blue and those under extension in red. The forces parallel and tangent to t are shown at two points, and push material points to regions of vanishing stress.

Here, we see that tangent forces couple nonequidistance to the stress tensor much like the Young-Laplace law couples in-plane stresses to normal forces in curved membranes [19]. This analogy becomes exact for zero twist textures: when $\mathbf{t} \cdot (\nabla \times \mathbf{t}) = 0$, filaments can be described by a set of surfaces normal to \mathbf{t} . In this case, it is possible to choose coordinates so that $\partial_{\alpha} \mathbf{r} \cdot \mathbf{t} = 0$, and tangential forces give the Young-Laplace force normal to each surface, with $h_{\alpha\beta}$ reducing to their second fundamental form. This illustrates the intuitive notion, shown schematically in Fig. 2, that columnar strain generates tangential body forces that push material points *towards* lower-stress locations in the array.

The bulk components of Eq. (5) perpendicular to \mathbf{t} give the transverse force

$$\mathbf{F}_{\perp} = -D_{\alpha} \left[S^{\alpha\beta} D_{\beta} \mathbf{r} \right] + D_{s} \left[S^{\alpha\beta} \frac{\mathbf{t} \cdot \partial_{\alpha} \mathbf{r}}{|\partial_{s} \mathbf{r}|} D_{\beta} \mathbf{r} \right].$$
(8)

This form captures the divergence of stress in the planes perpendicular to backbones. The second term accounts for the corrections arising from material derivatives that lie along the backbone, such as twisted textures, when $\partial_{\alpha} \mathbf{r} \cdot \mathbf{t} \neq 0$. Thus, the longitudinal derivatives in \mathbf{F}_{\perp} are needed to capture transverse mechanics of even equidistant twisted bundles beyond the lowest order geometric non-linearity [11].

We now illustrate the energetics of longitudinal frustration by considering a prototypical non-equidistant geometry: twisted toroidal bundles. Motivated in large part by the morphologies of condensed biopolymers [6, 7], theoretical models of twisted toroids have focused on their orientational elasticity costs [28–30], ignoring the unavoidable frustration of inter-filament spacing in this geometry. While satisfying force balance in non-equidistant bundles requires physical ingredients beyond the columnar strain energy, which we consider elsewhere [26], for the purposes of this Letter we take advantage of the full geometric-nonlinearity of Eq. (5) to explore the specific costs of longitudinal gradients in spacing required by simultaneous twist and bend.

We construct twisted toroids from equilibrium twisted

helical bundles of radius R and constant pitch, $2\pi/\Omega$ by bending them such that their central curve \mathbf{r}_0 is deformed from a straight line into a circle of radius κ_0^{-1} (see Fig. 1b). We then define perturbative displacements $\mathbf{r}(s + \delta s, \rho + \delta \rho, \phi + \delta \phi) \simeq \mathbf{r}_0 + \rho \hat{\rho} + \partial_s \mathbf{r}^{(0)} \delta s + \delta \rho \hat{\rho} + \delta \phi \hat{\phi}$ relative to the bent, pre-twisted bundles, where ρ and ϕ describe the (Eulerian) distance from the central curve and the angular position relative to its normal in the plane perpendicular to its tangent \mathbf{t}_0 , and where $\partial_s \mathbf{r}^{(0)} =$ $\mathbf{t}_0 + \Omega \rho \hat{\phi}$. The small- ρ limit of the force balance equations for the strain energy motivates the following displacements

$$\begin{pmatrix} \delta s \\ \delta \rho \\ \delta \phi \end{pmatrix} = \begin{pmatrix} a_s \Omega \kappa_0 \rho^3 \sin \phi \\ a_\rho \Omega^2 \kappa_0 \rho^4 \cos \phi \\ a_\phi \Omega^2 \kappa_0 \rho^3 \sin \phi \end{pmatrix}$$
(9)

where a_s , a_{ρ} , and a_{ϕ} are variational parameters. Notably, to linear order in curvature, these parameterize the "almost equidistant" ansatzes considered previously, including splay-free (tr(h) = 0) [28] configurations and det(h) = 0 [16] ansatzes.

We expand the energy to quadratic order in κ_0 , holding the center of area at \mathbf{r}_0 , which constrains $a_s = a_\rho - a_\phi$, then minimize Eq. (4) with respect to the displacements for a given ΩR . Examples of the distribution of pressure $P = -S_{\alpha}^{\alpha}/2$ are shown in Fig. 1b. Relative to the axisymmetric pressure induced by helical twist in the straight bundle, bending into a twisted toroid requires bunching (spreading) of the filaments at the inner (outer) positions in the toroid, leading to a polarization of the pressure towards the normal.

Because bending and twisting of bundles introduces longitudinal strain variation, bending pre-twisted bundle introduces additional stresses whose elastic cost can be characterized by an effective bending stiffness B, defined by $E_{\text{bend}} = \frac{B}{2} \int ds \kappa_0^2$, which derives purely from columnar strain, rather than intra-filament deformations. Fig. 3 shows that longitudinal frustration leads to a bending cost that increases with small twist as $B \sim (\Omega R)^4$, but eventually gives way to remarkable non-monotonic behavior at large pre-twist. We note further that the bending cost grows with the 2D Poisson ratio ν of the columnar array, highlighting the importance of local compressional deformations in optimal twisted toroids.

We analyze the optimal modes of deformation via the convective flow tensor, in particular, the trace $\operatorname{tr}(h)$ (splay) and deviatoric components $h^{\operatorname{dev}} = h_{\alpha\beta} - \operatorname{tr}(h)\delta_{\alpha\beta}/2$ (biaxial splay) [31], which characterize longitudinal gradients of dilatory and shear stress in the columnar array. In contrast to a heuristic view that optimal packings should favor the uniform area per filament of splay-free textures, the inset of Fig. 3 instead shows that optimal toroids incorporate a mixture of both splay and biaxial splay where we define the respective measures of average splay and biaxial splay, $\Psi \equiv \frac{1}{\kappa_{2}^{2}V} \int dV \operatorname{tr}(h)^{2}$



FIG. 3. The effective bending modulus B which results from the frustration of constant spacing in twisted-toroidal bundles. The inset shows dimensionless measures of mean splay (Ψ) and biaxial splay (Γ) defined in the text, for fixed twist $\Omega R = 0.8$ for a range of 2D Poisson ratios, from $\nu = 0.2$ to 1.0. Approaching incompressibility ($\nu \rightarrow 1$), Ψ goes to zero, maintaining uniform area-per-filament at the expense of expense of markedly increased strain energy.

and $\Gamma \equiv \frac{1}{\kappa_0^2 V} \int dV \operatorname{tr}(h_{\operatorname{dev}}^2)$. Only in the incompressible limit, as $\nu \to 1$, does the splay vanish, and only at the expense of additional biaxial splay and energetic cost, implying counterintuitively that splayed textures are in fact energetically favorable in longitudinally frustrated twisted toroids. Indeed, the energetic preference for splay in non-equidistant bundles, can be traced to force balance conditions in this geometry [26].

As an elastic theory, our treatment of reptation invariant energetics neglects the non-conservative forces of tangential friction in 2D ordered bundles. Such a description will apply, by definition, to equilibria of any system with columnar order (i.e. without long-range inter-axial correlations between filaments), conditions widely encountered in a range of molecular bundle-forming systems, from supermolecular fibers [32, 33] to nanotube ropes [34–36]. For example, simulations of nanotube ropes [34, 35], confirm that inter-filament cohesion is sufficient to maintain dense, 2D order while columns reptate under highly contorted, twisted and bent deformations. Condensed states of DNA and other polynucleotides, for example, exhibit columnar states [37], and notably exhibit twisted, contorted and 2D ordered packings within viral capsids [7, 38], although the existence of inter-helical correlations in such structure and their effects on observed packing remains a matter of open investigation [39]. The question of when the reptation symmetric elastic description captures goemetrically non-linear deformations of macroscopic bundles, in which static friction between filaments may transmit axial shears, is more complex. While some recent experiments [12, 40] with macroscopic bundles of densely-packed, 2D ordered and twisted bundles demonstrate that large degrees of distortion are possible without noticeable effects of sliding friction, prevailing theoretical models, backed up by finite-element simulations, of twisted cables, indicate that the effects of static friction lead to mechanical transitions between shear coupled and un-coupled regimes [41, 42]. Notably, models of bending in pretwisted cables show that above a threshold bending (which increases with static friction and pre-tension), deformation transitions to a slipping regime, [43, 44]. In this context, we might speculate that deformations in the post-slipping regimes of these models tends towards those captured by the minimal reptationinvarient elasticity. It may indeed by possible to explore this conjecture via a homogenization approach to a suitable constitutive model of fibered elastic composites [45–48], in which the inter-fiber matrix transmits strictly no elastic shears, but instead only an elastoplastic slipping of fibers determined by local transverse pressure (i.e. Coulomb friction).

In summary, we have shown that gauge-theoretic principles underlie the geometrically-nonlinear theory of columnar elasticity, providing a means to quantify the cost of longitudinal frustration in the mechanics of bundles. Unlike phase field models of nonlinear elasticity (such as [49, 50]), this description depends neither on the presence of a planar reference state, nor presupposes uniform crystalline order, allowing us to both accommodate the effective curvature of bundles of constant pitch helices [51], and providing a natural generalization to arbitrary target metrics [19]. Because this approach to elasticity relies only on the existence of local, continuous zero modes, we note that it can be generalized to other liquid crystals, like smectics, and anticipate that it may have applications beyond liquid crystals, including mechanical metamaterials.

The authors gratefully acknowledge useful discussions with R. Kusner, S. Zhou, and B. Davidovitch. M. Dimitriyev, D. Hall, and R. Kamien provided feedback on an early draft of this manuscript. This work was supported by the NSF under grants DMR-1608862 and NSF DMR-1822638; D.A. received additional funding from the Penn Provost's postdoctoral fellowship and NSF MRSEC DMR-1720530.

- * atkinsod@sas.upenn.edu, she/her or they/them
- X. Mao, N. Xu, and T. C. Lubensky, Physical Review Letters 104, 085504 (2010).
- [2] K. Sun, A. Souslov, X. Mao, and T. C. Lubensky, Proceedings of the National Academy of Sciences 109, 12369 (2012).
- [3] M. Warner, P. Bladon, and E. M. Terentjev, Journal de Physique II 4, 93 (1994).
- [4] Z. Xu, H. Sun, X. Zhao, and C. Gao, Advanced Materials 25, 188 (2013).
- [5] S. Poincloux, T. Chen, B. Audoly, and P. Reis, arXiv:2012.01335 [cond-mat] (2020), arXiv:2012.01335 [cond-mat].
- [6] N. V. Hud and K. H. Downing, Proceedings of the Na-

tional Academy of Sciences 98, 14925 (2001).

- [7] A. Leforestier and F. Livolant, Proceedings of the National Academy of Sciences 106, 9157 (2009).
- [8] M. Zhang, K. R. Atkinson, and R. H. Baughman, Science 306, 1358 (2004).
- [9] G. A. Costello, Theory of Wire Rope, Mechanical Engineering Series (Springer-Verlag, New York, 1990).
- [10] G. Grinstein and R. A. Pelcovits, Physical Review A 26, 915 (1982).
- [11] G. M. Grason, Physical Review E 85 (2012), 10.1103/PhysRevE.85.031603.
- [12] A. Panaitescu, G. M. Grason, and A. Kudrolli, Physical Review E 95 (2017), 10.1103/PhysRevE.95.052503.
- [13] G. M. Grason, Reviews of Modern Physics 87, 401 (2015).
- [14] Y. Bouligand, Journal de Physique 41, 1297 (1980).
- [15] M. Kléman, Journal de Physique **41**, 737 (1980).
- [16] D. W. Atkinson, C. D. Santangelo, and G. M. Grason, New Journal of Physics 21, 062001 (2019).
- [17] P. G. Ciarlet, Journal of Elasticity 78, 1 (2005).
- [18] Equidistant bundles—where components of g_{IJ}^{eff} are uniform along *s*—are known as Riemannian fibrations, see e.g. [52, 53], and the transverse structure can be mapped on to single 2D base space. Non-equidistant bundles correspond to Sub-Riemannian generalizations [54] where the inherited metric measures distances in the 2-planes perpendicular to **t** at each point.
- [19] E. Efrati, E. Sharon, and R. Kupferman, Journal of the Mechanics and Physics of Solids 57, 762 (2009).
- [20] M. A. Dias, J. A. Hanna, and C. D. Santangelo, Physical Review E 84, 036603 (2011).
- [21] J. V. Selinger and R. F. Bruinsma, Physical Review A 43, 2910 (1991).
- [22] G. M. Grason and R. F. Bruinsma, Physical Review Letters 99 (2007), 10.1103/PhysRevLett.99.098101.
- [23] See Supplemental Information, which includes [55, 56].
- [24] We raise and lower indices with $g^{\alpha\beta \text{tar}}$, the inverse of $g_{\alpha\beta}^{\text{tar}}$, rather than $g^{\alpha\beta\text{eff}}$, as a matter of convenience, and note that the difference in the elastic energy is higher order in the strain tensor [20].
- [25] The symmetry under local reptations in this theory is unlike a related treatment of both the nematic to smectic-A transition [57, 58] and its columnar analog [59–61] in which the gauge symmetry of a density-wave is explicitly broken by, e.g., the splay elasticity of the underlying nematic order.
- [26] D. W. Atkinson, C. D. Santangelo, and G. M. Grason, in preparation (2020).
- [27] R. S. Millman and G. D. Parker, <u>Elements of Differential Geometry</u> (Englewood Cliffs, N.J. : Prentice-Hall, 1977).
- [28] I. M. Kulić, D. Andrienko, and M. Deserno, Europhysics Letters 67, 418 (2004).
- [29] J. Charvolin and J. F. Sadoc, The European Physical Journal E 25, 335 (2008).
- [30] V. Koning, B. C. v. Zuiden, R. D. Kamien, and V. Vitelli, Soft Matter 10, 4192 (2014).
- [31] J. V. Selinger, Liquid Crystals Reviews 6, 129 (2018).
- [32] T. Kato, T. Yasuda, Y. Kamikawa, and M. Yoshio, Chem. Commun., 729 (2009).
- [33] K. Yano, Y. Itoh, F. Araoka, G. Watanabe, [5 T. Hikima, and T. Aida, Science 363, 161 (2019), [6 https://science.sciencemag.org/content/363/6423/161.full.pdf.

- [34] I. Ostanin, R. Ballarini, D. Potyondy, and T. Dumitric, Journal of the Mechanics and Physics of Solids 61, 762 (2013).
- [35] I. Ostanin, R. Ballarini, and T. Dumitrică, Journal of Applied Mechanics 81 (2014), 10.1115/1.4026484.
- [36] Y. Wang, I. Ostanin, C. Gaidu, and T. Dumitric, Langmuir **31**, 12323 (2015).
- [37] F. Livolant and A. Leforestier, Progress in Polymer Science 21, 1115 (1996), liquid Crystalline Polymers-Part 2.
- [38] S. L. Ilca, X. Sun, K. E. Omari, A. Kotecha, F. d. Haas, F. DiMaio, J. M. Grimes, D. I. Stuart, M. M. Poranen, and J. T. Huiskonen, Nature 570, 252 (2019), number: 7760 Publisher: Nature Publishing Group.
- [39] L. Barberi, F. Livolant, A. Leforestier, and M. Lenz, Nucleic Acids Research 49, 3709 (2021), https://academic.oup.com/nar/articlepdf/49/7/3709/37123994/gkab197.pdf.
- [40] A. Panaitescu, G. M. Grason, and A. Kudrolli, Physical Review Letters 120, 248002 (2018).
- [41] D. Liu, S. Zheng, and Y. He, Mathematics and Mechanics of Solids 24, 2154 (2019).
- [42] E. A. W. de Menezes and R. J. Marczak, Engineering Structures 241, 112487 (2021).
- [43] K.-J. Hong, A. Der Kiureghian, and J. L. Sackman, Journal of Engineering Mechanics 131, 500 (2005).
- [44] D. Zhang and M. Ostoja-Starzewski, Journal of Applied Mechanics 83 (2015), 10.1115/1.4032023.
- [45] K. D. Cherednichenko, V. P. Smyshlyaev, and V. V. Zhikov, Proceedings of the Royal Society of Edinburgh Section A: Mathematics 136, 87 (2006).
- [46] M. Bellieud, SIAM Journal on Mathematical Analysis 49, 2615 (2017).
- [47] D. Engl, C. Kreisbeck, and A. Ritorto, arXiv:2105.03971 [math] (2021), arXiv:2105.03971 [math].
- [48] E. Davoli and C. Kreisbeck, arXiv:2108.00038 [math] (2021), arXiv:2108.00038 [math].
- [49] P. M. Chaikin and T. C. Lubensky, <u>Principles of Condensed Matter Physics</u> <u>University Press, Cambridge</u>, 1995). (Cambridge
- [50] O. Stenull and T. C. Lubensky, private communication (2009).
- [51] I. R. Bruss and G. M. Grason, Proceedings of the National Academy of Sciences 109, 10781 (2012).
- [52] B. O'Neill, The Michigan Mathematical Journal 13, 459 (1966).
- [53] D. Gromoll and G. Walschap, "Submersions, foliations, and metrics," in <u>Metric Foliations and Curvature</u> (Birkhäuser Basel, Basel, 2009) pp. 1–44.
- [54] A. Agrachev, D. Barilari, and U. Boscain, <u>A Comprehensive Introduction to Sub-Riemannian Geometry</u>, Cambridge Studies in Advanced Mathematics (Cambridge University Press, 2019).
- [55] M. Kléman and P. Oswald, Journal de Physique 43, 655 (1982).
- [56] G. M. Grason, Soft Matter 9, 6761 (2013).
- [57] P. G. de Gennes, Solid State Communications 10, 753 (1972).
- [58] T. C. Lubensky, G. Grinstein, and R. A. Pelcovits, Physical Review B 25, 6022 (1982).
- [59] C. Giannessi, Physical Review A 34, 705 (1986).
- [60] R. D. Kamien and D. R. Nelson, Physical Review Letters
 f. 74, 2499 (1995).
- [61] R. D. Kamien, Journal de Physique II 6, 461 (1996).