

This is the accepted manuscript made available via CHORUS. The article has been published as:

Impact of Clustering on the math

$$\text{Li}^{8+}$$

$$\beta$$
 Decay and Recoil Form Factors

G. H. Sargsyan, K. D. Launey, M. T. Burkey, A. T. Gallant, N. D. Scielzo, G. Savard, A. Mercenne, T. Dytrych, D. Langr, L. Varriano, B. Longfellow, T. Y. Hirsh, and J. P. Draayer
Phys. Rev. Lett. **128**, 202503 — Published 19 May 2022

DOI: [10.1103/PhysRevLett.128.202503](https://doi.org/10.1103/PhysRevLett.128.202503)

Impact of clustering on the ${}^8\text{Li}$ β decay and recoil form factors

G. H. Sargsyan,¹ K. D. Launey,¹ M. T. Burkey,^{2,3,4} A. T. Gallant,⁴ N. D. Scielzo,⁴ G. Savard,³ A. Mercenne,^{1,5} T. Dytrych,^{1,6} D. Langr,⁷ L. Varriano,^{2,3} B. Longfellow,⁴ T. Y. Hirsh,⁸ and J. P. Draayer¹

¹*Department of Physics and Astronomy, Louisiana State University, Baton Rouge, LA 70803, USA*

²*Department of Physics, University of Chicago, Chicago, Illinois 60637, USA*

³*Physics Division, Argonne National Laboratory, Argonne, Illinois 60439, USA*

⁴*Lawrence Livermore National Laboratory, Livermore, California 94550, USA*

⁵*Center for Theoretical Physics, Sloane Physics Laboratory, Yale University, New Haven, Connecticut 06520, USA*

⁶*Nuclear Physics Institute of the Czech Academy of Sciences, 250 68 Rež, Czech Republic*

⁷*Department of Computer Systems, Faculty of Information Technology, Czech Technical University in Prague, Prague 16000, Czech Republic*

⁸*Soreq Nuclear Research Center, Yavne 81800, Israel*

We place unprecedented constraints on recoil corrections in the β decay of ${}^8\text{Li}$, by identifying a strong correlation between them and the ${}^8\text{Li}$ ground state quadrupole moment in large-scale *ab initio* calculations. The results are essential for improving the sensitivity of high-precision experiments that probe the weak interaction theory and test physics beyond the Standard Model (BSM). In addition, our calculations predict a 2^+ state of the $\alpha + \alpha$ system that is energetically accessible to β decay but has not been observed in the experimental ${}^8\text{Be}$ energy spectrum, and has an important effect on the recoil corrections and β decay for the $A = 8$ systems. This state and an associated 0^+ state are notoriously difficult to model due to their cluster structure and collective correlations, but become feasible for calculations in the *ab initio* symmetry-adapted no-core shell-model framework.

Introduction. – The left-handed vector minus axial-vector (V–A) structure of the weak interaction was postulated in late 1950’s and early 1960’s [1, 2] guided in large part by a series of β -decay experiments [3–5], and later was incorporated in the Standard Model of particle physics. However, in its most general form, the weak interaction can also have scalar, tensor, and pseudoscalar terms as well as right-handed currents.

Today, β -decay experiments continue to pursue increasingly sensitive searches for additional contributions to the weak interaction. Various experiments [5–8] have constrained the tensor part of the interaction, although the limits are less stringent compared to the other non-standard-model terms [9, 10]. While these experiments have achieved remarkable precision, further improvements require confronting the systematic uncertainties that stem from higher-order corrections (referred to as recoil-order terms) in nuclear β decay. These terms are inherently small compared to the allowed β decay terms; however, current experiments have reached a precision where even subtle distortions matter. Measurements of recoil-order terms are also interesting in their own right as they can test additional symmetries of the Standard Model, such as the existence of second-class currents [11–13] and the accuracy of the conserved vector current (CVC) hypothesis [12, 14–16].

The β decay of ${}^8\text{Li}$ to ${}^8\text{Be}$, which subsequently breaks up into two α particles, has long been recognized as an excellent testing ground to search for new physics [11, 14, 17] due to the high decay energy and the ease of detecting the β and two α particles. Recently, by taking advantage of ion-trapping techniques, high-precision measurements of β - $\bar{\nu}$ - α correlations [7, 8, 18] have been performed that set the most stringent limit on a tensor contribution to




FIG. 1. Decay scheme for β decay of ${}^8\text{Li}$ g.s. to the broad low-lying 2^+ state in ${}^8\text{Be}$. Energies, in MeV, are relative to ${}^8\text{Be}$ g.s. A small β -decay branching is observed to the doublet 2^+ states due to their resonant nature.

date [19]. However, in this type of experiment, one of the largest uncertainties comes from the several different recoil-order corrections that contribute to the decay. A number of other experiments have taken advantage of the presence of certain recoil-order terms in the β decay of ${}^8\text{Li}$ (Fig. 1) to perform CVC tests by studying β - α angular correlations [15, 20] and β -spin alignment correlations [13, 16]. In addition, these terms have been deduced from γ decays of the doublet 2^+ states near 17 MeV in the ${}^8\text{Be}$ spectrum, which contain the isobaric analogue of the ${}^8\text{Li}$ ground state (g.s.) [12]. Due to their small size, and the fact that there are several terms that contribute to decay observables, most of the experimentally extracted recoil-order terms have large uncertainties.

In this Letter, we report the first *ab initio* calculations

of recoil-order terms in the β decay of ^8Li . These calculations achieve highly reduced uncertainties compared to the experimentally extracted values of [16]. They help decrease the systematic uncertainties on the tensor-current estimates in the weak interaction reported in [19], and are of interest to experimental tests of the CVC hypothesis [12]. We also provide evidence that the β -transition strength of the ^8Li decay is affected by a disputed low-lying 2^+ state (sometimes referred to as an “intruder” state) below 16 MeV in the ^8Be spectrum. Our calculations in unprecedentedly large model spaces support the existence of low-lying states with a large overlap with the $\alpha + \alpha$ s - and d -waves. Indeed, a very broad 2^+ state along with a lower 0^+ were initially proposed by Barker from the R-matrix analysis of $\alpha + \alpha$ scattering and the β decays of ^8Li and ^8B [21–23]. Even though such states have not been directly observed experimentally, some earlier theoretical studies have predicted them in the low-lying spectrum of ^8Be [24–26]. Furthermore, there has been a recent experimental indication in favor of intruder states below 16 MeV [27].

SA-NCSM framework.— For this work, we employ the *ab initio* symmetry-adapted no-core shell model (SA-NCSM) [28–30]. The use of chiral effective-field-theory interactions [31–34] enables nuclear calculations informed by elementary particle physics, while the symmetry-adapted (SA) basis allows us to achieve ultra-large model spaces imperative for the description of challenging features in the ^8Be states, such as clustering and collectivity. It uses a harmonic oscillator (HO) basis with frequency $\hbar\Omega$ and a model space with an N_{max} cutoff, which is the maximum total HO excitation quanta above the lowest HO configuration for a given nucleus. These parameters are related to infrared (IR) and ultraviolet (UV) cutoffs [35], which can be understood as the effective size of the model space in which the nucleus resides, and its grid resolution, respectively. The calculations become independent of $\hbar\Omega$ at $N_{\text{max}} \rightarrow \infty$, providing a parameter-free *ab initio* prediction. The SA-NCSM results exactly reproduce those of the NCSM [36, 37] for the same nuclear interaction. However, by utilizing the emergent symplectic $\text{Sp}(3, \mathbb{R})$ -symmetry in nuclei [29], the SA-NCSM can expand the model space by a physically relevant subspace, which is only a fraction of the complete NCSM

space, thereby including localized- α degrees of freedom within the interaction effective range [38].

We adopt various chiral potentials without renormalization in nuclear medium: $\text{N}^3\text{LO-EM}$ [33], NNLO_{opt} [40], as well as NNLO_{sat} [41] with the three-nucleon (3N) forces, hierarchically smaller than their nucleon-nucleon (NN) forces, added as averages [30]. For comparison, we present results with the soft JISP16 phase-equivalent NN interaction [42]. We use $\hbar\Omega = 15\text{--}25$ MeV for $\text{N}^3\text{LO-EM}$, NNLO_{opt} , and JISP16, and $\hbar\Omega = 16$ and 22 MeV for NNLO_{sat} , unless otherwise stated. The NNLO_{opt} is used without 3N forces, which have been shown to contribute minimally to the 3- and 4-nucleon binding energy [40]. Furthermore, the NNLO_{opt} NN potential has been found to reproduce various observables, including the ^4He electric dipole polarizability [43]; the challenging analyzing power for elastic proton scattering on ^4He , ^{12}C , and ^{16}O [44]; along with B(E2) transition strengths for ^{21}Mg and ^{21}F [45] in the SA-NCSM without effective charges.

For the purposes of this study, the quadrupole moment of the ^8Li g.s., $Q(2^+_{\text{g.s.}})$, for which SA-NCSM calculations with the NNLO_{opt} NN are extrapolated to an infinite model-space size, is shown to reproduce the experimental value within the many-body model uncertainties (Fig. 2a). The result is in close agreement with the extrapolated value of [46] that uses renormalized NN+3N chiral potentials. The model uncertainties are based on variations in the model-space size and resolution, and extrapolations use the Shanks method [29].

Recoil-order corrections.— The recoil-order form factors are generally neglected in β -decay theory since they are of the order of q/m_N or higher, where q is the momentum transfer (typically several MeV/c) and m_N is the nucleon mass [14]. Thus, for most β decays, the recoil effects are typically less than a percent of the dominant Fermi and Gamow-Teller (GT) contributions (for an example see [47]). However, for measurements of sufficiently high precision, these terms must be included in the analysis especially when the leading contributions are suppressed or the recoil-order terms are unusually large. These recoil-order form factors include, the second forbidden axial vectors (j_2 and j_3), induced tensor (d), and weak magnetism (b), and along with the GT (c_0) are given in the impulse approximation (IA) as:

$$\begin{aligned}
 c_0(q^2) &= (-)^{(J'-J)} \frac{g_A(q^2)}{\sqrt{2J+1}} \langle J' || \sum_{i=1}^A \tau_i^\pm \sigma_i || J \rangle = (-)^{(J'-J)} \frac{g_A(q^2)}{\sqrt{2J+1}} M_{GT}, \\
 j_K(q^2) &= -(-)^{(J'-J)} \frac{2}{3} \frac{g_A(q^2)}{\sqrt{2J+1}} \frac{(Am_N c^2)^2}{(\hbar c)^2} \langle J' || \sum_{i=1}^A \tau_i^\pm [Q_i \times \sigma_i]^K || J \rangle, \text{ with } K = 2, 3, \\
 d(q^2) &= (-)^{(J'-J)} A \frac{g_A(q^2)}{\sqrt{2J+1}} \langle J' || \sum_{i=1}^A \tau_i^\pm \sqrt{2} [L_i \times \sigma_i]^1 || J \rangle, \\
 b(q^2) &= A \frac{(-)^{(J'-J)}}{\sqrt{2J+1}} \left[g_M(q^2) \langle J' || \sum_{i=1}^A \tau_i^\pm \sigma_i || J \rangle + g_V(q^2) \langle J' || \sum_{i=1}^A \tau_i^\pm L_i || J \rangle \right], \quad (1)
 \end{aligned}$$




FIG. 2. (a) *Ab initio* ^8Li g.s. quadrupole moment $Q(2^+)$ compared to experiment [39] (denoted as “Expt.”). Calculations use the NNLO_{opt} NN for different model-space sizes and resolutions (open symbols), along with the infinite-size extrapolated value (dashed line) and the corresponding many-body uncertainty (shaded area). (b) Calculated j_2/A^2c_0 and j_3/A^2c_0 (squares and triangles, respectively) and their predicted values (upper and lower horizontal lines, respectively) for the ^8Li β decay to 2_1^+ in ^8Be vs. the calculated ^8Li $Q(2^+)$. The first (second) set of the uncertainties in Eq. (2) is shown as the line thickness (gray bands). Calculations use the NNLO_{opt}, NNLO_{sat} and N³LO chiral potentials, and the JISP16 NN, in $N_{\text{max}} = 6$ -12 model spaces. Inset: subset of calculated j_2/A^2c_0 vs. $Q(2^+)$ for selected IR (in fm)/UV (in MeV) many-body cutoffs across all interactions, and for two interactions across several cutoffs.

where $g_V(0) = 1$, $g_A(0) \approx 1.27$ and $g_M(0) \approx 4.70$ are the vector, axial and weak magnetism coupling constants, A is the mass number and $J(J')$ is the total angular momentum of the initial (final) nucleus. The $\tau_i/2$, $\sigma_i/2$, $Q_i = \sqrt{16\pi/5} r_i^2 Y_{2\mu}(\hat{r}_i)$, and L_i are the isospin, intrinsic spin, quadrupole moment and angular momentum operators, respectively, of the i^{th} particle. M_{GT} is the conventional GT matrix element. The matrix elements in Eq. (1) are computed translationally invariant in the SA-NCSM. These recoil-order form factors, usually reported as the ratios $j_{2,3}/A^2c_0$, d/Ac_0 , and b/Ac_0 , enter into the expression of the β -decay rate for nuclei undergoing delayed α -particle emission [8, 14, 19, 48].

Remarkably, we identify a strong correlation between $j_{2,3}/A^2c_0$ and the ^8Li g.s. quadrupole moment based on calculations across several interactions, N_{max} and $\hbar\Omega$ parameters (Fig. 2b, using $N_{\text{max}} = 6$ to 12 for NNLO_{opt}, to 10 for N³LO-EM and JISP16, and to 8 for NNLO_{sat}). As can be seen in the Fig. 2b inset, the linear dependence is observed regardless of any errors that may arise from the many-body truncation and from the higher-order effects (e.g., [49, 50]) associated with various interactions. An identical spread is found for j_3/A^2c_0 due to the strong correlation between j_2 and j_3 (see Supplemental Material [51]). A linear regression along with the combination of the correlation to $Q(2^+_{\text{g.s.}})$ and its experimental value of $3.14(2) e \text{ fm}^2$ [39] lead to reduced uncertainties on our predictions:

$$\frac{j_2}{A^2c_0} = -966 \pm 13 \pm 33, \quad \frac{j_3}{A^2c_0} = -1546 \pm 19 \pm 40, \quad (2)$$

Here, the first set of uncertainties uses the quadrupole moment experimental uncertainties given the linear re-

gression slope, and the second set arises from the regression uncertainty using Student’s t -distribution and a 99% confidence level. This correlation is important, as we can reduce the problem of calculating a matrix element that depends on cluster physics in ^8Be to a bound state observable in ^8Li .

Most significantly, with the values in Eq. (2), the uncertainty from the recoil-order corrections on the tensor current contribution to the weak interaction presented in [48] is reduced by over 50% [19]. The recoil-order terms, including the b and d terms, for the lowest four SA-NCSM 2^+ states, are summarized in Table I. The d/Ac_0 prediction for 2_1^+ is based on a correlation similar to the one for $j_{2,3}/A^2c_0$ (see Supplemental Material [51]). These predictions can be used in future experiments to constrain BSM tensor currents, while these b weak magnetism predictions are of interest to experiments that test the CVC hypothesis and d is of importance to determining the existence of second-class currents [12].

TABLE I. The recoil-order terms from SA-NCSM. Results for the 2_1^+ $j_{2,3}/A^2c_0$ and d/Ac_0 are based on the correlation to $Q(2^+_{\text{g.s.}})$; all other calculations use NNLO_{opt} and have error bars from variations in $\hbar\Omega$ by 5 MeV and in model-space sizes up to $N_{\text{max}} = 16$ (12) for $j_{2,3}/A^2c_0$ (d/Ac_0 and b/Ac_0).

	j_2/A^2c_0	j_3/A^2c_0	d/Ac_0	b/Ac_0
2_1^+	-966 ± 36	-1546 ± 44	10.0 ± 1.0	6.0 ± 0.4
2_2^+ (new)	-10 ± 10	-80 ± 30	-0.5 ± 0.5	3.7 ± 0.4
2_3^+ (doublet 1)	12 ± 5	-60 ± 15	0.3 ± 0.2	3.8 ± 0.2
2_4^+ (doublet 2)	11 ± 3	-65 ± 11	0.2 ± 0.2	3.8 ± 0.2

New final state for β decays to ^8Be .— The experimentally

deduced values presented in [16], $j_2/A^2c_0 = -490 \pm 70$, $j_3/A^2c_0 = -980 \pm 280$, $d/Ac_0 = 5.5 \pm 1.7$ and $b/Ac_0 = 7.5 \pm 0.2$, are comparable but different from our predicted values. These experimental results were obtained through a global fit to β -spin alignment [16] and β - α angular correlation data [20] from ^8Li and ^8B β decays. Due to the small size of higher-order effects and relatively large statistical uncertainties, the $j_{2,3}/A^2c_0$ and d/Ac_0 were assumed in [16] to be independent of the ^8Be excitation energy. Thus, the results were averaged over the entire β decay spectrum. In contrast, the SA-NCSM wavefunctions are for individual states, hence, the predictions in Eq. (2) are for the lowest 2^+ state only, which is the dominant transition for the ^8Li β decay. The SA-NCSM reveals large differences between the recoil-order terms to the lowest 2_1^+ and higher-lying states, the most notable being for the j_K/A^2c_0 terms where the values differ by almost two orders of magnitude (see Table I). Hence, the angular-correlation experiment in [19] minimizes the sensitivity to the higher-lying states by restricting their analysis to decays centered on the broad 2_1^+ state.

Importantly, the SA-NCSM indicates the existence of another 2^+ state below 16 MeV – accessible to the ^8Li or ^8B β decays through allowed transitions – and a corresponding lower 0^+ state that largely overlaps with the $\alpha+\alpha$ system (Fig. 3, see also Supplemental Material [51]). In the SA-NCSM, these states quickly decrease in energy as the model space increases (Fig. 3a) regardless of the realistic interaction used, similar to the Hoyle-state rotational band in ^{12}C [58]. The extrapolations are performed using the three-parameter exponential formula from [59]. Notably, 0_3^+ converges to 20.1 ± 1.5 MeV and has a structure similar to the doublet 2^+ states and isospin $T = 1$. This state is not seen in the currently-available experimental spectrum and it is likely to be the isobaric analogue of the low-lying 0^+ state in ^8Li predicted by recent *ab initio* calculations [46, 60].

The calculated low-lying states in ^8Be are in good agreement with experiment (Fig. 3b). The NNLO_{sat} results include the average 3N contribution determined for a given isospin (for ^8Be , the contribution to the binding energy in $N_{\text{max}}=12$ is 1.51 MeV, resulting in a total extrapolated binding energy of 56.8 MeV). The extrapolations determine the energies of 0_2^+ and 2_2^+ between 5 and 15 MeV above the g.s., corroborating earlier estimates [21, 22, 24].

For NNLO_{opt} NN and the case of the fastest energy convergence of the 0_2^+ and 2_2^+ states ($\hbar\Omega=15$ MeV), we estimate α widths (Fig. 3b), by projecting the $N_{\text{max}}=16$ SA-NCSM wavefunctions onto $\alpha + \alpha$ cluster states and considering the exact continuum Coulomb wavefunctions outside of the interaction effective range, following the procedure of [38]. For this, the ^8Be and ^4He states are expressed in the $\text{Sp}(3, \mathbb{R})$ basis, associated with intrinsic shapes [29]. For ^8Be , we consider three dominant prolate shapes with contributions of 75%, 4%, and 3% to $0_{\text{g.s.}}^+$ (totaling 82%), and 46%, 15%, and 11% to 0_2^+ (totaling 72%), and similarly for the 2^+ states (see Supplemental

Material [51]). These shapes extend to 18 HO shells and start at the most deformed configurations among those in the valence shell: $2\hbar\Omega$ and $4\hbar\Omega$ excitations. Except for the $0_{\text{g.s.}}^+$ width that uses the experimental threshold of -92 keV relative to the ^8Be g.s., all the widths use the $\alpha+\alpha$ threshold estimated at -104 keV from the SA-NCSM extrapolations of the ^4He and ^8Be binding energies with NNLO_{opt} . These widths are in good agreement with experimentally deduced values [57] and earlier theoretical studies [61–63].

Intruder 0^+ and 2^+ states in the low-lying spectrum of ^8Be were proposed in the late 1960's by Barker from concurrent R-matrix fits to scattering, reaction, and decay data associated with the ^8Be nucleus [21, 22]. The inclusion of an intruder 2^+ state below 16 MeV in the R-matrix fits of β decays in [64] results in a decrease of the extracted M_{GT} for a decay to 2_1^+ by almost 1.5 times, which yields a closer agreement with the SA-NCSM M_{GT} (see Table II; the calculated M_{GT} are not used in the experimental analysis of [19]). Note that, depending on the interaction, two-body axial currents may significantly affect M_{GT} [65], however, here we are interested only in the IA part. Due to the large uncertainty on the 2_2^+ state in the calculations, we provide only the lower limits on the $\log(ft)$ based on the convergence pattern. The energies from Barker's R-matrix fits for the intruder 0^+ and 2^+ states are ~ 6 MeV and 9 MeV, respectively, with α widths > 7 MeV. These excitation energies agree with the SA-NCSM extrapolated results given the error bars (Fig. 3b), as well as with the predicted widths. The strong excitation-energy dependence of the recoil-order terms due to the presence of 2_2^+ has a small effect on the weak tensor currents constraints in the low excitation-energy range (see systematic uncertainty in Table I in [19]), but is imperative to consider in analyses over the entire β decay spectrum.

TABLE II. *Ab initio* M_{GT} , c_0 and $\log(ft)$ in IA, compared to the experimentally deduced values. Ref. [64] includes evaluations both with an intruder 2^+ state (denoted by *) around 8 MeV similar to Ref. [23], and without it.

	2_1^+		2_2^+	
	$ M_{\text{GT}} $	$ c_0 $	$\log(ft)$	$\log(ft)$
NNLO_{opt}	0.16(1)	0.09(1)	5.90	> 5.06
NNLO_{sat}	0.21(3)	0.12(2)	5.64	> 5.05
JISP16	0.23(4)	0.13(2)	5.54	> 4.28
Expt., Ref. [23]	0.190	0.108	5.72	5.27
Expt., Ref. [64]*	0.204	0.116	5.66	5.2
Expt., Ref. [64]	0.284	0.163	5.37	—

Summary.— The *ab initio* SA-NCSM has determined the size of the recoil-order form factors in the β decay of ^8Li . It has shown that states of the $\alpha + \alpha$ system not included in the evaluated ^8Be energy spectrum have an important effect on all $j_{2,3}/A^2c_0$, b/Ac_0 and d/Ac_0 terms, and can explain the M_{GT} discrepancy in the $A = 8$ systems.




FIG. 3. (a) Calculated ^8Be low-lying 0^+ state energies illustrated for the NNLO_{opt} chiral potential ($\hbar\Omega = 15$ MeV) vs. the model-space size, together with the extrapolation values (dotted lines) and uncertainties (bands). Extrapolations use complete model spaces up to $N_{\max} = 12$ and do not include the $N_{\max} = 14$ and 16 SA selected model spaces shown with uncertainties determined by the selection. The measured $0^+_{\text{g.s.}}$ energy is -56.5 MeV [56]. Inset: $\alpha + \alpha$ s -wave for $0^+_{\text{g.s.}}$ (blue solid) and 0^+_2 (orange dashed). (b) *Ab initio* low-lying states from extrapolations for ^8Be , compared to experiment (Expt.). The extrapolation uncertainties (error bars) for the 0^+_2 and 2^+_2 states (dashed levels) are based on variations in the model-space size and selection. For NNLO_{opt}, α width estimates (shaded areas) for the lowest two 0^+ and 2^+ states are shown with uncertainties (lighter shades) determined from the energy extrapolation uncertainties; the small $0^+_{\text{g.s.}}$ width (not shown) is estimated to be 5.7 eV, compared to 5.57 eV [57].

The outcomes reduce – by over 50% – the uncertainty on these recoil-order corrections, and help improve the sensitivity of high-precision β -decay experiments that probe the $V-A$ structure of the weak interaction [19]. Furthermore, our predicted b/Ac_0 and d/Ac_0 values are important for other investigations of the Standard Model symmetries, such as the CVC hypothesis and the existence of second-class currents.

Acknowledgments.– We thank Scott Marley and Konstantinos Kravvaris for useful discussions. This work was supported in part by the U.S. National Science Foundation (PHY-1913728), SURA, Czech Science Foundation (22-14497S), and the Czech Ministry

of Education, Youth and Sports under Grant No. CZ.02.1.01/0.0/0.0/16_019/0000765. It benefited from high performance computational resources provided by LSU (www.hpc.lsu.edu), the National Energy Research Scientific Computing Center (NERSC), a U.S. Department of Energy Office of Science User Facility operated under Contract No. DE-AC02-05CH11231, as well as the Frontera computing project at the Texas Advanced Computing Center, made possible by National Science Foundation award OAC-1818253. L. Varriano was supported by a National Science Foundation Graduate Research Fellowship under Grant No. DGE-1746045. This work was performed under the auspices of the U.S. Department of Energy by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.

-
- [1] R. P. Feynman and M. Gell-Mann, Phys. Rev. **109**, 193 (1958).
 - [2] E. C. Sudarshan and R. Marshak, Phys. Rev. **109**, 1860 (1958).
 - [3] C. S. Wu, E. Ambler, R. W. Hayward, D. D. Hoppes, and R. P. Hudson, Phys. Rev. **105**, 1413 (1957).
 - [4] W. B. Herrmannsfeldt, D. R. Maxson, P. Stähelin, and J. S. Allen, Phys. Rev. **107**, 641 (1957).
 - [5] C. Johnson, F. Pleasonton, and T. Carlson, Phys. Rev. **132**, 1149 (1963).
 - [6] X. Fléchar, P. Velten, E. Liénard, A. Méry, D. Rodríguez, G. Ban, D. Durand, F. Mauger, O. Naviliat-Cuncic, and J. Thomas, J. Phys. G: Nucl. Part. Phys. **38**, 055101 (2011).
 - [7] G. Li, R. Segel, N. D. Scielzo, P. F. Bertone, F. Buchinger, S. Caldwell, A. Chaudhuri, J. A. Clark, J. E. Crawford, C. M. Deibel, J. Fallis, S. Gulick, G. Gwinner, D. Lascar, A. F. Levand, M. Pedretti, G. Savard, K. S. Sharma, M. G. Sternberg, T. Sun, J. Van Schelt, R. M. Yee, and B. J. Zabransky, Phys. Rev. Lett. **110**, 092502 (2013).
 - [8] M. Sternberg, R. Segel, N. Scielzo, G. Savard, J. Clark, P. Bertone, F. Buchinger, M. Burkey, S. Caldwell, A. Chaudhuri, *et al.*, Phys. Rev. Lett. **115**, 182501 (2015).
 - [9] N. Severijns, M. Beck, and O. Naviliat-Cuncic, Rev. Mod. Phys. **78**, 991 (2006).
 - [10] F. Wauters, A. García, and R. Hong, Phys. Rev. C **89**, 025501 (2014).
 - [11] L. Grenacs, Annu. Rev. Nucl. Part. Sci. **35**, 455 (1985).
 - [12] L. De Braekeleer, E. G. Adelberger, J. H. Gundlach, M. Kaplan, D. Markoff, A. M. Nathan, W. Schieff, K. A.

- Snover, D. W. Storm, K. B. Swartz, D. Wright, and B. A. Brown, *Phys. Rev. C* **51**, 2778 (1995).
- [13] T. Sumikama, K. Matsuta, T. Nagatomo, M. Ogura, T. Iwakoshi, Y. Nakashima, H. Fujiwara, M. Fukuda, M. Mihara, K. Minamisono, *et al.*, *Phys. Lett. B* **664**, 235 (2008).
- [14] B. R. Holstein, *Rev. Mod. Phys.* **46**, 789 (1974).
- [15] R. E. Tribble and G. T. Garvey, *Phys. Rev. C* **12**, 967 (1975).
- [16] T. Sumikama, K. Matsuta, T. Nagatomo, M. Ogura, T. Iwakoshi, Y. Nakashima, H. Fujiwara, M. Fukuda, M. Mihara, K. Minamisono, T. Yamaguchi, and T. Minamisono, *Phys. Rev. C* **83**, 065501 (2011).
- [17] C. A. Barnes, W. A. Fowler, H. B. Greenstein, C. C. Lauritsen, and M. E. Nordberg, *Phys. Rev. Lett.* **1**, 328 (1958).
- [18] N. Scielzo, G. Li, M. Sternberg, G. Savard, P. Bertone, F. Buchinger, S. Caldwell, J. Clark, J. Crawford, C. Deibel, *et al.*, *Nucl. Instrum. Methods Phys. Res. A: Accel. Spectrom. Detect. Assoc. Equip.* **681**, 94 (2012).
- [19] M. Burkey, G. Savard, A. Gallant, N. Scielzo, J. Clark, T. Hirsh, L. Varriano, G. Sargsyan, K. Launey, M. Brodeur, D. Burdette, *et al.*, Submitted, *Phys. Rev. Lett.* (2021).
- [20] R. D. McKeown, G. T. Garvey, and C. A. Gagliardi, *Phys. Rev. C* **22**, 738 (1980).
- [21] F. Barker, H. Hay, and P. Treacy, *Aust. J. Phys.* **21**, 239 (1968).
- [22] F. Barker, *Aust. J. Phys.* **22**, 293 (1969).
- [23] F. Barker, *Aust. J. Phys.* **42**, 25 (1989).
- [24] E. Caurier, P. Navrátil, W. E. Ormand, and J. P. Vary, *Phys. Rev. C* **64**, 051301(R) (2001).
- [25] P. Maris, in *J. Phys. Conf. Ser.*, Vol. 445 (IOP Publishing, 2013) p. 012035.
- [26] D. Rodkin and Y. M. Tchuvil'sky, *Chin. Phys. C* **44**, 124105 (2020).
- [27] M. Munch, O. Sølund Kirsebom, J. A. Swartz, K. Risager, and H. O. U. Fynbo, *Phys. Lett. B* **782**, 779 (2018).
- [28] K. D. Launey, T. Dytrych, and J. P. Draayer, *Prog. Part. Nucl. Phys.* **89**, 101 (2016).
- [29] T. Dytrych, K. D. Launey, J. P. Draayer, D. J. Rowe, J. L. Wood, G. Rosensteel, C. Bahri, D. Langr, and R. B. Baker, *Phys. Rev. Lett.* **124**, 042501 (2020).
- [30] K. D. Launey, A. Mercenne, and T. Dytrych, *Annu. Rev. Nucl. Part. Sci.* **71** (2021).
- [31] P. F. Bedaque and U. van Kolck, *Annu. Rev. Nucl. Part. Sci.* **52**, 339 (2002).
- [32] E. Epelbaum, A. Nogga, W. Glöckle, H. Kamada, U.-G. Meißner, and H. Witala, *Phys. Rev. C* **66**, 064001 (2002).
- [33] D. R. Entem and R. Machleidt, *Phys. Rev. C* **68**, 041001(R) (2003).
- [34] E. Epelbaum, *Prog. Part. Nucl. Phys.* **57**, 654 (2006).
- [35] K. A. Wendt, C. Forssén, T. Papenbrock, and D. Sääf, *Phys. Rev. C* **91**, 061301(R) (2015).
- [36] P. Navrátil, J. P. Vary, and B. R. Barrett, *Phys. Rev. Lett.* **84**, 5728 (2000).
- [37] P. Navrátil, J. Vary, and B. Barrett, *Phys. Rev. C* **62**, 054311 (2000).
- [38] A. C. Dreyfuss, K. D. Launey, J. E. Escher, G. H. Sargsyan, R. B. Baker, T. Dytrych, and J. P. Draayer, *Phys. Rev. C* **102**, 044608 (2020).
- [39] D. Borremans, D. L. Balabanski, K. Blaum, W. Geithner, S. Gheysen, P. Himpe, M. Kowalska, J. Lassen, P. Lievens, S. Mallion, R. Neugart, G. Neyens, N. Vermeulen, and D. Yordanov, *Phys. Rev. C* **72**, 044309 (2005).
- [40] A. Ekström, G. Baardsen, C. Forssén, G. Hagen, M. Hjorth-Jensen, G. R. Jansen, R. Machleidt, W. Nazarewicz, *et al.*, *Phys. Rev. Lett.* **110**, 192502 (2013).
- [41] A. Ekström, G. R. Jansen, K. A. Wendt, G. Hagen, T. Papenbrock, B. D. Carlsson, C. Forssén, M. Hjorth-Jensen, P. Navrátil, and W. Nazarewicz, *Phys. Rev. C* **91**, 051301 (2015).
- [42] A. Shirokov, V. Kulikov, P. Maris, A. Mazur, E. Mazur, and J. Vary, in *EPJ Web of Conferences*, Vol. 3 (EDP Sciences, 2010) p. 05015.
- [43] R. B. Baker, K. D. Launey, S. Bacca, N. N. Dinur, and T. Dytrych, *Phys. Rev. C* **102**, 014320 (2020).
- [44] M. Burrows, C. Elster, S. P. Weppner, K. D. Launey, P. Maris, A. Nogga, and G. Popa, *Phys. Rev. C* **99**, 044603 (2019).
- [45] P. Ruotsalainen, J. Henderson, G. Hackman, G. H. Sargsyan, K. D. Launey, A. Saxena, P. C. Srivastava, S. R. Stroberg, T. Grahn, J. Pakarinen, G. C. Ball, R. Julin, P. T. Greenlees, J. Smallcombe, C. Andreoiu, N. Bernier, M. Bowry, M. Buckner, R. Caballero-Folch, A. Chester, S. Cruz, L. J. Evitts, R. Frederick, A. B. Garnsworthy, M. Holl, A. Kurkjian, D. Kisliuk, K. G. Leach, E. McGee, J. Measures, D. Mücher, J. Park, F. Sarazin, J. K. Smith, D. Southall, K. Starosta, C. E. Svensson, K. Whitmore, M. Williams, and C. Y. Wu, *Phys. Rev. C* **99**, 051301(R) (2019).
- [46] G. McCracken, P. Navrátil, A. McCoy, S. Quaglioni, and G. Hupin, *Phys. Rev. C* **103**, 035801 (2021).
- [47] A. Glick-Magid, C. Forssén, D. Gazda, D. Gazit, P. Gysbers, and P. Navrátil, arXiv preprint arXiv:2107.10212 (2021).
- [48] M. T. Burkey, *Searching for tensor currents in the weak interaction using lithium-8 β decay*, Ph.D. thesis, Chicago U. (2019).
- [49] A. A. Filin, D. Möller, V. Baru, E. Epelbaum, H. Krebs, and P. Reinert, *Phys. Rev. C* **103**, 024313 (2021).
- [50] P. Maris, E. Epelbaum, R. J. Furnstahl, J. Golak, K. Hebeler, T. Hübner, H. Kamada, H. Krebs, U.-G. Meißner, J. A. Melendez, A. Nogga, P. Reinert, R. Roth, R. Skibiński, V. Soloviov, K. Topolnicki, J. P. Vary, Y. Volkotrub, H. Witała, and T. Wolfgruber (LENPIC Collaboration), *Phys. Rev. C* **103**, 054001 (2021).
- [51] See Supplemental Material at [url], which includes Refs. [52–55], for additional correlation plots, intercluster wavefunctions used in calculations of widths, as well as the decomposition of the ground and intruder states in the SA basis.
- [52] J. Suhonen, *From nucleons to nucleus: concepts of microscopic nuclear theory* (Springer Science & Business Media, 2007).
- [53] P. Navrátil, *Phys. Rev. C* **70**, 054324 (2004).
- [54] O. Castaños, J. P. Draayer, and Y. Leschber, *Z. Phys. A* **329**, 33 (1988).
- [55] M. T. Mustonen, C. N. Gilbreth, Y. Alhassid, and G. F. Bertsch, *Phys. Rev. C* **98**, 034317 (2018).
- [56] W. Huang, M. Wang, F. Kondev, G. Audi, and S. Naimi, *Chin. Phys. C* **45**, 030002 (2021).
- [57] D. Tilley, J. Kelley, J. Godwin, D. Millener, J. Purcell, C. Sheu, and H. Weller, *Nucl. Phys. A* **745**, 155 (2004).

- [58] A. C. Dreyfuss, K. D. Launey, T. Dytrych, J. P. Draayer, and C. Bahri, Phys. Lett. B **727**, 511 (2013).
- [59] P. Maris, J. P. Vary, and A. M. Shirokov, Phys. Rev. C **79**, 014308 (2009).
- [60] P. Navrátil, R. Roth, and S. Quaglioni, Phys. Rev. C **82**, 034609 (2010).
- [61] K. Kravvaris and A. Volya, Phys. Rev. Lett. **119**, 062501 (2017).
- [62] K. Kravvaris, S. Quaglioni, G. Hupin, and P. Navratil, arXiv preprint arXiv:2012.00228 (2020).
- [63] S. Elhatisari, D. Lee, G. Rupak, E. Epelbaum, *et al.*, Nature **528**, 111 (2015).
- [64] E. K. Warburton, Phys. Rev. C **33**, 303 (1986).
- [65] G. B. King, L. Andreoli, S. Pastore, M. Piarulli, R. Schiavilla, R. B. Wiringa, J. Carlson, and S. Gandolfi, Phys. Rev. C **102**, 025501 (2020).