



# CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Boundary Criticality of the 3D  $O(N)$  Model:  
From Normal to Extraordinary

Francesco Parisen Toldin and Max A. Metlitski

Phys. Rev. Lett. **128**, 215701 — Published 27 May 2022

DOI: [10.1103/PhysRevLett.128.215701](https://doi.org/10.1103/PhysRevLett.128.215701)

# Boundary criticality of the 3d O(N) model: from normal to extraordinary

Francesco Parisen Toldin\*

*Institut für Theoretische Physik und Astrophysik, Universität Würzburg, Am Hubland, D-97074 Würzburg, Germany*

Max A. Metlitski†

*Department of Physics, Massachusetts Institute of Technology, Cambridge, MA 02139, USA*

It was recently realized that the three dimensional O(N) model possesses an extraordinary boundary universality class for a finite range of  $N \geq 2$ . For a given  $N$ , the existence and universal properties of this class are predicted to be controlled by certain amplitudes of the normal universality class, where one applies an explicit symmetry breaking field to the boundary. In this paper, we study the normal universality class for  $N = 2, 3$  using Monte Carlo simulations on an improved lattice model and extract these universal amplitudes. Our results are in good agreement with direct Monte Carlo studies of the extraordinary universality class serving as a non-trivial quantitative check of the connection between the normal and extraordinary classes.

*Introduction.*— When a physical system is in the vicinity of a continuous phase transition, various observables develop power-law singularities which have a character of universality: they are determined by the gross features of the system, such as dimensionality and symmetry, and not by the details of local interactions. Renormalization-Group (RG) theory allows to understand the emergence of universality as the result of the existence of fixed points in a suitably defined flow of Hamiltonians. Accordingly, systems exhibiting identical critical behavior define a given Universality Class (UC) [1]. The presence of a boundary gives rise to rich phenomena, which have attracted a large amount of experimental [2] and theoretical [3–5] studies. General RG arguments show that a given bulk UC class, describing the critical behavior far away from the boundary, potentially admits different surface UCs [1]. Further, surface critical exponents and other universal data generally differ from those of the bulk [3, 4]. Surface UCs also determine the critical Casimir force [6–12]. While boundary criticality is a mature subject, it has recently received renewed attention driven in part by advances in conformal field theory [13–23] and developments in topological phases of quantum matter. Many topological phases including quantum Hall states, topological insulators and certain quantum spin liquids possess protected boundary states. While it was initially thought that this protection relies on the presence of a bulk energy gap, examples where the boundary state survives in some form even as the bulk gap closes were later discovered [24–32]. The study of such “gapless topological states” and their boundaries lies in the domain of boundary critical phenomena. As gapless topological states were investigated in the context of quantum magnets [33–41], it was realized that even for the simplest model of *classical* magnets — the O(N) model — basic questions about the boundary phase diagram remain open [42–44].

The much investigated classical O(N) model [45] provides a prototypical example of boundary criticality. In three dimensions, for  $N = 1, 2$ , the bulk-surface phase diagram hosts a surface transition line, where the bulk is disordered and the surface critical behavior belongs to that of 2D O(N) UC. This line terminates at the bulk transition line dividing it into or-

inary and extraordinary surface UCs; the termination point is the so-called special UC [3, 4]. Surprisingly, the surface phase diagram for  $N > 2$  is still not fully settled. For  $d = 3$  and  $N > 2$  there is no surface transition for a disordered bulk [45], thus the topology of the phase diagram does not necessarily dictate the existence of the extraordinary UC or the special multicritical point [46, 47]. Yet, a recent field-theoretical analysis in Ref. [42] has pointed out that if one treats  $N$  as a continuous parameter, the extraordinary UC survives for a range  $2 < N < N_c$  where  $N_c$  is a currently unknown constant. Further, the extraordinary UC in the region  $2 \leq N < N_c$  exhibits a surface order parameter correlation function that falls off as:

$$\langle \vec{\phi}(\mathbf{x}) \cdot \vec{\phi}(0) \rangle \sim \frac{1}{(\log \mathbf{x})^q}, \quad (1)$$

thus, it was labeled the “extraordinary-log” UC in Ref. [42]; this should be contrasted to the extraordinary transition for  $N = 1$  or in  $d > 3$  where the above correlation function approaches a constant at large separation. In fact, for  $N = 3$  a recent numerical simulation [43] finds firm evidence of a special transition with exponents differing from those of the ordinary UC and a phase consistent with the extraordinary-log UC, implying  $N_c > 3$  [48]. For  $N = 2$ , the “logarithmic” character of the extraordinary phase was also verified numerically [44].

Ref. [42] showed that for a given  $N$  the existence of the extraordinary-log phase and its properties (such as the exponent  $q$  in Eq. (1)) are determined by certain universal amplitudes of the normal boundary UC. The latter is realized when an explicit symmetry breaking field is applied to the boundary [3, 4, 49, 50].

Motivated by these recent developments, in this letter we study the normal surface UC of the three dimensional O(N) model, for  $N = 2$  and  $N = 3$ , by means of Monte Carlo (MC) simulations of an *improved* lattice model [45], where the leading bulk irrelevant scaling field is suppressed. Through a finite-size scaling analysis of MC data we determine certain universal amplitudes of the normal UC. Such amplitudes are *per se* of interest, as they provide a quantitative description of the normal UC; for  $N = 1$  they have been studied in Ref. [51].

Furthermore, exploiting the analysis of Ref. [42], our results confirm the existence of the extraordinary-log UC for  $N = 3$ , and allow us to compute the universal exponent  $q$  in Eq. (1) for  $N = 2$  and  $N = 3$ . Our results are in good agreement with the value of  $q$  found in direct studies of the extraordinary phase in Refs. [43, 44].

*Model.*— We study the classical lattice  $\phi^4$  model by means of MC simulations. It is defined on a three-dimensional  $L_{\parallel} \times L_{\perp} \times L$  lattice, with periodic boundary conditions (BCs) along the lateral directions with size  $L_{\parallel}$ , and open BCs along the remaining direction. The reduced Hamiltonian  $\mathcal{H}$ , such that the Gibbs weight is  $\exp(-\mathcal{H})$ , is

$$\mathcal{H} = -\beta \sum_{\langle i j \rangle} \vec{\phi}_i \cdot \vec{\phi}_j - \beta_s \sum_{\langle i j \rangle_s} \vec{\phi}_i \cdot \vec{\phi}_j - \vec{h}_s \cdot \sum_{i \in s} \vec{\phi}_i + \sum_i [\phi_i^2 + \lambda(\phi_i^2 - 1)^2], \quad (2)$$

where  $\vec{\phi}_x$  is a  $N$ -components real field on the lattice site  $x$  and the first sum extends over the nearest-neighbor pairs where at least one site belongs to the inner bulk. The second and third sums extend over the lattice sites on the surface. The last term in Eq. (2) is summed over all lattice sites. In Eq. (2) the coupling constant  $\beta$  determines the critical behavior of the bulk, while  $\beta_s$  controls the surface coupling. Finally, we have introduced a symmetry-breaking boundary field  $\vec{h}_s = h_s \vec{e}_N$  along the  $N$ -th direction.

For  $\lambda \rightarrow \infty$ , the Hamiltonian (2) reduces to the hard spin  $O(N)$  model. In the  $(\beta, \lambda)$  plane, the bulk exhibits a second-order transition line in the  $O(N)$  UC [45, 52, 53]. For  $N = 2$  the model is *improved* for  $\lambda = 2.15(5)$  [53], i.e., the leading bulk irrelevant scaling field with dimension  $y_i = -0.789(4)$  [54] is suppressed. At  $\lambda = 2.15$  the model is critical for  $\beta = 0.50874988(6)$  [55]. For  $N = 3$  the model is improved for  $\lambda = 5.17(11)$  and the suppressed leading irrelevant scaling field has dimension  $y_i = -0.759(2)$  [56]. At  $\lambda = 5.2$ , the model is critical at  $\beta = 0.68798521(8)$  [56]. Improved models are instrumental to obtain accurate results in critical phenomena [45], in particular in boundary critical phenomena [43, 51, 57–65], because the broken translational invariance generically gives rise to additional scaling corrections, which cumulate to those arising from bulk irrelevant operators. The latter are suppressed for improved lattice models, hence enabling a more accurate analysis.

In the MC simulations presented here we set  $\beta$  and  $\lambda$  to the central value of the bulk critical point in the improved models, and  $\beta_s = \beta$ . Note that the boundary parameters  $\beta_s, h_s$  are chosen to be identical on the two surfaces: this realizes the normal UC on both surfaces, and allows us to compute improved estimators of surface observables by averaging them over the two surfaces. The geometry is fixed by  $L = L_{\parallel}$ . MC simulations are performed by combining Metropolis, overrelaxation, and Wolff single-cluster updates [66]; details of the algorithm are reported in Ref. [43], and the implementation of the Wolff algorithm in the presence of a symmetry-breaking surface field is discussed in Ref. [51]. The inclusion of a boundary field

breaks the  $O(N)$  symmetry to  $O(N-1)$ . Accordingly, we distinguish the components of  $\vec{\phi}$  defining  $\vec{\phi} \equiv (\vec{\varphi}, \sigma)$ , where  $\sigma$  is the component parallel to the surface field, and  $\vec{\varphi}$  is a  $(N-1)$ -component vector orthogonal to it. As discussed below, we measure the magnetization profile  $\langle \sigma \rangle$  and various surface-surface and surface-bulk two-point functions.

Besides the model realizing the normal UC, we also perform some MC simulations of the  $\phi^4$  model with periodic BCs, with the aim of determining the bulk field normalization. In this case, the Hamiltonian is as in Eq. (2), without the surface terms.

*Normal universality class.*— In this section, we discuss the normal surface UC of the  $O(N)$  model in  $d = 3$ . Unless otherwise stated, all operators in this section (e.g.  $\vec{\phi} = (\vec{\varphi}, \sigma)$ ) denote continuum fields; when referring to fields of the lattice model (2), we use the subscript “lat”. To leading order, the bulk field  $\vec{\phi}_{\text{lat}} \propto \vec{\phi}$ . The boundary operator spectrum contains two “protected” operators whose existence is mandated by bulk conservation laws and whose scaling dimensions are known exactly [49, 50, 67]:

- The “tilt” operator  $t^i$  of dimension  $\hat{\Delta}_t = d - 1 = 2$ , which is an  $O(N-1)$  vector ( $i = 1 \dots N - 1$ ). This operator is induced on the boundary when the symmetry breaking field  $\vec{h}_s$  is tilted — thus the nomenclature [68].
- The displacement operator  $D$  of dimension  $\hat{\Delta}_D = d = 3$ , which is an  $O(N-1)$  scalar. Perturbing the boundary with this operator is equivalent to moving the location of the boundary, justifying the name “displacement”.

These are believed to be the two lightest boundary operators. In particular, on the lattice the boundary field  $\varphi_{\text{lat}}^i \propto t^i$ . The boundary operator product expansion (OPE) holds for  $z \rightarrow 0$ :

$$\sigma(\mathbf{x}, z) = \frac{a_\sigma}{(2z)^{\Delta_\phi}} + b_D (2z)^{3-\Delta_\phi} D(\mathbf{x}) + \dots, \\ \varphi^i(\mathbf{x}, z) = b_t (2z)^{2-\Delta_\phi} t^i(\mathbf{x}) + \dots, \quad (3)$$

where  $\Delta_\phi$  is the bulk scaling dimension of  $\vec{\phi}$ . The coefficients  $a_\sigma, b_t, b_D$  are universal, assuming that the bulk and boundary operators are normalized.  $a_\sigma$  and  $b_t$  will be the main target of this paper — as was shown in Ref. [42], their ratio controls the existence and universal properties of the extraordinary-log phase (in the absence of a boundary magnetic field). Defining

$$\alpha \equiv \frac{\pi}{2} \left( \frac{a_\sigma}{4\pi b_t} \right)^2 - \frac{N-2}{2\pi}, \quad (4)$$

the extraordinary-log phase exists when  $\alpha > 0$ . Further, the exponent  $q$  in Eq. (1) is given by

$$q = \frac{N-1}{2\pi\alpha}. \quad (5)$$

We extract  $a_\sigma$  and  $b_t$  from the following correlators, which in a semi-infinite geometry take the form:

$$\langle \sigma(z) \rangle = \frac{a_\sigma}{(2z)^{\Delta_\phi}}, \quad \langle t^i(0) \varphi^j(\mathbf{x}, z) \rangle = \delta_{ij} b_t \frac{(2z)^{2-\Delta_\phi}}{(\mathbf{x}^2 + z^2)^2}. \quad (6)$$




FIG. 1. Bulk two-point function for (a)  $N = 2$  and (b)  $N = 3$ , rescaled to the large-distance decay exponent  $2\Delta_\phi$ . Error bars are smaller than symbol size.

The bulk field  $\phi^a$ ,  $a = 1 \dots N$ , is normalized so that in an infinite geometry  $\langle \phi^a(x) \phi^b(0) \rangle = \delta^{ab} x^{-2\Delta_\phi}$ , while  $t^i$  is normalized so that in a semi-infinite geometry  $\langle t^i(\mathbf{x}) t^j(0) \rangle = \delta^{ij} \mathbf{x}^{-4}$ . Thus, in a lattice model, to fix the normalizations above and to find  $a_\sigma$ ,  $b_t$  we will need to measure four different correlators. On the lattice, we expect both finite size scaling corrections and corrections to scaling [69].

*Results.*— We study first the XY UC. In order to determine the normalization of the bulk field, we have simulated the  $\phi^4$  model for  $N = 2$ , with periodic BCs and at the critical point, for lattice sizes  $L = 32 - 192$ . In Fig. 1(a) we show the two-point function  $\langle \vec{\phi}(x) \cdot \vec{\phi}(0) \rangle$ , rescaled to the expected decay  $x^{-2\Delta_\phi}$ . Here and below we use  $\Delta_\phi = 0.519088(22)$ ,  $\Delta_\epsilon = 1.51136(22)$  [70]. We fit the MC data to

$$\langle \vec{\phi}(x) \cdot \vec{\phi}(0) \rangle = \mathcal{N}_{\text{bulk}} x^{-2\Delta_\phi} \left[ 1 + B_\epsilon \left( \frac{x}{L} \right)^{\Delta_\epsilon} + C x^{-2} \right], \quad (7)$$

where the leading finite-size correction  $\propto L^{-\Delta_\epsilon}$  is due to the energy operator in the OPE of  $\phi^a \times \phi^a$ , while the correction  $\propto x^{-2}$  comes from the next-to-leading irrelevant operator in the action and from descendant operators in the expansion of the lattice field  $\vec{\phi}_i$  in terms of continuum fields [69]. Eq. (7) holds for  $(x/L) \ll 1$  and  $x \gtrsim x_0$ , with  $x_0$  a nonuniversal length governing the two-point function at short distance. The analysis of various fits to Eq. (7) allows us to infer [69]:

$$\mathcal{N}_{\text{bulk}} = 0.28152(15), \quad B_\epsilon = 2.758(10). \quad (8)$$

Next, we study the surface critical behavior. As discussed above, in order to implement the normal UC, we simulated the  $\phi^4$  model at the critical point with open BCs and a symmetry-breaking surface field. Preliminary MC data suggested a reduction of corrections to scaling for a surface field  $h_s = 1.5\beta_s$ . Our MC simulations reported below have thus been done at this value of  $h_s$ , for lattice sizes  $L = 32 - 192$ . To extract the normalization of the surface field component  $\varphi$ , we have computed its two-point function along the surface. We show it in Fig. 2(a), rescaled to its expected large-distance decay exponent 4. In this case finite-size corrections are rather small, such that we fit the data to [71]

$$\langle \vec{\varphi}(\mathbf{x}) \cdot \vec{\varphi}(0) \rangle = \mathcal{N}_\varphi \mathbf{x}^{-4} (1 + C \mathbf{x}^{-2}), \quad (9)$$




FIG. 2. Plots of surface observables. (a) and (b): two-point functions of the surface field component  $\varphi$ , for  $N = 2$  and  $N = 3$ , rescaled to the large-distance decay exponent 4. (c) and (d): order-parameter profile as a function of the distance from the surface, rescaled to the large-distance decay exponent  $\Delta_\phi$ . (e) and (f): surface-bulk correlation functions of the field component  $\varphi$ , rescaled to the large-distance decay exponent  $2 + \Delta_\phi$ .

where, analogous to Eq. (7), the leading correction to scaling  $\propto \mathbf{x}^{-2}$  originates from the expansion of the lattice operator in terms of continuum ones. Fits to Eq. (9) deliver [69]

$$\mathcal{N}_\varphi = 0.328(3). \quad (10)$$

In Fig. 2(c), we show the magnetization profile  $\langle \sigma(z) \rangle$  as a function of the distance from the surface  $z$ , and rescaled to its asymptotic decay exponent  $\Delta_\phi$ . For this quantity, scaling and finite-size corrections are relevant and we fit MC data to

$$\langle \sigma(z) \rangle = M_\sigma (z + z_0)^{-\Delta_\phi} \left[ 1 + B_\sigma \left( \frac{z + z_0}{L} \right)^3 \right], \quad (11)$$

where the perturbation of the surface action with the displacement operator  $D$  produces the replacement  $z \rightarrow z + z_0$ , with  $z_0$  a nonuniversal constant, and the leading finite-size correction originates from the OPE (3) [69]; the latter is also known as distant wall correction [6, 72, 73]. Fits to Eq. (11) allow us to estimate [69]

$$M_\sigma = 0.7540(3), \quad B_\sigma = 1.21(6), \quad z_0 = 1.018(6). \quad (12)$$

In Fig. 2(e) we show the surface-bulk correlation function  $\langle \varphi(0) \varphi(0, z) \rangle$  of the field component  $\varphi$ , where one point is

on the surface and the other a distance  $z$  away from the surface, so that the vector separating the two points is orthogonal to the surface; the correlations are rescaled to the large-distance decay exponent  $2 + \Delta_\phi$ . These correlations are affected by significant scaling corrections, while finite-size corrections, though not negligible, are smaller than in the case of  $\langle \sigma(z) \rangle$ . Together with the relatively fast large-distance decay,  $\sim z^{-2-\Delta_\phi}$ , this makes the analysis of the surface-bulk correlations more involved. A good Ansatz for the MC data is

$$\langle \vec{\varphi}(0) \cdot \vec{\varphi}(0, z) \rangle = M_\varphi (z + z_0)^{-2-\Delta_\phi} \left[ 1 + B_\varphi \left( \frac{z + z_0}{L} \right)^3 + C(z + z_0)^{-2} \right], \quad (13)$$

where we have included the corrections considered in Eq. (9) and Eq. (11) [69]. To avoid overfitting, in fits to Eq. (13) we plug in the result for  $z_0$  of Eq. (12), varying its value within one error bar quoted there. From the various fits we estimate

$$M_\varphi = 0.3146(8), \quad B_\varphi = -0.7(2). \quad (14)$$

In the analysis of the MC data for the Heisenberg normal UC, we proceed analogous to the case  $N = 2$ , using the critical exponents  $\Delta_\phi = 0.518920(25)$  and  $\Delta_\epsilon = 1.5948(2)$  [56]. To extract the normalization of the bulk field  $\phi$ , we simulated the  $\phi^4$  model for  $N = 3$ , periodic BCs, and at the critical point, for lattice sizes  $L = 32 - 192$ . In Fig. 1(b) we show the two-point function of  $\phi$ , rescaled to the expected decay  $x^{-2\Delta_\phi}$ . From fits of the correlations to Eq. (7) we obtain [69]

$$\mathcal{N}_{\text{bulk}} = 0.31230(15), \quad B_\epsilon = 2.432(7). \quad (15)$$

Concerning the surface critical behavior, preliminary MC simulations of the model (2) with  $N = 3$  suggested a reduction of subleading corrections for a surface field  $h_s = 1.4\beta_s$ . Here, we present results for this choice of  $h_s$ , and lattice sizes  $L = 32 - 192$ . In Fig. 2(b) we show the surface correlations. Fits of  $\langle \vec{\varphi}(\mathbf{x}) \cdot \vec{\varphi}(0) \rangle$  to Eq. (9) allow us to estimate [69]

$$\mathcal{N}_\varphi = 0.481(3). \quad (16)$$

Fits of the order-parameter profile  $\langle \sigma(z) \rangle$ , shown in Fig. 2(d), deliver the following results:

$$M_\sigma = 0.7062(2), \quad B_\sigma = 1.07(5), \quad z_0 = 1.031(4). \quad (17)$$

In Fig. 2(f) we show the surface-bulk correlation function  $\langle \vec{\varphi}(0) \cdot \vec{\varphi}(0, z) \rangle$ . We fit it to Eq. (13), employing the estimate of  $z_0$  given in Eq. (17). From the various fits we obtain [69]

$$M_\varphi = 0.4674(8), \quad B_\varphi = -0.7(1). \quad (18)$$

*Discussion.*— According to the discussion above of the normal UC and of the scaling forms in Ref. [69], the results of our scaling analysis of MC data allow us to extract universal amplitudes  $a_\sigma, b_t$  of the normal UC via

$$a_\sigma = \frac{2^{\Delta_\phi} M_\sigma}{\sqrt{\mathcal{N}_{\text{bulk}}/N}}, \quad b_t = \frac{2^{\Delta_\phi} M_\varphi}{4\sqrt{N-1}\sqrt{\mathcal{N}_{\text{bulk}}/N}\sqrt{\mathcal{N}_\varphi}}. \quad (19)$$

$N$	$a_\sigma$	$b_t$	$b_D$	$\alpha$	$\alpha_{\text{eo}}$
1	2.60(5)		0.244(8)		
2	2.880(2)	0.525(4)		0.300(5)	0.27(2)[44]
3	3.136(2)	0.529(3)		0.190(4)	0.15(2)[43]

TABLE I. Final results: universal amplitudes  $a_\sigma, b_t, b_D$  in Eq. (3) together with the corresponding value of  $\alpha$ , Eq. (4). We also tabulate  $\alpha_{\text{eo}}$  found in Refs. [43, 44] by direct MC simulations of the extraordinary region. For  $N = 1$  we present the values of  $a_\sigma, b_D$  extracted from MC results of Refs. [51, 58].

$a_\sigma$  is obtained from the amplitude of the order-parameter profile  $M_\sigma$  (11) and the normalization of the bulk field  $\mathcal{N}_{\text{bulk}}$  (7), while  $b_t$  is obtained in terms of the amplitude of the surface-bulk correlations  $M_\varphi$  (13), and the bulk and surface normalizations  $\mathcal{N}_{\text{bulk}}$ , (7),  $\mathcal{N}_\varphi$ , (9). We collect our results for  $a_\sigma, b_t$  in Table I, including the value of  $\alpha$  obtained from  $a_\sigma$  and  $b_t$  via Eq. (4). For both  $N = 2$  and  $N = 3$ ,  $\alpha > 0$ , which indicates that the extraordinary-log UC exists, in accord with MC results of Refs. [43, 44]. Ref. [42] predicts that  $\alpha$  controls various universal properties of the extraordinary-log phase, including the exponent  $q$  in Eq. (1), which is related to  $\alpha$  via Eq. (5).  $\alpha$  found here agrees well with  $\alpha$  extracted from direct MC simulations of the extraordinary region [43, 44], listed in Table I as  $\alpha_{\text{eo}}$ . This provides a highly non-trivial check of the theory in Ref. [42]. As pointed out in Ref. [43], the error bar on  $\alpha_{\text{eo}}$  should be taken with a grain of salt given the difficulty of fitting to the form in Eq. (1) and the presence of subleading logarithmic corrections. Thus, we expect the method for determining  $\alpha$  presented here to be more reliable than directly simulating the extraordinary-log phase. We also present results for the coefficients  $a_\sigma, b_D$  in Eq. (3) for the normal UC of the Ising model ( $N = 1$ ) [69], extracted from MC studies in Refs. [51, 58]. A numerical conformal bootstrap study of the normal UC of the O(N) model with  $N \geq 2$  was conducted in parallel to our work [68]. Our results for  $a_\sigma$  and  $b_t$  are within the bounds produced by positive bootstrap and agree reasonably well with the approximate truncated bootstrap results. For  $N = 1$ , both  $a_\sigma$  and  $b_D$  in Table I agree within error bars with the truncated bootstrap findings of Ref. [74].

We conclude by outlining some possible future directions. It will be interesting to extend the calculations presented here to the O(N) model with  $N > 3$ , with an eye to determining the critical value  $N_c$  where the extraordinary-log UC disappears.  $N = 4$  is a natural first target since bootstrap calculations [68], as well as previous MC simulations [75], suggest that the extraordinary transition still exists in this case. Another extension is to study the free energy density for the normal UC in the geometry considered here, which combined with the coefficient  $B_\sigma$  in Eq. (11) and  $a_\sigma$ , allows one to determine the OPE coefficient  $b_D$  in Eq. (3), as well as the universal coefficient  $C_D$ , characterizing the boundary OPE of the energy-momentum tensor  $T_{zz} \xrightarrow{z \rightarrow 0} -\sqrt{C_D} D$  [67]. (In fact, this is how for the Ising model  $b_D$  in Table I was obtained. [76]) Further interesting avenues for future research would be



to consider the  $N \rightarrow 0$  limit [77–82], which describes the physics of dilute polymers [1, 45, 83], and  $O(N)$  loop models [84, 85], which provide an extension of the standard  $O(N)$  model to noninteger values of  $N$ .

We are very grateful to Marco Meineri for discussions. M.M. thanks Ilya Gruzberg, Abijith Krishnan, Marco Meineri and Jay Padaysi for a collaboration on a related project. We thank Jian-Ping Lv for useful communications. F.P.T. is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)–Project No. 414456783. We gratefully acknowledge the Gauss Centre for Supercomputing e.V. for funding this project by providing computing time through the John von Neumann Institute for Computing (NIC) on the GCS Supercomputer JUWELS at Jülich Supercomputing Centre (JSC) [86]. M.M. is supported by the National Science Foundation under grant number DMR-1847861.

---

\* francesco.parisentoldin@physik.uni-wuerzburg.de

† mmetlits@mit.edu

- [1] J. Cardy, *Scaling and Renormalization in Statistical Physics* (Cambridge University Press, Cambridge, 1996).
- [2] H. Dosch, *Critical Phenomena at Surfaces and Interfaces: Evanescent X-Ray and Neutron Scattering*, Springer Tracts in Modern Physics (Springer Berlin Heidelberg, Berlin, 2006).
- [3] K. Binder, Critical behavior at surfaces, in *Phase Transitions and Critical Phenomena*, Phase Transitions and Critical Phenomena, Vol. 8, edited by C. Domb and J. L. Lebowitz (Academic Press, London, 1983) p. 1.
- [4] H. W. Diehl, Field-theoretical approach to critical behaviour at surfaces, in *Phase Transitions and Critical Phenomena*, Phase Transitions and Critical Phenomena, Vol. 10, edited by C. Domb and J. L. Lebowitz (Academic Press, London, 1986) p. 75.
- [5] M. Pleimling, Critical phenomena at perfect and non-perfect surfaces, *J. Phys. A: Math. Gen.* **37**, R79 (2004), [cond-mat/0402574](#).
- [6] M. E. Fisher and P.-G. de Gennes, Phénomènes aux parois dans un mélange binaire critique, *C. R. Acad. Sci. Paris Ser. B* **287**, 207 (1978).
- [7] M. Krech, *The Casimir Effect in Critical Systems* (World Scientific, London, 1994).
- [8] M. Krech, Fluctuation-induced forces in critical fluids, *J. Phys.: Condens. Matter* **11**, R391 (1999), [cond-mat/9909413](#).
- [9] Ī. Brankov, D. Danchev, and N. Tonchev, *Theory of Critical Phenomena in Finite-size Systems: Scaling and Quantum Effects*, Series in modern condensed matter physics (World Scientific, Singapore, 2000).
- [10] A. Gambassi, The Casimir effect: From quantum to critical fluctuations, *J. Phys.: Conf. Ser.* **161**, 012037 (2009), [arXiv:0812.0935 \[cond-mat.stat-mech\]](#).
- [11] A. Gambassi and S. Dietrich, Critical Casimir forces steered by patterned substrates, *Soft Matter* **7**, 1247 (2011), [arXiv:1011.1831 \[cond-mat.soft\]](#).
- [12] A. Maciołek and S. Dietrich, Collective behavior of colloids due to critical Casimir interactions, *Rev. Mod. Phys.* **90**, 045001 (2018), [arXiv:1712.06678 \[cond-mat.soft\]](#).
- [13] D. McAvity and H. Osborn, Conformal field theories near a boundary in general dimensions, *Nucl. Phys.* **B455**, 522 (1995), [arXiv:cond-mat/9505127 \[cond-mat\]](#).
- [14] P. Liendo, L. Rastelli, and B. C. van Rees, The bootstrap program for boundary  $CFT_d$ , *JHEP* **07**, 113, [arXiv:1210.4258 \[hep-th\]](#).
- [15] F. Gliozzi, P. Liendo, M. Meineri, and A. Rago, Boundary and Interface CFTs from the Conformal Bootstrap, *JHEP* **05**, 036, [arXiv:1502.07217 \[hep-th\]](#).
- [16] M. Billó, V. Gonçalves, E. Lauria, and M. Meineri, Defects in conformal field theory, *JHEP* **04**, 091, [arXiv:1601.02883 \[hep-th\]](#).
- [17] P. Liendo and C. Meneghelli, Bootstrap equations for  $\mathcal{N} = 4$  SYM with defects, *JHEP* **01**, 122, [arXiv:1608.05126 \[hep-th\]](#).
- [18] E. Lauria, M. Meineri, and E. Trevisani, Radial coordinates for defect CFTs, *JHEP* **11**, 148, [arXiv:1712.07668 \[hep-th\]](#).
- [19] D. Mazáč, L. Rastelli, and X. Zhou, An analytic approach to  $BCFT_d$ , *JHEP* **12**, 004, [arXiv:1812.09314 \[hep-th\]](#).
- [20] A. Kaviraj and M. F. Paulos, The functional bootstrap for boundary CFT, *JHEP* **04**, 135, [arXiv:1812.04034 \[hep-th\]](#).
- [21] P. Dey, T. Hansen, and M. Shpot, Operator expansions, layer susceptibility and two-point functions in BCFT, *JHEP* **12**, 051, [arXiv:2006.11253 \[hep-th\]](#).
- [22] C. Behan, L. Di Pietro, E. Lauria, and B. C. Van Rees, Bootstrapping boundary-localized interactions, *JHEP* **12**, 182, [arXiv:2009.03336 \[hep-th\]](#).
- [23] A. Gimenez-Grau, P. Liendo, and P. van Vliet, Superconformal boundaries in  $4 - \epsilon$  dimensions, *JHEP* **04**, 167, [arXiv:2012.00018 \[hep-th\]](#).
- [24] T. Grover and A. Vishwanath, Quantum criticality in topological insulators and superconductors: Emergence of strongly coupled majoranas and supersymmetry, arXiv e-prints (2012), [arXiv:1206.1332 \[cond-mat.str-el\]](#).
- [25] M. Barkeshli, M. Mulligan, and M. P. A. Fisher, Particle-hole symmetry and the composite fermi liquid, *Physical Review B* **92**, 165125 (2015), [arXiv:1502.05404 \[cond-mat.str-el\]](#).
- [26] M. Barkeshli and X.-L. Qi, Synthetic topological qubits in conventional bilayer quantum hall systems, *Physical Review X* **4**, 041035 (2014), [arXiv:1302.2673 \[cond-mat.mes-hall\]](#).
- [27] J. Cano, M. Cheng, M. Barkeshli, D. J. Clarke, and C. Nayak, Chirality-protected majorana zero modes at the gapless edge of abelian quantum hall states, *Physical Review B* **92**, 195152 (2015), [arXiv:1505.07825 \[cond-mat.str-el\]](#).
- [28] T. Scaffidi, D. E. Parker, and R. Vasseur, Gapless symmetry-protected topological order, *Phys. Rev. X* **7**, 041048 (2017), [arXiv:1705.01557 \[cond-mat.str-el\]](#).
- [29] D. E. Parker, T. Scaffidi, and R. Vasseur, Topological luttinger liquids from decorated domain walls, *Physical Review B* **97**, 165114 (2018), [arXiv:1711.09106 \[cond-mat.str-el\]](#).
- [30] R. Verresen, R. Thorngren, N. G. Jones, and F. Pollmann, Gapless topological phases and symmetry-enriched quantum criticality, *Phys. Rev. X* **11**, 041059 (2021), [arXiv:1905.06969 \[cond-mat.str-el\]](#).
- [31] R. Verresen, Topology and edge states survive quantum criticality between topological insulators, arXiv e-prints (2020), [arXiv:2003.05453 \[cond-mat.str-el\]](#).
- [32] R. Thorngren, A. Vishwanath, and R. Verresen, Intrinsically gapless topological phases, *Phys. Rev. B* **104**, 075132 (2021), [arXiv:2008.06638 \[cond-mat.str-el\]](#).
- [33] T. Suzuki and M. Sato, Gapless edge states and their stability in two-dimensional quantum magnets, *Phys. Rev. B* **86**, 224411 (2012), [arXiv:1209.3097 \[cond-mat.str-el\]](#).
- [34] L. Zhang and F. Wang, Unconventional Surface Critical Behavior Induced by a Quantum Phase Transition from the Two-Dimensional Affleck-Kennedy-Lieb-Tasaki Phase to a Néel-Ordered Phase, *Phys. Rev. Lett.* **118**, 087201 (2017),

- arXiv:1611.06477 [cond-mat.str-el].
- [35] C. Ding, L. Zhang, and W. Guo, Engineering Surface Critical Behavior of  $(2+1)$ -Dimensional  $O(3)$  Quantum Critical Points, *Phys. Rev. Lett.* **120**, 235701 (2018), arXiv:1801.10035 [cond-mat.str-el].
- [36] L. Weber, F. Parisen Toldin, and S. Wessel, Nonordinary edge criticality of two-dimensional quantum critical magnets, *Phys. Rev. B* **98**, 140403(R) (2018), arXiv:1804.06820 [cond-mat.str-el].
- [37] L. Weber and S. Wessel, Nonordinary criticality at the edges of planar spin-1 heisenberg antiferromagnets, *Phys. Rev. B* **100**, 054437 (2019), arXiv:1906.07051 [cond-mat.str-el].
- [38] C.-M. Jian, Y. Xu, X.-C. Wu, and C. Xu, Continuous Néel-VBS quantum phase transition in non-local one-dimensional systems with  $SO(3)$  symmetry, *SciPost Phys.* **10**, 033 (2021), arXiv:2004.07852 [cond-mat.str-el].
- [39] W. Zhu, C. Ding, L. Zhang, and W. Guo, Surface critical behavior of coupled Haldane chains, *Phys. Rev. B* **103**, 024412 (2021), arXiv:2010.10920 [cond-mat.str-el].
- [40] L. Weber and S. Wessel, Spin versus bond correlations along dangling edges of quantum critical magnets, *Phys. Rev. B* **103**, L020406 (2021), arXiv:2010.15691 [cond-mat.str-el].
- [41] C. Ding, W. Zhu, W. Guo, and L. Zhang, Special Transition and Extraordinary Phase on the Surface of a  $(2+1)$ -Dimensional Quantum Heisenberg Antiferromagnet, arXiv:2110.04762 [cond-mat.str-el].
- [42] M. A. Metlitski, Boundary criticality of the  $O(N)$  model in  $d = 3$  critically revisited, arXiv:2009.05119 [cond-mat.str-el].
- [43] F. Parisen Toldin, Boundary critical behavior of the three-dimensional Heisenberg universality class, *Phys. Rev. Lett.* **126**, 135701 (2021), arXiv:2012.00039 [cond-mat.stat-mech].
- [44] M. Hu, Y. Deng, and J.-P. Lv, Extraordinary-Log Surface Phase Transition in the Three-Dimensional X Y Model, *Phys. Rev. Lett.* **127**, 120603 (2021), arXiv:2104.05152 [cond-mat.stat-mech].
- [45] A. Pelissetto and E. Vicari, Critical phenomena and renormalization-group theory, *Phys. Rep.* **368**, 549 (2002), cond-mat/0012164.
- [46] M. Krech, Surface scaling behavior of isotropic heisenberg systems: Critical exponents, structure factor, and profiles, *Phys. Rev. B* **62**, 6360 (2000), arXiv:cond-mat/0006448 [cond-mat.stat-mech].
- [47] Y. Deng, H. W. J. Blöte, and M. P. Nightingale, Surface and bulk transitions in three-dimensional  $O(n)$  models, *Phys. Rev. E* **72**, 016128 (2005), cond-mat/0504173.
- [48] Earlier a hint of the special transition for  $N = 3$  was seen in Ref. [47].
- [49] A. J. Bray and M. A. Moore, Critical behaviour of semi-infinite systems, *J. Phys. A: Math. Gen.* **10**, 1927 (1977).
- [50] T. W. Burkhardt and J. L. Cardy, Surface critical behaviour and local operators with boundary-induced critical profiles, *J. Phys. A: Math. Gen.* **20**, L233 (1987).
- [51] F. Parisen Toldin and S. Dietrich, Critical Casimir forces and adsorption profiles in the presence of a chemically structured substrate, *J. Stat. Mech.* **P11003**, (2010), arXiv:1007.3913 [cond-mat.stat-mech].
- [52] M. Campostrini, M. Hasenbusch, A. Pelissetto, P. Rossi, and E. Vicari, Critical exponents and equation of state of the three-dimensional Heisenberg universality class, *Phys. Rev. B* **65**, 144520 (2002), cond-mat/0110336.
- [53] M. Campostrini, M. Hasenbusch, A. Pelissetto, and E. Vicari, Theoretical estimates of the critical exponents of the superfluid transition in  $^4\text{He}$  by lattice methods, *Phys. Rev. B* **74**, 144506 (2006), cond-mat/0605083.
- [54] M. Hasenbusch, Monte Carlo study of an improved clock model in three dimensions, *Phys. Rev. B* **100**, 224517 (2019), arXiv:1910.05916 [cond-mat.stat-mech].
- [55] F. Parisen Toldin, Finite-Size Scaling at fixed Renormalization-Group invariant, arXiv:2112.00392 [cond-mat.stat-mech].
- [56] M. Hasenbusch, Monte carlo study of a generalized icosahedral model on the simple cubic lattice, *Phys. Rev. B* **102**, 024406 (2020), arXiv:2005.04448 [cond-mat.stat-mech].
- [57] M. Hasenbusch, The thermodynamic Casimir effect in the neighbourhood of the  $\lambda$ -transition: a Monte Carlo study of an improved three-dimensional lattice model, *J. Stat. Mech.* **P07031**, (2009), arXiv:0905.2096 [cond-mat.stat-mech].
- [58] M. Hasenbusch, Thermodynamic Casimir effect for films in the three-dimensional Ising universality class: Symmetry-breaking boundary conditions, *Phys. Rev. B* **82**, 104425 (2010), arXiv:1005.4749 [cond-mat.stat-mech].
- [59] M. Hasenbusch, Thermodynamic Casimir force: A Monte Carlo study of the crossover between the ordinary and the normal surface universality class, *Phys. Rev. B* **83**, 134425 (2011), arXiv:1012.4986 [cond-mat.stat-mech].
- [60] M. Hasenbusch, Monte Carlo study of surface critical phenomena: The special point, *Phys. Rev. B* **84**, 134405 (2011), arXiv:1108.2425 [cond-mat.stat-mech].
- [61] M. Hasenbusch, Thermodynamic Casimir effect: Universality and corrections to scaling, *Phys. Rev. B* **85**, 174421 (2012), arXiv:1202.6206 [cond-mat.stat-mech].
- [62] F. Parisen Toldin, M. Tröndle, and S. Dietrich, Critical Casimir forces between homogeneous and chemically striped surfaces, *Phys. Rev. E* **88**, 052110 (2013), arXiv:1303.6104 [cond-mat.stat-mech].
- [63] F. Parisen Toldin, Critical Casimir force in the presence of random local adsorption preference, *Phys. Rev. E* **91**, 032105 (2015), arXiv:1308.5220 [cond-mat.stat-mech].
- [64] F. Parisen Toldin, M. Tröndle, and S. Dietrich, Line contribution to the critical Casimir force between a homogeneous and a chemically stepped surface, *J. Phys.: Condens. Matter* **27**, 214010 (2015), arXiv:1409.5536 [cond-mat.stat-mech].
- [65] F. Parisen Toldin, F. F. Assaad, and S. Wessel, Critical behavior in the presence of an order-parameter pinning field, *Phys. Rev. B* **95**, 014401 (2017), arXiv:1607.04270 [cond-mat.stat-mech].
- [66] U. Wolff, Collective Monte Carlo updating for spin systems, *Phys. Rev. Lett.* **62**, 361 (1989).
- [67] J. L. Cardy, Universal critical-point amplitudes in parallel-plate geometries, *Phys. Rev. Lett.* **65**, 1443 (1990).
- [68] J. Padayasi, A. Krishnan, M. Metlitski, I. Gruzberg, and M. Meineri, The extraordinary boundary transition in the 3d  $o(n)$  model via conformal bootstrap, arXiv:2111.03071 [cond-mat.str-el].
- [69] See Supplemental Material for a discussion of the scaling forms of various observables of the lattice model, technical details on the fits of statistically correlated data, and results of the fits of MC data.
- [70] S. M. Chester, W. Landry, J. Liu, D. Poland, D. Simmons-Duffin, N. Su, and A. Vichi, Carving out OPE space and precise  $O(2)$  model critical exponents, *J. High Energy Phys.* **2020**, 142 (2020), arXiv:1912.03324 [hep-th].
- [71] Even though  $\varphi$  has only a single component in the  $N = 2$  case, we write it as a vector here to set the normalization for  $N = 3$ .
- [72] E. Eisenriegler, M. Krech, and S. Dietrich, Absence of hyperuniversality in critical films, *Phys. Rev. Lett.* **70**, 619 (1993); Erratum: “Absence of hyperuniversality in critical films” [*Phys. Rev. Lett.* **70**, 619 (1993)], *Phys. Rev. Lett.* **70**, 2051(E) (1993).
- [73] E. Eisenriegler and M. Stapper, Critical behavior near a symmetry-breaking surface and the stress tensor, *Phys. Rev. B*

- 50**, 10009 (1994).
- [74] F. Gliozzi, P. Liendo, M. Meineri, and A. Rago, Boundary and interface CFTs from the conformal bootstrap, *J. High Energy Phys.* **5**, 36 (2015), [arXiv:1502.07217 \[hep-th\]](#).
- [75] Y. Deng, Bulk and surface phase transitions in the three-dimensional  $O(4)$  spin model, *Phys. Rev. E* **73**, 056116 (2006).
- [76] In principle, it is also possible to determine  $b_D$  by a method similar to the one we used to find  $b_t$ : one extracts the normalization of  $D$  from the connected two-point function of  $\sigma$  on the boundary, and the coefficient  $b_D$  from the connected bulk-boundary two point function of  $\sigma$ . However, we found that due to the large scaling dimension  $\hat{\Delta}_D = 3$ , the boundary two-point function  $\langle \sigma(\mathbf{x})\sigma(0) \rangle_{\text{conn}}$  falls off very fast, so that the normalization is difficult to extract reliably.
- [77] K. De’Bell, T. Lookman, and S. G. Whittington, Analysis of exact enumeration data for self-avoiding walks attached to a surface, *Phys. Rev. A* **41**, 682 (1990).
- [78] K. De’Bell and T. Lookman, Surface phase transitions in polymer systems, *Rev. Mod. Phys.* **65**, 87 (1993).
- [79] D. Zhao, T. Lookman, and K. De’Bell, Crossover behavior for self-avoiding walks interacting with a surface, *Phys. Rev. A* **42**, 4591 (1990).
- [80] E. Eisenriegler, *Polymers Near Surfaces* (World Scientific, 1993).
- [81] R. Hegger and P. Grassberger, Chain polymers near an adsorbing surface, *J. Phys. A: Math. Gen.* **27**, 4069 (1994).
- [82] M. T. Batchelor and J. Cardy, Extraordinary transition in the two-dimensional  $o(n)$  model, *Nuclear Physics B* **506**, 553–564 (1997).
- [83] P.-G. de Gennes, *Scaling Concepts in Polymer Physics* (Cornell University Press, Ithaca, NY, 1979).
- [84] Q. Liu, Y. Deng, T. M. Garoni, and H. W. J. Blöte, The  $O(n)$  loop model on a three-dimensional lattice, *Nucl. Phys. B* **859**, 107 (2012), [arXiv:1112.5647 \[cond-mat.stat-mech\]](#).
- [85] E. Domany, D. Mukamel, B. Nienhuis, and A. Schwimmer, Duality relations and equivalences for models with  $O(N)$  and cubic symmetry, *Nucl. Phys. B* **190**, 279 (1981).
- [86] Jülich Supercomputing Centre, JUWELS: Modular Tier-0/1 Supercomputer at the JülichSupercomputing Centre, *Journal of large-scale research facilities* **5**, A135 (2019).
- [87] D. Simmons-Duffin, The Lightcone Bootstrap and the Spectrum of the 3d Ising CFT, *JHEP* **03**, 086, [arXiv:1612.08471 \[hep-th\]](#).
- [88] M. Hasenbusch, Universal amplitude ratios in the three-dimensional Ising universality class, *Phys. Rev. B* **82**, 174434 (2010), [arXiv:1004.4983 \[cond-mat.stat-mech\]](#).
- [89] M. Campostrini, A. Pelissetto, P. Rossi, and E. Vicari, 25th-order high-temperature expansion results for three-dimensional Ising-like systems on the simple-cubic lattice, *Phys. Rev. E* **65**, 066127 (2002), [cond-mat/0201180](#).
- [90] A. J. Bray, Dispersion-theory approach to the correlation function for critical scattering, *Phys. Rev. B* **14**, 1248 (1976).
- [91] R. F. Chang, H. Burstyn, and J. V. Sengers, Correlation function near the critical mixing point of a binary liquid, *Phys. Rev. A* **19**, 866 (1979).
- [92] P. Damay, F. Leclercq, and P. Chieux, Critical scattering function in a binary fluid mixture: A study of sodium-deuteroammonia solution at the critical concentration by small-angle neutron scattering, *Phys. Rev. B* **40**, 4696 (1989).
- [93] P. Damay, F. Leclercq, R. Magli, F. Formisano, and P. Lindner, Universal critical-scattering function: An experimental approach, *Phys. Rev. B* **58**, 12038 (1998).
- [94] V. Martín-Mayor, A. Pelissetto, and E. Vicari, Critical structure factor in ising systems, *Phys. Rev. E* **66**, 026112 (2002).
- [95] A. P. Young, *Everything You Wanted to Know About Data Analysis and Fitting but Were Afraid to Ask*, Springer-Briefs in Physics (Springer International Publishing, 2015) [arXiv:1210.3781](#).
- [96] C. Michael, Fitting correlated data, *Phys. Rev. D* **49**, 2616 (1994), [arXiv:hep-lat/9310026 \[hep-lat\]](#).
- [97] D. Seibert, Undesirable effects of covariance matrix techniques for error analysis, *Phys. Rev. D* **49**, 6240 (1994), [arXiv:hep-lat/9305014 \[hep-lat\]](#).