



# CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

## Energy Spectrum of Two-Dimensional Acoustic Turbulence

Adam Griffin, Giorgio Krstulovic, Victor S. L'vov, and Sergey Nazarenko

Phys. Rev. Lett. **128**, 224501 — Published 3 June 2022

DOI: [10.1103/PhysRevLett.128.224501](https://doi.org/10.1103/PhysRevLett.128.224501)

# Energy spectrum of two-dimensional acoustic turbulence

Adam Griffin,<sup>1</sup> Giorgio Krstulovic,<sup>2,\*</sup> Victor S. L'vov,<sup>3</sup> and Sergey Nazarenko<sup>1</sup>

<sup>1</sup>*Institut de Physique de Nice, Université Côte d'Azur, CNRS, Nice, France*

<sup>2</sup>*Université Cte d'Azur, Observatoire de la Cte d'Azur, CNRS, Laboratoire Lagrange, Boulevard de l'Observatoire CS 34229 - F 06304 NICE Cedex 4, France.*

<sup>3</sup>*Dept. of Chemical and Biological Physics, Weizmann Institute of Science, Rehovot 76100, Israel*

(Dated: April 8, 2022)

We report an exact unique constant-flux power-law analytical solution of the wave kinetic equation for the turbulent energy spectrum,  $E(k) = C_1 \sqrt{\varepsilon} a c_s / k$ , of acoustic waves in 2D with almost linear dispersion law,  $\omega_k = c_s k [1 + (ak)^2]$ ,  $ak \ll 1$ . Here  $\varepsilon$  is the energy flux over scales, and  $C_1$  is the universal constant which was found analytically. Our theory describes, for example, acoustic turbulence in 2D Bose-Einstein condensates. The corresponding 3D counterpart of turbulent acoustic spectrum was found over half a century ago, however, due to the singularity in 2D, no solution has been obtained until now. We show the spectrum  $E(k)$  is realizable in direct numerical simulations of forced-dissipated Gross-Pitaevskii equation in the presence of strong condensate.

Waves in nonlinear systems interact and transfer energy along scales in a cascade process [with a constant flux](#), creating an out-of-equilibrium state known as wave turbulence. When nonlinearity is small, the Weak-Wave Turbulence (WWT) theory provides a mathematical description of the system [1, 2]. The most common applications of this theory are capillary-gravity waves [3], Alfvén waves in magnetohydrodynamics [4], Langmuir waves in plasmas [5], inertial and internal waves in rotating stratified fluids [6, 7], Kelvin waves in vortices [8], elastic plates [9], gravitational waves [10] and density waves in Bose-Einstein condensates [11].

Note, that acoustic waves in ideal compressible fluids, one of the most common examples in Nature, are non-dispersive: their frequency  $\omega_{\mathbf{k}} = c_s k$  is linear in wave number  $k \equiv |\mathbf{k}|$ . Accordingly, the resonance conditions

$$\omega_{\mathbf{k}} = \omega_{\mathbf{k}_1} + \omega_{\mathbf{k}_2}, \quad \mathbf{k} = \mathbf{k}_1 + \mathbf{k}_2, \quad (1)$$

allow for interaction of the waves with parallel wave vectors only:  $\mathbf{k} \parallel \mathbf{k}_1 \parallel \mathbf{k}_2$  [and in the same direction](#) [2]. Therefore in the reference frame moving with the speed of sound  $c_s$  in the direction of  $\mathbf{k}$ ,  $\mathbf{k}_1$  and  $\mathbf{k}_2$  all wave packets are at the rest and their overlapping time  $\tau_{\text{OVR}}$  goes to infinity, allowing for wave steepening and breaking effects, which requires finite shock creation time  $\tau_{\text{sh}}$ . In the other words, the main assumption of the WWT theory, roughly formulated as  $\tau_{\text{OVR}} \ll \tau_{\text{sh}}$ , fails for dispersionless acoustic waves even even at small nonlinearity.

This has caused a long-standing debate whether WWT theory is applicable for their description [12], or alternatively, if acoustic waves should be viewed as a random collection of weak shocks leading to the Kadomtsev-Petviashvili spectrum [13]. There is a handwaving argument—yet unsupported by rigorous proof—that the theory applies to 3D acoustic turbulence because the divergence of wave packets in 3D space plays a role similar to the wave dispersion in preventing wave breaking. It was argued first in [14] and later in [15], that the WWT description is indeed possible for 3D weak acoustic tur-

bulence. However, the respective kinetic equation for the spectrum has to be modified so that interactions of non-collinear waves are described correctly. Concerning the 2D case, it is evident that WWT theory is not applicable in its classical form. Indeed, the main equation of the theory, the wave-kinetic equation, is singular and meaningless in the 2D case. The possibility of an alternative statistical description of weak non-dispersive acoustic 2D sound was claimed in [14], but it remains an unfinished task. Fortunately, in some important physical applications, 2D sound is regularized by weak dispersion effects, and the use of the classical WWT theory becomes again possible. One of such examples, which we will use in the present Letter, is the acoustic turbulence in 2D Bose-Einstein condensates (BEC).

Recent experiments with 3D BECs have succeeded in creating wave-turbulence states where measurements can be explained using the WWT theory in the fully dispersive limit [16, 17]. On the other hand, much interest has been devoted to studying the dynamics of vortices experimentally in 2D BECs, as such states are close to hydrodynamic 2D classical turbulence [18, 19]. Unfortunately, experiments of acoustic 2D BECs still lack and no theoretical predictions are available.

In this Letter, we develop the theory of weak wave turbulence of weakly dispersive 2D sound and obtain a modified wave kinetic equation. We derive a new stationary power-law flux spectrum of Kolmogorov-Zakharov type, which corresponds to a cascade of energy from large to small spatial scales. We then determine the flux spectrum exponent and the value of the prefactor constant analytically. This prediction can not be naively guessed by dimensional arguments, unlike the existing 3D results. Our theory is then tested and validated with numerical simulations of weakly nonlinear sound in the 2D forced-dissipated Gross-Pitaevskii (GP) equation (17) which is a dynamical model for BEC.

Let us consider the classical wave-kinetic equation describing evolution of the wave action spectrum  $n_{\mathbf{k}} =$

$n(\mathbf{k}, t)$  (with  $\mathbf{k}$  and  $t$  being the wavenumber and time) driven by three-wave resonant interactions [1, 2],

$$\frac{\partial n_{\mathbf{k}}}{\partial t} = \text{St}_{\mathbf{k}} \quad (2)$$

with the wave-collision integral

$$\text{St}_{\mathbf{k}} = 2\pi \int (\mathcal{R}_{12}^{\mathbf{k}} - \mathcal{R}_{\mathbf{k}2}^1 - \mathcal{R}_{\mathbf{k}1}^2) d\mathbf{k}_1 d\mathbf{k}_2, \quad (3a)$$

$$\begin{aligned} \mathcal{R}_{12}^{\mathbf{k}} &= \mathcal{N}_{12}^{\mathbf{k}} \delta_{12}^{\mathbf{k}} \delta(\omega_{12}^{\mathbf{k}}), \\ \mathcal{N}_{12}^{\mathbf{k}} &= |V_{12}^{\mathbf{k}}|^2 [n_{\mathbf{k}_1} n_{\mathbf{k}_2} - n_{\mathbf{k}} n_{\mathbf{k}_1} - n_{\mathbf{k}} n_{\mathbf{k}_2}], \\ \delta_{12}^{\mathbf{k}} &= \delta(\mathbf{k} - \mathbf{k}_1 - \mathbf{k}_2), \quad \delta(\omega_{12}^{\mathbf{k}}) = \delta(\omega_{\mathbf{k}} - \omega_{\mathbf{k}_1} - \omega_{\mathbf{k}_2}), \end{aligned} \quad (3b)$$

$\omega_{\mathbf{k}} = \omega(\mathbf{k})$  and  $V_{12}^{\mathbf{k}} \equiv V(\mathbf{k}, \mathbf{k}_1, \mathbf{k}_2)$  is a three-wave interaction amplitude, that depends on the particular type of waves.

The central object in turbulence is the 1D energy spectrum defined as the distribution of energy in  $k = |\mathbf{k}|$ , so that the energy (per unit of mass) is defined as  $E = \int E(k) dk$ . It is related to the waveaction spectrum as  $E(k) = 4\pi\omega_k k^2 n_k$  in 3D and  $E(k) = 2\pi\omega_k k n_k$  in 2D.

In this Letter, we consider turbulence of weakly dispersive acoustic waves for which

$$\omega_k = c_s k [1 + (ak)^2]. \quad (4)$$

Here  $a = \text{const}$  is a dispersion length such that  $ak \ll 1$ . The interaction amplitude  $V_{12}^{\mathbf{k}}$  in case acoustic hydrodynamics [2, 11, 12, 20] is given by

$$V_{12}^{\mathbf{k}} = V_0 \sqrt{k k_1 k_2}, \quad V_0 = \text{const}. \quad (5)$$

To find the energy spectra, note that Eq. (3) conserves the total energy of the system and, therefore, it can be rewritten as the following continuity equation for  $E(k)$ ,

$$\frac{\partial E(k)}{\partial t} + \frac{\partial \varepsilon_k}{\partial k} = 0, \quad \text{where } \varepsilon_k = - \int_{k_1 < k} \text{St}_{\mathbf{k}_1} \omega_{k_1} d\mathbf{k}_1. \quad (6)$$

Dimensionally, the energy flux  $\varepsilon(k) \propto \text{St}_{\mathbf{k}} \propto n_k^2$ , so that  $n_k$  and  $E(k) \propto \sqrt{\varepsilon}$ . Assuming full self-similarity (i.e. no other dimensional parameters enter into the game), and that  $\varepsilon_k$  is independent of  $k$  in an inertial range of scales, one can reconstruct  $E(k)$  from the dimensional reasoning up to a dimensionless constant  $C$ ,

$$E(k) = C \sqrt{\varepsilon \omega_k} / k^2. \quad (7a)$$

For example, for non-dispersive 3D acoustic waves with  $\omega_k = c_s k$  we recover the Zakharov-Sagdeev spectrum [12],

$$E(k) \propto k^{-3/2}, \quad (7b)$$

which is a flux spectrum describing an energy cascade from large to small scales. It can also be obtained as an exact solution of Eq. (3) [12].

Recall that non-dispersive acoustic waves admit triad wave number and frequency resonances for colinear

wavevectors only, i.e.  $\mathbf{k} \parallel \mathbf{k}_1 \parallel \mathbf{k}_2$ . In this case the arguments of  $\delta_{12}^{\mathbf{k}}$  and  $\delta(\omega_{12}^{\mathbf{k}})$  in Eq. (3) for  $\text{St}_{\mathbf{k}}$  coincide which creates a singularity. Fortunately, in 3D, this singularity is integrable and  $\text{St}_{\mathbf{k}}$  remains finite. Therefore spectrum Eq. (7b) in 3D is a valid solution (see [14, 15, 20] for further discussions). In 2D, the singularity is NOT integrable and Eq. (3) for the non-dispersive case is therefore invalid. Furthermore, the energy spectrum (7b) obtained dimensionally is wrong in 2D because the extra dimensional parameter  $a$  becomes essential (unlike in 3D).

To solve the problem of acoustic turbulence in 2D, we take into account the dispersion correction in the frequency (4). We choose a reference system with  $\mathbf{k} \parallel \hat{\mathbf{x}}$ . Integrating in Eq. (3)  $\mathcal{R}_{12}^{\mathbf{k}}$  over  $\mathbf{k}_2$  with the help of  $\delta_{12}^{\mathbf{k}}$  and over  $k_{1y}$  using  $\delta(\omega_{12}^{\mathbf{k}})$ . We get

$$\begin{aligned} \Delta_{12}^{\mathbf{k}} &\stackrel{\text{def}}{=} \int_{\mathbf{k}_2, k_{1y}} \delta(\omega_{12}^{\mathbf{k}}) \delta_{12}^{\mathbf{k}} d\mathbf{k}_1 d\mathbf{k}_2 \\ &= |\partial_{k_{1y}} (\omega_{\mathbf{k}} - \omega_{\mathbf{k}_1} - \omega_{\mathbf{k}-\mathbf{k}_1})|^{-1} dk_{1x} \approx \frac{k_1 k_2 dk_{1x}}{c_s k |k_{1y}|}, \end{aligned} \quad (8a)$$

where we retained the leading order in  $ak \ll 1$  only. Analyzing resonance conditions (1) with the the dispersion law (4), for small  $ak$  we find  $k_{1y} \approx \sqrt{6} ak_1 k_2$  (see Supplemental Material). Substituting this into Eq. (8a), we have

$$\Delta_{12}^{\mathbf{k}} \approx dk_{1x} / (\sqrt{6} c_s ak). \quad (8b)$$

We can use this expression to modify the dimensional analysis, which now involves an extra dimensional quantity  $a$ . For this, it is only important to observe that  $\Delta_{12}^{\mathbf{k}} \propto 1/a$ . Therefore  $\text{St}_{\mathbf{k}} \propto n_k^2/a$  which, together with  $\text{St}_{\mathbf{k}} \propto \varepsilon$  (arising from Eq. (6)), gives  $n_k \propto \sqrt{\varepsilon a}$ . This leads to the following energy spectrum of weakly dispersive 2D acoustic turbulence,

$$E(k) = C_1 \sqrt{\varepsilon a \omega_k} / k^{3/2} = C_1 \sqrt{\varepsilon a c_s} / k, \quad (9)$$

where  $C_1$  is a dimensionless constant. Note that the spectrum  $E(k) \propto 1/k$  was suggested in [11] without proving that it is a solution of the kinetic equation. Below, we will prove that (9) is the unique stationary power-law flux solution of Eq. (3) and find  $C_1$  analytically. To do this, we compute  $\Delta_{\mathbf{k}2}^1$  and  $\Delta_{\mathbf{k}1}^2$ , similarly to Eqs. (8a) (see the Eqs. (27) in Supplemental Material) and present the collision term in (3) in simpler form,

$$\begin{aligned} \text{St}_{\mathbf{k}} &= \frac{2\pi}{\sqrt{6} c_s ak} \int dk_1 dk_2 \left[ \mathcal{N}_{12}^{\mathbf{k}} \delta(k - k_1 - k_2) \right. \\ &\quad \left. - \mathcal{N}_{\mathbf{k}2}^1 \delta(k_1 - k - k_2) - \mathcal{N}_{\mathbf{k}1}^2 \delta(k_2 - k - k_1) \right]. \end{aligned} \quad (10)$$

Let us seek a stationary solution of Eq. (3) in form

$$n_k = Ak^{-x}, \quad x = \text{const}. \quad (11)$$

There is, of course, a trivial solution with  $x = 1$  corresponding to the thermodynamic energy equipartition,

but we will be interested in non-zero flux states only. Assuming that the integrals in Eq. (10) converge (which is to be checked *a posteriori*), let us apply the Kraichnan-Zakharov transform

$$k_1 = \frac{k^2}{\tilde{k}_1}, \quad k_2 = \frac{\tilde{k}\tilde{k}_2}{\tilde{k}_1} \quad \text{and} \quad k_2 = \frac{k^2}{\tilde{k}_1}, \quad k_1 = \frac{\tilde{k}\tilde{k}_2}{\tilde{k}_1} \quad (12)$$

to the second and the third terms in Eq. (10) respectively. This gives (after dropping tildes on the integration variables for uniformity of notations):

$$\text{St}_{\mathbf{k}} = \frac{2\pi}{\sqrt{6}c_s ak} \int dk_1 dk_2 \mathcal{N}_{12}^{\mathbf{k}} \times \left[ 1 - \left(\frac{k_1}{k}\right)^{-y} - \left(\frac{k_2}{k}\right)^{-y} \right] \delta(k - k_1 - k_2), \quad (13)$$

with  $y = 5 - 2x$ . Thus,  $\text{St}_{\mathbf{k}} = 0$  if  $y = -1$ , so that  $x = 3$  and thus

$$n_k \propto \frac{1}{k^3}, \quad \text{so} \quad E(k) = 2\pi k \omega_k n_k \approx 2\pi k^2 c_s n_k \propto \frac{1}{k}. \quad (14)$$

This is an exact stationary solution of the Eqs. (3) and (10) coinciding with the dimensional prediction (9).

To demonstrate existence and uniqueness of solution (14), let us non-dimensionalize the collision integral for arbitrary  $x$ :

$$\text{St}_k = \frac{2\pi A^2 V_0^2}{\sqrt{6} a c_s} I(x) k^{3-2x}, \quad (15a)$$

$$I(x) = \int_0^1 q(1-q)(q^{-x}(1-q)^{-x} - q^{-x} - (1-q)^{-x}) \times (1 - q^{-y} - (1-q)^{-y}) dq. \quad (15b)$$

This integral converges if and only if  $2 < x < 4$  or  $x = 1$ , as we prove in the Supplemental Material. In Fig. 1 we show the plot  $I(x)$  obtained numerically for the convergence range  $2 < x < 4$ . As we see,  $x = 3$  is the only point at which  $I(x) = 0$ , which proves the uniqueness of the stationary power-law flux solution (9). The existence of the solution amounts to the finiteness of the constant  $C_1$ . We compute  $C_1$  in Sec. III.C of the Supplemental Material by substituting (13) into the definition of the flux (6). The result is as follows

$$\varepsilon_k = \frac{4\pi^4 A^2 V_0^2}{\sqrt{6} a}, \quad C_1 = \frac{6^{1/4} \sqrt{c_s}}{\pi V_0} \quad (16)$$

Notice that Eq. (3) is valid for sufficiently small nonlinearities and stochasticity of the phases [1, 2]. Let us define the interaction frequency  $\gamma_k$  as a frequency with which the wave packets are destroyed by the nonlinear interactions. It will also correspond to the nonlinear frequency broadening: a characteristic width of the time-Fourier spectrum at a fixed  $\mathbf{k}$ . Applicability of Eq. (3) requires frequency  $\gamma_k$  to be smaller than the characteristic frequency of interacting waves  $\delta\omega_k$ . For the 3D




FIG. 1.

Integral (13) in the window of convergence.

waves, we roughly (and definitely not rigorously) may take  $\delta\omega_k = \omega_k$ . However, due to the non-integrable singularity of the non-dispersive 2D system, in 2D we should take only the dispersive part of the frequency,  $\delta\omega_k = c_s a^2 k^3$ , ignoring its linear part  $c_s k$ , disappearing in the reference system, co-moving with velocity  $c_s$  in the  $\mathbf{k}$ -direction. On the other hand, for the wave turbulence to be considered acoustic, the dispersion must remain a small correction, i.e.  $ak \ll 1$ .

To test our theoretical predictions, we perform direct numerical simulations of the 2D GP equation for the complex wave function  $\psi$ . Written in terms of the healing length  $\xi$ , the speed of sound  $c_s$  and the bulk density  $\rho_0$ , this equation reads

$$i \frac{\partial \psi}{\partial t} = \frac{c_s}{\sqrt{2}\xi} \left[ -\xi^2 \nabla^2 + \frac{1}{\rho_0} |\psi|^2 - 1 \right] \psi + i D_c \nabla^6 \psi + F(\mathbf{r}, t), \quad (17)$$

where we have also included a large-scale forcing  $F$  and a hyper-viscous dissipation term. The healing length and the speed of sound both depend on the physical properties of the superfluid and on  $\rho_0$ , and they can be chosen arbitrarily in the dimensionless GP equation.

The GP equation is a well established model for BEC and it can be mapped to an effective compressible irrotational fluid flow via the Madelung transformation,

$$\psi(\mathbf{r}, t) = \sqrt{\rho(\mathbf{r}, t)} \exp[i\phi(\mathbf{r}, t)/\sqrt{2}c_s\xi],$$

with  $\rho(\mathbf{r}, t)$  and  $\phi(\mathbf{r}, t)$  being the fluid density and the velocity potential respectively. Perturbations about a still fluid with uniform density  $\rho(\mathbf{r}, t) \equiv \rho_0$  behave as a dispersive sound with frequency given by the Bogoliubov dispersion relation,  $\omega_k = c_s k \sqrt{1 + \xi^2 k^2/2}$ . In the weakly dispersive limit  $\xi k \ll 1$ , it becomes  $\omega_k = c_s k (1 + \xi^2 k^2/4)$  i.e. the dispersion relation (4) with  $a = \xi/2$ . In this limit, the three-wave interaction coefficient of Eq. (17) is of the form (5), although some ambiguities and discrepancies in the value of the coefficient  $V_0$  can be found in the previous works [2, 11, 12, 20]. In Sec.I.B of the Supplemental Material, we provide the corrected derivation which leads to  $V_0 = 3\sqrt{c_s}/4\sqrt{2}$ . Substituting this into Eq. (16) we have the following prediction for the pre-factor constant,

$$C_1 = \frac{2^{11/4}}{3^{3/4}\pi} \approx 0.94. \quad (18)$$

We simulate Eq.(17) using the standard pseudo-spectral code FROST [21] in a periodic domain of size  $L$  using  $N_c^2 = 1024^2$  and  $N_c^2 = 512^2$  collocation points, denoted by Run 1 and Run 2 respectively. The nonlinear term is de-aliased twice with 2/3–rule following the scheme introduced in [22] in order to conserve momentum (in addition to the energy and the number of particles) in the ideal case (with  $F = D_c = 0$ ). The Fourier transform of the forcing  $F$  obeys the OrnsteinUhlenbeck process  $dF_{\mathbf{k}} = -\alpha F_{\mathbf{k}}dt + f_0 dW_{\mathbf{k}}$ , where  $W_{\mathbf{k}}$  is the Wiener process. The forcing acts only on wavenumbers such that  $2\pi \leq kL \leq 3 \times 2\pi$ . In addition, the condensate amplitude is kept constant during the evolution. We set the initial data with uniform condensate with  $|\psi_0|^2 = \rho_0$ , the forcing then adds the acoustic disturbances, and we evolve the system until it reaches a steady-state. We then perform averages over time. In numerics, we have set  $c_s = 1$ ,  $\rho_0 = 1$  and  $\xi = 2L/N_c$ . For forcing and dissipation we set  $\alpha = 1$ ,  $f_0 = 1.25 \times 10^{-4}, 5 \times 10^{-4}$  and  $D_c = 4.1 \times 10^{-15}, 2.1 \times 10^{-11}$  for Run 1 and 2 respectively.

In absence of forcing and dissipation, Eq. (17) conserves the total energy (Hamiltonian) of the system. The energy per unit of mass, written in terms of the hydrodynamic variables, consists of the kinetic, internal and quantum energies [23]:

$$E = \frac{1}{L^2 \rho_0} \int \left[ \frac{\rho}{2} (\nabla \phi)^2 + \frac{c^2}{2\rho_0} (\rho - \rho_0)^2 + c^2 \xi^2 (\nabla \rho)^2 \right] d^2 \mathbf{r}. \quad (19)$$

In the dispersiveless limit ( $\xi \rightarrow 0$ ), we retrieve the standard energy for a compressible, isentropic, irrotational fluid [24]. The total energy spectrum is computed writing the energy as usual in quantum turbulence [23]. We also calculate the  $k$ -space energy flux  $\epsilon(k)$  directly using Eq. (17) (see the Eq.(31) of Supplemental Material for exact definitions).

In Fig. 2 we show the fluxes and spectra for Run 1 and Run 2. We see that the fluxes have a pronounced plateau which indicates the presence of an inertial range (free of forcing and dissipation effects). Both runs display a stationary power-law spectrum. Remarkably, both, the power-law exponent and the pre-factor  $C_1$  (calculated based on the averaged flux in the inertial range), closely agree with the theoretical predictions (9) and (16).

In Fig. 3 we show the spatio-temporal spectrum for Run1. We see that this spectrum follows closely the Bogoliubov dispersion law, which indicates that the nonlinearity is sufficiently weak. The  $\omega$ -width of the spectrum at each fixed  $k$  represents the nonlinear frequency broadening; we define it as

$$\gamma_k = \left[ \int_0^\infty (\omega - \omega_k)^2 |\hat{\psi}(\omega, k)|^2 d\omega / \int_0^\infty |\hat{\psi}(\omega, k)|^2 d\omega \right]^{1/2}.$$

In Fig. 4 we show the ratios  $\gamma_k / \delta\omega_k$ , where  $\delta\omega = c_s a^2 k^3$  is the dispersive correction. Recall that the WT theory is applicable when  $\gamma_k / \delta\omega_k \ll 1$ . We see that this quantity is indeed small in the scaling range in Run1, and




FIG. 2.

a) Energy flux. b) Energy spectra. Dashed lines correspond to theoretical prediction (9) and (18) using the corresponding flux values in the inertial range.




FIG. 3. Spatio-temporal spectrum  $S(\omega, k) = |\hat{\psi}(\omega, k)|^2$  normalized by the time-averaged spectra  $|\hat{\psi}(t, k)|^2$  of Run1 ( $1024^2$ ). The dashed and dot-dashed lines show the Bogoliubov and the pure acoustic dispersion relation.

only marginally small in a rather narrow range in Run2. This indicates that the WT theory has a good predictive power even when the formal applicability condition is on the borderline of validity.

The main result of our paper is the 1D energy spectrum of 2D weakly dispersive acoustic waves, Eq. (9), found as






FIG. 4.

Linear to nonlinear time ratios for Run1 ( $1024^2$ ) and Run2 ( $512^3$ ) computed from the STS in Fig. 3.

the unique stationary constant-flux solution of the kinetic equation (3) with convergent collision integral. From the physical view point such a convergence means that the main contribution to the energy balance of waves with wavenumber  $k$  comes from their energy exchange with the “neighbouring” waves with wavenumbers  $k'$  of the order of  $k$ . In the language of hydrodynamic turbulence we are dealing here with the step-by-step cascade energy transfer, local in the wavenumber space. We found the energy flux to be positive, meaning that the energy is transferred from small to large  $k$  i.e. it is a *direct energy cascade*.

We tested our analytical predictions by numerical simulations of the forced-dissipated GP equation (17) in the presence of a strong condensate. The analytically found spectrum (9) was confirmed by the numerics, including both the power-law exponent and the pre-factor  $C_1$  without any adjustable parameter. Such a double validation is a rare success in the theory of wave turbulence, where numerical tests were attempted by numerical simulations for various types of waves but, in most cases, only the spectrum exponent was confirmed. Wave turbulence is therefore a valid and productive approach for describing 2D superfluid BEC turbulence where interacting sound waves represent the principal mechanism of energy dissipation. **Since measurements of the spectrum are experimentally accessible in BEC [16, 17], our results present verifiable predictions which could guide future experiments.** The focus of the present Letter was on the weak turbulence of 2D acoustic waves, and the strong turbulence regimes would be an interesting subject for future studies.

This work was supported by the Agence Nationale de la Recherche through the project GIANTE ANR-18-CE30-0020-01, by the Simons Foundation Collaboration grant Wave Turbulence (Award ID 651471) and by NSF-BSF grant # 2020765. This work was granted access to the

HPC resources of CINES, IDRIS and TGCC under the allocation 2019-A0072A11003 made by GENCI. Computations were also carried out at the Msocentre SIGAMM hosted at the Observatoire de la Cte d’Azur.

\* krstulovic@oca.eu

- [1] V. Zakharov, V. L’vov, and G. Falkovich, *Kolmogorov Spectra of Turbulence, wave turbulence* (Springer, 2012).
- [2] S. Nazarenko, *Wave Turbulence* (Springer Berlin Heidelberg, series: Lecture Notes in Physics, 2011).
- [3] E. Falcon and N. Mordant, Annual Review of Fluid Mechanics **54**, annurev (2022), ISSN 0066-4189, 1545-4479.
- [4] S. Galtier, S. V. Nazarenko, A. C. Newell, and A. Pouquet, Journal of Plasma Physics **63**, 447 (2000), ISSN 0022-3778, 1469-7807.
- [5] V.E. Zakharov, Soviet PhysicsJETP **35**, 908-914 (1972).
- [6] P. Caillol and V. Zeitlin, Dynamics of Atmospheres and Oceans **32**, 81 (2000), ISSN 03770265.
- [7] S. Galtier, Physical Review E **68**, 015301 (2003), ISSN 1063-651X, 1095-3787.
- [8] V. S. L’vov and S. Nazarenko, Low Temperature Physics **36**, 785 (2010), ISSN 1063-777X, 1090-6517.
- [9] G. Düring, C. Josserand, and S. Rica, Physical review letters **97**, 025503 (2006).
- [10] S. Galtier and S. V. Nazarenko, Physical review letters **119**, 221101 (2017).
- [11] S. Dyachenko, A. C. Newell, A. Pushkarev, and V. E. Zakharov, Physica D: Nonlinear Phenomena **57**, 96 (1992).
- [12] V. E. Zakharov and R. Z. Sagdeev, Dokl. Akad. Nauk SSSR **192**, 297 (1970).
- [13] B. B. Kadomtsev and V. I. Petviashvili, Doklady Akademii Nauk SSSR **208**, 794 (1973).
- [14] A. C. Newell and P. J. Aucoin, Journal of Fluid Mechanics **49**, 593609 (1971).
- [15] V. S. L’vov, Y. L’vov, A. C. Newell, and V. Zakharov, Phys. Rev. E **56**, 390 (1997).
- [16] N. Navon, A. L. Gaunt, R. P. Smith, and Z. Hadzibabic, Nature **539**, 72 (2016), ISSN 0028-0836, 1476-4687.
- [17] N. Navon, C. Eigen, J. Zhang, R. Lopes, A. L. Gaunt, K. Fujimoto, M. Tsubota, R. P. Smith, and Z. Hadzibabic, Science **366**, 382 (2019), ISSN 0036-8075, 1095-9203.
- [18] S. P. Johnstone, A. J. Groszek, P. T. Starkey, C. J. Billington, T. P. Simula, and K. Helmerston, Science **364**, 1267 (2019), ISSN 0036-8075, 1095-9203.
- [19] G. Gauthier, M. T. Reeves, X. Yu, A. S. Bradley, M. A. Baker, T. A. Bell, H. Rubinsztein-Dunlop, M. J. Davis, and T. W. Neely, Science **364**, 1264 (2019), ISSN 0036-8075, 1095-9203.
- [20] V. E. Zakharov and S. V. Nazarenko, Physica D: Nonlinear Phenomena **201**, 203 (2005).
- [21] N. P. Müller, J. I. Polanco, and G. Krstulovic, Phys. Rev. X **11**, 011053 (2021), URL <https://link.aps.org/doi/10.1103/PhysRevX.11.011053>.
- [22] G. Krstulovic and M. Brachet, Phys. Rev. E **83**, 066311 (2011), URL <https://link.aps.org/doi/10.1103/PhysRevE.83.066311>.
- [23] C. Nore, M. Abid, and M. E. Brachet, Phys. Rev. Lett. **78**, 3896 (1997).
- [24] L. D. Landau and E. M. Lifshitz, *Fluid Mechanics, Sec-*

*2nd Edition: Volume 6 (Course of Theoretical Physics),*  
Course of theoretical physics / by L. D. Landau and E.

M. Lifshitz, Vol. 6 (Butterworth-Heinemann, 1987), 2nd  
ed., ISBN 0750627670.