

This is the accepted manuscript made available via CHORUS. The article has been published as:

Criticality and Phase Classification for Quadratic Open Quantum Many-Body Systems

Yikang Zhang and Thomas Barthel

Phys. Rev. Lett. **129**, 120401 — Published 12 September 2022

DOI: [10.1103/PhysRevLett.129.120401](https://doi.org/10.1103/PhysRevLett.129.120401)

Criticality and Phase Classification for Quadratic Open Quantum Many-Body Systems

Yikang Zhang and Thomas Barthel

Department of Physics, Duke University, Durham, North Carolina 27708, USA

(Dated: August 1, 2022)

We study the steady states of translation-invariant open quantum many-body systems governed by Lindblad master equations, where the Hamiltonian is quadratic in the ladder operators, and the Lindblad operators are either linear or quadratic and Hermitian. These systems are called quasi-free and quadratic, respectively. We find that steady states of one-dimensional systems with finite-range interactions necessarily have exponentially decaying Green's functions. For the quasi-free case without quadratic Lindblad operators, we show that fermionic systems with finite-range interactions are non-critical for any number of spatial dimensions and provide bounds on the correlation lengths. Quasi-free bosonic systems can be critical in $D > 1$ dimensions. Lastly, we address the question of phase transitions in quadratic systems and find that, without symmetry constraints beyond invariance under single-particle basis and particle-hole transformations, all gapped Liouvillians belong to the same phase.

Introduction. – For closed systems, criticality and quantum phase transitions have been studied extensively [1–4]. Particularly, for one-dimensional systems, we have obtained a thorough classification of gapped states using the tensor-network ansatz [5–8].

In practice, most quantum systems are not perfectly isolated from their environment. In addition to posing challenges for quantum technology, driving and dissipation in open systems could be designed to stabilize (novel) phases of matter or particular entangled states [9–11], e.g., to facilitate measurement-based quantum computation [12, 13], quantum phase estimation [14, 15], and quantum simulation [16–20]. For Markovian systems, the density matrix $\hat{\rho}$ evolves according to a Lindblad master equation [21–25]

$$\partial_t \hat{\rho} = \mathcal{L} \hat{\rho} = -i[\hat{H}, \hat{\rho}] + \sum_{\alpha} \left(\hat{L}_{\alpha} \hat{\rho} \hat{L}_{\alpha}^{\dagger} - \frac{1}{2} \{ \hat{L}_{\alpha}^{\dagger} \hat{L}_{\alpha}, \hat{\rho} \} \right).$$

In addition to the Hamiltonian part $-i[\hat{H}, \hat{\rho}]$, the Liouvillian superoperator \mathcal{L} captures decoherence processes with environment couplings described by the Lindblad operators \hat{L}_{α} .

In this work, we elucidate the occurrence of criticality and phase transitions in the steady states of open quasi-free and quadratic systems of fermions and bosons. Quasi-free open systems are characterized by Hamiltonians that are bilinear and Lindblad operators that are linear in ladder operators. Quadratic open systems may have additional bilinear self-adjoint Lindblad operators [26, 27]. A system is called *critical* if it has a unique steady state with algebraically decaying correlations. We establish that quadratic one-dimensional (1D) systems with finite-range interactions and unique steady states necessarily have exponentially decaying Green's functions. Next, we address quasi-free systems with finite-range interactions. Quasi-free fermionic systems are non-critical for any number of spatial dimensions [28]. Conversely, one can construct critical quasi-free bosonic systems for $D \geq 2$ dimensions. Gapped quasi-free systems

are always non-critical. Of course, the existence of critical steady states does not necessarily imply phase transitions. In fact we show that, without symmetry constraints beyond invariance under single-particle basis and particle-hole transformations, all gapped Liouvillians of quadratic open systems belong to the same phase.

Experimentally, systems of trapped ions [29, 30], Rydberg atoms [31, 32], ultracold atoms in optical lattices or tweezers [33–35], and superconducting circuits [36, 37] allow for the engineering of such dissipative systems [38–43]. In circuit QED systems [44–47], linear Lindblad operators arise naturally from photon loss and pump process, while the coupling of cavities can lead to bilinear Lindblad operators [48, 49].

Setup and covariance matrix. – Consider a system of identical bosons or fermions with ladder operators \hat{a}_j and \hat{a}_j^{\dagger} for modes $j = 1, \dots, N$. We employ Majorana operators $\hat{w}_{j+} := (\hat{a}_j + \hat{a}_j^{\dagger})/\sqrt{2}$ and $\hat{w}_{j-} := i(\hat{a}_j - \hat{a}_j^{\dagger})/\sqrt{2}$ which obey the (anti-)commutation relations

$$\begin{aligned} \{\hat{w}_{i\mu}, \hat{w}_{j\nu}\} &= \delta_{i,j} \delta_{\mu\nu} && \text{for fermions, and} \\ [\hat{w}_{i\mu}, \hat{w}_{j\nu}] &= -i\mu \delta_{i,j} \delta_{\mu,-\nu} && \text{for bosons.} \end{aligned}$$

We address Markovian systems with quadratic Hamiltonians $\hat{H} = \sum_{i\mu,j\nu} \hat{w}_{i\mu} H_{i\mu,j\nu} \hat{w}_{j\nu}$. Quasi-free systems only have linear Lindblad operators $\hat{L}_s = \sum_{j\nu} L_{s,j\nu} \hat{w}_{j\nu}$. Quadratic systems may feature additional bilinear self-adjoint Lindblad operators $\hat{M}_u = \hat{M}_u^{\dagger} = \sum_{i\mu,j\nu} \hat{w}_{i\mu} (M_u)_{i\mu,j\nu} \hat{w}_{j\nu}$. The $2N \times 2N$ covariance matrix

$$\Gamma_{i\mu,j\nu} := \begin{cases} \frac{i}{2} \langle \hat{w}_{i\mu} \hat{w}_{j\nu} - \hat{w}_{j\nu} \hat{w}_{i\mu} \rangle & \text{for fermions,} \\ \frac{1}{2} \langle \hat{w}_{i\mu} \hat{w}_{j\nu} + \hat{w}_{j\nu} \hat{w}_{i\mu} \rangle & \text{for bosons} \end{cases} \quad (1)$$

can be shown to evolve according to the equation of motion [26, 27]

$$\partial_t \Gamma = X \Gamma + \Gamma X^T + Y + \sum_u Z_u \Gamma Z_u^T, \quad (2)$$

where the real $2N \times 2N$ matrices X , Y , and Z_u depend on the coupling coefficients H , L_s , and M_u as detailed

in the Supplemental Material. The Z_u term vanishes for quasi-free systems.

For a translation-invariant system in D dimensions, each mode i is associated with a cell location $\mathbf{i} \in \mathbb{Z}^D$ and a crystal-basis index $c_i = 1, \dots, b$, where b is the number of bands. The covariance matrix elements and coupling coefficients are then functions of spatial distances such that

$$\Gamma_{i\mu, j\nu} =: \gamma_{c_i\mu, c_j\nu}(\mathbf{i} - \mathbf{j}), \quad X_{i\mu, j\nu} =: x_{c_i\mu, c_j\nu}(\mathbf{i} - \mathbf{j})$$

etc., and the equation of motion (2) takes the form

$$\begin{aligned} \partial_t \gamma(\mathbf{r}) = & \sum_{\mathbf{n}} [x(\mathbf{n})\gamma(\mathbf{r} - \mathbf{n}) + \gamma(\mathbf{r} + \mathbf{n})x^T(\mathbf{n})] + y(\mathbf{r}) \\ & + \sum_{u, \mathbf{n}, \mathbf{j}, \mathbf{l}} z_u(\mathbf{r} - \mathbf{n}, \mathbf{j} - \mathbf{n})\gamma(\mathbf{r} - \mathbf{l})z_u^T(-\mathbf{n}, \mathbf{l} - \mathbf{n}), \end{aligned} \quad (3)$$

where γ , x , y , and z_u are $2b \times 2b$ matrices depending on lattice translation vectors $\mathbf{r} \in \mathbb{Z}^D$.

Correlations in quadratic 1D systems. – As a first result, let us establish the following.

Proposition 1. *If a quadratic 1D system with translation-invariant finite-range couplings has a unique steady state, then its single-particle Green's function $\gamma(\mathbf{r})$ cannot follow a power-law decay with respect to the distance $|\mathbf{r}|$.*

For the steady-state covariance matrix $\gamma(\mathbf{r})$, the right-hand side of Eq. (3) needs to be zero. For distances r large enough such that the local z_u and y terms vanish, $\gamma(\mathbf{r})$ obeys a matrix difference equation of the form

$$C_0\gamma(\mathbf{r}) + C_1\gamma(\mathbf{r} + 1) + \dots + C_R\gamma(\mathbf{r} + R) = \mathbf{0}. \quad (4)$$

Here $\gamma(\mathbf{r})$ is the vectorization of $\gamma(\mathbf{r})$, the $4b^2 \times 4b^2$ matrices C_m are determined by the coupling matrices $x(m)$, and R denotes the interaction range [50].

In the simplest scenario, C_R is invertible such that we can solve Eq. (4) for $\gamma(\mathbf{r} + R)$ and

$$\mathbf{g}_{r+1} = \begin{bmatrix} A_{R-1} & A_{R-2} & \dots & A_1 & A_0 \\ \mathbb{1} & 0 & \dots & 0 & 0 \\ 0 & \mathbb{1} & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \dots & \mathbb{1} & 0 \end{bmatrix} \mathbf{g}_r \quad (5)$$

with $\mathbf{g}_r^T := (\gamma^T(\mathbf{r} + R - 1), \dots, \gamma^T(\mathbf{r}))$ and $A_m := -C_R^{-1}C_m$. The spectrum of the $4b^2 R \times 4b^2 R$ transfer matrix in Eq. (5) characterizes the spatial decay of $\gamma(\mathbf{r})$. As the spectrum is discrete, all elements of $\gamma(\mathbf{r})$ must decay exponentially, converge to a constant, or oscillate with constant amplitude. An algebraic decay that characterizes critical systems is not possible. The transfer matrix may have eigenvalues β with $|\beta| > 1$. These are however irrelevant as physical systems cannot feature indefinitely

growing $\gamma(\mathbf{r})$. For fermions, this is also prohibited by the constraint that all covariance matrix elements lie in the interval $[-1/2, 1/2]$ [26]. The Supplemental Material [50] gives a more general proof based on generating functions, which does not require invertibility of C_R .

Criticality in quasi-free systems. – Stronger results hold for the systems that have no quadratic Lindblad operators and, hence, no Z_u term in Eq. (2). Let us first consider *gapped* systems, where the Liouvillian \mathcal{L} has a single zero eigenvalue and the other eigenvalues λ have a nonzero *dissipative gap* $\Delta := -\max_{\lambda \neq 0} \text{Re } \lambda > 0$.

Proposition 2. *Gapped quasi-free systems with translation-invariant finite-range couplings are never critical.*

Note that, using quasi-locality [51], this proposition can be generalized to interacting systems. But quasi-free systems allow for a more direct proof that provides bounds on correlation lengths to be reused for Prop. 3:

Due to translation invariance, we can transform to a momentum-space representation with quasi-momenta $k_a = \frac{2\pi}{L}, \frac{4\pi}{L}, \dots, 2\pi$ for $a = 1, \dots, D$. With

$$\tilde{\gamma}(\mathbf{k}) := \sum_{\mathbf{r}} e^{-i\mathbf{k} \cdot \mathbf{r}} \gamma(\mathbf{r}), \quad \tilde{x}(\mathbf{k}) := \sum_{\mathbf{r}} e^{-i\mathbf{k} \cdot \mathbf{r}} x(\mathbf{r}) \quad (6)$$

and an analogous definition of \tilde{y} , according to Eq. (3), the steady state obeys the continuous Lyapunov equation

$$\tilde{x}(\mathbf{k})\tilde{\gamma}(\mathbf{k}) + \tilde{\gamma}(\mathbf{k})\tilde{x}^T(-\mathbf{k}) = -\tilde{y}(\mathbf{k}). \quad (7)$$

For a quasi-free system to be gapped, all eigenvalues of X in Eq. (2) or, equivalently, all eigenvalues of $\tilde{x}(\mathbf{k}) \forall \mathbf{k}$ in Eq. (7) need to have negative real-parts [26]. But this means that we can solve Eq. (7) for $\tilde{\gamma}(\mathbf{k})$ by inverting the matrix $\tilde{x}(\mathbf{k}) \otimes \mathbb{1} + \mathbb{1} \otimes \tilde{x}(-\mathbf{k})$. Due to the finite interaction range, $\tilde{x}(\mathbf{k})$ and $\tilde{y}(\mathbf{k})$ are polynomials in variables $z_a := e^{ik_a} \in \mathbb{C}$ and $1/z_a$. Hence, $\tilde{\gamma}(\mathbf{k})$ is a rational function of the z_a which, according to the invertibility of $\tilde{x}(\mathbf{k})$, has no poles on the manifold $|z_a| = 1$ that corresponds to real momenta $k_a \in (0, 2\pi]$ in the Brillouin zone. For concreteness, let us discuss $D = 2$ dimensions; the generalization to $D \neq 2$ is trivial. The established property of $\tilde{\gamma}(\mathbf{k}) =: \tilde{\gamma}(z_1, z_2)$ allows us to determine $\gamma(r_1, r_2)$ using Cauchy's residue theorem from complex analysis:

In the thermodynamic limit, the inverse of Eq. (6) is

$$\gamma(r_1, r_2) = - \oint_{|z_1|=|z_2|=1} \frac{dz_2}{(2\pi)^2} z_1^{r_1-1} z_2^{r_2-1} \tilde{\gamma}(z_1, z_2). \quad (8)$$

For fixed z_2 , let $\varrho(z_2) := i \sum_m \text{Res}(\tilde{\gamma}(\zeta_m(z_2), z_2))$ denote the sum over the residues of $\tilde{\gamma}$ at pole locations $z_1 = \zeta_m(z_2)$ inside the unit circle $|z_1| = 1$ [52]. With $|\zeta| := \max_{m, |z_2|=1} |\zeta_m(z_2)| < 1$, it follows that

$$|\gamma(r_1, r_2)| \leq |\zeta|^{r_1-1} \oint_{|z_2|=1} \frac{dz_2}{2\pi} |z_2^{r_2-1} \varrho(z_2)|. \quad (9)$$

As the contour integral is independent of r_1 , this bound establishes an exponential decay of $\gamma(r_1, r_2)$ with correlation length

$$\xi_1 \leq -1/\ln|\zeta| = -1/\max_{m, |z_2|=1} \ln|\zeta_m(z_2)| \quad (10)$$

in the positive r_1 direction. An exponential bound for negative r_1 is obtained by using $z_1 := e^{-ik_1}$ instead of e^{ik_1} , and the same arguments apply to r_2 or further dimensions.

The steady states of quasi-free systems are Gaussian [26]. Hence, according to Wick's theorem [53, 54], the steady state is fully characterized by $\gamma(\mathbf{r})$, and the exponential decay of $\gamma(\mathbf{r})$ implies the exponential decay of all connected real-space correlation functions which concludes the proof of Prop. 2. Let us now drop the constraint of a nonzero dissipative gap.

Proposition 3. *Quasi-free fermionic systems with translation-invariant finite-range couplings are never critical.*

For a unique steady state, the momentum-space covariance matrix $\tilde{\gamma}(\mathbf{k})$ solving Eq. (7) is again a rational function. Furthermore, it cannot have poles at real \mathbf{k} for any short-range fermionic system [28]: The covariance matrix Γ in Eq. (1) is real and anti-symmetric. Hence, there exists an orthogonal transformation $O \in O(2N)$ such that $\Gamma' := O\Gamma O^T = \begin{pmatrix} -\chi & \\ & \chi \end{pmatrix}$, where the elements χ_i of the $N \times N$ diagonal matrix χ correspond to the imaginary eigenvalue pairs $\pm i\chi_i$. The transformation defines an alternative set of Majorana operators $\hat{w}'_{i\mu} := \sum_{j\nu} O_{i\mu, j\nu} \hat{w}_{j\nu}$ with covariance matrix Γ' such that $\chi_i = i\langle \hat{w}'_{i+} \hat{w}'_{i-} \rangle$. As each fermionic occupation number operator $\hat{a}_j^\dagger \hat{a}_j$ has eigenvalues 0 and 1, the operators $i\hat{w}_{j+} \hat{w}_{j-} = 1/2 - \hat{a}_j^\dagger \hat{a}_j$ and the operators $i\hat{w}'_{i+} \hat{w}'_{i-}$ have eigenvalues $\pm 1/2$. Thus, all χ_i are in the interval $[-1/2, 1/2]$, and all covariance matrix elements obey $|\Gamma_{i,j}| \leq \|O^T \Gamma' O\| = \|\chi \oplus (-\chi)\| \leq \frac{1}{2}$. The Fourier transform (6) to momentum space just adds another unitary transformation. Hence, the elements of $\tilde{\gamma}(\mathbf{k})$ have modulus $\leq 1/2$, i.e., singularities can only occur at complex momenta. Their imaginary parts provide bounds on correlation lengths as in Eq. (10), and the system is not critical.

Proposition 3 is in stark contrast to closed fermionic systems, where tight-binding models have, for example, critical Fermi-sea ground states. The situation for open bosonic systems is different. Note that bosonic open systems can be unstable in the sense that the Liouvillian can have eigenvalues with positive real-parts that lead to unlimited absorption of energy and particles. In quasi-free systems, however, the existence of a steady state implies stability [26]. So, stability is implied in the following.

Proposition 4. *Quasi-free bosonic systems with translation-invariant finite-range couplings can be critical in $D \geq 2$ dimensions. 1D systems cannot be critical.*

The statement on 1D systems follows immediately from Prop. 1 and Wick's theorem. Furthermore, one can construct quasi-free bosonic models that are critical for $D \geq 2$ dimensions. Specifically, consider a purely dissipative model with one Lindblad operator $\hat{L}_j^{(1)} := \sqrt{2D\eta}(\hat{w}_{j+} - i\hat{w}_{j-}) = \sqrt{2D\eta}\hat{a}_j$ for every site $j \in \mathbb{Z}^D$ of the D -dimensional square lattice as well as four Lindblad operators $\hat{L}_{j,a}^{(2\pm)} := \hat{w}_{j+} + i\hat{w}_{(j\pm e_a)-}$ and $\hat{L}_{j,a}^{(3\pm)} := \hat{w}_{j+} \pm \hat{w}_{(j\pm e_a)-}$ for every edge, where e_a are the unit vectors for directions $a = 1, \dots, D$. One finds that $\tilde{x}(\mathbf{k}) = 2D(c_{\mathbf{k}} - \eta)\mathbb{1}_2$, where $c_{\mathbf{k}} := \sum_a \cos k_a/D$ [50]. The largest X eigenvalue real-part determines the dissipative gap Δ [26]. Here, $\tilde{x}(\mathbf{k})$ has the doubly degenerate eigenvalue $\xi(\mathbf{k}) = 2D(c_{\mathbf{k}} - \eta)$ and, hence, $\Delta = -\max_{\mathbf{k}} \text{Re } \xi(\mathbf{k}) = 2D(\eta - 1)$. So the model is stable for loss rates $\eta \geq 1$ and the gap closes for $\eta = 1$ at momentum $\mathbf{k} = \mathbf{0}$. Solving the Lyapunov equation (7) yields the covariance matrix $\tilde{\gamma}(\mathbf{k})$ with the diagonal and off-diagonal elements

$$\tilde{\gamma}_{\pm,\pm}(\mathbf{k}) = \frac{\eta + 2}{2(\eta - c_{\mathbf{k}})} \quad \text{and} \quad \tilde{\gamma}_{\pm,\mp}(\mathbf{k}) = \frac{\pm i s_{\mathbf{k}}}{2(\eta - c_{\mathbf{k}})}, \quad (11)$$

where $s_{\mathbf{k}} := \sum_a \sin k_a/D$. With a Fourier transform to $\gamma(\mathbf{r})$, one can assess criticality. For $D = 1$ dimensions, the Fourier integral can be evaluated exactly using the residue theorem. In agreement with Props. 1 and 2, we find an exponential decay of correlations if $\eta > 1$. The correlation length diverges for $\eta \rightarrow 1$, but there is no power-law decay. For dimensions $D \geq 2$, one can expand $\tilde{\gamma}(\mathbf{k})$ in a multipole series over hyperspherical harmonics [55] to reduce the Fourier transformation to a radial integral, which takes the form of a Hankel transform. The leading contributions to $\tilde{\gamma}_{\pm,\pm}$ are isotropic while those to $\tilde{\gamma}_{\pm,\mp}$ are anti-symmetric with respect to reflection. For $D = 2$ dimensions, the diagonal correlations $\gamma_{\pm,\pm}(\mathbf{r})$ decay logarithmically in $|\mathbf{r}|$ and the off-diagonal $\gamma_{\pm,\mp}(\mathbf{r})$ decay as $1/|\mathbf{r}|$. For $D = 3$, they decay as $1/|\mathbf{r}|$ and $1/|\mathbf{r}|^2$, respectively. A detailed discussion is given in the Supplemental Material.

Phase classification for quadratic systems. –

Like quantum phase transitions in closed systems [1–3], driven-dissipative phase transitions are characterized by a non-analytic dependence of steady-state expectation values on system parameters. This requires a non-analytic change in the steady-state density matrix and, hence, a level crossing [56]. So, the dissipative gap Δ needs to close at the transition point [57, 58]. As seen so far, there are some restrictions on criticality in quadratic open systems, but the gap can of course close. As another fundamental result, we will see why, here, closing the gap does generally not lead to phase transitions.

Proposition 5. *For quadratic systems without symmetry constraints beyond invariance under single-particle basis transformations and fermionic particle-hole symmetry, all gapped systems belong to the same phase. For any*

pair of gapped systems \mathcal{L}_1 and \mathcal{L}_2 , one can construct a continuous path of gapped Liouvillians that links the two.

In particular, we claim that for any quadratic Liouvillian \mathcal{L} with gap Δ , the auxiliary Liouvillian

$$\mathcal{L} + \kappa\mathcal{D} \quad \text{has a gap} \quad \Delta' \geq \Delta + \kappa. \quad (12)$$

For fermionic systems, the added dissipator \mathcal{D} comprises two linear Lindblad operators $\hat{L}_{i\pm} = \hat{w}_{i\pm}$ for every mode i [59]. For bosons, \mathcal{D} comprises one operator $\hat{L}_i = \hat{w}_{i+} - i\hat{w}_{i-} = \sqrt{2}\hat{a}_i$ per mode. With this choice and any $\kappa_0 > 0$, the gap stays nonzero, e.g., along the path $(1-g)\mathcal{L}_1 + g\mathcal{L}_2 + \kappa\mathcal{D}$, where the parameters are tuned as $(g, \kappa) : (0, 0) \rightarrow (0, \kappa_0) \rightarrow (1, \kappa_0) \rightarrow (1, 0)$ to connect \mathcal{L}_1 to \mathcal{L}_2 . Note that this proposition does not require short-range interactions.

Statement (12) can be proven by employing the third-quantization formalism [26, 60–62] as detailed in the companion paper [26]: (a) There exist ladder superoperators $a_{j\nu}$ and $a'_{j\nu}$ that obey canonical (anti-)commutation relations and form a basis for the superoperator algebra. (b) One can then construct a biorthogonal operator basis

$$|\mathbf{n}\rangle \quad \text{and} \quad \langle\mathbf{n}| \quad \text{with} \quad \langle\mathbf{n}|\mathbf{n}'\rangle = \delta_{\mathbf{n},\mathbf{n}'}, \quad (13)$$

occupation numbers $\mathbf{n}^T = (n_{1+}, \dots, n_{N+}, \dots, n_{N-})$, and $a'_{j\nu}a_{j\nu}|\mathbf{n}\rangle = n_{j\nu}|\mathbf{n}\rangle$. The Dirac notation with superbras $\langle\hat{A}|$ and superkets $|\hat{B}\rangle$, where \hat{A} and \hat{B} are operators on the Hilbert space, is based on the Hilbert-Schmidt inner product $\langle\hat{A}|\hat{B}\rangle \equiv \text{Tr}(\hat{A}^\dagger\hat{B})$. (c) The ladder superoperators can be chosen such that the matrix representation $\langle\mathbf{n}|\mathcal{L}|\mathbf{n}'\rangle$ of \mathcal{L} assumes a block-triangular form when ordering the basis (13) according to increasing eigenvalues $N_a \in \mathbb{N}$ of the number superoperator $\mathcal{N}_a := \sum_{j\nu} a'_{j\nu}a_{j\nu}$. The spectra of the blocks $\mathcal{L}|_{N_a}$ on the diagonal determine the full Liouvillian spectrum [63]. The only terms due to \mathcal{D} that affect the blocks $\mathcal{L}|_{N_a}$ are [26]

$$-\mathbf{a}'^T \frac{B+B^*}{2} \mathbf{a}, \quad -\mathbf{a}'^T \frac{B+B^*}{2} \mathbf{a}', \quad \mathbf{a}'^T U^\dagger \tau \frac{B-B^*}{2} U \mathbf{a} \quad (14)$$

for fermions with even N_a , fermions with odd N_a , and bosons, respectively. In Eq. (14), $\mathbf{a}^T = (a_{1+}, \dots, a_{N-})$ and $\mathbf{a}'^T = (a'_{1+}, \dots, a'_{N-})$ are vectors containing all ladder superoperators, U is a unitary matrix, $\tau = \begin{pmatrix} 0 & -i\mathbb{1}_N \\ i\mathbb{1}_N & 0 \end{pmatrix}$, and $B_{i\mu,j\nu} = \sum_s L_{s,i\mu} L_{s,j\nu}^*$ is a positive-semidefinite matrix, characterized by the expansion coefficients of linear Lindblad operators $\hat{L}_s = \sum_{j\nu} L_{s,j\nu} \hat{w}_{j\nu}$.

For fermions, the Lindblad operators of dissipator \mathcal{D} have coefficients $L_{i\pm,j\nu} = \delta_{i,j} \delta_{\pm,\nu}$ and, hence, $B = \mathbb{1}_{2N}$ such that the first two terms in Eq. (14) are simply $-\mathcal{N}_a$ and $\mathcal{N}_a - 2N$, respectively. This implies that the spectrum of block $\mathcal{L}|_{N_a}$ is shifted by $-N_a\kappa$ and $(N_a - 2N)\kappa$ for even and odd N_a , respectively. As the $N_a = 0$ block that contains the steady-state eigenvalue zero is one-dimensional, the spectral shifts due to $\kappa\mathcal{D}$ necessarily increase the gap to $\Delta' \geq \Delta + \kappa$. For bosons, we have




FIG. 1. The open fermionic model (15) with $\eta = 1$ and $\mu = 0$. Left: Both, in the quasi-free case ($\zeta = 0$) and quadratic case ($\zeta = 1/4$) with $\alpha = 1/5$ and $\phi = 2\pi/5$, correlations decay exponentially, where the asymptotic form $\sim \beta^r$ (dashed lines) is determined by an eigenvalue β of the transfer matrix in Eq. (5). Right: The dissipative gap Δ for $\zeta = 0$ and $\alpha = 1/2$ vanishes at $\phi = 0, \pi$. It can be increased using the additional dissipator $\kappa\mathcal{D}$ from Eq. (12).

$L_{i,j+} = \delta_{i,j}$ and $L_{i,j-} = -i\delta_{i,j}$. Hence, $B = \begin{pmatrix} \mathbb{1}_N & i\mathbb{1}_N \\ -i\mathbb{1}_N & \mathbb{1}_N \end{pmatrix}$ and $\tau(B - B^*)/2 = -\mathbb{1}_{2N}$ such that the third term in Eq. (14) reads $-\mathcal{N}_a$. Thus, also in the bosonic case, the gap increases at least by κ . For quasi-free fermionic and bosonic systems, the gap increases exactly by κ , i.e., $\Delta' = \Delta + \kappa$. Dissipator \mathcal{D} is invariant under single-particle basis transformations $\hat{a}_j \leftrightarrow \sum_i U_{ji} \hat{a}_i$ and also under particle-hole transformations $\hat{a}_j \leftrightarrow \hat{a}_j^\dagger$ for fermions. This completes the proof of Prop. 5.

Example. – To illustrate some of the above results, consider the quadratic fermionic 1D model with Hamiltonian

$$\hat{H} = \sum_j (\hat{a}_j^\dagger \hat{a}_{j+1} + \alpha \hat{a}_j^\dagger \hat{a}_{j+1}^\dagger + H.c.) - \mu \sum_j \hat{a}_j^\dagger \hat{a}_j, \quad (15)$$

corresponding to a spin-1/2 XY chain, and Lindblad operators $\hat{L}_j = \sqrt{\eta} (\hat{w}_{j+} + e^{i\phi} \hat{w}_{(j+1)+})$ as well as $\hat{M}_j = \sqrt{\zeta} (2\hat{a}_j^\dagger \hat{a}_j - 1)$. In accordance with Prop. 1, $\gamma(r)$ is always found to decay as β^r for an eigenvalue β of the transfer matrix in Eq. (5). Proposition 3 implies that the quasi-free model ($\zeta = 0$), considered in Ref. [64], is never critical, and $\tilde{x}(k)$ determines the full many-body spectrum [26]. In particular, if the Hamiltonian is gapped and $\eta > 0$, the dissipative gap Δ closes only at $\phi = 0$ and π . The correlation length diverges at those points ($\beta \rightarrow 1$) but, at the same time, $\gamma(r) \rightarrow 0$ for all r . Furthermore, employing the additional dissipator $\kappa\mathcal{D}$ from Eq. (12), any two gapped points can always be connected by a path of gapped Liouvillians as explained by Prop. 5 and illustrated in Fig. 1 for the points $\phi = \pi/4$ and $\phi = 9\pi/4$. So, the system is neither critical at $\phi = 0$ or π , nor does it undergo phase transitions. Details are presented in the Supplemental Material.

Discussion. – We have found fundamental prerequisites for criticality and phase transitions in driven-dissipative many-body systems that are in stark contrast to properties of closed systems. For any number

of spatial dimensions, there exist fermionic and bosonic closed systems with phase transitions and critical ground states, i.e., states featuring an algebraic decay of spatial correlations, even if the systems are quasi-free. In contrast, steady states of open 1D quasi-free systems as well as higher-dimensional quasi-free fermionic systems are never critical. For quadratic systems, we found that, while the dissipative gap may close and the system might even be critical for certain points in parameter space, all steady states basically belong to the same phase. The only way for realizing phase transitions in such systems is to impose symmetries on the considered Liouvillians that go beyond invariance under single-particle basis transformations (e.g., lattice symmetries) and fermionic particle-hole transformations or combinations thereof. A notable example are topological transitions in quasi-free systems, occurring under the (strong) restriction that the Lindblad operators form a complete anti-commuting set [65, 66]. The observation that dissipative phase transitions are, in the above sense, more rare than phase transitions in closed systems adds to the idea that steady states are in certain scenarios related to thermal states of closed systems [10, 67–69] such that continuous symmetries cannot be broken in $D \leq 2$ dimensions according to the Mermin-Wagner theorem [70, 71]. Interactions and more complex Lindblad operators can break the block-triangular Liouvillian structures [26, 63] that underlie our results on quasi-free and quadratic systems and can cause true phase transitions [68, 72–76] as long as we are below an upper critical dimension where all systems become effectively quasi-free.

We gratefully acknowledge discussions with Enej Ilievski, Israel Klich, Jianfeng Lu, and Tomaž Prosen as well as support through US Department of Energy grant DE-SC0019449.

-
- [1] S. Sachdev, *Quantum Phase Transitions*, 2nd ed. (Cambridge University Press, Cambridge, UK, 2011).
 - [2] M. Vojta, *Quantum phase transitions*, *Rep. Prog. Phys.* **66**, 2069 (2003).
 - [3] S. L. Sondhi, S. M. Girvin, J. P. Carini, and D. Shahar, *Continuous quantum phase transitions*, *Rev. Mod. Phys.* **69**, 315 (1997).
 - [4] B. Zeng, X. Chen, D.-L. Zhou, and X.-G. Wen, *Quantum Information Meets Quantum Matter*, *Quantum Science and Technology* (Springer, New York, 2019).
 - [5] X. Chen, Z.-C. Gu, and X.-G. Wen, *Classification of gapped symmetric phases in one-dimensional spin systems*, *Phys. Rev. B* **83**, 035107 (2011).
 - [6] N. Schuch, D. Pérez-García, and I. Cirac, *Classifying quantum phases using matrix product states and projected entangled pair states*, *Phys. Rev. B* **84**, 165139 (2011).
 - [7] X. Chen, Z.-C. Gu, and X.-G. Wen, *Complete classification of one-dimensional gapped quantum phases in interacting spin systems*, *Phys. Rev. B* **84**, 235128 (2011).
 - [8] J. I. Cirac, D. Pérez-García, N. Schuch, and F. Verstraete, *Matrix product states and projected entangled pair states: Concepts, symmetries, and theorems*, *Rev. Mod. Phys.* **93**, 045003 (2021).
 - [9] B. Kraus, H. P. Büchler, S. Diehl, A. Kantian, A. Micheli, and P. Zoller, *Preparation of entangled states by quantum Markov processes*, *Phys. Rev. A* **78**, 042307 (2008).
 - [10] S. Diehl, A. Micheli, A. Kantian, B. Kraus, H. P. Büchler, and P. Zoller, *Quantum states and phases in driven open quantum systems with cold atoms*, *Nat. Phys.* **4**, 878 (2008).
 - [11] F. Verstraete, M. M. Wolf, and J. I. Cirac, *Quantum computation and quantum-state engineering driven by dissipation*, *Nat. Phys.* **5**, 633 (2009).
 - [12] R. Raussendorf and H. J. Briegel, *A one-way quantum computer*, *Phys. Rev. Lett.* **86**, 5188 (2001).
 - [13] P. Walther, K. J. Resch, T. Rudolph, E. Schenck, H. Weinfurter, V. Vedral, M. Aspelmeyer, and A. Zeilinger, *Experimental one-way quantum computing*, *Nature* **434**, 169 (2005).
 - [14] D. S. Abrams and S. Lloyd, *Quantum algorithm providing exponential speed increase for finding eigenvalues and eigenvectors*, *Phys. Rev. Lett.* **83**, 5162 (1999).
 - [15] A. Aspuru-Guzik, A. D. Dutoi, P. J. Love, and M. Head-Gordon, *Simulated quantum computation of molecular energies*, *Science* **309**, 1704 (2005).
 - [16] R. P. Feynman, *Simulating physics with computers*, *Int. J. Theor. Phys.* **21**, 467 (1982).
 - [17] J. G. Bohnet, B. C. Sawyer, J. W. Britton, M. L. Wall, A. M. Rey, M. Foss-Feig, and J. J. Bollinger, *Quantum spin dynamics and entanglement generation with hundreds of trapped ions*, *Science* **352**, 1297 (2016).
 - [18] C. Gross and I. Bloch, *Quantum simulations with ultra-cold atoms in optical lattices*, *Science* **357**, 995 (2017).
 - [19] H. Bernien, S. Schwartz, A. Keesling, H. Levine, A. Omran, H. Pichler, S. Choi, A. S. Zibrov, M. Endres, M. Greiner, V. Vuletic, and M. D. Lukin, *Probing many-body dynamics on a 51-atom quantum simulator*, *Nature* **551**, 579 (2017).
 - [20] A. Browaeys and T. Lahaye, *Many-body physics with individually controlled Rydberg atoms*, *Nat. Phys.* **16**, 132 (2020).
 - [21] G. Lindblad, *On the generators of quantum dynamical semigroups*, *Commun. Math. Phys.* **48**, 119 (1976).
 - [22] V. Gorini, A. Kossakowski, and E. C. G. Sudarshan, *Completely positive dynamical semigroups of N-level systems*, *J. Math. Phys.* **17**, 821 (1976).
 - [23] H.-P. Breuer and F. Petruccione, *The Theory of Open Quantum Systems* (Oxford University Press, Oxford, 2007).
 - [24] Á. Rivas and S. F. Huelga, *Open Quantum Systems: An Introduction* (Springer, Heidelberg, 2012).
 - [25] M. Wolf and J. I. Cirac, *Dividing quantum channels*, *Commun. Math. Phys.* **279**, 147 (2008).
 - [26] T. Barthel and Y. Zhang, *Solving quasi-free and quadratic Lindblad master equations for open fermionic and bosonic systems*, *arXiv:2112.08344* (2021).
 - [27] B. Horstmann, J. I. Cirac, and G. Giedke, *Noise-driven dynamics and phase transitions in fermionic systems*, *Phys. Rev. A* **87**, 012108 (2013).
 - [28] Ref. [64] contains an incomplete argument for 1D quasi-free fermionic systems.
 - [29] J. I. Cirac and P. Zoller, *Quantum computations with cold trapped ions*, *Phys. Rev. Lett.* **74**, 4091 (1995).

- [30] R. Blatt and D. Wineland, *Entangled states of trapped atomic ions*, *Nature* **453**, 1008 (2008).
- [31] D. Jaksch, J. I. Cirac, P. Zoller, S. L. Rolston, R. Côté, and M. D. Lukin, *Fast quantum gates for neutral atoms*, *Phys. Rev. Lett.* **85**, 2208 (2000).
- [32] M. D. Lukin, M. Fleischhauer, R. Cote, L. M. Duan, D. Jaksch, J. I. Cirac, and P. Zoller, *Dipole blockade and quantum information processing in mesoscopic atomic ensembles*, *Phys. Rev. Lett.* **87**, 037901 (2001).
- [33] I. Bloch, J. Dalibard, and W. Zwerger, *Many-body physics with ultracold gases*, *Rev. Mod. Phys.* **80**, 885 (2008).
- [34] M. A. Norcia, A. W. Young, and A. M. Kaufman, *Microscopic control and detection of ultracold strontium in optical-tweezer arrays*, *Phys. Rev. X* **8**, 041054 (2018).
- [35] A. Cooper, J. P. Covey, I. S. Madjarov, S. G. Porsev, M. S. Safronova, and M. Endres, *Alkaline-earth atoms in optical tweezers*, *Phys. Rev. X* **8**, 041055 (2018).
- [36] R. J. Schoelkopf and S. M. Girvin, *Wiring up quantum systems*, *Nature* **451**, 664 (2008).
- [37] M. H. Devoret and R. J. Schoelkopf, *Superconducting circuits for quantum information: An outlook*, *Science* **339**, 1169 (2013).
- [38] H. Weimer, M. Müller, I. Lesanovsky, P. Zoller, and H. P. Büchler, *A Rydberg quantum simulator*, *Nat. Phys.* **6**, 382 (2010).
- [39] J. T. Barreiro, M. Müller, P. Schindler, D. Nigg, T. Monz, M. Chwalla, M. Hennrich, C. F. Roos, P. Zoller, and R. Blatt, *An open-system quantum simulator with trapped ions*, *Nature* **470**, 486 (2011).
- [40] M. Müller, S. Diehl, G. Pupillo, and P. Zoller, *Engineered open systems and quantum simulations with atoms and ions*, *Advances in Atomic, Molecular, and Optical Physics* **61**, 1 (2012).
- [41] Z. Leghtas, U. Vool, S. Shankar, M. Hatridge, S. M. Girvin, M. H. Devoret, and M. Mirrahimi, *Stabilizing a Bell state of two superconducting qubits by dissipation engineering*, *Phys. Rev. A* **88**, 023849 (2013).
- [42] A. W. Carr and M. Saffman, *Preparation of entangled and antiferromagnetic states by dissipative Rydberg pumping*, *Phys. Rev. Lett.* **111**, 033607 (2013).
- [43] D. D. Bhaktavatsala Rao and K. Mølmer, *Dark entangled steady states of interacting Rydberg atoms*, *Phys. Rev. Lett.* **111**, 033606 (2013).
- [44] M. J. Hartmann, F. G. S. L. Brandão, and M. B. Plenio, *Strongly interacting polaritons in coupled arrays of cavities*, *Nat. Phys.* **2**, 849 (2006).
- [45] D. G. Angelakis, M. F. Santos, and S. Bose, *Photon-blockade-induced Mott transitions and XY spin models in coupled cavity arrays*, *Phys. Rev. A* **76**, 031805 (2007).
- [46] M. J. Hartmann, *Quantum simulation with interacting photons*, *J. Opt.* **18**, 104005 (2016).
- [47] M. Fitzpatrick, N. M. Sundaresan, A. C. Y. Li, J. Koch, and A. A. Houck, *Observation of a dissipative phase transition in a one-dimensional circuit QED lattice*, *Phys. Rev. X* **7**, 011016 (2017).
- [48] D. Marcos, A. Tomadin, S. Diehl, and P. Rabl, *Photon condensation in circuit quantum electrodynamics by engineered dissipation*, *New J. Phys.* **14**, 055005 (2012).
- [49] A. Tomadin, S. Diehl, M. D. Lukin, P. Rabl, and P. Zoller, *Reservoir engineering and dynamical phase transitions in optomechanical arrays*, *Phys. Rev. A* **86**, 033821 (2012).
- [50] The Supplemental Material provides details on the steady-state equation for the covariance matrix, completes the proof of Prop. 1 for non-invertible C_R in Eq. (4), establishes the criticality of the $D > 2$ dimensional bosonic models for Prop. 4, and discusses an illustrative quadratic fermionic model.
- [51] T. Barthel and M. Kliesch, *Quasi-locality and efficient simulation of Markovian quantum dynamics*, *Phys. Rev. Lett.* **108**, 230504 (2012).
- [52] This statement and the following considerations actually concern matrix elements of γ and $\tilde{\gamma}$, but we drop the matrix indices for brevity of notation.
- [53] G. C. Wick, *The evaluation of the collision matrix*, *Phys. Rev.* **80**, 268 (1950).
- [54] J. W. Negele and H. Orland, *Quantum Many-Particle Systems* (Perseus Books, Reading, MA, 1988).
- [55] J. E. Avery and J. S. Avery, *Hyperspherical Harmonics and Their Physical Applications* (World Scientific, Singapore, 2017).
- [56] T. Kato, *Perturbation Theory for Linear Operators, Classics in Mathematics*, 2nd ed. (Springer, Berlin, 1995).
- [57] E. M. Kessler, G. Giedke, A. Imamoglu, S. F. Yelin, M. D. Lukin, and J. I. Cirac, *Dissipative phase transition in a central spin system*, *Phys. Rev. A* **86**, 012116 (2012).
- [58] F. Minganti, A. Biella, N. Bartolo, and C. Ciuti, *Spectral theory of Liouvillians for dissipative phase transitions*, *Phys. Rev. A* **98**, 042118 (2018).
- [59] A dissipator \mathcal{D} comprising Lindblad operators $\hat{L}_1, \dots, \hat{L}_n$ acts as $\mathcal{D}\hat{\rho} = \sum_{\alpha=1}^n (\hat{L}_{\alpha}\hat{\rho}\hat{L}_{\alpha}^{\dagger} - \frac{1}{2}\{\hat{L}_{\alpha}^{\dagger}\hat{L}_{\alpha}, \hat{\rho}\})$.
- [60] T. Prosen, *Third quantization: A general method to solve master equations for quadratic open Fermi systems*, *New J. Phys.* **10**, 043026 (2008).
- [61] T. Prosen, *Spectral theorem for the Lindblad equation for quadratic open fermionic systems*, *J. Stat. Mech.* **2010**, P07020 (2010).
- [62] T. Prosen and T. H. Seligman, *Quantization over boson operator spaces*, *J. Phys. A: Math. Theor.* **43**, 392004 (2010).
- [63] T. Barthel and Y. Zhang, *Superoperator structures and no-go theorems for dissipative quantum phase transitions*, *Phys. Rev. A* **105**, 052224 (2022).
- [64] J. Eisert and T. Prosen, *Noise-driven quantum criticality*, *arXiv:1012.5013* (2010).
- [65] C.-E. Bardyn, M. A. Baranov, C. V. Kraus, E. Rico, A. Imamoglu, P. Zoller, and S. Diehl, *Topology by dissipation*, *New J. Phys.* **15**, 085001 (2013).
- [66] J. C. Budich, P. Zoller, and S. Diehl, *Dissipative preparation of Chern insulators*, *Phys. Rev. A* **91**, 042117 (2015).
- [67] A. Mitra, S. Takei, Y. B. Kim, and A. J. Millis, *Nonequilibrium quantum criticality in open electronic systems*, *Phys. Rev. Lett.* **97**, 236808 (2006).
- [68] R. Rota, F. Minganti, C. Ciuti, and V. Savona, *Quantum critical regime in a quadratically driven nonlinear photonic lattice*, *Phys. Rev. Lett.* **122**, 110405 (2019).
- [69] J. T. Young, A. V. Gorshkov, M. Foss-Feig, and M. F. Maghrebi, *Nonequilibrium fixed points of coupled Ising models*, *Phys. Rev. X* **10**, 011039 (2020).
- [70] P. C. Hohenberg, *Existence of long-range order in one and two dimensions*, *Phys. Rev.* **158**, 383 (1967).
- [71] N. D. Mermin and H. Wagner, *Absence of ferromagnetism or antiferromagnetism in one- or two-dimensional isotropic Heisenberg models*, *Phys. Rev. Lett.* **17**, 1133 (1966).
- [72] A. Le Boité, G. Orso, and C. Ciuti, *Steady-state phases and tunneling-induced instabilities in the driven dissipa-*

- tive Bose-Hubbard model*, [Phys. Rev. Lett. **110**, 233601 \(2013\)](#).
- [73] A. Le Boité, G. Orso, and C. Ciuti, *Bose-Hubbard model: Relation between driven-dissipative steady states and equilibrium quantum phases*, [Phys. Rev. A **90**, 063821 \(2014\)](#).
- [74] R. M. Wilson, K. W. Mahmud, A. Hu, A. V. Gorshkov, M. Hafezi, and M. Foss-Feig, *Collective phases of strongly interacting cavity photons*, [Phys. Rev. A **94**, 033801 \(2016\)](#).
- [75] V. Savona, *Spontaneous symmetry breaking in a quadratically driven nonlinear photonic lattice*, [Phys. Rev. A **96**, 033826 \(2017\)](#).
- [76] W. Verstraelen, R. Rota, V. Savona, and M. Wouters, *Gaussian trajectory approach to dissipative phase transitions: The case of quadratically driven photonic lattices*, [Phys. Rev. Research **2**, 022037 \(2020\)](#).
- [77] P. C. Parks, *A. M. Lyapunov's stability theory – 100 years on*, [IMA J. Math. Control Inf. **9**, 275 \(1992\)](#).
- [78] V. Simoncini, *Computational methods for linear matrix equations*, [SIAM Rev. **58**, 377 \(2016\)](#).
- [79] S. Sastry, *Nonlinear Systems: Analysis, Stability, and Control*, Vol. 10 of *Interdisciplinary Applied Mathematics* (Springer, New York, 1999).
- [80] H. K. Khalil, *Nonlinear Systems*, 3rd ed. (Prentice Hall, Upper Saddle River, NJ, 2002).
- [81] H. S. Wilf, *Generatingfunctionology*, 3rd ed. (A. K. Peters, Ltd., USA, 2006).
- [82] P. Jordan and E. Wigner, *About the Pauli exclusion principle*, [Z. Phys. **47**, 631 \(1928\)](#).
- [83] E. H. Lieb, T. Schultz, and D. Mattis, *Two soluble models of an antiferromagnetic chain*, [Ann. Phys. **16**, 407 \(1961\)](#).