



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Simulability of High-Dimensional Quantum Measurements

Marie Ioannou, Pavel Sekatski, Sébastien Designolle, Benjamin D. M. Jones, Roope Uola,
and Nicolas Brunner

Phys. Rev. Lett. **129**, 190401 — Published 4 November 2022

DOI: [10.1103/PhysRevLett.129.190401](https://doi.org/10.1103/PhysRevLett.129.190401)

Simulability of high-dimensional quantum measurements

Marie Ioannou,^{1,*} Pavel Sekatski,^{1,*} Sébastien Designolle,¹
Benjamin D.M. Jones,^{1,2,3} Roope Uola,¹ and Nicolas Brunner¹

¹*Department of Applied Physics University of Geneva, 1211 Geneva, Switzerland*

²*H. H. Wills Physics Laboratory, University of Bristol, Bristol, BS8 1TL, UK*

³*Quantum Engineering Centre for Doctoral Training, University of Bristol, Bristol, BS8 1FD UK*

We investigate the compression of quantum information with respect to a given set \mathcal{M} of high-dimensional measurements. This leads to a notion of simulability, where we demand that the statistics obtained from \mathcal{M} and an arbitrary quantum state ρ are recovered exactly by first compressing ρ into a lower dimensional space, followed by some quantum measurements. A full quantum compression is possible, i.e., leaving only classical information, if and only if the set \mathcal{M} is jointly measurable. Our notion of simulability can thus be seen as a quantification of measurement incompatibility in terms of dimension. After defining these concepts, we provide an illustrative examples involving mutually unbiased basis, and develop a method based on semi-definite programming for constructing simulation models. In turn we analytically construct optimal simulation models for all projective measurements subjected to white noise or losses. Finally, we discuss how our approach connects with other concepts introduced in the context of quantum channels and quantum correlations.

Introduction.— Quantum measurements play a fundamental role in quantum theory and its applications, notably in quantum information processing and metrology. Indeed measurements represent the bridge between a quantum system and an external observer, hence essentially any quantum experiment relies on a quantum measurement process. More recently, the role of quantum measurements as a resource was clarified in the context of quantum information processing tasks, see [1, 2] for a review. At the formal level, the concept of joint measurability [3] provides a framework for characterisation and quantification of the incompatibility of quantum measurements, which can be connected to their usefulness in, e.g., state-discrimination problems [4–9] and quantum steering [10–13].

A natural question is whether quantum measurement incompatibility can also be quantified in terms of dimension. Intuitively, a set of quantum measurement defined on a high-dimensional Hilbert space may feature a stronger form of incompatibility than what is possible for lower dimensions.

In this work we address this question and propose a notion of dimensionality for measurement incompatibility. This notion can be understood, and naturally motivated, in a scenario involving the compression of quantum information. Loosely speaking, we ask whether the statistics of a set of d -dimensional positive operator-valued measures (POVMs) \mathcal{M} (considering any possible quantum state ρ) can be exactly recovered from first projecting ρ onto a n -dimensional space (with $1 \leq n < d$) and then performing POVMs in this lower-dimensional space. If such a compression is possible, we say the set \mathcal{M} is *n -dimensional simulable*. Note that the case $n = 1$ exactly corresponds to the notion of joint measurability; indeed, in this case, the full quantum information can be compressed to classical information, see also [14, 15]. However, as we will see below, there exist sets of POVMs

\mathcal{M} that are incompatible, but yet simulable with n -dimensional measurements with $1 < n < d$. The notion of *n -dimensional simulability* can thus be seen as an extension of the concept of joint measurability, providing a quantification of the incompatibility of quantum measurements in terms of dimension.

After introducing these ideas more formally, we provide illustrative examples based on sets of mutually unbiased measurements. Then, we present a method based on semi-definite programming (SDP) to show that a set of measurements is *n -dimensional simulable*. In turn we consider the continuous sets of all projective measurements subjected to noise or losses, and construct optimal n -simulation models. Finally, we establish a link to partially entanglement-breaking channels [16], and discuss connections to other concepts as well as open questions.

Scenario and definition.— Consider the following task: a sender (Alice) is located on the moon and wants to transmit an (arbitrary) d -dimensional quantum state ρ to a receiver (Bob) located on earth. Upon receiving ρ , Bob will perform a set of measurements $\mathcal{M} = \{M_{a|x}\}_{a,x}$, where $\{M_{a|x}\}_a$ denotes a POVM, i.e., $M_{a|x} \geq 0$ and $\sum_a M_{a|x} = 1 \forall a, x$. This leads to the following statistics termed the target data: $p(a|x, \rho) = \text{Tr}(M_{a|x}\rho)$.

So far, we assumed that the channel between Alice and Bob is ideal, i.e., a d -dimensional identity channel. In the following, however, we will consider the dimensionality of the quantum channel as a resource, which we will aim to minimise. Note that a classical channel (of arbitrary capacity) is always available for free.

We now ask if the target data could be reproduced by using a lower-dimensional quantum channel. That is, we look for a quantum instrument $\{\mathcal{E}_\lambda\}_\lambda$, where each $\mathcal{E}_\lambda : B(\mathbb{C}^d) \rightarrow B(\mathbb{C}^n)$ is a completely positive (CP) map, λ a classical outcome, and the map $\sum_\lambda \mathcal{E}_\lambda$ is trace-preserving. Alice would then send the compressed n -dimensional state $\mathcal{E}_\lambda(\rho)$ to Bob, together with the clas-




FIG. 1. A measurement assemblage $\{M_{a|x}\}$ in dimension d is said to be **n -dimensional simulable** if it can be replicated by first compressing the measured system down to dimension $n < d$ with a “parent” instrument $\{\mathcal{E}_\lambda\}$ independent of the setting x , and then performing some measurements $\{N_{a|x,\lambda}\}$ on the n -dimensional system.

sical outcome λ . Bob finally chooses from a set of measurements $\mathcal{N} = \{N_{a|x,\lambda}\}_{a,x,\lambda}$. The protocol is successful if we recover the target data, i.e., if

$$\sum_\lambda \text{Tr}[N_{a|x,\lambda} \mathcal{E}_\lambda(\rho)] = \text{Tr}(M_{a|x} \rho) \quad \forall \rho, \quad (1)$$

or equivalently if $M_{a|x} = \sum_\lambda \mathcal{E}_\lambda^*(N_{a|x,\lambda})$, where \mathcal{E}_λ^* is the adjoint map to \mathcal{E}_λ (for continuous instruments one replaces the sum with an integral, see examples below). In this case, we say that \mathcal{M} is **n -dimensional simulable**, as illustrated in Fig. 1. Clearly, in the case $n = 1$, the instrument corresponds to a POVM and the measurements afterwards form classical post-processings of the classical output. This coincides with the concept of joint measurability: $M_{a|x} = \sum_\lambda G_\lambda p(a|x, \lambda)$, where $\{G_\lambda\}$ represents the joint (or parent) POVM [1, 2].

Illustrative example.— To illustrate the concept, we present an example involving mutually unbiased bases (MUBs). Recall that two bases $\{|\varphi_i^1\rangle\}_i$ and $\{|\varphi_j^2\rangle\}_j$ are termed mutually unbiased if $|\langle \varphi_i^1 | \varphi_j^2 \rangle|^2 = 1/d$ for all i and j . There are at most $d + 1$ MUBs in dimension d , and a construction for the complete set of $d + 1$ MUBs is only known when d is a power of a prime number [17].

Here we consider a set \mathcal{M} consisting of m measurements in MUBs subject to white noise. That is, projection-valued measures (PVMs) composed of projectors $P_{a|x} = |\varphi_a^x\rangle\langle \varphi_a^x|$ preceded by a white noise channel. The resulting POVMs read

$$M_{a|x}^\eta = \eta P_{a|x} + (1 - \eta) \frac{\mathbb{1}}{d}, \quad (2)$$

where $x \in \{1, \dots, m\}$ and $a \in \{1, \dots, d\}$. Note that we use the standard construction of MUBs [17] with $\{|\varphi_i^1\rangle\}_i$ being the computational basis.

We now show that, depending on the amount of noise, $1 - \eta$, the set \mathcal{M} becomes **n -dimensional simulable**. Let us first construct an appropriate map, implementing the compression from dimension d to n . For a given basis $\{|\psi_i\rangle\}_i$ we consider the set of $\binom{d}{n}$ projectors Π_λ onto an n -dimensional subspace of \mathbb{C}^d spanned by the vectors of the basis.

This defines an instrument $\{\mathcal{E}_\lambda\}$, with each \mathcal{E}_λ given by a Kraus operator $K_\lambda = \sqrt{\binom{d}{n}^{-1} \frac{d}{n}} \Pi_\lambda$, compressing from dimension d to n . Here $\binom{d}{n}$ is simply the number of projectors Π_λ with n ones on the diagonal, and the term $\sqrt{\frac{d}{n}}$ is due to normalisation. Next, for any POVM P_a acting on \mathbb{C}^d , define its restriction to the subspace labelled by λ as

$$N_{a|\lambda} = \Pi_\lambda P_a \Pi_\lambda, \quad (3)$$

which is a POVM on \mathbb{C}^n since $N_{a|\lambda} \geq 0$ and $\sum_a N_{a|\lambda} = \Pi_\lambda \mathbb{1}_d \Pi_\lambda = \mathbb{1}_n$. We can now compute

$$\sum_\lambda \mathcal{E}_\lambda^*(N_{a|\lambda}) = \frac{n-1}{d-1} P_a + \left(1 - \frac{n-1}{d-1}\right) \mathcal{T}_{\{|\psi_i\rangle\}}[P_a] \quad (4)$$

where $\mathcal{T}_{\{|\psi_i\rangle\}}[P_a] = \sum_i |\psi_i\rangle\langle \psi_i| P_a |\psi_i\rangle\langle \psi_i|$ is the twirling map in the basis $\{|\psi_i\rangle\}$ used to define the instrument. In particular, if the eigenbasis of P_a and $\{|\psi_i\rangle\}$ are mutually unbiased, one obtains $\mathcal{T}_{\{|\psi_i\rangle\}}[P_a] = \frac{\mathbb{1}}{d}$. While if they coincide, one trivially gets $\mathcal{T}_{\{|\psi_i\rangle\}}[P_a] = P_a$.

If the set \mathcal{M} is composed of m MUBs, we construct an instrument that chooses one of the bases $y = 1, \dots, m$ randomly and performs $\{\mathcal{E}_{\lambda|y}\}$ as defined above. Then, for the setting x Bob does the measurement $N_{a|\lambda,x,y}$ as defined in Eq. (3). This construction results in

$$\sum_{\lambda,y} \frac{1}{m} \mathcal{E}_{\lambda|y}^*(N_{a|\lambda,x,y}) = \frac{1}{m} P_{a|x} + \frac{m-1}{m} \left[\frac{n-1}{d-1} P_{a|x} + \left(1 - \frac{n-1}{d-1}\right) \frac{\mathbb{1}}{d} \right], \quad (5)$$

which equals $M_{a|x}^\eta$ for $1 - \eta = \frac{m-1}{m} \left(1 - \frac{n-1}{d-1}\right)$, and implies the following observation.

Result 1. *The set \mathcal{M} in Eq. (2) of m noisy projective measurements in MUBs on \mathbb{C}^d is **n -dimensional simulable** if*

$$\eta \leq 1 - \frac{m-1}{m} \cdot \frac{d-n}{d-1}. \quad (6)$$

Let us first discuss the case of a pair of MUBs, i.e., $m = 2$. It is known that the pair is jointly measurable (i.e., **1-dimensional simulable**) if and only if $\eta \leq \eta^* = \frac{1}{2} \left(1 + \frac{1}{1+\sqrt{d}}\right)$ [18, 19]. Hence it follows that for a noise parameter satisfying $1 - \frac{1}{2} \frac{d-n}{d-1} \geq \eta > \eta^*$ (which is possible for all $d > 2$) we get a set of measurements that is incompatible but nevertheless **n -dimensional simulable** (for some $n > 1$). Alternatively, consider the full set of $m = 4$ MUBs in $d = 3$. This set is incompatible if and only if $\eta > (1 + 3\sqrt{5}) \approx 0.4818$ [19] while it is **2-dimensional simulable** for $\eta \leq \frac{5}{8} = 0.625$.

Finally, note that the above construction for **n -dimensional simulability** uses a heuristic choice of bases for the compression instrument. We verify below that this choice is sub-optimal.

SDP method.— In order to explore more general schemes, involving arbitrary bases for the compression, we now present a numerical approach based on semi-definite programming (SDP). For the example of MUBs, this shows that better schemes are indeed possible.

Consider a set of measurements $M_{a|x}$, to which we add noise. For clarity we focus here on the case of white noise, but the technique applies in general. Formally, we consider sets of POVMs of the form:

$$M_{a|x}^\eta = \eta M_{a|x} + (1 - \eta) \text{Tr}[M_{a|x}] \frac{\mathbb{1}_d}{d}.$$

Our goal now is to derive a lower bound on the critical value η^* , below which the set is n -dimensional simulable. In particular, if the lower bound is found to be trivial, i.e., $\eta \geq 1$, we conclude that the original measurements $M_{a|x}$ are n -dimensional simulable.

We first choose a compression map, i.e an instrument $\{\mathcal{E}_\lambda\}$ consisting of a finite number of CP maps \mathcal{E}_λ . For fixed compression map $\{\mathcal{E}_\lambda\}$ and measurements $M_{a|x}^\eta$ we can now find the maximal value of η (i.e., minimise the noise) while ensuring n -dimensional simulability, optimising over Bob's final measurements, via the following SDP:

$$\begin{aligned} \max_{\eta, \{\tilde{N}_{a|x,\lambda}\}} \quad & \eta \\ \text{s.t.} \quad & \sum_{\lambda} \mathcal{E}_\lambda \tilde{N}_{a|x,\lambda} \mathcal{E}_\lambda^\dagger = M_{a|x}^\eta \quad \forall a, x \\ & \sum_a \tilde{N}_{a|x,\lambda} \leq \mathbb{1}_n \quad \forall x, \lambda \\ & \tilde{N}_{a|x,\lambda} \geq 0 \quad \forall a, x, \lambda, \end{aligned} \quad (7)$$

where $\{\tilde{N}_{a|x,\lambda}\}_{a,x,\lambda}$ is the set of subnormalized measurements in dimension n . To illustrate the relevance of this method, let us consider again the case of a pair of MUBs in dimension $d = 3$, i.e., setting $M_{a|x} = |\varphi_a^x\rangle\langle\varphi_a^x|$, and compressing to $n = 2$. Here we choose a simple compression channel: we first choose a basis (via a unitary U_μ) and perform a projection onto the $\binom{d}{n}$ n -dimensional subspaces, denoted by projectors Π_i . Hence we set $\mathcal{E}_\lambda = U_\mu \Pi_i$ with $\mu \in 0, \dots, |\mu| - 1$, $i \in \{0, \dots, \binom{d}{n} - 1\}$ and hence $\lambda \in \{0, \dots, |\mu| \binom{d}{n} - 1\}$. Optimising over choices of $|\mu| = 2$ and $|\mu| = 3$ basis, we find noise thresholds of $\eta \approx 0.7803$ and $\eta \approx 0.8281$, respectively, hence clearly improving upon the bound of $\eta = 5/8$ we got in the analytical construction, see Eq. (6). Moreover, when allowing for more basis (up to $|\mu| = 5$), we could find no improvement. The Matlab code can be found in the Supplementary Material [20].

The case of all projective measurements.— So far we analysed sets \mathcal{M} with finitely many measurements. Now we turn our attention to continuous sets of measurements. Precisely, we consider assemblages \mathcal{M}_{PVM}^η made

of all rank-1 projective measurements subjected to white noise. Note that this automatically extends to all projective measurements which can be obtained from rank-1 projective measurements by post-processing. Assuming an isotropic noise, our set is made of all POVMs

$$\mathcal{M}_{PVM}^\eta = \{M_{a|U}^\eta\}_U \quad \text{with} \quad M_{a|U}^\eta = U^\dagger M_a^\eta U, \quad (8)$$

where U runs through all unitary operators on \mathbb{C}^d and M_a^η is the noisy (or lossy) measurement in the computational basis. On the one hand, the continuous case may seem more complicated as the infinite number of possible measurements cannot be tackled with the SDP of Eq. (7). On the other hand, the symmetry of the set helps simplifying the optimal compression scheme as we shall see now.

The set \mathcal{M}_{PVM}^η is a particular case of what we call an *invariant assemblage*. This is a set \mathcal{M} such that $M_a \in \mathcal{M}$ implies that $U^\dagger M_a U$ is also in \mathcal{M} for all unitaries U . For such an invariant assemblage the choice of the compression bases does not play any role, which leads to the following observation.

Result 2. *An n -dimensional simulable invariant assemblage \mathcal{M} can be compressed with the continuous instrument $\{\mathcal{E}_V\}_V$ with CP map density \mathcal{E}_V , such that $\mathcal{E}_V(\rho) = K_V \rho K_V^\dagger$ with*

$$K_V = \sqrt{\frac{d}{n}} \Pi_n V, \quad (9)$$

with V running through all unitary operators on \mathbb{C}^d and Π_n is a projection onto a fixed n -dimensional subspace of \mathbb{C}^d . In addition, if there exists a measurement $M_a \in \mathcal{M}$ such that $U^\dagger M_a U$ generates all of \mathcal{M} (i.e. the action is transitive), then it suffices to find $N_{a|V}$ satisfying $M_a = \int dV \mathcal{E}_V^*(N_{a|V})$, as then $U^\dagger M_a U = \int dV \mathcal{E}_V^*(N_{a|VU^\dagger})$.

Here and below $dV = d\mu(V)$ is the Haar measure. If there are multiple orbits, that is, a family of measurements \mathcal{N} such that $U^\dagger \mathcal{N} U$ generates all of \mathcal{M} , then it suffices to find a measurement $N_{a|V}$ as above for each measurement in \mathcal{N} . A detailed proof of the result can be found in the Supplementary Material [20] and we explain only the intuition here. First, because of the invariance of \mathcal{M} one can freely apply any (random) unitary V to the state before the compression. Thus if the set is compressible with some instrument $\{\mathcal{E}_\lambda\}$ it is also compressible with an instrument where a random V is applied before $\{\mathcal{E}_\lambda\}$. Second, one shows that any such instrument can be obtained by post-processing of $\{\mathcal{E}_V\}$ defined in the statement of the result. Finally, the relation between the compressed POVMs $N_{a|V}$ and $N_{a|VU^\dagger}$ is a simple consequence of Haar measure invariance.

All noisy projective measurements.— We now consider the set of all projective measurements \mathcal{M}_{PVM}^η in Eq. (8) subject to white noise

$$M_a^\eta = \eta |a\rangle\langle a| + \frac{1-\eta}{d} \mathbb{1}, \quad (10)$$

where $|a\rangle$ denotes the d vectors of the computational basis. We now want to find the $N_{a|V}$ such that $\int dV \mathcal{E}_V^*(N_{a|V})$ equals M_a^η for the highest value of η possible. In fact, the optimal compressed POVM here is given by

$$N_{a|V} = \operatorname{argmax}_{\tilde{N}_a: \text{POVM on } \mathbb{C}^n} \sum_{a=1}^d \langle a|V^\dagger \tilde{N}_a V|a\rangle \quad (11)$$

with \tilde{N}_a embedded in \mathbb{C}^d , which can be seen in two steps, c.f. Supplementary Material [20] for details. First, this choice does result in an operator of the desired form $M'_a = \eta(x) |a\rangle\langle a| + \frac{1-\eta(x)}{d} \mathbb{1}$ with $\eta(x) = \frac{dx-1}{d-1}$ and

$$x = \frac{1}{n} \int dV \max_{\tilde{N}_a: \text{POVM on } \mathbb{C}^n} \sum_{a=1}^d \langle a|V^\dagger \tilde{N}_a V|a\rangle. \quad (12)$$

Second, by construction with the max inside the integral this yields the highest value of x , and thus η , possible (this is precisely the intuition behind the definition (11)). This leads to the following full characterisation of ***n-dimensional simulability*** of noisy PVMs in any finite dimensional system.

Result 3. *The set of all noisy (white noise) projective measurements \mathcal{M}_{PVM}^η in dimension d is ***n-dimensional simulable*** if and only if*

$$\eta \leq \eta_{d \rightarrow n} = \frac{dx-1}{d-1} \quad (13)$$

with x given in Eq. (12) where dV is the Haar measure on the unitary operators V on \mathbb{C}^d , $\{|a\rangle\}_{a=1}^d$ is the computational basis of \mathbb{C}^d , and \mathbb{C}^n is any n -dimensional subspace of \mathbb{C}^d .

$n \setminus d$	2	3	4	5	6
1	<i>0.5</i>	<i>0.42</i>	<i>0.36</i>	<i>0.32</i>	<i>0.29</i>
2		0.70	0.56	0.48	0.42
3			0.77	0.64	0.56
4				0.81	0.70
5					0.84

TABLE I. Some values of the white noise threshold $\eta_{d \rightarrow n}$ for the ***n-dimensional simulability*** of \mathcal{M}_{PVM}^η – the set of all projective measurements in dimension d . We computed the values with Wolfram Mathematica using the build-in function `CircularUnitaryMatrixDistribution` to sample from the Haar measure (5000 points in each case), and `SemidefiniteOptimization` to solve the optimal POVMs \tilde{N}_a in Eq. (12), the code can be found in the Supplementary Material [20]. The values in italic give the known white noise threshold for the incompatibility of all PVMs and equal to $\eta_{d \rightarrow 1} = (\sum_{k=1}^d \frac{1}{k} - 1)/(d-1)$, cf. [11, 21].

The POVM maximisation Eq. (12) is a simple SDP, so the threshold $\eta_{d \rightarrow n}$ can be computed numerically by sampling from the Haar measure (or integrating numerically) and solving the SDP for each V . We report some values in Table I.

An upper bound on the threshold can be obtained by applying the Cauchy-Schwarz inequality to the sum in Eq. (12), and leads to $\eta_{d \rightarrow n} \leq \frac{d\sqrt{\frac{n+1}{d+1}}-1}{d-1}$, as we show in the Supplementary Material [20] by using results from Refs. [22, 23]. There we also derive a lower bound $\eta_{d \rightarrow (d-1)} \geq \frac{d^2-d(1+\sum_{k=1}^d \frac{1}{k})+1}{(d-1)^2}$ for the case $n = d-1$.

For $n = 1$ our considerations reduce to the joint measurability of noisy projective measurements, where the white noise threshold is known $\eta_{d \rightarrow 1} = (\sum_{k=1}^d \frac{1}{k} - 1)/(d-1)$ [21]. In this case the subspace \mathbb{C}^n contains a single state $|\Psi\rangle = V|1\rangle$. Accordingly to Eq. (12) $\eta_{d \rightarrow 1}$ can be computed from $x = \int d\Psi \max_a |\langle a|\Psi\rangle|^2 = \frac{1}{d} \sum_{k=1}^d \frac{1}{k}$, where $d\Psi$ is the uniform measure over states in \mathbb{C}^d (invariant under unitaries), which is equivalent to the derivation in [21].

All lossy projective measurements. — We now briefly analyse the set of all projective measurements \mathcal{M}_{PVM}^η in Eq. (8) subject to loss

$$M_a^\eta = \eta |a\rangle\langle a|; \quad M_\emptyset^\eta = (1-\eta)\mathbb{1} \quad (14)$$

This set describes measurements with a limited efficiency – the additional element M_\emptyset^η corresponds to the "no-click" outcome. The required rank-1 form of the operators $M_a^\eta \propto |a\rangle\langle a|$ implies that for $a \neq \emptyset$ the POVM element $N_{a|V}$ can only be nonzero if $\mathcal{E}_V^*(\tilde{N}_{a|V}) \propto |a\rangle\langle a|$. In other words, the vector $|a\rangle$ has to be in the n -dimensional subspace selected by the instrument, i.e.

$$|a\rangle \in \operatorname{span}\{V|1\rangle, \dots, V|n\rangle\}. \quad (15)$$

This condition is only fulfilled for a set of V of measure zero, hence $M'_a \propto |a\rangle\langle a| \implies M'_a = 0$. We can thus conclude that

Result 4. *For any positive efficiency $\eta > 0$ the set of all lossy projective measurements \mathcal{M}_{PVM}^η is not ***n-dimensional simulable*** for any $n < d$.*

An equivalent definition. — The notion of joint measurability has a direct connection to entanglement-breaking properties of quantum channels, cf. [13] and [24, 25]. As a final point we now discuss how this connection naturally extends to ***n-dimensional simulability***.

A quantum channel Λ is said to be *n-partially entanglement breaking* (*n*-PEB) if, for all states ρ , the Schmidt number of the state $(\Lambda \otimes \mathbb{1})[\rho]$ is less than or equal to n [16]. In the finite-dimensional setting, this is equivalent to the existence of Kraus operators $\{K_\lambda\}$ of Λ each with $\operatorname{rank}(K_\lambda) \leq n$ [16]. This notion provides an

alternative way to define *n-dimensional simulability*:

Claim 5. *A measurement assemblage $M_{a|x}$ is *n-dimensional simulable* if and only if there exists a quantum channel Λ that is *n-PEB* and a measurement assemblage $N_{a|x}$ such that*

$$M_{a|x} = \Lambda^*(N_{a|x}). \quad (16)$$

For the if direction one uses the singular value decomposition of the rank- n Kraus operators $\{K_\lambda\}$ and $N_{a|x}$ to define the instrument $\{\mathcal{E}_\lambda\}$ with n -dimensional output and the subsequent measurement $N_{a|x,\lambda}$. For the only if direction one can simply view any compression instrument $\{\mathcal{E}_\lambda\}$ as a channel Λ that outputs a quantum system of dimension n and a classical register encoding λ . This channel is manifestly *n-PEB*. The details of the proof can be found in the Supplementary Material [20]. Note that we prove the result for finite amount of classical communication λ , respectively a channel Λ with a finite dimensional output and conjecture it to be true in general.

Related concepts.— We now compare the idea of *n-dimensional simulability* with some previously introduced notions.

First, in [26] a related notion of “*n-compressibility*” of a set of quantum measurements has been proposed. Similarly to our definition a set of measurements is said to be *n-compressible* if $M_{a|x} = \sum_\lambda \mathcal{E}_\lambda^*(N_{a|x,\lambda})$ ¹, but in sharp contrast the compressed measurement has to take the form $N_{a|x,\lambda} = \tilde{\mathcal{E}}_\lambda^*(M_{a|x})$. Here, $\tilde{\mathcal{E}}_\lambda$ is a set of CPTP maps decompressing the system back into dimension d , and the same measurement $M_{a|x}$ has to be used after the decompression. Clearly, a set of *n-compressible* measurements is *n-dimensional simulable* by construction, but not the other way around. In particular, it is known that single measurements can have a larger compression dimension than one [26].

Other works have investigated measurement (in)compatibility in subspaces [27–29], while Ref. [30] defined a concept of *n-compatibility* considering a scenario where a set of measurements is performed on n copies of a state. As far as we can say, these concepts are unrelated to *n-dimensional simulability*.

Finally, the problem of implementing POVMs with given resources has been investigated [31–33], for example the implementation of POVMs using only projective measurements of the same dimension [34–36]. This question has been connected to the compression of quantum information [31].

¹ Notably the authors of [26] also consider the possibility to bound the amount of classical communication via the number of different instrument outcomes λ (the dimension of the classical register output by the instrument).

Conclusion.— We have introduced the concept of *n-dimensional simulability* of a set of measurements, motivated by a scenario of compression of quantum information. When full compression is possible (i.e. $n=1$), our notion corresponds to joint measurability. Hence our approach can be used as a quantification of measurement incompatibility in terms of dimension. We discussed a number of examples, providing analytical constructions as well as numerical methods.

More generally, the concept of *n-dimensional simulability* turns out to be connected to several other relevant notions of quantum information theory. First, as we showed above, *n-dimensional simulability* relates to partially entanglement-breaking channels. Second, there is a direct connection between *n-dimensional simulability* and the notion of genuine high-dimensional steering [37], which will be presented in detail in a companion article [38]. These links generalise the well-known connection between joint measurability, steering and quantum channels [10–12]. They also open new questions, for example, whether the dimensionality of a quantum channel could be tested in a partially device-independent manner.

Beyond quantum compression, our approach also has implications for device-independent quantum information processing. For example, it is clear that a set of measurements that is *n-dimensional simulable* is of limited use for randomness certification, as it can lead to (at most) $2 \log(n)$ bits of randomness in a black-box setting.

Finally, an intriguing question is whether our approach could be generalised to the case of continuous variable measurements. It turns out that, in the infinite-dimensional case, the Kraus operators’ ranks and entanglement breaking properties of a channel are not anymore tied together [39, 40], opening different possibilities to extend our concept.

Acknowledgments.— We thank Denis Rosset, Costantino Budroni, Paul Skrzypczyk, Alex Little, Michalis Skotiniotis, and Thomas Cope for discussions. We acknowledge financial support from the Swiss National Science Foundation (projects 192244, Ambizione PZ00P2-202179, and NCCR SwissMAP). BDMJ acknowledges support from UK EPSRC (EP/SO23607/1).

* These authors contributed equally to this work

- [1] Teiko Heinosaari, Takayuki Miyadera, and Mário Ziman. An invitation to quantum incompatibility. *Journal of Physics A: Mathematical and Theoretical*, 49(12):123001, Feb 2016.
- [2] Otfried Gühne, Erkkka Haapasalo, Tristan Kraft, Juha-Pekka Pellonpää, and Roope Uola. Incompatible measurements in quantum information science. *arXiv:2112.06784*, 2021.
- [3] P. Busch, P. Lahti, J.-P. Pellonpää, and K.Ylinen. *Quantum Measurement (Theoretical and Mathematical*

- Physics*). Springer, 2016.
- [4] C. Carmeli, T. Heinosaari, and A. Toigo. Quantum incompatibility witnesses. *Phys. Rev. Lett.*, 122:130402, 2019.
 - [5] P. Skrzypczyk, I. Šupić, and D. Cavalcanti. All sets of incompatible measurements give an advantage in quantum state discrimination. *Phys. Rev. Lett.*, 122:130403, 2019.
 - [6] Michał Oszmaniec and Tanmoy Biswas. Operational relevance of resource theories of quantum measurements. *Quantum*, 3:133, Apr 2019.
 - [7] Roope Uola, Tristan Kraft, Jiangwei Shang, Xiao-Dong Yu, and Otfried Gühne. Quantifying quantum resources with conic programming. *Phys. Rev. Lett.*, 122:130404, 2019.
 - [8] Roope Uola, Tom Bullock, Tristan Kraft, Juha-Pekka Pellonpää, and Nicolas Brunner. All quantum resources provide an advantage in exclusion tasks. *Phys. Rev. Lett.*, 125:110402, 2020.
 - [9] Andrés F. Ducuara and Paul Skrzypczyk. Operational interpretation of weight-based resource quantifiers in convex quantum resource theories. *Phys. Rev. Lett.*, 125:110401, Sep 2020.
 - [10] Marco Túlio Quintino, Tamás Vértesi, and Nicolas Brunner. Joint measurability, einstein-podolsky-rosen steering, and bell nonlocality. *Phys. Rev. Lett.*, 113:160402, 2014.
 - [11] Roope Uola, Tobias Moroder, and Otfried Gühne. Joint measurability of generalized measurements implies classicality. *Phys. Rev. Lett.*, 113:160403, 2014.
 - [12] Roope Uola, Costantino Budroni, Otfried Gühne, and Juha-Pekka Pellonpää. One-to-one mapping between steering and joint measurability problems. *Phys. Rev. Lett.*, 115:230402, 2015.
 - [13] J. Kiukas, C. Budroni, R. Uola, and J.-P. Pellonpää. Continuous-variable steering and incompatibility via state-channel duality. *Phys. Rev. A*, 96:042331, 2017.
 - [14] Teiko Heinosaari and Takayuki Miyadera. Incompatibility of quantum channels. *Journal of Physics A: Mathematical and Theoretical*, 50(13):135302, Mar 2017.
 - [15] Leonardo Guerini, Marco Túlio Quintino, and Leandro Aolita. Distributed sampling, quantum communication witnesses, and measurement incompatibility. *Physical Review A*, 100(4):042308, 2019.
 - [16] Dariusz Chruściński and Andrzej Kossakowski. On partially entanglement breaking channels. *Open Systems & Information Dynamics*, 13(1):17–26, 2006.
 - [17] William K Wootters and Brian D Fields. Optimal state-determination by mutually unbiased measurements. *Ann. Phys.*, 191(2):363–381, 1989.
 - [18] Roope Uola, Kimmo Luoma, Tobias Moroder, and Teiko Heinosaari. Adaptive strategy for joint measurements. *Physical Review A*, 94(2), Aug 2016.
 - [19] Sébastien Designolle, Paul Skrzypczyk, Florian Fröwis, and Nicolas Brunner. Quantifying measurement incompatibility of mutually unbiased bases. *Phys. Rev. Lett.*, 122:050402, 2019.
 - [20] See Supplemental Material at [url will be inserted by publisher] for the proof of Result 2 (Section A), the proof of Result 3 (Section B), an upper bound on the n -simulability of all noisy PVMs (Section C), a lower bound for the simulability of noisy PVMs of $n = d - 1$ (Section D), and the proof of Result 5 (Section E). DiscreteSDP.zip contains the Matlab code used to obtain the numerical results discussed after equation (7). Table1.nb contains the Mathematica code that was used to produce the results of Table 1.
 - [21] Howard M Wiseman, Steve James Jones, and Andrew C Doherty. Steering, entanglement, nonlocality, and the einstein-podolsky-rosen paradox. *Physical review letters*, 98(14):140402, 2007.
 - [22] Aram W Harrow. The church of the symmetric subspace. *arXiv preprint arXiv:1308.6595*, 2013.
 - [23] Christoph Spengler, Marcus Huber, and Beatrix C Hiesmayr. Composite parameterization and haar measure for all unitary and special unitary groups. *Journal of mathematical physics*, 53(1):013501, 2012.
 - [24] Ioannis Kogias, Paul Skrzypczyk, Daniel Cavalcanti, Antonio Acín, and Gerardo Adesso. Hierarchy of steering criteria based on moments for all bipartite quantum systems. *Physical Review Letters*, 115(21), Nov 2015.
 - [25] Tobias Moroder, Oleg Gittsovich, Marcus Huber, Roope Uola, and Otfried Gühne. Steering maps and their application to dimension-bounded steering. *Physical Review Letters*, 116(9), Mar 2016.
 - [26] Andreas Bluhm, Lukas Rauber, and Michael M Wolf. Quantum compression relative to a set of measurements. In *Annales Henri Poincaré*, volume 19, pages 1891–1937. Springer, 2018.
 - [27] Jukka Kiukas. Subspace constraints for joint measurability. *Journal of Physics: Conference Series*, 1638(1):012003, oct 2020.
 - [28] Faedi Loulidi and Ion Nechita. The compatibility dimension of quantum measurements. *Journal of Mathematical Physics*, 62(4):042205, Apr 2021.
 - [29] Roope Uola, Tristan Kraft, Sébastien Designolle, Nikolai Miklin, Armin Tavakoli, Juha-Pekka Pellonpää, Otfried Gühne, and Nicolas Brunner. Quantum measurement incompatibility in subspaces. *Physical Review A*, 103(2), Feb 2021.
 - [30] Claudio Carmeli, Teiko Heinosaari, Daniel Reitzner, Jussi Schultz, and Alessandro Toigo. Quantum incompatibility in collective measurements. *Mathematics*, 4(3):54, Sep 2016.
 - [31] Andreas Winter. “extrinsic” and “intrinsic” data in quantum measurements: Asymptotic convex decomposition of positive operator valued measures. *Communications in Mathematical Physics*, 244(1):157–185, 2004.
 - [32] G Sentís, B Gendra, S D Bartlett, and A C Doherty. Decomposition of any quantum measurement into extremals. *Journal of Physics A: Mathematical and Theoretical*, 46(37):375302, aug 2013.
 - [33] Tanmay Singal, Filip B. Maciejewski, and Michał Oszmaniec. Implementation of quantum measurements using classical resources and only a single ancillary qubit. *npj Quantum Information*, 8(1):82, 2022.
 - [34] Michał Oszmaniec, Leonardo Guerini, Peter Wittek, and Antonio Acín. Simulating positive-operator-valued measures with projective measurements. *Physical Review Letters*, 119(19), Nov 2017.
 - [35] Leonardo Guerini, Jessica Bavaresco, Marcelo Terra Cunha, and Antonio Acín. Operational framework for quantum measurement simulability. *Journal of Mathematical Physics*, 58(9):092102, Sep 2017.
 - [36] Flavien Hirsch, Marco Túlio Quintino, Tamás Vértesi, Miguel Navascués, and Nicolas Brunner. Better local hidden variable models for two-qubit Werner states and an upper bound on the Grothendieck constant $K_G(3)$.

- Quantum*, 1:3, April 2017.
- [37] Sébastien Designolle, Vatshal Srivastav, Roope Uola, Natalia Herrera Valencia, Will McCutcheon, Mehul Malik, and Nicolas Brunner. Genuine high-dimensional quantum steering. *Physical Review Letters*, 126(20):200404, 2021.
- [38] B. Jones et al. to appear. 2022.
- [39] A. S. Holevo, M. E. Shirokov, and R. F. Werner. Separability and entanglement-breaking in infinite dimensions, 2005.
- [40] M. E. Shirokov. The schmidt number and partially entanglement breaking channels in infinite dimensions, 2011.