



This is the accepted manuscript made available via CHORUS. The article has been published as:

Emergent Glassy Behavior in a Kagome Rydberg Atom Array

Zheng Yan, Yan-Cheng Wang, Rhine Samajdar, Subir Sachdev, and Zi Yang Meng
Phys. Rev. Lett. **130**, 206501 — Published 18 May 2023

DOI: [10.1103/PhysRevLett.130.206501](https://doi.org/10.1103/PhysRevLett.130.206501)

Emergent glassy behavior in a kagome Rydberg atom array

Zheng Yan,¹ Yan-Cheng Wang,² Rhine Samajdar,^{3,4} Subir Sachdev,^{5,*} and Zi Yang Meng^{1,†}

¹*Department of Physics and HKU-UCAS Joint Institute of Theoretical and Computational Physics, The University of Hong Kong, Pokfulam Road, Hong Kong SAR, China*

²*Beihang Hangzhou Innovation Institute Yuhang, Hangzhou 310023, China*

³*Department of Physics, Princeton University, Princeton, NJ, 08544, USA*

⁴*Princeton Center for Theoretical Science, Princeton University, Princeton, NJ, 08544, USA*

⁵*Department of Physics, Harvard University, Cambridge, MA 02138, USA*

(Dated: April 11, 2023)

We present large-scale quantum Monte Carlo simulation results on a realistic Hamiltonian of kagome-lattice Rydberg atom arrays. Although the system has no intrinsic disorder, intriguingly, our analyses of static and dynamic properties on large system sizes reveal *emergent* glassy behavior in a region of parameter space located between two valence bond solid phases. The extent of this glassy region is demarcated using the Edwards-Anderson order parameter, and its phase transitions to the two proximate valence bond solids—as well as the crossover towards a trivial paramagnetic phase—are identified. We demonstrate the intrinsically slow (imaginary) time dynamics deep inside the glassy phase and discuss experimental considerations for detecting such a quantum disordered phase with numerous nearly degenerate local minima. Our proposal paves a new route to the study of real-time glassy phenomena and highlights the potential for quantum simulation of a distinct phase of quantum matter beyond solids and liquids in current-generation Rydberg platforms.

Introduction.—Over the last decade, quantum simulators based on programmable Rydberg atom arrays [1–5] have emerged as powerful platforms for the investigation of highly correlated quantum matter. These systems have opened up new avenues to study interesting many-body states [6–9], quantum dynamics [10–12], gauge theories [13–16], and even combinatorial optimization problems [17–19].

An especially promising direction that has recently attracted much attention is the simulation of quantum phases of matter in these tunable atomic setups. Such phases and the transitions between them have been intensely studied for Rydberg atoms arrayed in one spatial dimension [20–23] as well as in various two-dimensional geometries, including on the square [24–28], triangular [29], honeycomb [30, 31], kagome [32], and ruby [33–35] lattices. In particular, Ref. 32 identified an intriguing highly correlated regime in the phase diagram of the kagome-lattice Rydberg atom array characterized by a lack of symmetry-breaking solid order and a large entanglement entropy. The correlations in this region were found to be “liquid-like” in that the density of excitations is limited by the strong Rydberg-Rydberg interactions (as opposed to a weakly interacting gas wherein the laser drive induces independent atomic excitations). Mapping this system to a quantum dimer model [36] on the triangular lattice [37–39] raises the possibility of a spin liquid phase with \mathbb{Z}_2 topological order [32, 40]. However, since the precise microscopic interactions differ between the Rydberg [32] and dimer [40] models, it is crucial to *independently* establish the properties of the former in the thermodynamic limit. This poses a challenging problem for numerical techniques such as the density-matrix renormalization group (DMRG), which is not only hindered by the geometrically frustrated and long-ranged




FIG. 1. Phase diagram, spanned by the δ/Ω and R_b axes, obtained from QMC simulations. The four stars mark the points in each phase for which the equal-time dimer structure factor is presented in Fig. 2. The white dashed lines indicate the cuts in parameter space along which the quantum phase transitions are studied in Figs. 3 and S2. The colored background shows the Edwards-Anderson order parameter as obtained in Fig. 3(c), with the colorbar on top denoting the scale of q_{EA} . The two insets schematically sketch the two crystalline phases (nematic and staggered). Each red (white) circle representing an atom in the Rydberg (ground) state on the kagome lattice [black] can be mapped to the presence (absence) of a dimer on the medial triangular lattice [white].

nature of the Hamiltonian in Eq. (1) but is also limited to relatively small system sizes on cylinders.

In light of the situation, here, we overcome this obstacle and present large-scale quantum Monte Carlo (QMC) simulation results on the realistic Hamiltonian of kagome-lattice Rydberg arrays in Eq. (1) below. Surprisingly,

even though the Hamiltonian is translationally invariant and has no disorder, our unbiased numerical results for large system sizes and dynamic and static data reveal *emergent* glassy behavior [41] in the region located between the so-called “nematic” [38, 42] and “staggered” [43] valence bond solid (VBS) phases [32]. Although disorder-free glassiness was observed in previous results obtained in extended Heisenberg models [44], it is novel in the context of the realistic Rydberg arrays. Moreover, we emphasize that unlike previous work that identified glassy behavior in open dissipative Rydberg gases [45, 46], our findings here apply to an isolated closed quantum system. We utilize the Edwards-Anderson order parameter to map out the extent of the glassy region in the phase diagram. Furthermore, the phase transitions between the glassy phase and the two valence bond phases as well as the crossover towards the paramagnetic phase are identified. Our results highlight the intrinsically slow (imaginary) time dynamics deep inside the glassy phase, and we suggest experimental protocols to detect such a quantum disordered phase with numerous nearly degenerate local minima in its energy landscape.

Rydberg Hamiltonian on the kagome lattice.—We investigate the following realistic Hamiltonian describing Rydberg arrays on the kagome lattice,

$$H = \sum_{i=1}^N \left[\frac{\Omega}{2} (|g\rangle_i \langle r| + |r\rangle_i \langle g|) - \delta |r\rangle_i \langle r| \right] + \sum_{i,j=1}^N \frac{V_{ij}}{2} (|r\rangle_i \langle r| \otimes |r\rangle_j \langle r|), \quad (1)$$

where the sum on i runs over all N sites of the kagome lattice. The ket $|g\rangle$ ($|r\rangle$) represents the ground (Rydberg) state, while Ω (δ) stands for the Rabi frequency (detuning) of the laser drive, which can be mapped to a transverse (longitudinal) field in the language of quantum Ising model. The repulsive interaction is of the van der Waals form $V_{ij} = \Omega R_b^6 / R_{ij}^6$, where R_{ij} is the distance between the sites i and j , and R_b defines the Rydberg blockade radius (within which no two atoms can be simultaneously excited to the Rydberg state). Note that we are implicitly working in units where the lattice spacing is set to one. Since V_{ij} falls off rapidly with the sixth power of the interatomic distance, we truncate the interactions beyond a cutoff of third-nearest neighbours in our simulations, akin to Ref. 32. We set $\Omega = 1$ and scan the parameters δ and R_b to explore the phase diagram, paying particular attention to the previously identified correlated region between the solid phases.

To solve the model in Eq. (1) in an unbiased manner, we modify and employ several stochastic series expansion (SSE) QMC schemes [40, 47–53] to deal with such Rydberg arrays. By monitoring the behavior of various physical observables, e.g., correlation functions and structure factors, we map out the detailed phase di-

agram in Fig. 1. Our simulations are performed on the kagome lattice with periodic boundary conditions and system sizes $N = 3L^2$ for linear dimensions $L = 6, 8, 12$, while setting the inverse temperature $\beta = L$ to scale to the ground state. Besides the conventional observables employed in recent QMC work [40], here, to reveal the intricate nature of the glassy phase and its transitions or crossovers to the neighboring phases, we employ different annealing, quench, and parallel tempering schemes [19, 54–62]. Scanning along parameter paths in the phase diagram, we compute the Edwards-Anderson order parameter to detect the spin-glass behavior [63–66]. More details about these schemes can be found in the Supplemental Material (SM) [67].

Phase diagram.—The phase diagram thus obtained is illustrated in Fig. 1. When δ/Ω is small but positive, we observe a disordered paramagnetic (PM) phase. Once δ/Ω is tuned to larger values, we find two symmetry-breaking VBS phases, in agreement with previous DMRG results [32] but with slightly shifted phase boundaries. These solid phases, termed nematic and staggered, correspond to (approximately) 1/3 and 1/6 filling of Rydberg excitations, respectively. The schematic plots of these crystalline phases are sketched in the insets of Fig. 1, both on the direct kagome lattice in the Rydberg basis and on the medial triangular lattice in the dimer basis. The exact manner in which they connect to each other, in the thermodynamic limit, is an interesting open question, with possibilities including topologically ordered even or




FIG. 2. Equal-time dimer structure factors $S(\mathbf{k}, \tau=0)$ in the Brillouin zone for the (a) 1/6 staggered ($R_b = 2.3$), (b) glass ($R_b = 2.1$), (c) nematic ($R_b = 1.9$) and (d) PM ($R_b = 1.05$) phases at $\delta/\Omega = 3.3$ (vertical cut in the phase diagram of Fig. 1). The data shown here is simulated for $\beta = L = 12$. The number in the upper-right corner shows the enlargement factor of the color bar; e.g., $\times 10$ changes the color bar from $[0, 0.1]$ to $[0, 1]$.




FIG. 3. (a) Energy per site plotted along the cut with $\delta/\Omega = 3.3$ in Fig. 1 using different initial states for the QMC simulations, showing that the nematic–glass phase transition is first-order while the nematic–PM one is also weakly first-order. (b) The loop order parameter (green dashed string), which can be used to distinguish the even/odd \mathbb{Z}_2 QSL ($\langle loop \rangle \neq 0$) from the trivial disordered phase without topological order ($\langle loop \rangle = 0$), is always close to zero, suggesting that the phase is not a pure even or odd \mathbb{Z}_2 QSL. (c) The Edwards-Anderson order parameter increases sharply upon going from the PM to the glass phase as δ/Ω increases at different R_b .

odd quantum spin liquids (QSLs), an intervening trivial disordered phase, or some new emergent intermediate phase [32, 40].

Interestingly, we discover that a glassy disordered phase—which can be distinguished from the PM by the magnitude of the Edwards-Anderson order parameter—exists in the central region between the two VBSs. The phase boundaries between this region and proximate phases are determined by examining various parameter points and paths scanning through the phase diagram, as denoted by the red stars and dashed lines in Fig. 1, and addressed in detail below.

To characterize the variety of phases, we first compute the equal-time ($\tau = 0$) structure factor (see Fig. 2) as

$$S(\mathbf{k}, \tau) = \frac{1}{N} \sum_{\substack{i,j \\ \alpha=1,2,3}}^{L^3} e^{i\mathbf{k}\cdot\mathbf{r}_{ij}} \left(\langle n_{i,\alpha}(\tau) n_{j,\alpha}(0) \rangle - \langle n_{i,\alpha} \rangle \langle n_{j,\alpha} \rangle \right), \quad (2)$$

where n_i is the density operator on site i and α stands for the three sublattices of the kagome lattice, at four representative parameter points corresponding to the four distinct phases in the phase diagram. Figures 2 (b) and (d) show $S(\mathbf{k}, 0)$ inside the glass and PM phases, respectively. In the hexagonal Brillouin zone, we observe that there are no peaks signifying long-range order but only broad profiles associated with different short-range density correlation patterns in real space. In contrast, Figs. 2 (a) and (c) present the structure factors inside the staggered and nematic phases, respectively, where one now clearly sees the Bragg peaks at the relevant ordering wavevectors.

Quantum phase transitions.—Having established the lack of long-range density correlations in both the PM and glass phases, we move on to study the associated quantum phase transitions [68]. Since the glass phase is expected to have many degenerate energy minima and very long autocorrelation times (which render the QMC simulation difficult), special care needs to be taken in de-

termining its phase boundaries. Our results in this regard are summarized in Fig. 3, which shows the data along several parameter scans in the phase diagram (dashed lines in Fig. 1).

First, in Fig. 3(a), we illustrate the energy density along the line $\delta/\Omega = 3.3$, computed with different initial states and annealing/quench schemes, to find the phase transition between the nematic and PM phases. The red curve shows the energy density simulated from random initial configurations with thermal annealing (by decreasing the temperature slowly) [55, 56]. On the other hand, the data plotted in blue is simulated from nematic configurations by quenching (i.e., starting at a very low temperature). Deep in the nematic phase, the two energy lines are clearly distinct. The difference between the two becomes small on progressing towards the transition point, where the two energy lines cross and then split weakly in the PM phase. Thus, the phase transition between the PM and nematic phases, which belongs to the (2+1)D three-state Potts universality class [32], is seen to be weakly first-order in consistency with prior findings [69]. This first-order phase transition can also be detected from the order parameter of the nematic phase, as detailed in the SM [67]. By the same logic, Fig. 3(a) also conveys that the transition between the glass and the nematic phase, which occurs at $R_b \sim 2$ in Fig. 1, is first-order as well. Scanning the energy density along the line $R_b = 1.9$, as shown in the SM [67], similarly manifests a first-order phase transition.

We now study the central disordered region between the two VBS phases, considering, in particular, the possible even and odd QSLs, or PM phases that emerge in an approximate quantum dimer model [40]. To this end, we define a nonlocal loop operator [9]—schematically shown by the green dashed loop in the inset of Fig. 3(b)—as $\langle loop \rangle = \langle (-1)^{\# \text{ cut dimers}} \rangle$, which measures the parity of the number of dimers intersected along a rhomboid with odd linear size on the medial dimer lattice. This

operator can be used to distinguish the two QSLs and the PM phase [9, 40]. In an odd (even) \mathbb{Z}_2 QSL without spin excitations, the value of $\langle loop \rangle$ is pinned to -1 ($+1$) because of the exact constraint requiring one (two) dimer(s) per site of the triangular lattice; this operator continues to be well-defined for a small density of matter fields [16]. From Fig. 3(b), we see that $\langle loop \rangle$ remains close to zero in the central correlated region, indicating that the ground state is not a pure even or odd \mathbb{Z}_2 QSL. As shown below, we further find that this region is also not a trivial PM phase, but, perhaps surprisingly for a homogeneous Hamiltonian, an emergent glass phase. This finding also underscores that the low-energy effective theory—a triangular lattice quantum dimer model with variable dimer density [40]—used to describe the physics proximate to the VBS phases departs from the realistic Rydberg Hamiltonian in this part of the phase diagram, where the Rydberg excitation density differs significantly from $1/6$ or $1/3$ (corresponding to the limit of one or two dimers per site, respectively). Moreover, the snapshots of the sampled configurations drawn in the SM [67] also demonstrate that apart from the Rydberg blockade, all local constraints (associated with the \mathbb{Z}_2 topological order) are relaxed in this glass region.

To differentiate between the PM and glass phases, we utilize the Edwards-Anderson order parameter [63–66, 70] $q_{EA} = \sum_{i=1}^N \langle n_i - \rho \rangle^2 / [N\rho(1 - \rho)]$, where $n_i \equiv |r\rangle_i \langle r|$, ρ is the average density defined as $\rho \equiv \sum_{i=1}^N \langle n_i \rangle / N$, and $\langle \dots \rangle$ indicates a statistical average over Monte Carlo snapshots [44, 71]. The spin-glass order is characterized by the breaking of translational invariance and the Edwards-Anderson order parameter $q_{EA} \in [0, 1]$, which captures the on-site deviation from the average density, is a measure of the glassy behavior. The large magnitude of q_{EA} shown in Fig. 3(c) demonstrates the emergent glassy nature of the disordered region amid the VBS phases. While q_{EA} clearly tells the PM and glass phases apart, whether the two are separated by a phase transition or by a crossover is an interesting open question. It is also noteworthy that this Edwards-Anderson order parameter decays extremely slow with increasing the number of Monte Carlo steps [see Fig. 4(b)]; this is another signature of the slow dynamics in the glass phase [72], which is consistent with the small and nearly degenerate gaps that we will now establish.

Glassy dynamics.—One of the hallmarks of a quantum glass is its ability to support nearly gapless excitations at all momenta due to the existence of exponentially many local minima in its energy landscape [72–75]. Meanwhile, the PM phase is obviously gapped and dispersionless with a short correlation length. In this section, we focus on the measurement of imaginary-time correlations $S(\mathbf{k}, \tau)$ at different momenta, deep inside the PM ($\delta/\Omega = 3.3$, $R_b = 1.0$) and glass ($\delta/\Omega = 3.3$, $R_b = 2.1$) phases, as marked by the red stars (d) and (b) in Fig. 1,




FIG. 4. (a) Dynamical correlations $S(\mathbf{k}, \tau)$ deep in the glass ($\delta/\Omega = 3.3$, $R_b = 2.1$) and PM ($\delta/\Omega = 3.3$, $R_b = 1$) phases. The correlations decay quickly with similar slopes in the PM phase, indicating a large gap and a flat band. Meanwhile, the correlations in the glass phase decay very slowly, which suggests that there are many nearly degenerate local minima in the energy landscape. (b) The Edwards-Anderson order parameter q_{EA} also decays much more slowly with the increasing number of Monte Carlo steps for the glass than for the QSL. The data in the glass phase is computed for the model (1) at $\delta/\Omega = 3.55$, $R_b = 2.05$, while the QSL’s data is obtained from the triangular-lattice quantum loop model at $V/t = 0.9$. All the data are simulated at $\beta = 6$ on a $L = 6$ lattice.

respectively. Figure 4(a) highlights a striking distinction in $S(\mathbf{k}, \tau)$ between these two regions. The upper ones, which are measured inside the glass phase, decay slowly, indicating that their gaps are all very small and almost equal. On the other hand, the lower ones measuring the correlations in PM phase decay much more quickly with similar slopes. These two kinds of correlations are consistent with the respective features of the glass phase, which possesses nearly gapless excitations, as well as the PM phase, which hosts gapped flat bands.

Lastly, we also compare the behavior of the Edwards-Anderson order parameter q_{EA} , as a function of the number of Monte Carlo steps, inside the glass phase to that in an even QSL [39]. Since the representative parameter point chosen for the spin glass ($\delta/\Omega = 3.55$, $R_b = 2.05$) lies close to the boundary of the nematic phase (which, recall, is a VBS with two dimers per site), the low-energy effective model describing the QSL is taken to be the triangular-lattice quantum loop model close to the Rokhsar-Kivelson point [39, 40] ($V/t = 0.9$, where V is the interaction between parallel dimers and $-t$ is the dimer resonance energy). As shown in Fig. 4(b), in the QSL phase, q_{EA} is not only an order-of-magnitude smaller but also decays much faster than in the glass. The Monte Carlo *dynamics* of q_{EA} thus complement the *static* results of Fig. 3(b) and together, highlight the distinction between the QSL and glass phases.

Discussion.—In this work, we investigated a realistic Rydberg Hamiltonian on the kagome lattice with large-scale QMC simulations and uncover—besides two VBSs and a PM phase—an emergent glass phase in the phase diagram. The origin of glassiness is likely due to the kinetic constraints associated with the Rydberg blockade that forbid several hopping processes [41, 45, 76]. Such a glass phase constitutes a new addition to the growing list of correlated quantum many-body states that can be studied on current-generation Rydberg platforms. Through detailed QMC analyses, we explore the subtle behavior of this glass phase with its intricate degenerate energy landscapes, establish its unique static and dynamic fingerprints, and compare and contrast its properties to those of competing QSL candidates. However, we note that the observation of a glassy *ground state* does not necessarily rule out the dynamical preparation of QSL states, which may be obtained as a macroscopic superposition of dimer configurations during quasiadiabatic sweeps in experiments [9, 34, 35].

Experimentally, the spin glass phase can be detected by preparing a (deterministic) far-from-equilibrium initial state, quenching to the glassy region, measuring snapshots in the occupation basis, and repeating the protocol, but stopping at different points in the temporal evolution each time. This would allow one to observe the anomalously slow relaxation dynamics, as recently demonstrated in experiments studying a disordered XXZ model on a Rydberg quantum simulator [77]. It is also possible to use Bragg spectroscopy to measure the dynamics of the glass phase in cold-atom systems held in optical lattices [78]. Finally, even though the Edwards-Anderson order parameter can be challenging to measure in quantum simulators (since it requires knowledge of two-time correlation functions), one can probe an efficient proxy for q_{EA} by measuring the eigenstate spin-glass order parameter [79] constructed from two-site reduced density matrices, which can be accessed by a variety of methods including quantum state tomography [80].

Acknowledgments.—Z.Y. and Z.Y.M. acknowledge useful discussions with Adriano Angelone, Xue-Feng Zhang and Zheng Zhou. R.S. and S.S. thank their coauthors in earlier collaborations [9, 32] for helpful discussions. R.S. is supported by the Princeton Quantum Initiative Fellowship. S.S. is supported by the U.S. Department of Energy under Grant DE-SC0019030. Z.Y. and Z.Y.M. acknowledge support from the RGC of Hong Kong SAR of China (Grant Nos. 17301420, 17301721, AoE/P-701/20, 17309822, A.HKU703/22 and C7037-22G), the K. C. Wong Education Foundation (Grant No. GJTD-2020-01), and the Seed Funding “Quantum-Inspired explainable-AI” at the HKU-TCL Joint Research Centre for Artificial Intelligence. Y.C.W. acknowledges support from Zhejiang Provincial Natural Science Foundation of China (Grant Nos. LZ23A040003) and the Beihang Hangzhou Innovation Institute Yuhang. We

thank Beijing PARATERA Tech CO.,Ltd., the High-performance Computing Centre of Beihang Hangzhou Innovation Institute Yuhang, HPC2021 system under the Information Technology Services, and the Blackbody HPC system at the Department of Physics, University of Hong Kong for providing computational resources that have contributed to the research results in this paper.

* sachdev@g.harvard.edu

† zymeng@hku.hk

- [1] M. Endres, H. Bernien, A. Keesling, H. Levine, E. R. Anschuetz, A. Krajenbrink, C. Senko, V. Vuletić, M. Greiner, and M. D. Lukin, Atom-by-atom assembly of defect-free one-dimensional cold atom arrays, *Science* **354**, 1024 (2016).
- [2] H. Bernien, S. Schwartz, A. Keesling, H. Levine, A. Omran, H. Pichler, S. Choi, A. S. Zibrov, M. Endres, M. Greiner, V. Vuletić, and M. D. Lukin, Probing many-body dynamics on a 51-atom quantum simulator, *Nature* **551**, 579 (2017).
- [3] A. Browaeys and T. Lahaye, Many-body physics with individually controlled Rydberg atoms, *Nat. Phys.* **16**, 132 (2020).
- [4] M. Morgado and S. Whitlock, Quantum simulation and computing with Rydberg-interacting qubits, *AVS Quantum Sci.* **3**, 023501 (2021).
- [5] D. Bluvstein, H. Levine, G. Semeghini, T. T. Wang, S. Ebadi, M. Kalinowski, A. Keesling, N. Maskara, H. Pichler, M. Greiner, V. Vuletic, and M. D. Lukin, A quantum processor based on coherent transport of entangled atom arrays, *Nature* **604**, 451 (2022).
- [6] S. de Léséleuc, V. Lienhard, P. Scholl, D. Barredo, S. Weber, N. Lang, H. P. Büchler, T. Lahaye, and A. Browaeys, Observation of a symmetry-protected topological phase of interacting bosons with Rydberg atoms, *Science* **365**, 775 (2019).
- [7] S. Ebadi, T. T. Wang, H. Levine, A. Keesling, G. Semeghini, A. Omran, D. Bluvstein, R. Samajdar, H. Pichler, W. W. Ho, S. Choi, S. Sachdev, M. Greiner, V. Vuletić, and M. D. Lukin, Quantum phases of matter on a 256-atom programmable quantum simulator, *Nature* **595**, 227 (2021).
- [8] P. Scholl, M. Schuler, H. J. Williams, A. A. Eberharter, D. Barredo, K.-N. Schymik, V. Lienhard, L.-P. Henry, T. C. Lang, T. Lahaye, A. M. Läuchli, and A. Browaeys, Quantum simulation of 2D antiferromagnets with hundreds of Rydberg atoms, *Nature* **595**, 233 (2021).
- [9] G. Semeghini, H. Levine, A. Keesling, S. Ebadi, T. T. Wang, D. Bluvstein, R. Verresen, H. Pichler, M. Kalinowski, R. Samajdar, A. Omran, S. Sachdev, A. Vishwanath, M. Greiner, V. Vuletić, and M. D. Lukin, Probing topological spin liquids on a programmable quantum simulator, *Science* **374**, 1242 (2021).
- [10] C. J. Turner, A. A. Michailidis, D. A. Abanin, M. Serbyn, and Z. Papić, Weak ergodicity breaking from quantum many-body scars, *Nature Phys.* **14**, 745 (2018).
- [11] A. Keesling, A. Omran, H. Levine, H. Bernien, H. Pichler, S. Choi, R. Samajdar, S. Schwartz, P. Silvi, S. Sachdev, P. Zoller, M. Endres, M. Greiner, V. Vuletić,

- and M. D. Lukin, Quantum Kibble–Zurek mechanism and critical dynamics on a programmable Rydberg simulator, *Nature* **568**, 207 (2019).
- [12] D. Bluvstein, A. Omran, H. Levine, A. Keesling, G. Semeghini, S. Ebadi, T. T. Wang, A. A. Michailidis, N. Maskara, W. W. Ho, S. Choi, M. Serbyn, M. Greiner, V. Vuletić, and M. D. Lukin, Controlling quantum many-body dynamics in driven Rydberg atom arrays, *Science* **371**, 1355 (2021).
- [13] A. Celi, B. Vermersch, O. Viyuela, H. Pichler, M. D. Lukin, and P. Zoller, Emerging Two-Dimensional Gauge Theories in Rydberg Configurable Arrays, *Phys. Rev. X* **10**, 021057 (2020).
- [14] F. M. Surace, P. P. Mazza, G. Giudici, A. Lerose, A. Gambassi, and M. Dalmonte, Lattice Gauge Theories and String Dynamics in Rydberg Atom Quantum Simulators, *Phys. Rev. X* **10**, 021041 (2020).
- [15] D. González-Cuadra, T. V. Zache, J. Carrasco, B. Kraus, and P. Zoller, Hardware Efficient Quantum Simulation of Non-Abelian Gauge Theories with Qudits on Rydberg Platforms, *Phys. Rev. Lett.* **129**, 160501 (2022).
- [16] R. Samajdar, D. G. Joshi, Y. Teng, and S. Sachdev, Emergent \mathbb{Z}_2 Gauge Theories and Topological Excitations in Rydberg Atom Arrays, *Phys. Rev. Lett.* **130**, 043601 (2023).
- [17] H. Pichler, S.-T. Wang, L. Zhou, S. Choi, and M. D. Lukin, Quantum optimization for maximum independent set using Rydberg atom arrays, [arXiv:1808.10816 \[quant-ph\]](https://arxiv.org/abs/1808.10816) (2018).
- [18] S. Ebadi, A. Keesling, M. Cain, T. T. Wang, H. Levine, D. Bluvstein, G. Semeghini, A. Omran, J.-G. Liu, R. Samajdar, X.-Z. Luo, B. Nash, X. Gao, B. Barak, E. Farhi, S. Sachdev, N. Gemelke, L. Zhou, S. Choi, H. Pichler, S.-T. Wang, M. Greiner, V. Vuletic, and M. D. Lukin, Quantum optimization of maximum independent set using rydberg atom arrays, *Science* **376**, 1209 (2022).
- [19] Z. Yan, Z. Zhou, Y.-C. Wang, Z. Y. Meng, and X.-F. Zhang, Preparing state within target topological sector of lattice gauge theory model on quantum simulator, [arXiv:2105.07134 \[quant-ph\]](https://arxiv.org/abs/2105.07134) (2021).
- [20] R. Samajdar, S. Choi, H. Pichler, M. D. Lukin, and S. Sachdev, Numerical study of the chiral \mathbb{Z}_3 quantum phase transition in one spatial dimension, *Phys. Rev. A* **98**, 023614 (2018).
- [21] S. Whitsitt, R. Samajdar, and S. Sachdev, Quantum field theory for the chiral clock transition in one spatial dimension, *Phys. Rev. B* **98**, 205118 (2018).
- [22] N. Chepiga and F. Mila, Floating Phase versus Chiral Transition in a 1D Hard-Boson Model, *Phys. Rev. Lett.* **122**, 017205 (2019).
- [23] N. Chepiga and F. Mila, Kibble-Zurek exponent and chiral transition of the period-4 phase of Rydberg chains, *Nat. Commun.* **12**, 1 (2021).
- [24] R. Samajdar, W. W. Ho, H. Pichler, M. D. Lukin, and S. Sachdev, Complex Density Wave Orders and Quantum Phase Transitions in a Model of Square-Lattice Rydberg Atom Arrays, *Phys. Rev. Lett.* **124**, 103601 (2020).
- [25] T. Felser, S. Notarnicola, and S. Montangero, Efficient Tensor Network Ansatz for High-Dimensional Quantum Many-Body Problems, *Phys. Rev. Lett.* **126**, 170603 (2021).
- [26] M. Kalinowski, R. Samajdar, R. G. Melko, M. D. Lukin, S. Sachdev, and S. Choi, Bulk and boundary quantum phase transitions in a square Rydberg atom array, *Phys. Rev. B* **105**, 174417 (2022).
- [27] C. Miles, R. Samajdar, S. Ebadi, T. T. Wang, H. Pichler, S. Sachdev, M. D. Lukin, M. Greiner, K. Q. Weinberger, and E.-A. Kim, Machine learning discovery of new phases in programmable quantum simulator snapshots, *Phys. Rev. Res.* **5**, 013026 (2023).
- [28] M. J. O’Rourke and G. K.-L. Chan, Entanglement in the quantum phases of an unfrustrated Rydberg atom array, [arXiv:2201.03189 \[cond-mat.str-el\]](https://arxiv.org/abs/2201.03189) (2022).
- [29] C.-X. Li, S. Yang, and J.-B. Xu, Quantum phases of Rydberg atoms on a frustrated triangular-lattice array, *Opt. Lett.* **47**, 1093 (2022).
- [30] S. Yang and J.-B. Xu, Density-wave-ordered phases of rydberg atoms on a honeycomb lattice, *Phys. Rev. E* **106**, 034121 (2022).
- [31] M. Kornjača, R. Samajdar, T. Macrì, N. Gemelke, S.-T. Wang, and F. Liu, Trimer quantum spin liquid in a honeycomb array of Rydberg atoms, [arXiv:2211.00653 \[cond-mat.quant-gas\]](https://arxiv.org/abs/2211.00653) (2022).
- [32] R. Samajdar, W. W. Ho, H. Pichler, M. D. Lukin, and S. Sachdev, Quantum phases of Rydberg atoms on a kagome lattice, *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2015785118 (2021), 2011.12295.
- [33] R. Verresen, M. D. Lukin, and A. Vishwanath, Prediction of Toric Code Topological Order from Rydberg Blockade, *Phys. Rev. X* **11**, 031005 (2021).
- [34] G. Giudici, M. D. Lukin, and H. Pichler, Dynamical Preparation of Quantum Spin Liquids in Rydberg Atom Arrays, *Phys. Rev. Lett.* **129**, 090401 (2022).
- [35] Y. Cheng, C. Li, and H. Zhai, Variational Approach to Quantum Spin Liquid in a Rydberg Atom Simulator, [arXiv:2112.13688 \[cond-mat.quant-gas\]](https://arxiv.org/abs/2112.13688) (2021).
- [36] R. Moessner and K. S. Raman, Quantum dimer models, in *Introduction to Frustrated Magnetism* (Springer, 2011) pp. 437–479.
- [37] R. Moessner and S. L. Sondhi, Ising models of quantum frustration, *Phys. Rev. B* **63**, 224401 (2001).
- [38] K. Roychowdhury, S. Bhattacharjee, and F. Pollmann, \mathbb{Z}_2 topological liquid of hard-core bosons on a kagome lattice at 1/3 filling, *Phys. Rev. B* **92**, 075141 (2015).
- [39] Z. Yan, X. Ran, Y.-C. Wang, R. Samajdar, J. Rong, S. Sachdev, Y. Qi, and Z. Y. Meng, Fully packed quantum loop model on the triangular lattice: Hidden vison plaquette phase and cubic phase transitions, [arXiv:2205.04472 \[cond-mat.str-el\]](https://arxiv.org/abs/2205.04472) (2022).
- [40] Z. Yan, R. Samajdar, Y.-C. Wang, S. Sachdev, and Z. Y. Meng, Triangular lattice quantum dimer model with variable dimer density, *Nat. Commun.* **13**, 5799 (2022).
- [41] J. Feldmeier, F. Pollmann, and M. Knap, Emergent Glassy Dynamics in a Quantum Dimer Model, *Phys. Rev. Lett.* **123**, 040601 (2019).
- [42] X. Plat, F. Alet, S. Capponi, and K. Totsuka, Magnetization plateaus of an easy-axis kagome antiferromagnet with extended interactions, *Phys. Rev. B* **92**, 174402 (2015).
- [43] R. Moessner and S. L. Sondhi, Resonating Valence Bond Phase in the Triangular Lattice Quantum Dimer Model, *Phys. Rev. Lett.* **86**, 1881 (2001).
- [44] A. Angelone, F. Mezzacapo, and G. Pupillo, Superglass phase of interaction-blockaded gases on a triangular lattice, *Phys. Rev. Lett.* **116**, 135303 (2016).
- [45] I. Lesanovsky and J. P. Garrahan, Kinetic constraints, hierarchical relaxation, and onset of glassiness in strongly interacting and dissipative Rydberg gases, *Phys. Rev.*

- Lett. **111**, 215305 (2013).
- [46] C. Pérez-Espigares, I. Lesanovsky, J. P. Garrahan, and R. Gutiérrez, Glassy dynamics due to a trajectory phase transition in dissipative Rydberg gases, *Phys. Rev. A* **98**, 021804 (2018).
- [47] A. W. Sandvik, Stochastic series expansion method for quantum Ising models with arbitrary interactions, *Phys. Rev. E* **68**, 056701 (2003).
- [48] Z. Yan, Y. Wu, C. Liu, O. F. Syljuåsen, J. Lou, and Y. Chen, Sweeping cluster algorithm for quantum spin systems with strong geometric restrictions, *Phys. Rev. B* **99**, 165135 (2019).
- [49] Z. Yan, Global scheme of sweeping cluster algorithm to sample among topological sectors, *Phys. Rev. B* **105**, 184432 (2022).
- [50] Z. Yan, Y.-C. Wang, N. Ma, Y. Qi, and Z. Y. Meng, Topological phase transition and single/multi anyon dynamics of Z_2 spin liquid, *npj Quantum Mater.* **6**, 39 (2021).
- [51] Z. Yan, Z. Zhou, O. F. Syljuåsen, J. Zhang, T. Yuan, J. Lou, and Y. Chen, Widely existing mixed phase structure of the quantum dimer model on a square lattice, *Phys. Rev. B* **103**, 094421 (2021).
- [52] Y. Da Liao, H. Li, Z. Yan, H.-T. Wei, W. Li, Y. Qi, and Z. Y. Meng, Phase diagram of the quantum Ising model on a triangular lattice under external field, *Phys. Rev. B* **103**, 104416 (2021).
- [53] E. Merali, I. J. De Vlugt, and R. G. Melko, Stochastic Series Expansion Quantum Monte Carlo for Rydberg Arrays, [arXiv:2107.00766 \[cond-mat.str-el\]](https://arxiv.org/abs/2107.00766) (2021).
- [54] K. Hukushima and K. Nemoto, Exchange Monte Carlo method and application to spin glass simulations, *J. Phys. Soc. Japan* **65**, 1604 (1996).
- [55] J. Brooke, D. Bitko, F. T. Rosenbaum, and G. Aeppli, Quantum annealing of a disordered magnet, *Science* **284**, 779 (1999).
- [56] J. Brooke, T. Rosenbaum, and G. Aeppli, Tunable quantum tunnelling of magnetic domain walls, *Nature* **413**, 610 (2001).
- [57] E. Farhi, J. Goldstone, S. Gutmann, and M. Sipser, Quantum computation by adiabatic evolution (2000), [arXiv:quant-ph/0001106](https://arxiv.org/abs/quant-ph/0001106).
- [58] E. Farhi, J. Goldstone, S. Gutmann, J. Lapan, A. Lundgren, and D. Preda, A quantum adiabatic evolution algorithm applied to random instances of an NP-complete problem, *Science* **292**, 472 (2001).
- [59] R. G. Melko, Simulations of quantum XXZ models on two-dimensional frustrated lattices, *J. Phys.: Condens. Matter* **19**, 145203 (2007).
- [60] A. Das and B. K. Chakrabarti, Colloquium: Quantum annealing and analog quantum computation, *Rev. Mod. Phys.* **80**, 1061 (2008).
- [61] I. S. Aranson, N. B. Kopnin, and V. M. Vinokur, Dynamics of vortex nucleation by rapid thermal quench, *Phys. Rev. B* **63**, 184501 (2001).
- [62] A. Mitra, Quantum quench dynamics, *Annu. Rev. Condens. Matter Phys.* **9**, 245 (2018).
- [63] S. F. Edwards and P. W. Anderson, Theory of spin glasses, *J. Phys. F* **5**, 965 (1975).
- [64] P. M. Richards, Spin-glass order parameter of the random-field Ising model, *Phys. Rev. B* **30**, 2955 (1984).
- [65] A. Georges, O. Parcollet, and S. Sachdev, Quantum fluctuations of a nearly critical Heisenberg spin glass, *Phys. Rev. B* **63**, 134406 (2001).
- [66] K. Binder and A. P. Young, Spin glasses: Experimental facts, theoretical concepts, and open questions, *Rev. Mod. Phys.* **58**, 801 (1986).
- [67] The implementation of the different simulating schemes (annealing, quenching and parallel tempering), a brief description of the Edwards-Anderson order parameter, the phase transition between the nematic and paramagnetic phases, and the snapshots in the glassy region are presented in the Supplemental Material, which includes Refs. 81–83.
- [68] S. Sachdev, *Quantum Phase Transitions* (Cambridge University Press, New York, 2011).
- [69] W. Janke and R. Villanova, Three-dimensional 3-state Potts model revisited with new techniques, *Nucl. Phys. B* **489**, 679 (1997).
- [70] A. D. King, J. Raymond, T. Lanting, R. Harris, A. Zucca, F. Altomare, A. J. Berkley, K. Boothby, S. Ejtemaee, C. Enderud, E. Hoskinson, S. Huang, E. Ladizinsky, A. J. R. MacDonald, G. Marsden, R. Molavi, T. Oh, G. Poulin-Lamarre, M. Reis, C. Rich, Y. Sato, N. Tsai, M. Volkmann, J. D. Whittaker, J. Yao, A. W. Sandvik, and M. H. Amin, Quantum critical dynamics in a 5000-qubit programmable spin glass, [arXiv:2207.13800 \[quant-ph\]](https://arxiv.org/abs/2207.13800) (2022).
- [71] G. Carleo, M. Tarzia, and F. Zamponi, Bose-Einstein Condensation in Quantum Glasses, *Phys. Rev. Lett.* **103**, 215302 (2009).
- [72] A. Keren, P. Mendels, I. A. Campbell, and J. Lord, Probing the Spin-Spin Dynamical Autocorrelation Function in a Spin Glass above T_g via Muon Spin Relaxation, *Phys. Rev. Lett.* **77**, 1386 (1996).
- [73] V. K. Malinovsky, V. N. Novikov, and A. P. Sokolov, Log-normal spectrum of low-energy vibrational excitations in glasses, *Phys. Lett. A* **153**, 63 (1991).
- [74] M. J. Rozenberg and D. R. Grempel, Dynamics of the Infinite-Range Ising Spin-Glass Model in a Transverse Field, *Phys. Rev. Lett.* **81**, 2550 (1998).
- [75] Z. Ma, J. Wang, Z.-Y. Dong, J. Zhang, S. Li, S.-H. Zheng, Y. Yu, W. Wang, L. Che, K. Ran, S. Bao, Z. Cai, P. Čermák, A. Schneidewind, S. Yano, J. S. Gardner, X. Lu, S.-L. Yu, J.-M. Liu, S. Li, J.-X. Li, and J. Wen, Spin-Glass Ground State in a Triangular-Lattice Compound YbZnGaO_4 , *Phys. Rev. Lett.* **120**, 087201 (2018).
- [76] F. Ritort and P. Sollich, Glassy dynamics of kinetically constrained models, *Advances in physics* **52**, 219 (2003).
- [77] A. Signoles, T. Franz, R. Ferracini Alves, M. Gärttner, S. Whitlock, G. Zürn, and M. Weidemüller, Glassy Dynamics in a Disordered Heisenberg Quantum Spin System, *Phys. Rev. X* **11**, 011011 (2021).
- [78] R. Senaratne, D. Cavazos-Cavazos, S. Wang, F. He, Y.-T. Chang, A. Kafle, H. Pu, X.-W. Guan, and R. G. Hulet, Spin-charge separation in a one-dimensional Fermi gas with tunable interactions, *Science* **376**, 1305 (2022).
- [79] Y. Javanmard, S. Bera, and M. Heyl, Accessing eigenstate spin-glass order from reduced density matrices, *Phys. Rev. B* **99**, 144201 (2019).
- [80] H.-Y. Huang, R. Kueng, and J. Preskill, Predicting many properties of a quantum system from very few measurements, *Nature Phys.* **16**, 1050 (2020).
- [81] A. W. Sandvik and J. Kurkijärvi, Quantum Monte Carlo simulation method for spin systems, *Phys. Rev. B* **43**, 5950 (1991).
- [82] A. W. Sandvik, Stochastic series expansion method with operator-loop update, *Phys. Rev. B* **59**, R14157 (1999).

- [83] Z. Zhou, X.-F. Zhang, F. Pollmann, and Y. You, Fractal quantum phase transitions: Critical phenomena beyond renormalization, [arXiv:2105.05851](https://arxiv.org/abs/2105.05851) [cond-mat.str-el] (2021).