



This is the accepted manuscript made available via CHORUS. The article has been published as:

## Robust Fermi-Liquid Instabilities in Sign Problem-Free Models

Ori Grossman and Erez Berg

Phys. Rev. Lett. **131**, 056501 — Published 1 August 2023

DOI: [10.1103/PhysRevLett.131.056501](https://doi.org/10.1103/PhysRevLett.131.056501)

# Robust Fermi liquid instabilities in sign problem-free models

Ori Grossman\* and Erez Berg

*Department of Condensed Matter Physics, Weizmann Institute of Science, Rehovot, 76100, Israel*

Determinant Quantum Monte Carlo (DQMC) is a powerful numerical technique to study many-body fermionic systems. In recent years, several classes of sign-free (SF) models have been discovered, where the notorious sign problem can be circumvented. However, it is not clear what are the inherent physical characteristics and limitations of SF models. In particular, which zero-temperature quantum phases of matter are accessible within such models, and which are fundamentally inaccessible? Here, we show that a model belonging to any of the known SF classes within DQMC cannot have a stable Fermi liquid ground state in spatial dimension  $d \geq 2$ , unless the anti-unitary symmetry that prevents the sign problem is spontaneously broken (for which there are currently no known examples in SF models). For SF models belonging to one of the symmetry classes (where the absence of the sign problem follows from a combination of non-unitary symmetries of the fermionic action), any putative Fermi liquid fixed point generically includes an attractive Cooper-like interaction that destabilizes it. In the recently discovered lower-symmetry classes of SF models, the Fermi surface is generically unstable even at the level of the quadratic action. Our results suggest a fundamental link between Fermi liquids and the fermion sign problem. Interestingly, our results do not rule out a non-Fermi liquid ground state with a Fermi surface in a sign-free model.

*Introduction.*— The importance of reliable and practical simulations of strongly correlated fermionic systems cannot be overstated. In recent decades, substantial progress was made thanks to the development of the Determinant Quantum Monte Carlo (DQMC) technique [1–6]. However, this technique is often hindered by the sign problem [7, 8], associated with negative or complex amplitudes in the quantum partition sum. The sign problem results in inefficient simulations, generally scaling exponentially with system size and inverse temperature [6, 9] (although interesting exceptions exist [10, 11]).

Interestingly, certain classes of fermionic models do not suffer from the sign problem [12–18], and hence can be solved at polynomial cost. In these models, the absence of the sign problem is guaranteed by a combination of symmetries of the fermionic action. Sign free (SF) models were used to simulate a plethora of interesting phenomena, including fermionic quantum criticality [19–26] and unconventional superconductivity [27–30]. These developments raise the question of the physical properties and intrinsic limitations of sign problem free models [11, 31–34]. In particular, which quantum phases of matter can be accessed within SF models, and which are fundamentally inaccessible?

In this work, we show that all the currently known classes of sign problem free models within DQMC cannot support a stable Fermi liquid (FL) ground state. This is shown by demonstrating that any putative Fermi surface in a SF model is generically unstable in the presence of interactions at zero temperature. Depending on the model, the instability may be towards a fully gapped superconducting or density wave state, or towards a Dirac semi-metal [35, 36].

It is important to note that we cannot rule out a fine-tuned SF model with a Fermi surface (e.g., a model of

Table I. The known sign-free (SF) classes (see text for a detailed description), and the origin of the Fermi surface instability in each class. In the symmetry classes (Symm. SF), there is an interaction-induced instability of the Fermi surface that opens a gap through spontaneous symmetry breaking. In the lower-symmetry classes (lower-symm. SF), the Fermi surface is not protected (in general) by symmetry, and a gap generically opens even at the single-particle level.

	Kramers' class $\{T_1^-, T_2^-\} = 0$	Majorana class $\{T_1^-, T_2^+\} = 0$
Symm. SF	$U(1)$ breaking	$\mathbb{Z}_2$ breaking
Lower-symm. SF	Single-particle gap	Single-particle gap

non-interacting electrons). Our argument shows that in such cases, the Fermi surface is unstable, in the sense that a generic, arbitrarily small perturbation can destroy it [37]. In addition, a non-FL state with a Fermi surface (of the type that arises, e.g., at certain quantum critical points) is not ruled out.

Our main findings are summarized in Table I. In the symmetric SF classes [18] the SF property is guaranteed by the combination of two anti-unitary time-reversal symmetries (TRSs) of the fermionic action matrix. A detailed description of the different SF classes is given below. The product of the two TRSs defines a unitary symmetry, which is either a  $U(1)$  or a  $\mathbb{Z}_2$  symmetry, depending on the class. Interactions generically lead to spontaneous breaking of the unitary symmetry, gapping out the Fermi surface. There are many known examples of these phenomena (see Suppl. Material [38]); our work shows that these are general properties of SF models. There are also lower-symmetry classes where the SF is due to less strict conditions [15, 16]. We show that in these cases,

\* ori.grossman@weizmann.ac.il

generically, there is no FS even at the single-particle level, since there is no symmetry to protect it. If a symmetry that protects a FS is imposed, the FS is unstable in the presence of interactions, as in the symmetric classes.

*Sign-free DQMC.*— DQMC is based on introducing a bosonic field  $\phi$  via a Hubbard-Stratonovich (HS) transformation, deriving an effective action for  $\phi$  by integrating out the fermions, and averaging stochastically over  $\phi$ . The field  $\phi$  mediates the fermionic interaction [6]. Alternatively,  $\phi$  may represent a physical boson (such as a phonon). A typical action has the form of  $S = S_F + S_\phi + S_{\text{Int}}$  where  $S_F$  is the non-interacting fermionic action,  $S_\phi$  is the bosonic action, and  $S_{\text{Int}}$  is a Yukawa-like interaction. We assume that  $S_\phi \in \mathbb{R}$ . Upon integrating out the fermions, an effective bosonic action is obtained:  $S'_\phi = S_\phi + \ln \det M_\phi$ , where  $M_\phi$  is the ( $\phi$  dependent) quadratic fermionic action matrix,  $\bar{\psi} M_\phi \psi = S_{\text{Int}} + S_F$ . The fermionic problem has therefore been mapped to a classical statistical mechanical problem in  $d + 1$  dimensions for the field  $\phi(\mathbf{r}, \tau)$ .

However, it is not guaranteed that the statistical weights in the partition sum can be treated as probabilities, since they are not necessarily real and non-negative. This is known as the sign problem. Models that satisfy  $S'_\phi \in \mathbb{R}$  (or equivalently  $\det M_\phi \geq 0$ ) for any configuration  $\phi(\mathbf{r}, \tau)$  are known as SF models.

A set of sufficient conditions is known to guarantee the absence of the sign problem [18]. These conditions are most conveniently stated using a Majorana representation, writing the complex fermion field in terms of two real (Majorana) fields:  $\psi = \frac{1}{2}(\gamma_1 + i\gamma_2)$ . The fermionic bilinear action takes the form  $\gamma^T \tilde{M}_\phi \gamma$  where  $\gamma = (\gamma_{1,1}, \dots, \gamma_{1,N}, \dots, \gamma_{2,N})$  (for  $N$  Dirac fermions) and  $\tilde{M}_\phi$  is a  $2N \times 2N$  skew-symmetric matrix. In addition, a Majorana TRS,  $T$ , is defined as an anti-unitary operator that satisfies  $[\tilde{M}_\phi, T] = 0$  for any  $\phi(\mathbf{r}, \tau)$ . In this framework, one can distinguish between two fundamental SF classes:

- **Kramers' class:**  $\tilde{M}_\phi$  has two mutually anti-commuting TRSs, satisfying  $(T_1^-)^2 = (T_2^-)^2 = -1$ . Since  $(iT_1 T_2)^2 = 1$ , models of this class have a conserved  $U(1)$  charge,  $\hat{Q} = \gamma^T iT_1 T_2 \gamma$ . They can therefore be represented by Dirac fermions with a  $U(1)$  symmetry ( $\gamma^T \tilde{M}_\phi \gamma \rightarrow \bar{\psi} M_\phi \psi$ ). By Kramer's theorem, the eigenvalues of  $M_\phi$  come in complex conjugate pairs.
- **Majorana class:**  $\tilde{M}_\phi$  has two mutually anti-commuting TRSs, satisfying  $(T_1^-)^2 = -1, (T_2^+)^2 = 1$ . Since  $U = T_1^- T_2^+$  is a  $\mathbb{Z}_2$  unitary symmetry,  $\tilde{M}_\phi$  can be brought to the block diagonal form

$$\tilde{M}_\phi = \begin{bmatrix} B & 0 \\ 0 & B^* \end{bmatrix} \quad (1)$$

in the eigenbasis of  $U$ . Integrating out the fermions yields  $\text{Pf}(\tilde{M}_\phi) = \text{Pf}(B)\text{Pf}(B^*) \geq 0$  (where  $\text{Pf}(B)$  is the Pfaffian of the skew-symmetric matrix  $B$ ). Models of this type may not have a  $U(1)$  symmetry,

but can still have a FS of the zero energy Bogoliubov-like excitations, protected at the single-particle level by the  $\mathbb{Z}_2$  symmetry  $U$ .

The conditions above can be somewhat relaxed [15, 16]. In the so called lower-symmetry classes, we keep the requirement  $[\tilde{M}_\phi, T_2^\pm] = 0$ , but the second condition becomes  $iK [T_1^-, \tilde{M}_\phi] \leq 0$  (i.e., the left-hand side is a negative semi-definite matrix), where  $K$  is complex conjugation. As before  $\{T_1, T_2\} = 0$ . These requirements are sufficient to guarantee  $\det M_\phi \geq 0$  [16]. The limiting case where  $[\tilde{M}_\phi, T_1^-] = 0$  corresponds to the symmetry classes discussed above. In case of a strict inequality, we have two lower-symmetry SF classes: the Majorana class ( $T_2^+$ ) and the Kramers' class ( $T_2^-$ ).

*Fermi surface instability.*— We consider a general, translationally invariant model in  $d \geq 2$  spatial dimension, that belongs to one of the SF classes. The model includes Majorana fermions interacting with a bosonic field via a Yukawa coupling (note that any model of fermions with a quartic interaction can be recast in this form via a HS transformation). The lattice-scale, ultra-violet (UV) action is given by

$$S_{\text{UV}} = \int dk \bar{\gamma}_{\alpha,k} \frac{(i\omega \delta_{\alpha,\alpha'} - h_{\alpha,\alpha',\mathbf{k}})}{2} \gamma_{\alpha',k} + S_\phi + S_{\text{Int}}, \quad (2)$$

where  $\alpha$  denotes a general fermionic flavor such as spin or Majorana flavor ( $\gamma_{1,2}$ ), and summation over repeated indices is assumed. We have defined  $\bar{\gamma}_k \equiv \gamma_{-k}$ , and used the notation  $k = \{i\omega, \mathbf{k}\}$ , where  $\mathbf{k}$  is the spatial momentum.  $S_\phi$  is a bosonic action, and  $S_{\text{Int}}$  is given by

$$S_{\text{Int}} = \int dk dk' \frac{1}{2} \lambda_{\alpha,\alpha',k,k'}^\nu \phi_{k,k'}^\nu \bar{\gamma}_{\alpha,k} \gamma_{\alpha',k'}. \quad (3)$$

Here  $\nu$  is the index of the auxiliary field and  $\lambda$  is the coupling function (which depends on momenta, bosonic index, and fermionic flavor). Note that the bosonic field  $\phi^\nu$  may also be complex, but  $S_\phi \in \mathbb{R}$  as the model is SF. In addition, we assume that our action corresponds to a physical Hamiltonian. I.e., Eq.(2) corresponds to a hermitian Hamiltonian  $\hat{H}(\hat{\gamma}, \hat{\phi}, \hat{\pi})$ , where  $\hat{\phi}, \hat{\pi}$  are canonically conjugate operators.

We use proof by contradiction, assuming that the ground state is a FL, and showing that the putative FL phase is unstable. The proof proceeds in two steps: (1) Obtaining the low-energy FL effective action, (2) Showing that the SF requirement necessitates the existence of an instability of the Fermi surface.

**Step 1:** We divide the fermionic modes into slow modes residing within a thin shell of thickness  $2\Lambda$  around the FS (shaded region in Fig. 1), and fast modes residing outside of the shell. The fast modes are integrated out, obtaining the infra-red effective theory ( $S_{\text{IR}}$ ) close to the FS. Note that  $S_{\text{IR}}$  is still quadratic in the fermions at this stage, as we do not integrate over  $\phi$  at this stage [38].

The infra-red action is written as

$$S_{\text{IR}} = \int_{|\epsilon_{\mathbf{k},\alpha}| < \Lambda} dk \bar{\gamma}_{\alpha,k} \frac{(i\omega - h_{\alpha,\alpha',\mathbf{k}})}{2} \gamma_{\alpha',k} + S'_\phi + S'_{\text{Int}} \quad (4)$$




Figure 1. A schematic plot of the fast modes integration out for a time-reversal symmetric model. The shaded region around the Fermi surface (FS) represents the slow momenta while all the rest of the Brillouin zone (fast modes) is integrated out. In **(a)** it is shown for the simple case of circular FS, **(b)** is for a generic case in which  $\epsilon_{\mathbf{k}} = \epsilon_{-\mathbf{k}}$ .

with

$$S'_{\text{Int}} = \int dk dk' \bar{\gamma}_{\alpha,k} \Sigma_{\alpha,\alpha',k,k'}(\phi) \gamma_{\alpha',k'} . \quad (5)$$

Here,  $\Sigma(\phi)$  is the fermionic self-energy obtained from integrating out the fast modes, including all  $\lambda$  dependent terms (we suppress the indices of  $\phi^\nu$  for brevity).  $S'_\phi$  is the renormalized bosonic action. More details concerning the diagrammatic representation of  $\Sigma(\phi)$  and  $S'_\phi$  can be found in the SI [38]. Importantly,  $\epsilon_{\mathbf{k}}$  is the “true” (renormalized) dispersion, given by the eigenvalues of the matrix  $h_{\alpha,\alpha',\mathbf{k}} + \langle \Sigma_{\alpha,\alpha',k,k} \rangle_{S'_\phi}$  (with  $\omega = 0$  within the self-energy). Hence, the exact FS is given by  $\epsilon_{\mathbf{k}} = 0$ .

Next, we integrate out  $\phi$  in order to obtain a purely fermionic low-energy action. By our assumption, this action has a FL form. We assume that we are not exactly at a quantum critical point (QCP); at a QCP, singular interactions between the low-energy fermions arise, violating the FL assumption.

In the resulting fermionic action  $S_{\text{eff}}(\gamma, \bar{\gamma})$ , we are interested only in the quartic terms, as in a FL, higher order terms are irrelevant in the renormalization group (RG) sense [39, 40]. The quartic part of  $-S_{\text{eff}}(\gamma, \bar{\gamma})$  is written as

$$\Gamma_{k_1, k_2, k'_1, k'_2}^{\alpha, \beta, \alpha', \beta'} \bar{\gamma}_{\alpha, k_1} \bar{\gamma}_{\beta, k_2} \gamma_{\beta', k'_2} \gamma_{\alpha', k'_1} \delta(\{k\}), \quad (6)$$

where  $\delta(\{k\}) = \delta_{k_1+k_2-k'_1-k'_2}$ , and we can neglect the dependence of  $\Gamma$  on both frequency and the momentum perpendicular to the FS, since both dependencies are irrelevant [39, 40]. Using the cumulant expansion, we can express  $\Gamma$  as

$$\Gamma_{\mathbf{k}_1, \mathbf{k}_2, \mathbf{k}'_1, \mathbf{k}'_2}^{\alpha, \beta, \alpha', \beta'} = \left\langle \Sigma_{\alpha, \alpha', \mathbf{k}_1, \mathbf{k}'_1} \Sigma_{\beta, \beta', \mathbf{k}_2, \mathbf{k}'_2} \right\rangle_{S'_\phi} - \left\langle \Sigma_{\alpha, \alpha', \mathbf{k}_1, \mathbf{k}'_1} \right\rangle_{S'_\phi} \left\langle \Sigma_{\beta, \beta', \mathbf{k}_2, \mathbf{k}'_2} \right\rangle_{S'_\phi} . \quad (7)$$

Note that, due to the SF property of the model,  $S'_\phi \in \mathbb{R}$  (see [38]).

**Step 2:** We now perform a stability analysis of the putative FL fixed point. The analysis follows closely the

standard RG procedure for interacting fermions [39]. In the limit  $\Lambda \rightarrow 0$ , forward scattering processes are exactly marginal. Since the model is SF, it has at least one TRS, and hence  $\epsilon_{\mathbf{k}} = \epsilon_{-\mathbf{k}}$ . Then, FS instabilities may arise in the Cooper channel, corresponding to  $k_1 = -k_2, k'_1 = -k'_2$ . To obtain a compact form of the RG equations, it is convenient to define

$$\tilde{\Gamma}_{\mathbf{k}, -\mathbf{k}, \mathbf{k}', -\mathbf{k}'}^{\alpha, \beta, \alpha', \beta'} \equiv \frac{1}{v_{\mathbf{k}} v_{\mathbf{k}'}} \Gamma_{\mathbf{k}, -\mathbf{k}, \mathbf{k}', -\mathbf{k}'}^{\alpha, \beta, \alpha', \beta'} \quad (8)$$

where  $v_{\mathbf{k}} = |\nabla_{\mathbf{k}} \epsilon_{\mathbf{k}}|$ . We treat  $\tilde{\Gamma}_{\mathbf{k}, -\mathbf{k}, \mathbf{k}', -\mathbf{k}'}^{\alpha, \beta, \alpha', \beta'}$  as a matrix  $\tilde{\Gamma}$ , where the first (second) index corresponds to the set  $\{\alpha, \beta, k\}$  ( $\{\alpha', \beta', k'\}$ ), respectively. The hermiticity of the effective FL Hamiltonian implies that the matrix  $\tilde{\Gamma}$  is hermitian.

The one-loop RG equations for  $\tilde{\Gamma}$  take the simple form

$$\frac{d\tilde{\Gamma}}{dl} = \frac{1}{4\pi} \tilde{\Gamma}^2, \quad (9)$$

where  $dl = -d\Lambda/\Lambda$  is the infinitesimal scaling factor. Diagonalizing  $\tilde{\Gamma}$ , we obtain a set of differential equations for the eigenvalues (denoted by  $\lambda_i$ ):

$$\frac{d\lambda_i}{dl} = \frac{1}{4\pi} \lambda_i^2. \quad (10)$$

A positive  $\lambda_i$  (corresponding to attraction in a certain channel) grows under RG, destroying the FL. Thus, a stable FL phase requires that  $\lambda_i < 0$  for all  $i$ . Conversely, if there is at least one positive eigenvalue, there is no FS at  $T = 0$ . If  $\lambda_i \leq 0$  for all  $i$ , but there is at least one zero eigenvalue, there can be a FS, but it is unstable to the addition of an infinitesimal perturbation that makes one of the eigenvalues positive. (The latter case corresponds, e.g., to a non-interacting electron gas).

We now show that there exists a vector  $\tilde{w}$  for which  $\tilde{w}^T \tilde{\Gamma} \tilde{w} \geq 0$ , and hence  $\tilde{\Gamma}$  has at least one non-negative eigenvalue, and the FL phase cannot be stable. As mentioned above, the model has at least one TRS. We write the corresponding TRS operator as  $T = OK$ , where  $O$  is an orthogonal matrix ( $O$  is real since we are dealing with Majorana fermions), and  $K$  denotes complex conjugation. Under TRS,  $\gamma_{\alpha, k} \rightarrow O_{\alpha\beta} \gamma_{\beta, -k}$ . Setting  $w_{\mathbf{k}, \alpha, \beta} = O_{\alpha\beta} \delta_{\mathbf{k}, \mathbf{k}_0}$  with an arbitrarily chosen  $\mathbf{k}_0$ , and using the identity

$$\sum_{\alpha, \alpha'} O_{\alpha\beta} \Sigma_{\mathbf{k}_0, \mathbf{k}_0}^{\alpha, \alpha'} O_{\alpha'\beta'} = \left( \Sigma_{-\mathbf{k}_0, -\mathbf{k}_0}^{\beta, \beta'} \right)^*, \quad (11)$$

which follows from the time-reversal invariance of the fermionic action under  $T$ , we obtain

$$\tilde{w}^T \tilde{\Gamma} \tilde{w} = \frac{1}{v_{\mathbf{k}_0}} \sum_{\beta, \beta'} \left[ \left| \Sigma_{\beta, \beta', -\mathbf{k}_0, -\mathbf{k}_0} \right|^2 \right]_{S'_\phi} - \left\langle \Sigma_{\beta, \beta', -\mathbf{k}_0, -\mathbf{k}_0} \right\rangle_{S'_\phi} \left\langle \Sigma_{\beta, \beta', -\mathbf{k}_0, -\mathbf{k}_0}^* \right\rangle_{S'_\phi} \right] \geq 0. \quad (12)$$

Thus, the putative FL phase is either intrinsically unstable, or can be destabilized by adding an infinitesimal attractive interaction. This is our main result.




Figure 2. Schematic description of Cooper channel instability in the sign-free classes. We demonstrate it for the simple case of 2 flavors. (a) The time-reversal mapping between the two bands. (b) The obtained nesting between the Fermi surfaces.

It is worth examining the key elements required for our proof. In essence, the TRS (which all presently known SF classes require) ensures that the FS has a Cooper-like instability, with states at opposite momenta being degenerate (see Fig. 2). In addition, the SF property guarantees that the effective interaction is attractive in some channel (although it may be repulsive in other channels). Therefore, the FS cannot be stable. It is natural to expect a gapped, spontaneously broken ground state as a result.

We further stress that we cannot rule out the possibility of a non-FL metal with a FS. We also do not rule out a manifold of co-dimension  $d - 2$  or lower of zero energy excitations. SF models can have a stable Dirac semi-metal ground state in  $d = 2$  [36, 41–43], or a nodal line semimetal in  $d \geq 3$ .

Importantly, within our proof, we have implicitly assumed that TRS is not broken *spontaneously* [44]. If such spontaneous symmetry breaking had occurred, the FS could be stable. Ref. [31] conjectured, and showed explicitly for some cases, that an anti-unitary symmetry cannot be spontaneously broken in a bosonic SF model. We do not know of cases where TRS is spontaneously broken in a SF model containing fermions; whether such TRS breaking is fundamentally possible remains to be seen.

*The gapped phase.*— While for all SF classes, the ground state is not a stable FL, its exact nature is class and model-dependent. However, in both symmetry classes, it is natural to expect that as a result of the FS instability, the unitary  $T_1 T_2$  symmetry is spontaneously broken, and the FS is gapped out. In Kramers’ class, this typically results in a superconducting ground state, as  $\gamma^T i T_1 T_2 \gamma$  is the generator of a  $U(1)$  symmetry. The angular momentum of the order parameter is model specific, and both  $s$ -wave [45–51] and nodal or nodeless  $d$ -wave [27, 28, 30] superconductivity were found within SF DQMC. In the Majorana class,  $T_1 T_2$  is a unitary  $\mathbb{Z}_2$  symmetry. Its spontaneous breaking results in a 2-fold degenerate ground state. In certain physical realizations, the symmetry broken phase may correspond to a charge ordered state [13, 15, 47, 50, 52, 53].

It is important to note that the spontaneous breaking of  $T_1 T_2$  does not always occur; if the fermions form a band insulator or a Dirac semi-metal, the symmetry unbroken phase may extend down to  $T = 0$ . We provide detailed

examples of realizations in different symmetry SF classes and their symmetry broken phases in [38].

In the lower-symmetry SF classes, as we only have a single TRS and not a unitary symmetry, there is generically no FS already at the level of the quadratic (non-interacting) part of the action. The single-particle Hamiltonian is analogous to a Bogoliubov-de Gennes (BdG) Hamiltonian with TRS, where real off-diagonal pairing terms are allowed. Consequently, the co-dimension of the zero energy modes is at most  $d - 2$  (see [38]). If we add a symmetry (beyond those required by the SF property) that protects the FS at the quadratic level (e.g., a  $U(1)$  or  $\mathbb{Z}_2$  symmetry), the FS is still generically unstable in the presence of interactions, just as in the symmetry classes. This is because our proof requires only a single TRS, which is present in the lower-symmetry classes. We summarize our conclusions for all currently known SF classes in Table I.

*Comment regarding the Kohn-Luttinger mechanism.*— We have shown that SF models cannot have a stable FL ground state, due to TRS and the presence of attraction in the Cooper channel. In this context, it is important to address the question whether, conversely, non-SF models with TRS can support a stable Fermi surface at  $T = 0$ . It is well known that under a wide range of circumstances, bare repulsive interactions can lead to superconductivity at high angular momentum channel. This is the Kohn-Luttinger (KL) mechanism [54].

In general, however, the KL mechanism relies on the bare repulsion being short-ranged, i.e., exponentially decaying as a function of distance (see [38] for details). The Fermi surface can be stabilized at  $T = 0$  by adding a small power-law repulsive interaction. In contrast, our proof shows that in SF models, the Fermi surface is generically unstable, even in the presence of arbitrary long-ranged interactions.

*Concluding remarks.*— Our arguments apply to all SF classes that are currently known within DQMC. However, these observations naturally lead to the stronger conjecture that a stable FL phase cannot be realized in any SF model. As new classes of SF models are found, this conjecture will be put to the test. For example, there are models [55] that do not have a known SF DQMC formulation, but the sign problem can be solved in a continuous time QMC [14, 56–58] (or the “fermion bag” approach [59, 60]). None of these models have a FL ground state, as far as we know; however, our proof does not formally encompass these cases. In addition, we note that it is possible to get stable metallic phases in “mixed dimensionality” systems [61, 62], where the interaction terms are sub-extensive. These lie beyond the scope of our results since they are not fully translationally invariant.

Looking ahead, it should be straightforward to extend our results to include quenched disorder. In this case, we expect SF models to obey a version of Anderson’s theorem [63], i.e., disorder that preserves the SF property does not suppress the superconducting instability at the mean-field level.



*Acknowledgments*– We are grateful to A. Chubukov, R. Shankar, S. Gazit, O. Golan, S. Kivelson, D. Mross, Z. Ringel, J. S. Hofmann and S. Trebst for illuminating discussions. This work was supported by the European Research Council (ERC) under the European Union’s

Horizon 2020 research and innovation programme (grant agreement No 817799), the Israel-USA Binational Science Foundation (BSF), and the ISF Quantum Science and Technology grant no. 2074/19.

- 
- [1] D. J. Scalapino and R. L. Sugar, Method for Performing Monte Carlo Calculations for Systems with Fermions, *Phys. Rev. Lett.* **46**, 519 (1981).
- [2] R. Blankenbecler, D. J. Scalapino, and R. L. Sugar, Monte Carlo calculations of coupled boson-fermion systems. I, *Phys. Rev. D* **24**, 2278 (1981).
- [3] J. E. Hirsch, D. J. Scalapino, R. L. Sugar, and R. Blankenbecler, Efficient Monte Carlo Procedure for Systems with Fermions, *Phys. Rev. Lett.* **47**, 1628 (1981).
- [4] J. E. Hirsch, Discrete Hubbard-Stratonovich transformation for fermion lattice models, *Phys. Rev. B* **28**, 4059 (1983).
- [5] S. R. White, D. J. Scalapino, R. L. Sugar, N. E. Bickers, and R. T. Scalettar, Attractive and repulsive pairing interaction vertices for the two-dimensional Hubbard model, *Physical Review B* **39**, 839 (1989).
- [6] F. F. Assaad and H. G. Evertz, World-line and Determinantal Quantum Monte Carlo Methods for Spins, Phonons and Electrons, *Lecture Notes in Physics* **739**, 277 (2008).
- [7] E. Y. Loh, J. E. Gubernatis, R. T. Scalettar, S. R. White, D. J. Scalapino, and R. L. Sugar, Sign problem in the numerical simulation of many-electron systems, *Physical Review B* **41**, 9301 (1990).
- [8] G. Pan and Z. Y. Meng, Sign problem in quantum monte carlo simulation, arXiv preprint arXiv:2204.08777 (2022).
- [9] M. Troyer and U.-J. Wiese, Computational complexity and fundamental limitations to fermionic quantum monte carlo simulations, *Phys. Rev. Lett.* **94**, 170201 (2005).
- [10] X. Zhang, G. Pan, X. Y. Xu, and Z. Y. Meng, Fermion sign bounds theory in quantum monte carlo simulation, *Phys. Rev. B* **106**, 035121 (2022).
- [11] Z.X Li, Z.Q Wan, and H.Yao, Asymptotic sign free in interacting fermion models, arXiv 10.48550/ARXIV.2211.00663 (2022).
- [12] C. Wu and S.-C. Zhang, Sufficient condition for absence of the sign problem in the fermionic quantum Monte Carlo algorithm, *Phys. Rev. B* **71**, 155115 (2005).
- [13] Z. X. Li, Y. F. Jiang, and H. Yao, Solving the fermion sign problem in quantum Monte Carlo simulations by Majorana representation, *Physical Review B* **91**, 241117 (2015).
- [14] L. Wang, Y.-H. Liu, M. Iazzi, M. Troyer, and G. Harcos, Split orthogonal group: A guiding principle for sign-problem-free fermionic simulations, *Phys. Rev. Lett.* **115**, 250601 (2015).
- [15] Z. C. Wei, C. Wu, Y. Li, S. Zhang, and T. Xiang, Majorana Positivity and the Fermion Sign Problem of Quantum Monte Carlo Simulations, *Physical Review Letters* **116**, 250601 (2016).
- [16] Z.-C. Wei, Semigroup Approach to the Sign Problem in Quantum Monte Carlo Simulations, arXiv:1712.09412 10.48550/arxiv.1712.09412 (2017).
- [17] Z. X. Li, Y. F. Jiang, and H. Yao, Majorana-Time-Reversal Symmetries: A Fundamental Principle for Sign-Problem-Free Quantum Monte Carlo Simulations, *Physical Review Letters* **117**, 267002 (2016).
- [18] Z. X. Li and H. Yao, Sign-Problem-Free Fermionic Quantum Monte Carlo: Developments and Applications, <https://doi.org/10.1146/annurev-conmatphys-033117-054307> **10**, 337 (2019).
- [19] F. F. Assaad and I. F. Herbut, Pinning the order: The nature of quantum criticality in the hubbard model on honeycomb lattice, *Phys. Rev. X* **3**, 031010 (2013).
- [20] Z.-X. Li, Y.-F. Jiang, and H. Yao, Fermion-sign-free majorana-quantum-monte-carlo studies of quantum critical phenomena of dirac fermions in two dimensions, *New Journal of Physics* **17**, 085003 (2015).
- [21] M. H. Gerlach, Y. Schattner, E. Berg, and S. Trebst, Quantum critical properties of a metallic spin-density-wave transition, *Physical Review B* **95**, 035124 (2017).
- [22] E. Berg, S. Lederer, Y. Schattner, and S. Trebst, Monte Carlo Studies of Quantum Critical Metals, <https://doi.org/10.1146/annurev-conmatphys-031218-013339> **10**, 63 (2019).
- [23] O. Grossman, J. S. Hofmann, T. Holder, and E. Berg, Specific heat of a quantum critical metal, *Phys. Rev. Lett.* **127**, 017601 (2021).
- [24] Y. Liu, W. Jiang, A. Klein, Y. Wang, K. Sun, A. V. Chubukov, and Z. Y. Meng, Dynamical exponent of a quantum critical itinerant ferromagnet: A monte carlo study, *Phys. Rev. B* **105**, L041111 (2022).
- [25] J. Schwab, L. Janssen, K. Sun, Z. Y. Meng, I. F. Herbut, M. Vojta, and F. F. Assaad, Nematic quantum criticality in dirac systems, *Phys. Rev. Lett.* **128**, 157203 (2022).
- [26] P. Lunts, M. S. Albergo, and M. Lindsey, Non-hertz-millis scaling of the antiferromagnetic quantum critical metal via scalable hybrid monte carlo, arXiv:2204.14241 10.48550/ARXIV.2204.14241 (2022).
- [27] F. F. Assaad, M. Imada, and D. J. Scalapino, Quantum transition between an antiferromagnetic mott insulator and  $d_{x^2-y^2}$  superconductor in two dimensions, *Phys. Rev. Lett.* **77**, 4592 (1996).
- [28] Y. Schattner, M. H. Gerlach, S. Trebst, and E. Berg, Competing orders in a nearly antiferromagnetic metal, *Phys. Rev. Lett.* **117**, 097002 (2016).
- [29] Z.-X. Li, F. Wang, H. Yao, and D.-H. Lee, Nature of the effective interaction in electron-doped cuprate superconductors: A sign-problem-free quantum monte carlo study, *Phys. Rev. B* **95**, 214505 (2017).
- [30] X. Y. Xu and T. Grover, Competing Nodal  $d$ -Wave Superconductivity and Antiferromagnetism, *Phys. Rev. Lett.* **126**, 217002 (2021).
- [31] Z. Ringel and D. L. Kovrizhin, Quantized gravitational responses, the sign problem, and quantum complexity, *Science Advances* **3**, e1701758 (2017).
- [32] O. Golan, A. Smith, and Z. Ringel, Intrinsic sign problem

- in fermionic and bosonic chiral topological matter, *Phys. Rev. Res.* **2**, 043032 (2020).
- [33] A. Smith, O. Golan, and Z. Ringel, Intrinsic sign problems in topological quantum field theories, *Phys. Rev. Res.* **2**, 033515 (2020).
- [34] R. Mondaini, S. Tarat, and R. T. Scalettar, Quantum critical points and the sign problem, *Science* **375**, 418 (2022).
- [35] S. Pujari, T. C. Lang, G. Murthy, and R. K. Kaul, Interaction-Induced Dirac Fermions from Quadratic Band Touching in Bilayer Graphene, *Physical Review Letters* **117**, 086404 (2016).
- [36] S. Gazit, M. Randeria, and A. Vishwanath, Emergent Dirac fermions and broken symmetries in confined and deconfined phases of Z2 gauge theories, *Nature Physics* **2017** 13:5 **13**, 484 (2017).
- [37] This is in contrast, e.g., to a superfluid phase, which is realizable within a SF model and is robust against arbitrary small perturbations (both ones that preserve the SF property and ones that do not).
- [38] See Supplementary Material, where we provide detailed examples and explain proof steps that require further clarification.
- [39] R. Shankar, Renormalization-group approach to interacting fermions, *Reviews of Modern Physics* **66**, 129 (1994).
- [40] J. Polchinski, Effective field theory and the fermi surface, arXiv [10.48550/arxiv.HEP-TH/9210046](https://arxiv.org/abs/10.48550/arxiv.HEP-TH/9210046) (1992).
- [41] Z. Meng, T. Lang, S. Wessel, F. Assaad, and A. Muramatsu, Quantum spin liquid emerging in two-dimensional correlated dirac fermions, *Nature* **464**, 847 (2010).
- [42] S. Sorella, Y. Otsuka, and S. Yunoki, Absence of a spin liquid phase in the hubbard model on the honeycomb lattice, *Scientific reports* **2**, 1 (2012).
- [43] H.-K. Tang, J. N. Leaw, J. N. B. Rodrigues, I. F. Herbut, P. Sengupta, F. F. Assaad, and S. Adam, The role of electron-electron interactions in two-dimensional dirac fermions, *Science* **361**, 570 (2018).
- [44] Importantly, here we refer to breaking of all anti-unitary symmetries. In the symmetric SF classes, it is possible that one of the protecting TRSs is spontaneously broken, while the other is preserved.
- [45] R. T. Scalettar, E. Y. Loh, J. E. Gubernatis, A. Moreo, S. R. White, D. J. Scalapino, R. L. Sugar, and E. Dagotto, Phase diagram of the two-dimensional negative-u hubbard model, *Phys. Rev. Lett.* **62**, 1407 (1989).
- [46] A. Moreo and D. J. Scalapino, Two-dimensional negative-u hubbard model, *Phys. Rev. Lett.* **66**, 946 (1991).
- [47] R. M. Noack, D. J. Scalapino, and R. T. Scalettar, Charge-density-wave and pairing susceptibilities in a two-dimensional electron-phonon model, *Phys. Rev. Lett.* **66**, 778 (1991).
- [48] Y. Schattner, S. Lederer, S. A. Kivelson, and E. Berg, Ising Nematic Quantum Critical Point in a Metal: A Monte Carlo Study, *Physical Review X* **6**, 031028 (2016).
- [49] Y. L. Loh, M. Randeria, N. Trivedi, C.-C. Chang, and R. Scalettar, Superconductor-Insulator Transition and Fermi-Bose Crossovers, *Phys. Rev. X* **6**, 021029 (2016).
- [50] I. Esterlis, B. Nosarzewski, E. W. Huang, B. Moritz, T. P. Devereaux, D. J. Scalapino, and S. A. Kivelson, Breakdown of the migdal-eliasberg theory: A determinant quantum monte carlo study, *Phys. Rev. B* **97**, 140501 (2018).
- [51] J. S. Hofmann, E. Berg, and D. Chowdhury, Superconductivity, pseudogap, and phase separation in topological flat bands, *Phys. Rev. B* **102**, 201112 (2020).
- [52] S. Johnston, E. A. Nowadnick, Y. F. Kung, B. Moritz, R. T. Scalettar, and T. P. Devereaux, Determinant quantum monte carlo study of the two-dimensional single-band hubbard-holstein model, *Phys. Rev. B* **87**, 235133 (2013).
- [53] Y.-F. Jiang, Z.-X. Li, S. A. Kivelson, and H. Yao, Charge-4e superconductors: A majorana quantum monte carlo study, *Phys. Rev. B* **95**, 241103 (2017).
- [54] W. Kohn and J. M. Luttinger, New Mechanism for Superconductivity, *Physical Review Letters* **15**, 524 (1965).
- [55] E. Huffman and S. Chandrasekharan, Solution to sign problems in models of interacting fermions and quantum spins, *Phys. Rev. E* **94**, 043311 (2016).
- [56] E. Gull, A. J. Millis, A. I. Lichtenstein, A. N. Rubtsov, M. Troyer, and P. Werner, Continuous-time monte carlo methods for quantum impurity models, *Rev. Mod. Phys.* **83**, 349 (2011).
- [57] L. Wang, P. Corboz, and M. Troyer, Fermionic quantum critical point of spinless fermions on a honeycomb lattice, *New Journal of Physics* **16**, 103008 (2014).
- [58] E. F. Huffman and S. Chandrasekharan, Solution to sign problems in half-filled spin-polarized electronic systems, *Phys. Rev. B* **89**, 111101 (2014).
- [59] S. Chandrasekharan, Fermion bag approach to lattice field theories, *Phys. Rev. D* **82**, 025007 (2010).
- [60] S. Chandrasekharan, Fermion bag approach to fermion sign problems: New opportunities in lattice fermion field theories, *The European Physical Journal A* **49**, 1 (2013).
- [61] Z. H. Liu, B. Frank, L. Janssen, M. Vojta, and F. F. Assaad, Magnetic quantum phase transition in a metallic kondo heterostructure, arXiv.2211.02071 [10.48550/ARXIV.2211.02071](https://arxiv.org/abs/10.48550/ARXIV.2211.02071) (2022).
- [62] B. Frank, Z. H. Liu, F. F. Assaad, M. Vojta, and L. Janssen, Marginal fermi liquid at magnetic quantum criticality from dimensional confinement, arXiv.2212.11283 [10.48550/ARXIV.2212.11283](https://arxiv.org/abs/10.48550/ARXIV.2212.11283) (2022).
- [63] P. Anderson, Theory of dirty superconductors, *Journal of Physics and Chemistry of Solids* **11**, 26 (1959).