



# CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

## Quasi-Floquet Prethermalization in a Disordered Dipolar Spin Ensemble in Diamond

Guanghai He, Bingtian Ye, Ruotian Gong, Zhongyuan Liu, Kater W. Murch, Norman Y. Yao, and Chong Zu

Phys. Rev. Lett. **131**, 130401 — Published 27 September 2023

DOI: [10.1103/PhysRevLett.131.130401](https://doi.org/10.1103/PhysRevLett.131.130401)

# Quasi-Floquet prethermalization in a disordered dipolar spin ensemble in diamond

Guanghui He,<sup>1,\*</sup> Bingtian Ye,<sup>2,3,\*†</sup> Ruotian Gong,<sup>1</sup> Zhongyuan Liu,<sup>1</sup>

Kater W. Murch,<sup>1,4</sup> Norman Y. Yao,<sup>2,3,5</sup> Chong Zu,<sup>1,4,‡</sup>

<sup>1</sup>Department of Physics, Washington University, St. Louis, MO 63130, USA

<sup>2</sup>Department of Physics, Harvard University, Cambridge, MA 02138, USA

<sup>3</sup>Department of Physics, University of California, Berkeley, CA 94720, USA

<sup>4</sup>Institute of Materials Science and Engineering, Washington University, St. Louis, MO 63130, USA

<sup>5</sup>Materials Science Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

\*These authors contributed equally to this work.

†To whom correspondence should be addressed; E-mail: bingtian\_ye@berkeley.edu

‡To whom correspondence should be addressed; E-mail: zu@wustl.edu

(Dated: August 21, 2023)

Floquet (periodic) driving has recently emerged as a powerful technique for engineering quantum systems and realizing non-equilibrium phases of matter [1–22]. A central challenge to stabilizing quantum phenomena in such systems is the need to prevent energy absorption from the driving field. Fortunately, when the frequency of the drive is significantly larger than the local energy scales of the many-body system, energy absorption is suppressed [23–32]. The existence of this so-called prethermal regime depends sensitively on the range of interactions and the presence of multiple driving frequencies [14, 33–36]. Here, we report the observation of Floquet prethermalization in a strongly interacting dipolar spin ensemble in diamond, where the angular dependence of the dipolar coupling helps to mitigate the long-ranged nature of the interaction. Moreover, we extend our experimental observation to quasi-Floquet drives with *multiple* incommensurate frequencies. In contrast to a *single*-frequency drive, we find that the existence of prethermalization is extremely sensitive to the smoothness of the applied field. Our results open the door to stabilizing and characterizing non-equilibrium phenomena in quasi-periodically driven systems.

Floquet theory describes the dynamics of a system whose Hamiltonian exhibits a *single* time-translation symmetry. Often used as a tool to control quantum systems, Floquet engineering (i.e. periodic driving) can help to prevent environment-induced decoherence and more recently, has enabled the study of novel quantum dynamical phenomena [8, 37–42]. In particular, many-body Floquet systems can host intrinsically non-equilibrium phases of matter, ranging from discrete time crystals [6, 7, 10–12, 18, 19, 43, 44] to Floquet topological states [15, 20, 22, 45–51]. Even richer non-equilibrium behaviors can arise in “quasi-Floquet” systems, where a *single* time-translation symmetry is replaced by *multiple* time-translation symmetries [21, 35, 52–55]. For instance, the spontaneous breaking of the latter can result in time quasi-crystalline order, which features a subharmonic response that is fundamentally distinct from conventional time crystals [21, 35].

A critical obstacle to stabilizing and observing such phenomena in driven quantum systems is Floquet heating: the inevitable absorption of energy from the driving field. One potential solution arises when the driving frequency,  $\omega$ , is significantly larger than the local energy scale,  $J$ , of the many-body system; in this case, Floquet prethermalization occurs and there exists an exponentially long-lived preheating regime described by an effective time-independent Hamiltonian,  $\mathcal{H}_{\text{eff}}$  [23–30]. The intuition underlying Floquet prethermalization is simple — in order to absorb a single photon from the drive, the

system must undergo  $\sim \omega/J$  off-resonant rearrangements of its local degrees of freedom. This higher-order process leads to an exponentially slow heating rate  $\sim \mathcal{O}(e^{-\omega/J})$ .

Despite this promise, there are two natural scenarios where prethermalization can break down: (i) systems with long-range, power-law interactions [14, 33] and (ii) quasi-Floquet systems where multi-photon processes can enable resonant energy absorption [35].

Here, we report the experimental observation of Floquet prethermalization in a long-range interacting quantum system under quasi-periodic driving. Our experimental platform consists of a dense ensemble of dipolar interacting nitrogen-vacancy (NV) centers in diamond (Fig. 1) [7, 8, 40, 57–59]. With single-frequency modulation, we observe that the heating time,  $\tau^*$ , is consistent with an exponential scaling with increasing driving frequency. In contrast, by driving quasi-periodically with two frequencies, we find that Floquet heating can be fitted to a stretched exponential profile with  $\tau^* \sim \mathcal{O}(e^{\omega^{\frac{1}{2}}})$  [35]. Interestingly, in the quasi-periodic case, the heating is extremely sensitive to the *smoothness* of the drive; indeed, when the system is driven via rectangular pulses (as opposed to sinusoidal pulses), we observe a significant enhancement in the heating rate (Fig. 3) [35]. We remark that the presence of slow heating is reminiscent of classic results from the NMR literature detailing the observation of long-lived dynamics in driven systems [38, 60–64]. However, the origin of the long-lived dynamics are either expected to be independent of the




FIG. 1. **Quasi-Floquet prethermalization in a dipolar spin system.** (a) Quasi-periodic driving of a strongly interacting NV ensemble in diamond with two incommensurate frequencies  $\omega$  and  $\varphi\omega$ , where  $\varphi$  is the golden ratio. Typical thermalization dynamics of the spin polarization  $\langle S^x(t) \rangle$  exhibit an initial fast decay followed by a late-time slow relaxation. Top: Level structure of NV center. Without an external field,  $|m_s = \pm 1\rangle$  sublevels are degenerate and sit  $D_{gs} = (2\pi) \times 2.87$  GHz above  $|m_s = 0\rangle$ . A magnetic field  $B \sim 350$  G along the NV axis splits  $|m_s = \pm 1\rangle$ , enabling the isolation of a two-level system. (b) Experimental sequence. The dynamical decoupling sequence eliminates the on-site random fields induced by the environmental bath spins. The sequence includes a series of fast  $\pi$ -pulses with alternating phases along  $\hat{x}$  and  $-\hat{x}$  axes to compensate the pulse errors. The inter-pulse spacing is fixed at  $\tau = 0.1 \mu\text{s}$ , much smaller than the interaction timescale between NV centers. A static microwave  $\Omega \sum_i S_i^x$  together with dipolar interaction serves as the static Hamiltonian  $\mathcal{H}_0$ , and a time-dependent microwave  $\Omega f(t) \sum_i S_i^x$  serves as the Floquet and quasi-Floquet drives. A final  $\frac{\pi}{2}$ -pulse along  $\mp \hat{y}$  axis rotates the spins back to  $\hat{z}$  for detection [56].

driving frequency or exhibits a lifetime that scales as a power-law of the driving frequency [56]. These scenarios are markedly distinct from the context of Floquet prethermalization, which exhibits an exponentially long lifetime in  $\omega$ . Despite such distinction, how to unequivocally demonstrate exponential over power-law scaling from real experiment remains a challenging task [56].

*Experimental system*—We choose to work with a diamond sample containing a dense ensemble of spin-1 NV centers with concentration,  $\rho \sim 4.5$  ppm [56, 57]. The NV centers can be optically initialized and read out using green laser. In the presence of an external magnetic field  $\sim 350$  G, the  $|m_s = \pm 1\rangle$  sublevels are Zeeman split, allowing us to isolate an effective two-level system,  $\{|m_s = 0\rangle, |m_s = -1\rangle\}$  (Fig. 1a). By applying a resonant microwave field with Rabi frequency  $\Omega$ , the effective Hamiltonian governing the system (in the rotating frame) is [7, 8, 58]:

$$\mathcal{H}_0 = - \sum_{i < j} \frac{J_0 \mathcal{A}_{i,j}}{r_{i,j}^3} (S_i^z S_j^z - S_i^x S_j^x - S_i^y S_j^y) + \Omega \sum_i S_i^x, \quad (1)$$

where  $J_0 = (2\pi) \times 52$  MHz·nm<sup>3</sup>,  $\mathcal{A}_{i,j}$  characterizes the angular dependence of the dipolar interaction,  $r_{i,j}$  is the distance between the  $i^{\text{th}}$  and  $j^{\text{th}}$  NV centers, and  $\hat{S}$  is spin operator.

We note that  $\mathcal{H}_0$  contains only the energy-conserving terms of the dipolar interaction under the rotating-wave approximation. The approximation holds because the NV transition frequency is more than three orders of magnitude larger than any other terms in the interacting Hamiltonian [56]. The presence of other paramagnetic spins in the diamond lattice [65] leads to an additional

on-site random field at each NV center which is eliminated using dynamical decoupling (Fig. 1b) [56].

Let us begin by characterizing the dynamics of the NV ensemble under the static Hamiltonian  $\mathcal{H}_0$ . We set  $\Omega = (2\pi) \times 0.05$  MHz, comparable to the average dipolar interaction strength. After optically initializing the NV spins to  $|m_s = 0\rangle$ , we then prepare a product state,  $\otimes_i \frac{|0\rangle_i + |-1\rangle_i}{\sqrt{2}}$ , by applying a global  $\pi/2$ -pulse around the  $\hat{y}$  axis. We let the system evolve under  $\mathcal{H}_0$  for a time  $t$ , before measuring the final NV polarization,  $\langle S^x(t) \rangle$ , along the  $\hat{x}$  direction.

The polarization dynamics proceed in two steps. At early times,  $t \lesssim 100 \mu\text{s}$ , the polarization exhibits rapid decay toward a plateau value, reflecting local equilibration under  $\mathcal{H}_0$  (Fig. 2a). Following these initial dynamics, the system exhibits a slow exponential decay  $\sim A_0 e^{-t/T_0}$  with  $A_0 = (0.43 \pm 0.01)$  and a time-scale  $T_0 = (0.82 \pm 0.03)$  ms that is consistent with spin-phonon relaxation [66]. To ensure that the observed spin dynamics does not come from the incorporated dynamical decoupling pulses, we also investigate the corresponding spin dynamics at  $\Omega = 0$ . The measured NV polarization quickly decays to zero, in agreement with the expectation of thermalization behavior (Fig. 2a Inset).

*Floquet prethermalization*—To probe the nature and existence of Floquet prethermalization, we modulate the Rabi frequency as  $\Omega(t) = \Omega[1 + f(t)]$ , where  $f(t) = \sin(\omega t)$  (Fig. 1b) [67]. Starting with a driving frequency  $\omega = (2\pi) \times 0.07$  MHz, which is comparable to energy scales within  $\mathcal{H}_0$ , we perform the same spin polarization measurement (light blue curve in Fig. 2a). The measured spin dynamics at stroboscopic times,  $t = 2\pi N/\omega$  (where  $N$  is an integer), exhibit an initial relaxation, which is




FIG. 2. **Probing the spin dynamics under periodic and quasi-periodic drives.** (a) The measured spin polarization  $\langle S^x(t) \rangle$  under Floquet drive. After an initial fast decay (gray shaded area), a long-lived prethermal regime persists. Dashed lines are fits of the late-time dynamics using single exponential decay. Inset: Measured initial spin dynamics for the undriven case (red): After the initial fast relaxation, the polarization decays to an equilibration plateau (dashed line), due to the finite static field,  $\Omega \sum_i S_i^x$ . In contrast, the polarization with only dynamical decoupling ( $\Omega = 0$ ) quickly decays to zero. (b) Measured prethermal equilibrium value,  $A(\omega)$ , as a function of  $\omega^{-1}$ . Dashed line is a linear fit with the gray shaded area representing 95% confidence interval. The red diamond marks the measured amplitude for the undriven case. (c) Heating timescale  $\tau^*$  as a function of the driving frequency  $\omega$ . For both sinusoidal and rectangular single-frequency drives,  $\tau^* \sim \mathcal{O}(e^{\omega/J})$ . For sinusoidal quasi-periodic drive,  $\tau^* \sim \mathcal{O}(e^{\omega^{\frac{1}{2}}})$ ; while for rectangular quasi-periodic drive,  $\tau^* \sim \mathcal{O}(\omega^{\frac{1}{2}})$ . (d) Measured spin dynamics under quasi-periodic drive [ $\omega = (2\pi) \times 0.103$  MHz]. We observe an additional small time-quasiperiodic micromotion on top of a slow relaxation. Left inset: zoom-in of the micromotion. Right inset: relative amplitude of the micromotion,  $W$ , scales quadratically with  $\omega^{-1}$ . (e) Using rolling-average to remove the micromotion [56], we observe a quasi-Floquet prethermal regime, whose lifetime  $\tau^*$  increases with  $\omega$ . Dashed lines are fits using single exponential decay. Errorbars on the spin polarization represent 1 s.d. accounting statistical uncertainties, and errorbars on the prethermal plateau and timescale represent 1 s.d. from the fitting.

qualitatively similar to the undriven case. However, the late-time dynamics exhibit a significantly faster polarization decay, arising from Floquet heating. To obtain the heating timescale  $\tau^*$ , we fit the experimental data to  $\sim Ae^{-(\frac{t}{\tau^*} + \frac{t}{T_0})}$ , where  $T_0$  is the previously extracted spin-phonon lifetime. As shown in Fig. 2a, by increasing the driving frequency, one can extract the frequency dependence of the amplitude,  $A(\omega)$ , and the heating time-scale,  $\tau^*(\omega)$ ; both are crucial for understanding the nature of Floquet prethermalization.

Focusing first on the heating time-scale, we find that  $\tau^*$  increases exponentially with  $\omega$  for more than an order of magnitude, demonstrating the existence of a parametrically long-lived prethermal regime (Fig. 2c). The observed exponential scaling also allows us to extract a phenomenological local energy scale of the NV many-body system,  $J_{\text{exp}} = (2\pi) \times (0.032 \pm 0.006)$  MHz.

Intuitively, this  $J_{\text{exp}}$  extracted from Floquet heating process is expected to agree with the local energy scale

of the system. However, for systems with power-law interaction as ours ( $\sim 1/r^3$  in 3D), a naive estimation of the local energy scale  $J \approx \int \frac{J_0}{r^3} \rho d^3\mathbf{r}$  is divergent, and thus, one should expect the prethermalization to not exist. Nevertheless, an important missing piece to this puzzle is the angular dependence of the dipolar interaction,  $\mathcal{A}_{i,j}$  [33]. Crucially, the combination of this angular dependence and the NV's random positioning in the diamond lattice ensures that the average,  $\bar{\mathcal{A}}_{i,j} = 0$ , which helps to mitigate the divergence of the above integral [68].

A more careful analysis reveals that the relevant local energy scale is the variance of the interaction,  $\tilde{J} \approx [f(\frac{J_0 \mathcal{A}_{i,j}}{r^3})^2 \rho d^3\mathbf{r}]^{\frac{1}{2}} = \sqrt{\frac{16\pi}{15}} J_0 \rho$  [56]. As long as the driving frequency  $\omega > \tilde{J}$ , one should expect the presence of prethermalization, in agreement with previous theoretical studies [33]. Using the independently calibrated NV density,  $\rho$ , we estimate the local energy scale  $\tilde{J} \approx (2\pi) \times 0.02$  MHz, which is indeed comparable with

$J_{\text{exp}}$  extracted from Floquet heating.

Next, let us turn to analyzing the frequency dependence of the amplitude,  $A(\omega)$ . One can think of  $A(\omega)$  as the value of the prethermal plateau. In general, for short-range interactions, it is expected that  $A(\omega)$  is determined by a time-independent effective Hamiltonian,  $\mathcal{H}_{\text{eff}}(\omega) = \mathcal{H}_0 + \mathcal{O}(\frac{\tilde{J}}{\omega})$ , which can be calculated order-by-order via a Magnus expansion [69]. In this case,  $A(\omega) = \text{Tr}[\sum_i S_i^x e^{-\beta H_{\text{eff}}(\omega)}] = A_0 + \mathcal{O}(\frac{\tilde{J}}{\omega})$ , where the inverse temperature  $\beta$ , is set by the energy density of the initial state.

For sufficiently long-range interactions (such as dipolar interactions in 3D), the existence of  $\mathcal{H}_{\text{eff}}$  is unproven [14, 26, 69–71]. However, by probing the functional form of  $A(\omega)$  and its extrapolated value as  $\omega \rightarrow \infty$ , one can provide experimental evidence for the existence of  $\mathcal{H}_{\text{eff}}$ . As depicted in Fig. 2b, we find that the frequency dependence of  $A(\omega)$  is linear in  $\omega^{-1}$ , allowing us to extrapolate  $A(\omega \rightarrow \infty) = (0.47 \pm 0.06)$ . This is consistent with the measured value in the undriven case,  $A_0 = (0.43 \pm 0.01)$ , suggesting that despite the presence of strong long-range interactions, the effective Hamiltonian exists and can be well-approximated by  $\mathcal{H}_0$  at leading order [56].

*Quasi-Floquet prethermalization*— We now turn to the quasi-Floquet setting. Specifically, we choose  $f(t) = \frac{1}{2}[\sin(\omega t) + \sin(\varphi\omega t)]$ , where  $\varphi = (\sqrt{5} - 1)/2 \approx 0.618$  is the golden ratio, so that the system is driven by two incommensurate frequencies. From the perspective of Floquet heating, the situation is significantly more complex. In particular, recall that within Fermi’s golden rule, the heating rate can be estimated from the overlap between the Fourier spectrum of the drive,  $F(\nu) = \int f(t)e^{i\nu t} dt$ , and the local spectral function of the spin ensemble,  $S(\nu) = \sum_{i,j} \delta(E_i - E_j - \nu) |\langle i | S^x | j \rangle|^2$  [72].

This picture immediately provides a more formal intuition for the exponentially slow heating observed in the context of single-frequency driving (Fig. 3c). In particular, for  $f(t) = \sin(\omega t)$ ,  $F(\nu)$  exhibits a cut-off at frequency  $\omega$ . Meanwhile, as aforementioned,  $S(\nu)$  exhibits an exponentially small tail for frequencies  $\nu > \tilde{J}$  [73]. In combination, this implies that for a single driving frequency,  $\omega > \tilde{J}$ , energy absorption is strongly suppressed leading to  $\tau^* \sim e^{\omega/\tilde{J}}$ .

For driving with two incommensurate frequencies, even when  $\omega > \tilde{J}$ , there are multi-photon processes that are effectively resonant within the local energy scale; these processes correspond, for example, to the absorption of  $n_1$  photons of energy  $\omega$  and the emission of  $n_2$  photons of energy  $\varphi\omega$ . Thus, there is no strict cut-off for  $F(\nu)$ , and the drive spectrum exhibits a non-zero amplitude for all frequencies  $\nu = |n_1\omega - n_2\varphi\omega| < \tilde{J}$ . Interestingly, despite this, for sufficiently large driving frequencies, seminal results have proven that the quasi-Floquet heating timescale remains extremely slow, exhibiting a stretched exponential lower-bound [74].

In contrast to the Floquet case, we measure the dynamics at evenly spaced time points, since there does not exist a stroboscopic time which is an integer multiple of *both* drives. Much like the single-frequency drive, after an early-time transient, the spin polarization exhibits a slow decay. However, we observe small oscillations scaling as  $\sim \omega^{-2}$  on top of the decay (Fig. 2d), corresponding to the micromotion of the quasi-Floquet system [75–78]. To reliably extract a heating timescale  $\tau^*$  from our quasi-Floquet measurements, we perform a rolling average to obtain the overall decay profile (Fig. 2e) [56]. By varying the driving frequency, we extract a heating time-scale,  $\tau^* \sim e^{\omega^{\frac{1}{2}}}$  (Fig. 2c), which is consistent with the theoretically predicted stretched exponential form [35].

*Robustness of quasi-Floquet prethermalization*—The stability of slow prethermal heating is quite different depending on whether one considers the Floquet or quasi-Floquet setting. For the Floquet setting, the exponential behavior of  $\tau^*$  is robust to the functional form of the drive amplitude  $f(t)$ . However, in the quasi-Floquet setting, the stretched exponential behavior of  $\tau^*$  is *only* predicted to hold when  $f(t)$  is smooth. In particular, when  $f(t)$  is smooth, even though  $F(\nu)$  does not exhibit a cut-off for small  $\nu$ , its amplitude is exponentially small in this regime (Fig. 3c) [35, 36].

These expectations are indeed borne out by the data (Fig. 2c, Fig. 3). Using a rectangular wave  $f(t) = \text{Sgn}[\frac{1}{2}\sin(\omega t) + \frac{1}{2}\sin(\varphi\omega t)]$ , we observe that the heating timescale is significantly shortened and scales as a power-law with increasing driving frequency  $\sim \omega^{\frac{1}{2}}$ , as opposed to a stretched exponential. In contrast, for a single-frequency drive, the smoothness of the driving field is not critical: the Floquet heating time-scale exhibits an exponential scaling for both sinusoidal and rectangular forms of  $f(t)$ .

*Outlook*—Looking forward, our work opens the door to a number of intriguing future directions. First, it is interesting to ask whether the restriction on  $\bar{A}_{i,j} = 0$  is essential for realizing prethermalization in long-range interacting systems [14, 33, 34]. Second, the observed long-lived quasi-Floquet prethermal regime can enable the experimental realization of novel non-equilibrium phases of matter [15, 20, 22, 45–51]. Finally, while our experiments suggest the presence of power-law-slow-heating in the case of a quasi-Floquet, rectangular-wave drive, the precise frequency dependence of the heating rate remains unknown and requires future study.

*Acknowledgements:* We gratefully acknowledge the insights of and discussions with E. Henriksen, P. Y. Hou, A. Jayich, F. Machado, P. Peng, G. Refael, and W. Wu. We thank C. Gaikwad, D. Kowsari, and K. Zheng for their assistance in setting up the experiment. This work is supported by the Startup Fund, the Center for Quantum Leaps, the Institute of Materials Science and Engineering and the OVCR Seed Grant from Washington University.




FIG. 3. **Comparison between sinusoidal and rectangular driving fields.** (a) A representative late-time dynamics of the spin ensemble under single-frequency (Floquet) drive [ $\omega = (2\pi) \times 0.151$  MHz]. Here the plotted polarization is normalized by the intrinsic decay timescale,  $T_0$ , measured without the driving field, i.e. normalized  $\langle S^x(t) \rangle \sim e^{-t/\tau^*}$ . (b) Under quasi-periodic drive [ $\omega = (2\pi) \times 0.143$  MHz], the spin dynamics is extremely sensitive to the smoothness of the drive, and displays a significantly faster decay under a rectangular drive. Errorbars represent 1 s.d. accounting statistical uncertainties. (c) Two-dimensional Fourier spectrum  $F_{n_1, n_2}$  of the driving field  $f(t)$  [56]. The horizontal (vertical) axis corresponds to the absorption of  $n_1$  ( $n_2$ ) photons with energy  $\omega$  ( $\varphi\omega$ ) from the drive, where  $n_1, n_2 = 0, \pm 1, \dots$ . The color of each green dot represents the amplitude of  $F_{n_1, n_2}$  at frequency  $\nu = n_1\omega + n_2\varphi\omega$  (darker represents larger amplitude). The Fourier spectrum  $F(\nu)$  can be thought of as a projection of green dots onto a line with slope  $\varphi$ . Right panel: Fourier spectrum  $F(\nu)$  of quasi-periodic and periodic drive with sinusoidal and rectangular amplitude. For the quasi-Floquet scenario, the spectrum with rectangular drive exhibits resonances at the small frequency range, which overlaps with the system's local spectral function  $S(\nu)$  (black) and leads to energy absorption.

B.Y. acknowledges support from the U.S. Department of Energy (BES grant No. DE-SC0019241). K.W.M. acknowledges support from NSF Grant No. PHY-1752844 (CAREER). N.Y.Y. acknowledges support from the U.S. Department of Energy, Office of Science, through the Quantum Systems Accelerator (QSA), a National Quantum Information Science Research Center and the David and Lucile Packard foundation.

---

[1] Eckardt, A. Colloquium: Atomic quantum gases in periodically driven optical lattices. *Reviews of Modern Physics* **89**, 011004 (2017).

[2] Oka, T. & Kitamura, S. Floquet engineering of quantum materials. *Annual Review of Condensed Matter Physics* **10**, 387–408 (2019).

[3] Harper, F., Roy, R., Rudner, M. S. & Sondhi, S. Topology and broken symmetry in floquet systems. *Annual Review of Condensed Matter Physics* **11**, 345–368 (2020).

[4] Moessner, R. & Sondhi, S. L. Equilibration and order in quantum floquet matter. *Nature Physics* **13**, 424–428 (2017).

[5] Bukov, M., D'Alessio, L. & Polkovnikov, A. Universal high-frequency behavior of periodically driven systems: from dynamical stabilization to floquet engineering. *Advances in Physics* **64**, 139–226 (2015).

[6] Zhang, J. *et al.* Observation of a discrete time crystal. *Nature* **543**, 217–220 (2017). URL <https://doi.org/10.1038/nature21413>.

[7] Choi, S. *et al.* Observation of discrete time-crystalline order in a disordered dipolar many-body system. *Nature* **543**, 221–225 (2017). URL <https://doi.org/10.1038/nature21426>.

[8] Choi, J. *et al.* Robust dynamic hamiltonian engineering of many-body spin systems. *Phys. Rev. X* **10**, 031002 (2020). URL <https://link.aps.org/doi/10.1103/PhysRevX.10.031002>.

[9] Geier, S. *et al.* Floquet hamiltonian engineering of an isolated many-body spin system. *Science* **374**, 1149–1152 (2021). URL <https://www.science.org/doi/abs/10.1126/science.abd9547>. <https://www.science.org/doi/pdf/10.1126/science.abd9547>.

[10] Mi, X. *et al.* Time-crystalline eigenstate order on a quantum processor. *Nature* **601**, 531–536 (2022).

[11] Yao, N. Y., Potter, A. C., Potirniche, I.-D. & Vishwanath, A. Discrete time crystals: Rigidity, criticality, and realizations. *Physical review letters* **118**, 030401 (2017).

[12] Else, D. V., Bauer, B. & Nayak, C. Floquet time crystals. *Physical review letters* **117**, 090402 (2016).

[13] Khemani, V., Lazarides, A., Moessner, R. & Sondhi, S. L. Phase structure of driven quantum systems. *Physical review letters* **116**, 250401 (2016).

[14] Machado, F., Else, D. V., Kahanamoku-Meyer, G. D., Nayak, C. & Yao, N. Y. Long-range prethermal phases of nonequilibrium matter. *Physical Review X* **10**, 011043 (2020).

[15] Potirniche, I.-D., Potter, A. C., Schleier-Smith, M., Vishwanath, A. & Yao, N. Y. Floquet symmetry-protected topological phases in cold-atom systems. *Physical review letters* **119**, 123601 (2017).

[16] Eisert, J., Friesdorf, M. & Gogolin, C. Quantum many-body systems out of equilibrium. *Nature Physics* **11**, 124–130 (2015). URL <https://doi.org/10.1038/nphys3215>.

[17] Ye, B., Machado, F. & Yao, N. Y. Floquet phases of matter via classical prethermalization. *Physical Review*

- Letters* **127**, 140603 (2021).
- [18] Kyprianidis, A. *et al.* Observation of a prethermal discrete time crystal. *Science* **372**, 1192–1196 (2021).
- [19] Randall, J. *et al.* Many-body-localized discrete time crystal with a programmable spin-based quantum simulator. *Science* **374**, 1474–1478 (2021).
- [20] Rechtsman, M. C. *et al.* Photonic floquet topological insulators. *Nature* **496**, 196–200 (2013).
- [21] Dumitrescu, P. T., Vasseur, R. & Potter, A. C. Logarithmically slow relaxation in quasiperiodically driven random spin chains. *Physical review letters* **120**, 070602 (2018).
- [22] Long, D. M., Crowley, P. J. & Chandran, A. Nonadiabatic topological energy pumps with quasiperiodic driving. *Physical Review Letters* **126**, 106805 (2021).
- [23] Peng, P., Yin, C., Huang, X., Ramanathan, C. & Cappellaro, P. Floquet prethermalization in dipolar spin chains. *Nature Physics* **17**, 444–447 (2021). URL <https://doi.org/10.1038/s41567-020-01120-z>.
- [24] Rubio-Abadal, A. *et al.* Floquet prethermalization in a bose-hubbard system. *Physical Review X* **10**, 021044 (2020).
- [25] Beatrez, W. *et al.* Floquet prethermalization with lifetime exceeding 90 s in a bulk hyperpolarized solid. *Physical review letters* **127**, 170603 (2021).
- [26] Abanin, D., De Roeck, W., Ho, W. W. & Huvneers, F. A rigorous theory of many-body prethermalization for periodically driven and closed quantum systems. *Communications in Mathematical Physics* **354**, 809–827 (2017).
- [27] Weidinger, S. A. & Knap, M. Floquet prethermalization and regimes of heating in a periodically driven, interacting quantum system. *Scientific reports* **7**, 1–10 (2017).
- [28] D’Alessio, L. & Rigol, M. Long-time behavior of isolated periodically driven interacting lattice systems. *Physical Review X* **4**, 041048 (2014).
- [29] Ho, W. W., Mori, T., Abanin, D. A. & Dalla Torre, E. G. Quantum and classical floquet prethermalization. *arXiv preprint arXiv:2212.00041* (2022).
- [30] Ye, B., Machado, F., White, C. D., Mong, R. S. K. & Yao, N. Y. Emergent hydrodynamics in nonequilibrium quantum systems. *Phys. Rev. Lett.* **125**, 030601 (2020). URL <https://link.aps.org/doi/10.1103/PhysRevLett.125.030601>.
- [31] Singh, K. *et al.* Quantifying and controlling prethermal nonergodicity in interacting floquet matter. *Phys. Rev. X* **9**, 041021 (2019). URL <https://link.aps.org/doi/10.1103/PhysRevX.9.041021>.
- [32] Viebahn, K. *et al.* Suppressing dissipation in a floquet-hubbard system. *Phys. Rev. X* **11**, 011057 (2021). URL <https://link.aps.org/doi/10.1103/PhysRevX.11.011057>.
- [33] Ho, W. W., Protopopov, I. & Abanin, D. A. Bounds on energy absorption and prethermalization in quantum systems with long-range interactions. *Physical review letters* **120**, 200601 (2018).
- [34] Machado, F., Kahanamoku-Meyer, G. D., Else, D. V., Nayak, C. & Yao, N. Y. Exponentially slow heating in short and long-range interacting floquet systems. *Physical Review Research* **1**, 033202 (2019).
- [35] Else, D. V., Ho, W. W. & Dumitrescu, P. T. Long-lived interacting phases of matter protected by multiple time-translation symmetries in quasiperiodically driven systems. *Phys. Rev. X* **10**, 021032 (2020). URL <https://link.aps.org/doi/10.1103/PhysRevX.10.021032>.
- [36] Mori, T., Zhao, H., Mintert, F., Knolle, J. & Moessner, R. Rigorous bounds on the heating rate in thue-morse quasiperiodically and randomly driven quantum many-body systems. *Physical Review Letters* **127**, 050602 (2021).
- [37] Hahn, E. L. Spin echoes. *Physical review* **80**, 580 (1950).
- [38] Waugh, J. S., Huber, L. M. & Haeberlen, U. Approach to high-resolution nmr in solids. *Physical Review Letters* **20**, 180 (1968).
- [39] Viola, L., Knill, E. & Lloyd, S. Dynamical decoupling of open quantum systems. *Physical Review Letters* **82**, 2417 (1999).
- [40] Zhou, H. *et al.* Quantum metrology with strongly interacting spin systems. *Physical review X* **10**, 031003 (2020).
- [41] Gong, R. *et al.* Coherent dynamics of strongly interacting electronic spin defects in hexagonal boron nitride. *Nature Communications* **14**, 3299 (2023).
- [42] Hahn, W. & Dobrovitski, V. Long-lived coherence in driven many-spin systems: from two to infinite spatial dimensions. *New Journal of Physics* **23**, 073029 (2021).
- [43] Rovny, J., Blum, R. L. & Barrett, S. E. Observation of discrete-time-crystal signatures in an ordered dipolar many-body system. *Physical review letters* **120**, 180603 (2018).
- [44] Beatrez, W. *et al.* Critical prethermal discrete time crystal created by two-frequency driving. *Nature Physics* **1–7** (2023).
- [45] Lindner, N. H., Refael, G. & Galitski, V. Floquet topological insulator in semiconductor quantum wells. *Nature Physics* **7**, 490–495 (2011).
- [46] Wang, Y. H., Steinberg, H., Jarillo-Herrero, P. & Gedik, N. Observation of floquet-bloch states on the surface of a topological insulator. *Science* **342**, 453–457 (2013). URL <https://www.science.org/doi/abs/10.1126/science.1239834>. <https://www.science.org/doi/pdf/10.1126/science.1239834>.
- [47] Po, H. C., Fidkowski, L., Morimoto, T., Potter, A. C. & Vishwanath, A. Chiral floquet phases of many-body localized bosons. *Physical Review X* **6**, 041070 (2016).
- [48] Po, H. C., Fidkowski, L., Vishwanath, A. & Potter, A. C. Radical chiral floquet phases in a periodically driven kitaev model and beyond. *Physical Review B* **96**, 245116 (2017).
- [49] Zhang, X. *et al.* Digital quantum simulation of floquet symmetry-protected topological phases. *Nature* **607**, 468–473 (2022).
- [50] Dumitrescu, P. T. *et al.* Dynamical topological phase realized in a trapped-ion quantum simulator. *Nature* **607**, 463–467 (2022).
- [51] Xu, P., Zheng, W. & Zhai, H. Topological micromotion of floquet quantum systems. *Physical Review B* **105**, 045139 (2022).
- [52] Qi, Z., Refael, G. & Peng, Y. Universal nonadiabatic energy pumping in a quasiperiodically driven extended system. *Physical Review B* **104**, 224301 (2021).
- [53] Peng, Y. & Refael, G. Time-quasiperiodic topological superconductors with majorana multiplexing. *Physical Review B* **98**, 220509 (2018).
- [54] Blekher, P., Jauslin, H. & Lebowitz, J. Floquet spectrum for two-level systems in quasiperiodic time-dependent fields. *Journal of statistical physics* **68**, 271–310 (1992).
- [55] Zhao, H., Mintert, F., Moessner, R. & Knolle, J. Random multipolar driving: tunably slow heating through spec-

- tral engineering. *Physical Review Letters* **126**, 040601 (2021).
- [56] See Supplementary Materials, which includes [79–90].
- [57] Doherty, M. W. *et al.* The nitrogen-vacancy colour centre in diamond. *Physics Reports* **528**, 1–45 (2013).
- [58] Zu, C. *et al.* Emergent hydrodynamics in a strongly interacting dipolar spin ensemble. *Nature* **597**, 45–50 (2021).
- [59] Davis, E. J. *et al.* Probing many-body noise in a strongly interacting two-dimensional dipolar spin system. *Nature Physics* **19**, 836–844 (2023).
- [60] Maricq, M. Matti. Spin thermodynamics of periodically time-dependent systems: The quasistationary state and its decay. *Phys. Rev. B* **36**, 516 (1987).
- [61] M. Matti Maricq. Long-time limitations of the average hamiltonian theory: A dressed-states viewpoint (1990).
- [62] Dimitris Sakellariou and Paul Hodgkinson and Sabine Hediger and Lyndon Emsley. Experimental observation of periodic quasi-equilibria in solid-state nmr. *Chemical Physics Letters* **308**, 381 (1999).
- [63] Carravetta, M. & Levitt, M. H. Long-lived nuclear spin states in high-field solution nmr. *Journal of the American Chemical Society* **126**, 6228–6229 (2004).
- [64] Redfield, A. G. Nuclear spin thermodynamics in the rotating frame: Spins in solids interact weakly and behave irreversibly, yet retain vestiges of isolated-spin dynamics. *Science* **164**, 1015–1023 (1969).
- [65] Such as  $^{13}\text{C}$  nuclear spins and substitutional nitrogen impurities.
- [66] Jarmola, A., Acosta, V., Jensen, K., Chemerisov, S. & Budker, D. Temperature-and magnetic-field-dependent longitudinal spin relaxation in nitrogen-vacancy ensembles in diamond. *Physical review letters* **108**, 197601 (2012).
- [67] We note that  $\Omega(t)$  contains two fundamentally different components: the constant field,  $\Omega \sum_i S_i^x$ , is a part of the previous undriven Hamiltonian  $\mathcal{H}_0$ , while the time-dependent component  $\Omega f(t) \sum_i S_i^x$  acts as the Floquet drive.
- [68] Even if the NV centers are located on a three-dimensional regular lattice, we expect the angular dependence should also average to zero over a large lengthscale, so the positional disorder is not necessary here.
- [69] Kuwahara, T., Mori, T. & Saito, K. Floquet–magnus theory and generic transient dynamics in periodically driven many-body quantum systems. *Annals of Physics* **367**, 96–124 (2016).
- [70] Else, D. V., Bauer, B. & Nayak, C. Prethermal phases of matter protected by time-translation symmetry. *Physical Review X* **7**, 011026 (2017).
- [71] Mori, T., Kuwahara, T. & Saito, K. Rigorous bound on energy absorption and generic relaxation in periodically driven quantum systems. *Physical review letters* **116**, 120401 (2016).
- [72] Here  $E_i$  and  $|i\rangle$  are the eigenenergy and eigenstate of the spin system.
- [73] This is specifically for our case where we need the dipolar interaction to average to zero, in general this is true for the local energy scale  $J$ .
- [74] The predicted lower bound is  $\mathcal{O}(e^{C(\omega/\tilde{J})^{\frac{1}{m}}})$ , where  $m$  is the number of incommensurate driving frequencies and  $C$  is a dimensionless factor [35].
- [75] Sambe, H. Steady states and quasienergies of a quantum-mechanical system in an oscillating field. *Physical Review A* **7**, 2203 (1973).
- [76] Goldman, N. & Dalibard, J. Periodically driven quantum systems: effective hamiltonians and engineered gauge fields. *Physical review X* **4**, 031027 (2014).
- [77] Eckardt, A. & Anisimovas, E. High-frequency approximation for periodically driven quantum systems from a floquet-space perspective. *New journal of physics* **17**, 093039 (2015).
- [78] We note that for a single-frequency drive, similar micro-motion will also emerge if one does not measure the spin dynamics at stroboscopic times [56]. Intuitively, such micro-motion arises from the time-dependent portion of the Hamiltonian which only averages to zero for each complete Floquet cycle.
- [79] Edmonds, A. M. *et al.* Characterisation of cvd diamond with high concentrations of nitrogen for magnetic-field sensing applications. *Materials for Quantum Technology* **1**, 025001 (2021).
- [80] DNV Series Diamond. <https://e6cvd.com/uk/material/single-crystalline/dnv-b14-%203-0mmx3-0mm-0-5mm.html>.
- [81] Jacques, V. *et al.* Dynamic polarization of single nuclear spins by optical pumping of nitrogen-vacancy color centers in diamond at room temperature. *Physical review letters* **102**, 057403 (2009).
- [82] Fischer, R., Jarmola, A., Kehayias, P. & Budker, D. Optical polarization of nuclear ensembles in diamond. *Physical Review B* **87**, 125207 (2013).
- [83] Zu, C. *et al.* Experimental realization of universal geometric quantum gates with solid-state spins. *Nature* **514**, 72–75 (2014).
- [84] Mrózek, M. *et al.* Longitudinal spin relaxation in nitrogen-vacancy ensembles in diamond. *EPJ Quantum Technology* **2**, 22 (2015).
- [85] Choi, J. *et al.* Depolarization dynamics in a strongly interacting solid-state spin ensemble. *Physical review letters* **118**, 093601 (2017).
- [86] Mittiga, T. *et al.* Imaging the local charge environment of nitrogen-vacancy centers in diamond. *Physical review letters* **121**, 246402 (2018).
- [87] Aslam, N., Waldherr, G., Neumann, P., Jelezko, F. & Wrachtrup, J. Photo-induced ionization dynamics of the nitrogen vacancy defect in diamond investigated by single-shot charge state detection. *New Journal of Physics* **15**, 013064 (2013).
- [88] Manson, N. B. *et al.* Nv—n+ pair centre in 1b diamond. *New Journal of Physics* **20**, 113037 (2018).
- [89] Jayakumar, H. *et al.* Optical patterning of trapped charge in nitrogen-doped diamond. *Nature communications* **7**, 1–8 (2016).
- [90] Bluvstein, D., Zhang, Z. & Jayich, A. C. B. Identifying and mitigating charge instabilities in shallow diamond nitrogen-vacancy centers. *Physical review letters* **122**, 076101 (2019).