



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Quantum Phase Transitions in Optomechanical Systems

Bo Wang, Franco Nori, and Ze-Liang Xiang

Phys. Rev. Lett. **132**, 053601 — Published 30 January 2024

DOI: [10.1103/PhysRevLett.132.053601](https://doi.org/10.1103/PhysRevLett.132.053601)

Quantum Phase Transitions in Optomechanical Systems

Bo Wang,¹ Franco Nori,^{2,3,4} and Ze-Liang Xiang^{1,*}

¹*School of Physics, Sun Yat-sen University, Guangzhou 510275, China*

²*Theoretical Quantum Physics Laboratory, Cluster for Pioneering Research, RIKEN, Wako-shi, Saitama 351-0198, Japan*

³*Center for Quantum Computing, RIKEN, Wako-shi, Saitama 351-0198, Japan*

⁴*Department of Physics, University of Michigan, Ann Arbor, Michigan 48109-1040, USA*

In this letter, we investigate the ground state properties of an optomechanical system consisting of a coupled cavity and mechanical modes. An exact solution is given when the ratio η between the cavity and mechanical frequencies tends to infinity. This solution reveals a coherent photon occupation in the ground state by breaking continuous or discrete symmetries, exhibiting an equilibrium quantum phase transition (QPT). In the $U(1)$ -broken phase, an unstable Goldstone mode can be excited. In the model featuring Z_2 symmetry, we discover the mutually (in the finite η) or unidirectionally (in $\eta \rightarrow \infty$) dependent relation between the squeezed vacuum of the cavity and mechanical modes. In particular, when the cavity is driven by a squeezed field along the required squeezing parameter, it enables modifying the region of Z_2 -broken phase and significantly reducing the coupling strength to reach QPTs. Furthermore, by coupling atoms to the cavity mode, the hybrid system can undergo a QPT at a hybrid critical point, which is cooperatively determined by the optomechanical and light-atom systems. These results suggest that this optomechanical system complements other phase transition models for exploring novel critical phenomena.

Introduction.—Quantum phase transitions have garnered significant attention for their crucial role in comprehending the various intricacies involved in the evolution of matter phases [1–3]. Equilibrium QPTs can be characterized by the closing spectral gap and the emergence of degenerate ground states due to the spontaneous breaking of a continuous (or discrete) symmetry. For systems possessing continuous $U(1)$ symmetry, such as the Bose-Hubbard model, their phase transitions [4, 5] are attracting intense interest due to their involvement in exhibiting notable physics of the Anderson-Higgs mechanism [6, 7].

In quantum optics, the well-known Dicke model, describing the coupling between a cavity mode and N two-level systems, can exhibit a superradiant QPT in the thermodynamic limit, $N \rightarrow \infty$, where the bosonic mode gains the occupation of macroscopic coherence in the ground state [8–10]. Recently, this QPT was also predicted in the Rabi model in the classical oscillator limit [11, 12], where the ratio between frequencies of the atomic transition and the cavity mode approaches infinity. These two fundamental spin-boson models can be reduced to a model with a $U(1)$ symmetry by dropping the so-called counter-rotating terms, resulting in the emergence of Goldstone modes [13, 14]. In the ultrastrong-coupling regime, these terms usually cannot be ignored [15, 16]; however, recent progress indicated that this requirement could be realized by engineering the light-matter interaction in circuit-QED systems [14] or through quantum simulation techniques [17–19].

Previous investigations of the superradiant QPT have predominantly focused on the spin-boson model, with either infinite [20–31] or finite components [32–39]. This naturally leads to an intriguing question of whether the boson-boson model can similarly exhibit a rich range of

quantum phases, thereby improving our understanding of QPTs. The optomechanical system considered here is a boson-boson model, describing the interaction between a cavity mode and a mechanical oscillator through radiation pressure [40, 41]. With recent developments of strong single-photon optomechanical couplings [42–51], it enables the investigation of quantum nonlinear effects in optomechanical systems, such as preparing non-classical states [52–57] and observing the dynamical Casimir effect [58–62]. These nonlinear effects lead to energy-level repulsion and attraction that can cause the degeneration of the lowest levels, implying the emergence of QPTs.

In this letter, we study the ground state properties of optomechanical systems based on models with either $U(1)$ symmetry or Z_2 symmetry and give the analytical solutions of their equilibrium QPTs. The exact solution for a model possessing $U(1)$ symmetry reveals the instability of the ground state and indicates the emergence of a Goldstone mode. For the model with Z_2 -symmetry, in the finite η , displacing the phonon space can generate a pair of mutually dependent squeezed vacuum between the cavity and mechanical modes via radiation pressure; however, in the limit, $\eta \rightarrow \infty$, such dependence is unidirectional. Interestingly, applying a squeezed field to drive the cavity with the required squeezing parameter, we find that the features of QPTs can be remarkably influenced: the region where the Z_2 -broken phase occurs is alterable, and the coupling strength to reach the critical point can be significantly reduced. In addition, interacting with atoms, the hybrid system will have a hybrid critical point, where the optomechanical and light-atom components of the system could cooperatively determine the critical phenomena.

Model.—We consider a typical optomechanical system

consisting of a cavity with a movable mirror. The system Hamiltonian [63] can be written as

$$H = \omega_c a^\dagger a + \omega_m b^\dagger b + g(a + a^\dagger)^2(b + b^\dagger). \quad (1)$$

where the annihilation operator $a(b)$ denotes the optical (mechanical) mode with the resonant frequency $\omega_c(\omega_m)$, and g is the strength of single-photon optomechanical coupling. For most experiments to date [40], the Hamiltonian $H_{\text{om}} = \omega_c a^\dagger a + \omega_m b^\dagger b + 2ga^\dagger a(b + b^\dagger)$ is a sufficient good approximation. While the single-photon coupling strength is increasing, the terms $g(a^2 + a^{\dagger 2})(b + b^\dagger)$, describing the creation and annihilation of photon pairs [58, 59], become considerable. Obviously, $2ga^\dagger a(b + b^\dagger)$ and $g(a^2 + a^{\dagger 2})(b + b^\dagger)$ have distinct effects on the photon occupation in the ground state due to the different symmetries of the cavity mode.

Photon occupations in ground state.— First we focus on the simplest model described by the Hamiltonian H_{om} . Let us denote $|n\rangle$ as an n -photon Fock state and $|k\rangle$ as an k -phonon Fock state, respectively. By performing a unitary transformation with $U = \exp[-(g/\omega_m)a^\dagger a(b^\dagger - b)]$, the Hamiltonian becomes $\tilde{H}_{\text{om}} = (U^\dagger H_{\text{om}} U)/\omega_c = a^\dagger a + \eta^{-1} b^\dagger b - (1/\kappa^2) a^\dagger a a^\dagger a$ with a dimensionless coupling strength $\kappa = \sqrt{\omega_c \omega_m}/2g$ and a frequency ratio $\eta = \omega_c/\omega_m$, where the eigenstate and the eigenvalue can be described by the ket $|n, k\rangle$ and $E_{\text{om}} = n - (n^2/\kappa^2) + \eta^{-1}k$, respectively. In the displaced basis, \tilde{H}_{om} has a conserved operator, $\tilde{P} = \exp[i\theta N]$ with $\theta \in [0, 2\pi]$ and $N = a^\dagger a + b^\dagger b$, such that $[\tilde{H}_{\text{om}}, \tilde{P}] = 0$, satisfying a $U(1)$ -continuous symmetry. Going back to the original basis, the corresponding conserved operator and eigenstates can be written as

$$P = U \tilde{P} U^\dagger = e^{i\theta(a^\dagger a + b^\dagger b + \frac{2g}{\omega_m} a^\dagger a(b + b^\dagger) + \frac{4g^2}{\omega_m^2} a^\dagger a a^\dagger a)} \quad (2)$$

and $|\psi\rangle = |n, k_n\rangle = D(n\sqrt{\eta}/\kappa)|n, k\rangle$, with $D(x) = \exp[x(b - b^\dagger)]$, respectively.

Now we consider a particular limit, i.e., $\eta \rightarrow \infty$. The Hamiltonian can be rewritten as

$$\tilde{H}_{\text{om}} = a^\dagger a - \frac{1}{\kappa^2} a^\dagger a a^\dagger a \quad (3)$$

with eigenvalues $\tilde{E}_{\text{om}} = n[1 - (n/\kappa^2)]$, exhibiting an anharmonic spectrum. To obtain a well-defined ground state in Eq. (3), $a^\dagger a$ needs to be treated as a perturbation, which means the system energy is dominated by the negative nonlinear term with $(-1/\kappa^2)n^2$ [64]. Therefore, we can perform the transformation $\tilde{E}_{\text{om}} \rightarrow -\tilde{E}_{\text{om}}$ shown in Fig. 1 (or $\bar{E}_G \rightarrow -\bar{E}_G$ shown in Fig. 2), which only changes the reference frame but allows us to conveniently capture the nature of the ground state.

For $\kappa < 1$, the well-defined ground state of \tilde{H}_{om} is the vacuum state $|0\rangle$, until at $\kappa = 1$ there occurs a level crossing between $|0\rangle$ and $|1\rangle$. After that, the ground state meets a series of level crossings between the states $|n\rangle$ and




FIG. 1. Analytic solution of the Hamiltonian H_{om} . (a) Level crossings of the low-energy state near the ground state for a frequency ratio $\eta \rightarrow \infty$. (b) The total number of excitations in the ground state for a frequency ratio $\eta \rightarrow \infty$.

$|n+1\rangle$, as shown in Fig. 1(a). From $\tilde{E}_{\text{om}}^{(n+1)} - \tilde{E}_{\text{om}}^{(n)} = 0$, the number of photons occupying the ground state, denoted as n_G , can be described by

$$n_G = \frac{\kappa^2 - 1}{2}. \quad (4)$$

Obviously, as κ increases, the quantity $[n_G]$, which counts the number of photons using the ceiling function $[\]$, experiences a discrete step-wise ascent, as shown in Fig. 1(b), showing the trend of photon occupations in the ground state.

Results in mean-field approach.—Next we use the mean-field approach, a semiclassical approximation, to observe the distribution of the ground-state energy. By displacing both modes a and b in \tilde{H}_{om} , with respect to their corresponding mean values α and β , and neglecting the fluctuations, we can obtain the mean value of the ground state energy in terms of α and β ,

$$\bar{E}_G = |\alpha|^2 + \eta^{-1}|\beta|^2 - \frac{1}{\kappa^2}(|\alpha|^2 + |\alpha|^4). \quad (5)$$

When $\kappa < 1$, the ground state energy has a single minimum at $\alpha = \beta = 0$, where the energy is $\bar{E}_G = 0$, as shown in Fig. 2(a). For $\kappa > 1$, the distribution of the ground-state energy abruptly changes as a profile of Mexican hat shown in Fig. 2(b), where the ground state in $\alpha = 0$ becomes unstable and will fall into the stable one with the energy minima occurring at

$$\alpha = \pm e^{i\theta} \sqrt{\frac{1}{2}(\kappa^2 - 1)} \text{ and } \beta = 0. \quad (6)$$

Because the phase θ can be an arbitrary value from 0 to 2π , the ground states are infinitely degenerate with the energy minima lying on a circle, which is known as Goldstone mode [65] with broken $U(1)$ -symmetry.

For $\kappa > 1$, the ground state energy becomes $\bar{E}_G(\kappa) = (1/4)(\kappa^2 + \kappa^{-2} - 2)$. When the ground state energy is continuous, there is a discontinuity in the second derivative of $\bar{E}_G(\kappa)$ at $\kappa = 1$, revealing a second-order




FIG. 2. Mean-field solution of the Hamiltonian \bar{H}_{om} . Mean-field energy of the ground state when $\kappa = 0.5$ (a) and when $\kappa = 2$ (b). (c) Ground state energy \bar{E}_G (solid blue line) and its second derivative $d^2 \bar{E}_G / d\kappa^2$ (red dashed line). (d) Coherence $\langle a \rangle$ of the cavity field in its ground state.

phase transition [Fig. 2(c)]. In Eq. (6), the nonzero values of α mean that the ground state in $\kappa > 1$ has a nonzero coherence of the cavity mode, which is an order parameter of the phase transition [Fig. 2(d)] and indicates the spontaneous breaking of $U(1)$ symmetry. Based on nonzero coherence, we can obtain the photon occupation in the ground state, namely, $\langle a^\dagger a \rangle_G = (\kappa^2 - 1)/2$, which is consistent with the solution of Eq. (4).

Although the results in Fig. 2 can significantly indicate that the model H_{om} can undergo a phase transition through the critical point $\kappa = 1$, the lack of thermodynamic limit in Eq. (6) implies that spontaneous symmetry breaking would be restored by quantum fluctuations. Nevertheless, for a finite-component system [10] or finite-frequency system [12], quantum fluctuations still require a finite time to restore the symmetry of the real ground state [66]. Therefore, it is possible to observe the above results, even though the symmetry-breaking ground state is not stable.

Squeezed vacuum in photon and phonon modes.— When the Hamiltonian includes $g(a^2 + a^{\dagger 2})(b + b^\dagger)$, it is difficult to determine the symmetry of the system. However, we find that the Hamiltonian H commutes with the operator $P_c = \exp(i\pi a^\dagger a)$, indicating that the cavity mode possesses an independent Z_2 -parity symmetry. Accordingly, one may execute a displacement transformation on the mechanical mode with the unitary operator, $U_1 = \exp[-(g/\omega_m)(b^\dagger - b)]$,

and the transformed Hamiltonian $U_1^\dagger H U_1$ becomes

$$\begin{aligned} \bar{H} = & \omega_c a^\dagger a + \omega_m b^\dagger b - \frac{2g^2}{\omega_m} (a + a^\dagger)^2 + \frac{g^2}{\omega_m} \\ & + g(a^2 + a^{\dagger 2})(b + b^\dagger) + 2ga^\dagger a(b + b^\dagger). \end{aligned} \quad (7)$$

Note that this displacement elicits an anti-squeezing term, the third one in Eq. (7), which could induce non-analytic behavior of the cavity mode at one point upon changing the coupling strength [64]. Such behavior will give feedback to the mechanical oscillator by radiation pressure and potentially influence the phonon mode's behavior. To gain insight into the underlying physics, one can perform a transformation on the Hamiltonian \bar{H} [64] with the unitary operator, $U_2 = \exp[-(g/2\omega_c)(b^\dagger + b)(a^{\dagger 2} - a^2)]$. This transformed Hamiltonian, $\tilde{H} = U_2^\dagger \bar{H} U_2$, although not yet able to be diagonalized, may be analyzed via a variational method.

We propose a trial wave function $|\psi(r, s)\rangle = \mathcal{S}_a(r)\mathcal{S}_b(s)|0_a, 0_b\rangle$ for \tilde{H} , where $\mathcal{S}_y(x) = \exp[(x^2/2)(y^{\dagger 2} - y^2)]$ is the squeezing operator with a bosonic operator $y \in [a, b]$ and a variational parameter $x \in [r, s]$ providing the energy function $\tilde{E}(r, s)$ [64]. Without loss of generality, we here require that the energy function is up to fourth order in γ in the following analysis, where $\gamma = 2\sqrt{2}g/\sqrt{\omega_c\omega_m}$ is a dimensionless coupling strength. By minimizing the energy with respect to r and s , we can obtain

$$e^{4r} = \frac{1 + \frac{1}{4}\gamma^2\eta^{-1}e^{2s} + \frac{1}{32}\gamma^4\eta^{-2}e^{4s}}{1 - \gamma^2(1 + \frac{3}{4}\eta^{-1}e^{2s} + \frac{1}{4}\gamma^2\eta^{-1}e^{2s} + \frac{7}{32}\gamma^2\eta^{-2}e^{4s})} \quad (8)$$

and

$$e^{4s} = \frac{[\frac{3}{32}\gamma^4\eta^{-1}(8\sinh^2 r + 4) + \frac{1}{2}\gamma^4\eta^{-1}\sinh 2r]e^{6s} + 1}{[1 - \gamma^2(\sinh^2 r + \sinh 2r + \frac{1}{2} + \frac{1}{4}\gamma^2 e^{2r})]} \quad (9)$$

When η is finite, nonzero and correlated solutions for the squeezing parameters r and s always exist if $\gamma < 1$. This reflects the inseparable relation between the squeezed vacuum states of the cavity and mechanical modes induced by the radiation pressure.

When $\eta \rightarrow \infty$, Eqs. (8,9) can be rewritten as $e^{4r} = 1/(1 - \gamma^2)$ and $e^{4s} = 1/[1 - \gamma^2(\sinh^2 r + \sinh 2r + \frac{1}{2} + \frac{1}{4}\gamma^2 e^{2r})]$, respectively. In this case, r no longer relies on s , but s still depends on r , indicating that the cavity mode carries an irreversible feedback to the mechanical mode. This arises from the deterministic symmetry of the cavity mode itself, independent of the mechanical mode, namely, $[H, P_c] = 0$. Moreover, as $\gamma \rightarrow \gamma_c = 1$, a transition arises, signifying a profound divergence in both r and s . These divergences in the ground state imply the phase transitions of the system [10, 12], where the two-fold degeneracy occurs. Interestingly, the phase transition of the mechanical mode would suggest a hidden symmetry within H , which is cooperatively determined

by the cavity and mechanical modes and would be broken in the ground state.

Superradiant phase.—To address the superradiant phase in the Hamiltonian H , it is necessary to recognize that $H_I = g(a + a^\dagger)^2(b + b^\dagger)$ is unbounded, which is an obstacle in locating the position of macroscopic coherence of the bosonic mode in the ground state [67]. One feasible way is to introduce a nonlinear quartic term to each bosonic mode, thereby obtaining a Hamiltonian

$$H_{\text{op}} = N\omega_c a^\dagger a + \omega_m b^\dagger b + g(a^2 + a^{\dagger 2} + 2a^\dagger a)(b + b^\dagger) + Ng(b + b^\dagger) + \frac{\epsilon_1}{N^2} a^\dagger a^\dagger a a + \frac{\epsilon_2}{N^2} b^\dagger b^\dagger b b, \quad (10)$$

where N is a macroscopic factor. Given the factor N , the detuning between the two oscillators is determined. As N becomes large, the resulting Hamiltonian in Eq. (10) is equivalent to the one in Eq. (1), faithfully illustrating that the two nonlinear quartic terms are vanishingly small.

Next, we adopt a mean-field approach to examine the ground state energy of the Hamiltonian H_{op} by displacing both modes a and b from their mean values, denoted as α and β . The expression of the ground state energy is given by

$$E_G = N\omega_c \alpha^2 + \omega_m \beta^2 + 2g\beta(N + 4\alpha^2) + \frac{\epsilon_1}{N^2} \alpha^4 + \frac{\epsilon_2}{N^2} \beta^4. \quad (11)$$

To minimize the energy E_G , we let $\beta/\alpha^2 \sim 1$ and $\alpha \sim \sqrt{N}$, which yields

$$\alpha^2 = \frac{N}{4}(\gamma^2 - 1) \text{ and } \beta = -\frac{N\omega_c}{8g}, \quad (12)$$

where $\epsilon_1 = \epsilon_2 = (4\omega_m^2/\omega_c)(\gamma^6 - \gamma^2)$. For $\gamma > 1$, the nonzero coherence of the mode $\langle a \rangle = \pm\alpha$ indicates the superradiant phase with a spontaneously broken-parity symmetry P_c .

Modification to features of QPT.—We will now show that by employing a controllable squeezed field of the cavity mode to drive the optomechanical system, it is possible to modify the characteristics of the QPT. To illustrate this point, we apply the transformation with the unitary operator, $S_\zeta = \exp[(1/2)(\zeta^* a^2 - \zeta a^{\dagger 2})]$, where $\zeta = \xi \exp(i\theta)$ represents the squeezed parameter, to the system Hamiltonian H with a squeezing-driven term $\xi(a^{\dagger 2} e^{-i\theta} + a^2 e^{i\theta})$. The transformed Hamiltonian depends on the squeezing direction θ and the squeezing amplitude ξ [64], providing a channel to modify the ground state properties of the cavity mode.

First, we consider the squeezing direction $\theta = 0$, where the Hamiltonian $H_{(\theta=0,\xi)}$ exhibits a similar structure to H . By performing a displacement transformation with $\tilde{U} = \exp[-(ge^{-2\xi}/\omega_m)(b^\dagger - b)]$, and taking the limit $\eta \rightarrow \infty$, the Hamiltonian $H_{(\theta=0,\xi)}$ can be

diagonalized as $\tilde{H}_{(\theta=0,\xi)} = 2\varepsilon_\xi d^\dagger d + \tilde{E}_{G(\xi)}$, with $\varepsilon_\xi = \sqrt{(1/4)(1 - \gamma^2 e^{-2\xi})}$. For $\xi = 2 \ln(\gamma)$, ε_ξ simplifies to

$$\varepsilon_{\xi \rightarrow 2 \ln(\gamma)} = \sqrt{(1/4)(1 - \gamma^{-2})}, \quad (13)$$

which is real only for $\gamma \geq 1$ and vanishes at $\gamma = 1$. However, when $\xi = 0$, i.e., without the squeezing-driven field, the eigenvalue is given by $\varepsilon_{\xi \rightarrow 0} = \sqrt{(1/4)(1 - \gamma^2)}$, indicating the occurrence of the phase for $\gamma < 1$. This result demonstrates that the region where the quantum phase occurs can be altered by driving the system with a squeezed field of the squeezing direction $\theta = 0$ and an appropriate squeezing amplitude. This feature is remarkably unusual within the field of phase transitions, indicating the modifying capability on phase diagrams.

Furthermore, for the squeezing direction $\theta = \pi$, the corresponding eigenvalue is given by $\varepsilon_\xi = \sqrt{(1/4)(1 - \gamma^2 e^{2\xi})}$, indicating that the optomechanical coupling strength required to reach the critical point can be exponentially reduced by increasing the squeezing amplitude ξ . This result suggests the possibility of the phase transition in the optomechanical system even with a common coupling strength. Of note, such a way cannot apply to the Rabi and Dicke models, as these models lack the inherent symmetry of the cavity mode itself.

QPTs in hybrid systems.—We now turn to an optomechanical system interacting with atoms in the cavity, as shown in Fig. 3. The results reveal that QPT can emerge in such a system, which has a *hybrid* critical point cooperatively determined by the optomechanical system and the light-atom system. Here, we employ the Dicke model to describe the light-atom interaction. The Hamiltonian of this hybrid system [68] can be written as

$$H_h = H + \omega_a J_z + \frac{\lambda}{\sqrt{N_a}}(a + a^\dagger)(J_+ + J_-) + \chi(a + a^\dagger)^2 \quad (14)$$

with the angular momentum operator $J_z = (1/2) \sum_{i=1}^{N_a} \sigma_z^{(i)}$ and $J_\pm = \sum_{i=1}^{N_a} \sigma_\pm^{(i)}$, satisfying the commutation relation $[J_-, J_+] = -2J_z$. The last term in Eq. (14) is so-called A^2 term with $\chi = \alpha\lambda^2/\omega_a$ [69]. The coefficient $\alpha \geq 1$ ensures that the Dicke model has a precisely gauge-invariant Hamiltonian satisfying the TRK sum rule [70, 71], while $\alpha = 0$ corresponds to the standard Dicke model [10]. After diagonalizing the Hamiltonian H_h [64], the spectrum shows that the excitation energy of the lowest branch ϵ_- vanishes at

$$\mu^2(1 - \alpha) + \gamma^2 = 1, \quad (15)$$

indicating a quantum critical point of the hybrid system, where $\mu = 2\lambda/\sqrt{\omega_a\omega_c}$ is a dimensionless coupling strength of the light-atoms interacting system.

When neglecting the A^2 term, i.e., $\alpha = 0$, the critical point can be expressed as $\mu^2 + \gamma^2 = 1$. This formula contains two distinct types of dimensionless coupling strength, μ and γ , which correspond to the critical




FIG. 3. Schematic of a hybrid system consisting of a cavity coupled to both a movable mirror and an atomic ensemble.

points in the standard Dicke model ($\mu^2 = 1$) and the optomechanical model ($\gamma^2 = 1$), respectively. Thus, the hybrid quantum system features a *hybrid* critical point that separates the normal and superradiant phases, where the critical phenomena are dominated by both the light-atom and optomechanical systems.

When $\alpha = 1$, the *hybrid* critical point becomes $\gamma^2 = 1$. In this scenario, one trend of the closing spectrum gap, induced by the light-atom interaction, is completely suppressed by the \mathbf{A}^2 term, in agreement with the no-go theorem [70]. However, the other trend, caused by the interaction between the cavity mode and the mechanical oscillator, results in a gapless spectrum, where the critical point only depends on γ . Nevertheless, the energy spectrum for the superradiant phase remains influenced by μ [64].

Conclusion and outlook.—We demonstrate that several optomechanical systems can exhibit distinct spontaneous broken-symmetry phases, either continuous or discrete, yielding different coherent photon occupations in the ground state. By interacting with the two-level atoms, we find that the closing spectrum gap of the hybrid quantum system is determined by two distinct types of coupling degrees of freedom, which results in the emergence of a hybrid critical point.

In addition, the hybrid critical point may display anomalous behavior when a drive with squeezed light is applied to the cavity. With appropriate squeezed parameters, it is possible to find two superradiant phases separated by this critical point. Considering that these two phases are independently induced by the two subsystems, respectively, they should be characterized by the two corresponding thermodynamic limits (η and N_a), yielding two distinct order parameters. In this scenario, exploring whether such a system can undergo a second-order QPT between the two ordered phases with different broken symmetries would be an interesting topic beyond the Landau-Ginzburg-Wilson paradigm [72–74]. With the impressive ongoing advancements in technology that allow for the realization of strong and even ultra-strong coupling between the cavity and a movable mirror, cavity optomechanical systems will become a potential platform for investigating critical phenomena.

Acknowledgments.—We thank S. Yin and S. Ashhab for critical reading and stimulating discussions. This

work was supported by the National Natural Science Foundation of China (Grant No. 11874432), the National Key R&D Program of China (Grant No. 2019YFA0308200), and the China Postdoctoral Science Foundation (Grant No. 2021M693682). F. N. is supported in part by Nippon Telegraph and Telephone Corporation (NTT) Research, the Japan Science and Technology Agency (JST) [via the Quantum Leap Flagship Program (Q-LEAP) and the Moonshot R&D Grant No. JPMJMS2061], the Asian Office of Aerospace Research and Development (AOARD) (via Grant No. FA2386-20-1-4069), and the Foundational Questions Institute Fund (FQXi) via Grant No. FQXiIAF19-06.

* xiangzliang@mail.sysu.edu.cn

- [1] S. Sachdev, *Quantum Phase Transitions*, 2nd ed. (Cambridge University Press, Cambridge, England, 2011).
- [2] S. L. Sondhi, S. M. Girvin, J. P. Carini, and D. Shahar, Continuous quantum phase transitions, *Rev. Mod. Phys.* **69**, 315 (1997).
- [3] M. Vojta, Quantum phase transitions, *Reports on Progress in Physics* **66**, 2069 (2003).
- [4] M. Endres, T. Fukuhara, D. Pekker, M. Cheneau, P. Schauß, C. Gross, E. Demler, S. Kuhr, and I. Bloch, The ‘Higgs’ amplitude mode at the two-dimensional superfluid/Mott insulator transition, *Nature* **487**, 454 (2012).
- [5] S.-P. Wang, A. Ridolfo, T. Li, S. Savasta, F. Nori, Y. Nakamura, and J. Q. You, Probing the symmetry breaking of a light-matter system by an ancillary qubit, *Nature Communications* **14**, 4397 (2023).
- [6] P. W. Anderson, Plasmons, gauge invariance, and mass, *Phys. Rev.* **130**, 439 (1963).
- [7] P. W. Higgs, Broken symmetries and the masses of gauge bosons, *Phys. Rev. Lett.* **13**, 508 (1964).
- [8] K. Hepp and E. H. Lieb, On the superradiant phase transition for molecules in a quantized radiation field: the Dicke maser model, *Annals of Physics* **76**, 360 (1973).
- [9] Y. K. Wang and F. T. Hioe, Phase transition in the Dicke model of superradiance, *Phys. Rev. A* **7**, 831 (1973).
- [10] C. Emary and T. Brandes, Chaos and the quantum phase transition in the Dicke model, *Phys. Rev. E* **67**, 066203 (2003).
- [11] S. Ashhab, Superradiance transition in a system with a single qubit and a single oscillator, *Phys. Rev. A* **87**, 013826 (2013).
- [12] M.-J. Hwang, R. Puebla, and M. B. Plenio, Quantum phase transition and universal dynamics in the Rabi model, *Phys. Rev. Lett.* **115**, 180404 (2015).
- [13] M.-J. Hwang and M. B. Plenio, Quantum phase transition in the finite Jaynes-Cummings lattice systems, *Phys. Rev. Lett.* **117**, 123602 (2016).
- [14] A. Baksic and C. Ciuti, Controlling discrete and continuous symmetries in “superradiant” phase transitions with circuit QED systems, *Phys. Rev. Lett.* **112**, 173601 (2014).
- [15] A. Frisk Kockum, A. Miranowicz, S. De Liberato, S. Savasta, and F. Nori, Ultrastrong coupling between

- light and matter, *Nature Reviews Physics* **1**, 19 (2019).
- [16] P. Forn-Díaz, L. Lamata, E. Rico, J. Kono, and E. Solano, Ultrastrong coupling regimes of light-matter interaction, *Rev. Mod. Phys.* **91**, 025005 (2019).
- [17] N. K. Langford, R. Sagastizabal, M. Kounalakis, C. Dickel, A. Bruno, F. Luthi, D. J. Thoen, A. Endo, and L. DiCarlo, Experimentally simulating the dynamics of quantum light and matter at deep-strong coupling, *Nature communications* **8**, 1715 (2017).
- [18] A. Mezzacapo, U. Las Heras, J. Pedernales, L. DiCarlo, E. Solano, and L. Lamata, Digital quantum Rabi and Dicke models in superconducting circuits, *Scientific Reports* **4**, 7482 (2014).
- [19] I. M. Georgescu, S. Ashhab, and F. Nori, Quantum simulation, *Rev. Mod. Phys.* **86**, 153 (2014).
- [20] F. Dimer, B. Estienne, A. S. Parkins, and H. J. Carmichael, Proposed realization of the Dicke-model quantum phase transition in an optical cavity qed system, *Phys. Rev. A* **75**, 013804 (2007).
- [21] K. Baumann, C. Guerlin, F. Brennecke, and T. Esslinger, Dicke quantum phase transition with a superfluid gas in an optical cavity, *Nature* **464**, 1301 (2010).
- [22] K. Baumann, R. Mottl, F. Brennecke, and T. Esslinger, Exploring symmetry breaking at the Dicke quantum phase transition, *Phys. Rev. Lett.* **107**, 140402 (2011).
- [23] D. Nagy, G. Kónya, G. Szirmai, and P. Domokos, Dicke-model phase transition in the quantum motion of a Bose-Einstein condensate in an optical cavity, *Phys. Rev. Lett.* **104**, 130401 (2010).
- [24] J. Klinder, H. Keßler, M. Wolke, L. Mathey, and A. Hemmerich, Dynamical phase transition in the open Dicke model, *Proceedings of the National Academy of Sciences* **112**, 3290 (2015).
- [25] J. Vidal and S. Dusuel, Finite-size scaling exponents in the Dicke model, *Europhysics Letters* **74**, 817 (2006).
- [26] L. Bakemeier, A. Alvermann, and H. Fehske, Quantum phase transition in the Dicke model with critical and noncritical entanglement, *Phys. Rev. A* **85**, 043821 (2012).
- [27] N. Lambert, C. Emary, and T. Brandes, Entanglement and the phase transition in single-mode superradiance, *Phys. Rev. Lett.* **92**, 073602 (2004).
- [28] D. Guerci, P. Simon, and C. Mora, Superradiant phase transition in electronic systems and emergent topological phases, *Phys. Rev. Lett.* **125**, 257604 (2020).
- [29] G. Mazza and A. Georges, Superradiant quantum materials, *Phys. Rev. Lett.* **122**, 017401 (2019).
- [30] G. Chiriacò, M. Dalmonte, and T. Chanda, Critical light-matter entanglement at cavity mediated phase transitions, *Phys. Rev. B* **106**, 155113 (2022).
- [31] J. Zhao and M.-J. Hwang, Frustrated superradiant phase transition, *Phys. Rev. Lett.* **128**, 163601 (2022).
- [32] R. Puebla, M.-J. Hwang, J. Casanova, and M. B. Plenio, Probing the dynamics of a superradiant quantum phase transition with a single trapped ion, *Phys. Rev. Lett.* **118**, 073001 (2017).
- [33] M. Liu, S. Chesi, Z.-J. Ying, X. Chen, H.-G. Luo, and H.-Q. Lin, Universal scaling and critical exponents of the anisotropic quantum Rabi model, *Phys. Rev. Lett.* **119**, 220601 (2017).
- [34] L. Garbe, M. Bina, A. Keller, M. G. A. Paris, and S. Felicetti, Critical quantum metrology with a finite-component quantum phase transition, *Phys. Rev. Lett.* **124**, 120504 (2020).
- [35] H.-J. Zhu, K. Xu, G.-F. Zhang, and W.-M. Liu, Finite-component multicriticality at the superradiant quantum phase transition, *Phys. Rev. Lett.* **125**, 050402 (2020).
- [36] X. Chen, Z. Wu, M. Jiang, X.-Y. Lü, X. Peng, and J.-F. Du, Experimental quantum simulation of superradiant phase transition beyond no-go theorem via antisqueezing, *Nature communications* **12**, 6281 (2021).
- [37] M.-L. Cai, Z.-D. Liu, W.-D. Zhao, Y.-K. Wu, Q.-X. Mei, Y. Jiang, L. He, X. Zhang, Z.-C. Zhou, and L.-M. Duan, Observation of a quantum phase transition in the quantum Rabi model with a single trapped ion, *Nature communications* **12**, 1126 (2021).
- [38] Y.-Y. Zhang, Z.-X. Hu, L. Fu, H.-G. Luo, H. Pu, and X.-F. Zhang, Quantum phases in a quantum Rabi triangle, *Phys. Rev. Lett.* **127**, 063602 (2021).
- [39] D. Fallas Padilla, H. Pu, G.-J. Cheng, and Y.-Y. Zhang, Understanding the quantum Rabi ring using analogies to quantum magnetism, *Phys. Rev. Lett.* **129**, 183602 (2022).
- [40] M. Aspelmeyer, T. J. Kippenberg, and F. Marquardt, Cavity optomechanics, *Rev. Mod. Phys.* **86**, 1391 (2014).
- [41] S. Barzanjeh, A. Xuereb, S. Gröblacher, M. Paternostro, C. A. Regal, and E. M. Weig, Optomechanics for quantum technologies, *Nature Physics* **18**, 15 (2022).
- [42] T. T. Heikkilä, F. Massel, J. Tuorila, R. Khan, and M. A. Sillanpää, Enhancing optomechanical coupling via the Josephson effect, *Phys. Rev. Lett.* **112**, 203603 (2014).
- [43] G. Via, G. Kirchmair, and O. Romero-Isart, Strong single-photon coupling in superconducting quantum magnetomechanics, *Phys. Rev. Lett.* **114**, 143602 (2015).
- [44] X.-Y. Lü, Y. Wu, J. R. Johansson, H. Jing, J. Zhang, and F. Nori, Squeezed optomechanics with phase-matched amplification and dissipation, *Phys. Rev. Lett.* **114**, 093602 (2015).
- [45] O. Shevchuk, G. A. Steele, and Y. M. Blanter, Strong and tunable couplings in flux-mediated optomechanics, *Phys. Rev. B* **96**, 014508 (2017).
- [46] L. Neumeier, T. E. Northup, and D. E. Chang, Reaching the optomechanical strong-coupling regime with a single atom in a cavity, *Phys. Rev. A* **97**, 063857 (2018).
- [47] M. Kounalakis, Y. M. Blanter, and G. A. Steele, Flux-mediated optomechanics with a transmon qubit in the single-photon ultrastrong-coupling regime, *Phys. Rev. Res.* **2**, 023335 (2020).
- [48] J. Manninen, M. T. Haque, D. Vitali, and P. Hakonen, Enhancement of the optomechanical coupling and Kerr nonlinearity using the Josephson capacitance of a cooper-pair box, *Phys. Rev. B* **105**, 144508 (2022).
- [49] J.-M. Pirkkalainen, S. Cho, F. Massel, J. Tuorila, T. Heikkilä, P. Hakonen, and M. Sillanpää, Cavity optomechanics mediated by a quantum two-level system, *Nature communications* **6**, 6981 (2015).
- [50] N. Carlon Zambon, Z. Denis, R. De Oliveira, S. Ravets, C. Ciuti, I. Favero, and J. Bloch, Enhanced cavity optomechanics with quantum-well exciton polaritons, *Phys. Rev. Lett.* **129**, 093603 (2022).
- [51] I. Rodrigues, D. Bothner, and G. Steele, Coupling microwave photons to a mechanical resonator using quantum interference, *Nature communications* **10**, 5359 (2019).
- [52] S. Bose, K. Jacobs, and P. L. Knight, Preparation of nonclassical states in cavities with a moving mirror, *Phys. Rev. A* **56**, 4175 (1997).
- [53] A. Nunnenkamp, K. Børkje, and S. M. Girvin, Single-

- photon optomechanics, Phys. Rev. Lett. **107**, 063602 (2011).
- [54] P. Rabl, Photon blockade effect in optomechanical systems, Phys. Rev. Lett. **107**, 063601 (2011).
- [55] J.-Q. Liao and F. Nori, Photon blockade in quadratically coupled optomechanical systems, Phys. Rev. A **88**, 023853 (2013).
- [56] J. Qian, A. A. Clerk, K. Hammerer, and F. Marquardt, Quantum signatures of the optomechanical instability, Phys. Rev. Lett. **109**, 253601 (2012).
- [57] R. Riedinger, S. Hong, R. A. Norte, J. A. Slater, J. Shang, A. G. Krause, V. Anant, M. Aspelmeyer, and S. Gröblacher, Non-classical correlations between single photons and phonons from a mechanical oscillator, Nature **530**, 313 (2016).
- [58] V. Macrì, A. Ridolfo, O. Di Stefano, A. F. Kockum, F. Nori, and S. Savasta, Nonperturbative dynamical Casimir effect in optomechanical systems: Vacuum Casimir-Rabi splittings, Phys. Rev. X **8**, 011031 (2018).
- [59] B. Wang, J.-M. Hu, V. Macrì, Z.-L. Xiang, and F. Nori, Coherent resonant coupling between atoms and a mechanical oscillator mediated by cavity-vacuum fluctuations, Phys. Rev. Res. **5**, 013075 (2023).
- [60] O. Di Stefano, A. Settineri, V. Macrì, A. Ridolfo, R. Stassi, A. F. Kockum, S. Savasta, and F. Nori, Interaction of mechanical oscillators mediated by the exchange of virtual photon pairs, Phys. Rev. Lett. **122**, 030402 (2019).
- [61] J. R. Johansson, G. Johansson, C. M. Wilson, and F. Nori, Dynamical Casimir effect in a superconducting coplanar waveguide, Phys. Rev. Lett. **103**, 147003 (2009).
- [62] P. D. Nation, J. R. Johansson, M. P. Blencowe, and F. Nori, Colloquium: Stimulating uncertainty: Amplifying the quantum vacuum with superconducting circuits, Rev. Mod. Phys. **84**, 1 (2012).
- [63] C. K. Law, Interaction between a moving mirror and radiation pressure: A Hamiltonian formulation, Phys. Rev. A **51**, 2537 (1995).
- [64] See supplemental material.
- [65] J. Goldstone, A. Salam, and S. Weinberg, Broken symmetries, Phys. Rev. **127**, 965 (1962).
- [66] X.-G. Wen, *Quantum field theory of many-body systems: from the origin of sound to an origin of light and electrons* (Oxford University Press, New York, 2004).
- [67] S. Felicetti and A. Le Boité, Universal spectral features of ultrastrongly coupled systems, Phys. Rev. Lett. **124**, 040404 (2020).
- [68] We consider the resonant relationship between the cavity frequency and the atomic transition frequency. Moreover, using Hamiltonian H_{op} in Eq. (10) instead of H to describe the optomechanical system can also obtain the same conclusions.
- [69] It naturally arises by applying minimal coupling $\mathbf{p} \rightarrow \mathbf{p} - q\mathbf{A}/c$ to the electron kinetic energy $\mathbf{p}^2/(2m)$ [70, 75–77].
- [70] K. Rzażewski, K. Wódkiewicz, and W. Żakowicz, Phase transitions, two-level atoms, and the A^2 term, Phys. Rev. Lett. **35**, 432 (1975).
- [71] S. Savasta, O. Di Stefano, and F. Nori, Thomas–Reiche–Kuhn (TRK) sum rule for interacting photons, Nanophotonics **10**, 465 (2020).
- [72] T. Senthil, A. Vishwanath, L. Balents, S. Sachdev, and M. P. Fisher, Deconfined quantum critical points, Science **303**, 1490 (2004).
- [73] T. Senthil, L. Balents, S. Sachdev, A. Vishwanath, and M. P. A. Fisher, Quantum criticality beyond the Landau-Ginzburg-Wilson paradigm, Phys. Rev. B **70**, 144407 (2004).
- [74] C. Wang, A. Nahum, M. A. Metlitski, C. Xu, and T. Senthil, Deconfined quantum critical points: Symmetries and dualities, Phys. Rev. X **7**, 031051 (2017).
- [75] O. Di Stefano, A. Settineri, V. Macrì, L. Garziano, R. Stassi, S. Savasta, and F. Nori, Resolution of gauge ambiguities in ultrastrong-coupling cavity quantum electrodynamics, Nature Physics **15**, 803 (2019).
- [76] L. Garziano, A. Settineri, O. Di Stefano, S. Savasta, and F. Nori, Gauge invariance of the Dicke and Hopfield models, Phys. Rev. A **102**, 023718 (2020).
- [77] A. Settineri, O. Di Stefano, D. Zueco, S. Hughes, S. Savasta, and F. Nori, Gauge freedom, quantum measurements, and time-dependent interactions in cavity QED, Phys. Rev. Res. **3**, 023079 (2021).