



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Topological photonics

Tomoki Ozawa, Hannah M. Price, Alberto Amo, Nathan Goldman, Mohammad Hafezi, Ling Lu, Mikael C. Rechtsman, David Schuster, Jonathan Simon, Oded Zilberberg, and Iacopo Carusotto

Rev. Mod. Phys. **91**, 015006 — Published 25 March 2019

DOI: [10.1103/RevModPhys.91.015006](https://doi.org/10.1103/RevModPhys.91.015006)

Topological Photonics

Tomoki Ozawa

*Interdisciplinary Theoretical and Mathematical Sciences Program (iTHEMS),
RIKEN, Wako, Saitama 351-0198,
Japan
Center for Nonlinear Phenomena and Complex Systems,
Université Libre de Bruxelles,
CP 231, Campus Plaine,
B-1050 Brussels,
Belgium
INO-CNR BEC Center and Dipartimento di Fisica,
Università di Trento, I-38123 Povo,
Italy*

Hannah M. Price

*School of Physics and Astronomy,
University of Birmingham,
Edgbaston, Birmingham B15 2TT,
United Kingdom
INO-CNR BEC Center and Dipartimento di Fisica,
Università di Trento, I-38123 Povo,
Italy*

Alberto Amo

*Université de Lille, CNRS,
UMR 8523 - PhLAM - Laboratoire de Physique des Lasers Atomes et Molécules, F-59000 Lille,
France*

Nathan Goldman

*Center for Nonlinear Phenomena and Complex Systems,
Université Libre de Bruxelles,
CP 231, Campus Plaine,
B-1050 Brussels,
Belgium*

Mohammad Hafezi

*Joint Quantum Institute,
NIST/University of Maryland,
College Park, MD 20742,
USA
Department of Electrical and Computer Engineering and Institute for Research in Electronics and Applied Physics,
University of Maryland,
College Park, MD 20742,
USA*

Ling Lu

*Institute of Physics,
Chinese Academy of Sciences/Beijing National Laboratory for Condensed Matter Physics, Beijing 100190,
China*

Mikael C. Rechtsman

*Department of Physics,
The Pennsylvania State University,
University Park, PA 16802,
USA*

David Schuster

*The James Franck Institute and Department of Physics,
University of Chicago,*

Chicago, Illinois 60637,
USA

Jonathan Simon

*The James Franck Institute and Department of Physics,
University of Chicago,
Chicago, Illinois 60637,
USA*

Oded Zilberberg

*Institute for Theoretical Physics,
ETH Zurich, 8093 Zurich,
Switzerland*

Iacopo Carusotto

*INO-CNR BEC Center and Dipartimento di Fisica,
Università di Trento, I-38123 Povo,
Italy*

(Dated: October 4, 2018)

Topological photonics is a rapidly-emerging field of research in which geometrical and topological ideas are exploited to design and control the behavior of light. Drawing inspiration from the discovery of the quantum Hall effects and topological insulators in condensed matter, recent advances have shown how to engineer analogous effects also for photons, leading to remarkable phenomena such as the robust unidirectional propagation of light, which hold great promise for applications. Thanks to the flexibility and diversity of photonics systems, this field is also opening up new opportunities to realise exotic topological models and to probe and exploit topological effects in new ways. In this article, we review experimental and theoretical developments in topological photonics across a wide-range of experimental platforms, including photonic crystals, waveguides, metamaterials, cavities, optomechanics, silicon photonics and circuit-QED. We discuss how changing the dimensionality and symmetries of photonics systems has allowed for the realization of different topological phases, and we review progress in understanding the interplay of topology with non-Hermitian effects, such as dissipation. As an exciting perspective, topological photonics can be combined with optical nonlinearities, leading towards new collective phenomena and novel strongly-correlated states of light, such as an analogue of the fractional quantum Hall effect.

CONTENTS

I. Introduction	3	2. Topological RF Circuits	36
II. Basic concepts	5	3. Twisted Optical Resonators	38
A. Topological phases of matter	5	4. Intrinsic spin-orbit interactions for light	39
1. Integer quantum Hall effect	6	5. Bianisotropic Metamaterials	40
2. Quantum spin-Hall system	10	6. Photonic structures with crystalline symmetries	42
3. Topological phases in other dimensions	11	7. Other proposals	43
4. Topological Pumping	13	C. Anomalous Floquet topological insulators	43
5. Floquet engineering: Topology through time-periodic modulations	16	D. Topology in gapless photonic systems	45
B. Features of photonic systems	17	IV. Topological photonics in one dimension	49
1. Bosonic nature	18	A. Topology in 1D chiral systems	49
2. Non-equilibrium nature	18	B. Topological pumps	50
3. Basics of nonlinear optics	22	V. Topological photonics in higher dimensions	52
III. Topological photonics in two dimensions	25	A. Three-dimensional gapless phases	52
A. Analogue quantum Hall systems in photonics	25	1. Weyl points and helicoid surface states	52
1. Gyro-magnetic photonic crystals	25	2. Multi-Weyl and Dirac points	53
2. Propagating geometries	27	3. Nodal lines and surface	53
3. Optomechanics	30	B. Three-dimensional gapped phases	53
4. Cavity- and Circuit-QED systems	30	1. 3D Chern Insulator	53
5. Other proposals	31	2. One-way fibers	54
B. Analogue quantum spin Hall systems in photonics	32	3. Single surface Dirac cone	54
1. Silicon ring resonator arrays	33	4. Non-magnetic designs	54
		C. Towards even higher dimensions	55
		1. Synthetic dimensions	55
		2. Four-dimensional quantum Hall effect	56

VI. Gain and loss in topological photonics	56
A. Non-Hermitian topological photonics	56
B. Emergent topology of Bogoliubov modes	57
VII. Topological effects for interacting photons	58
A. Weak nonlinearities	59
B. Strong nonlinearities	60
VIII. Conclusion and perspectives	62
A. Optical isolation and robust transport	62
B. Quantum emitters and topological laser	63
1. Topological lasers: theory	63
2. Topological lasers: experiments	63
C. Measurement of bulk topological and geometrical properties	64
D. Topological quantum computing	65
Acknowledgments	65
References	66

I. INTRODUCTION

Over the last decade, *topological photonics* appeared as a rapidly growing field of study. This field aims to explore the physics of topological phases of matter, originally discovered in solid-state electron systems, in a novel optical context. In this review, we attempt to cover the main achievements of topological photonics, beginning from the basic concepts of topological phases of matter and of photonics, so that readers can follow our discussion independently of their background.

Topological photonics is rooted in ideas that were first developed to understand topological phases of matter in solid state physics. This field of research began with the discovery of the integer quantum Hall effect in 1980 (Klitzing *et al.*, 1980; von Klitzing, 1986). In this effect, a two-dimensional electron gas in the presence of a strong perpendicular magnetic field was found to exhibit robust plateaus in the Hall conductance as a function of the magnetic field, at values equal to integer multiples of the fundamental constant e^2/h . The far-reaching conceptual consequences of this integer quantum Hall effect were soon highlighted in (Kohmoto, 1985; Thouless *et al.*, 1982). These works related the integer appearing in the Hall conductance to a *topological invariant* of the system, the Chern number, that is an integer-valued quantity which describes the global structure of the wavefunction in momentum space over the Brillouin zone.

An important insight into the physical meaning of the topological invariant is given by the *bulk-edge correspondence* (Hatsugai, 1993a,b; Jackiw and Rebbi, 1976; Qi *et al.*, 2006): when two materials with different topological invariants are put in contact, there must exist edge states that are spatially localized at the interface at energies that lie within the energy gap of the surrounding bulk materials.

The bulk-edge correspondence can be heuristically understood in the following way: an integer topological in-

variant of a gapped system cannot change its value under perturbations or deformations of the system, unless the energy gap to excited states is somewhere closed. This implies that when two materials with different topological invariants are put in contact, the energy gap must close somewhere in the interface region, which leads to the appearance in this region of localized states. In a finite-size sample of a topologically non-trivial material, the physical edge of the sample can be considered as an interface between a region with a non-zero topological invariant and the topologically trivial vacuum, guaranteeing the existence of localized states at the system boundary.

In the quantum Hall effect, these edge modes display *chiral* properties, in the sense that they can only propagate in one direction along the sample boundary but not in the opposite direction. The number of such chiral edge modes that are available at the Fermi energy for electric conduction is proportional to the Hall conductance. Because of the unidirectional nature of the edge states, the edge currents are immune to backscattering, resulting in the precise and robust quantization of the measured Hall conductance (Büttiker, 1988; Halperin, 1982; MacDonald and Středa, 1984).

Interest in the topological physics of electronic systems surged further when a different class of topological phases of matter, now known as the quantum spin-Hall systems or \mathbb{Z}_2 topological insulators, was discovered in 2005 (Bernevig *et al.*, 2006; Bernevig and Zhang, 2006; Kane and Mele, 2005a,b; König *et al.*, 2007). In these systems, the Chern number is zero but the wavefunction is characterised by a binary (\mathbb{Z}_2) topological invariant, that can be non-zero and robust in the presence of time-reversal symmetry. Since then, there has been intense investigation in condensed matter physics into what different topological phases of matter are possible under various symmetries, and what are the physical consequences of this physics (Bernevig and Hughes, 2013; Chiu *et al.*, 2016; Hasan and Kane, 2010; Qi and Zhang, 2011). Besides electronic systems in solid-state materials, topological phases of matter are also being actively studied in other quantum many-body systems, in particular liquid helium (Volovik, 2009) and ultracold atomic gases (Cooper, 2008; Dalibard *et al.*, 2011; Goldman *et al.*, 2016a, 2014).

Parallel to the growth in the study of topological phases of matter in condensed matter systems, Haldane and Raghunathan made the crucial observation that topological band structures are, in fact, a ubiquitous property of waves inside a periodic medium, regardless of the classical or quantum nature of the waves. In their seminal works (Haldane and Raghunathan, 2008; Raghunathan and Haldane, 2008), they considered electromagnetic waves in two-dimensional spatially periodic devices embedding time-reversal-breaking magneto-optical elements, and showed that the resulting photonic bands would have nontrivial




FIG. 1 (Color online) Panel (a1): sketch of the gyro-magnetic photonic crystal slab used in the experiments of (Wang *et al.*, 2009). The blue dots indicate the ferrite rods which are organized in a two-dimensional square lattice along the xy plane and are subject to a magnetic field of 0.2 T. The structure is sandwiched between two parallel copper plates providing confinement along z . The chiral edge state is located at the boundary of the photonic crystal next to the metal wall. Two dipole antennas A and B serve as feeds and/or probes. Back-scattering is investigated by inserting a variable-length metal obstacle between the antennas. Panel (a2) shows a top view photograph of the actual waveguide with the top plate removed. Panels (b): Theoretical calculation of light propagation on edge states: subpanels (b1,b3) show unidirectional, non-reciprocal propagation from the antennas A and B, respectively. Subpanel (b2) shows the immunity to backscattering against a defect. Panel (c): reciprocal transmission when the two antennas are located in the bulk. Panel (d): non-reciprocal transmission via the chiral edge state. Blue and red curves refer to transmission from antenna A to antenna B and viceversa. Panel (e): projected dispersion of the allowed photonic bands in the bulk (blue and gray) and of the chiral edge state (red). The white numbers indicate the Chern number of each band. All panels adapted from (Wang *et al.*, 2009).

topological invariants. Consequently, they predicted that such photonic systems would support robust chiral states propagating along the edge of the system at frequencies inside the photonic band gap.

Shortly afterwards, following a realistic proposal of Ref. (Wang *et al.*, 2008), the idea of Haldane and Raghu was experimentally implemented using the two-dimensional magneto-optical photonic crystal structure in the microwave domain sketched in Fig. 1(a1-a2) (Wang *et al.*, 2009): a clear signature of the non-trivial band

topology was indeed found in the unidirectionally propagating edge states and in the corresponding non-reciprocal behaviour, as illustrated in the simulations of Fig. 1(b) and in the experimental data of Fig. 1(c,d). More details on this and related following experiments are given in Sec.III.A.

Further progress towards the implementation of such a model in the optical domain and the exploration of other topological models remained however elusive. One major challenge was the absence of large magneto-optical response in the optical domain. One way to overcome this difficulty is to consider internal degrees of freedom of photons as pseudospins and look for an analogy of quantum spin-Hall systems, namely where the overall time-reversal symmetry is not broken but each pseudo-spin feels an artificial magnetic field (Hafezi *et al.*, 2011; Khanikaev *et al.*, 2013; Umucalilar and Carusotto, 2011). A second way is to use ideas from the Floquet topological insulators (Kitagawa *et al.*, 2010a; Lindner *et al.*, 2011; Oka and Aoki, 2009) known in condensed matter physics, where temporal modulation is applied to the system to generate an effective time-independent Hamiltonian which breaks time-reversal symmetry (Fang *et al.*, 2012b). A third way is to employ time-dependent modulation to implement a “topological pump” (Thouless, 1983); this last approach was realized experimentally in photonics in 2012 (Kraus *et al.*, 2012), while the previous two ideas have been realized in 2013 by two concurrent experiments (Hafezi *et al.*, 2013b; Rechtsman *et al.*, 2013b).

Since then, there has been great activity in the study of a variety of photonic systems realizing band structures with non-trivial topological invariants, leading to the emerging research field of *topological photonics* (Khanikaev and Shvets, 2017; Lu *et al.*, 2014, 2016c; Sun *et al.*, 2017b). Along similar lines, intense theoretical and experimental work has also been devoted to related topological effects in other areas of classical physics, such as in mechanical and acoustic systems. Reviews of the advances of these other fields can be found in (Fleury *et al.*, 2015; Huber, 2016).

The present review is focused on the recent developments in the study of topological phases of matter in the photonics context. As we shall see in the following, in the last decade, topological ideas have successfully permeated the field of photonics, having been applied to a wide range of different material platforms, arranged in lattices of various dimensionalities, and operating in different regions of the electromagnetic spectrum, from radio- and micro-waves up to visible light. One long term goal of topological photonics is to achieve and control strongly correlated states of photons with topological features such as fractional quantum Hall states. In addition to opening up perspectives for exploring the fundamental physics of topological phases of matter beyond solid-state systems, topological photonics also offers rich potential applications of these concepts to a novel gener-

ation of optoelectronic devices, such as optical isolators and topological lasers.

The structure of this review article is the following. In Sec. II.A, we offer a general review of the main geometrical and topological concepts that have been developed in the study of solid-state electronic systems and that are commonly used in topological photonics. The following Sec. II.B gives a general overview of the specific features that characterize photonic systems in contrast to electronic topological insulators. The readers familiar with basic concepts of topological physics can safely skip Sec. II.A, and similarly those who are from photonic backgrounds can safely skip Sec. II.B.

In Sec. III, we discuss two dimensional photonic systems which show topological features. This section is divided into four subsections. In Sec. III.A, we discuss two dimensional photonic structures which break time-reversal symmetry, and hence display physics analogous to integer quantum Hall systems. Relevant photonic systems include gyro-magnetic photonic crystals, waveguide arrays, optomechanics, cavity-QED, and circuit-QED. The following Sec. III.B deals with two dimensional photonic systems which do not break time-reversal symmetry, and hence can be considered as analogs of the quantum spin-Hall systems. Systems discussed in this subsection include silicon ring resonator arrays, radio-frequency circuits, twisted optical resonators, and photonic metamaterials. In Sec. III.C, we review photonic realizations of anomalous Floquet topological systems with waveguide arrays, namely temporally modulated systems displaying topological features that do not have an analog in static Hamiltonians. Sec. III.D discusses two dimensional gapless systems such as honeycomb lattices, whose features can also be understood from topological considerations. Systems discussed in this subsection include microwave resonators, photorefractive crystals, coupled microlasers, exciton-polariton lattices, and waveguide arrays.

Sec. IV is devoted to photonic realizations of one-dimensional topological systems. In the first subsection Sec. IV.A we will concentrate on systems with chiral symmetry, such as the Su-Schrieffer-Heeger (SSH) model. In Sec. IV.B we will review photonic realizations of topological pumping. Systems of higher dimensionality will then be considered in Sec. V. Three dimensional gapless phases with features originating from topological charges in momentum space, such as Weyl points, are discussed in Sec. V.A. In Sec. V.B, we discuss gapped three dimensional phases and their topological interface states. The following Sec. V.C will present the concept of synthetic dimensions, which could be used to realize models with an effective spatial dimensionality higher than three, e.g., four dimensional quantum Hall systems.

In Sec. VI, we discuss photonic systems where gain and loss play an essential role. Such systems are described by non-Hermitian Hamiltonians and do not find a direct

counterpart in electronic topological insulators. The first subsection Sec. VI.A will focus on the interplay of gain and loss, while the following Sec. VI.B will review the emergent topology of Bogoliubov modes that arises from parametric down conversion processes.

Sec. VII is devoted to an overview of the interplay between topology and optical nonlinearities. Theoretical work on nonlinear effects stemming from weak nonlinearities will be reviewed in Sec. VII.A, while the following Sec. VII.B will highlight the prospect of strong photon-photon interactions mediated by strong nonlinearities to realize topologically nontrivial strongly correlated states of photons. Some of the future perspectives of the field of topological photonics are finally illustrated in Sec. VIII.

Compared to earlier reviews on topological photonics (Khanikaev and Shvets, 2017; Lu *et al.*, 2014, 2016c; Sun *et al.*, 2017b), this review aims to be comprehensive, starting from the very basics of topological phases of matter and trying to cover most of the works that have appeared in the last years in relation to topological phases of matter in optical systems in any dimensionality. However, we need to warn the readers that space restrictions force us to leave out many other fields of the optical sciences that relate to topological concepts, e.g. the rich dynamics of optical vortices in singular optics (Dennis *et al.*, 2009; Gbur, 2016), the topology underlying knots in complex electromagnetic fields (Arrayás *et al.*, 2017), and the topological ideas underlying bound states in the continuum (Hsu *et al.*, 2016). For all these advances, we refer the readers to the rich specific literature that is available on each of them.

II. BASIC CONCEPTS

In this section, we introduce general concepts of topological phases of matter and of optical and photonic systems that will be needed in the following sections. In Sec. II.A, we briefly review the paradigm of topological phases of matter, as it has been originally developed in the context of electronic systems in solid state materials, and illustrate the basic technical and mathematical tools to describe them. Then, in Sec. II.B, we review the principal features of photonic systems used for topological photonics, with a special emphasis on their differences and peculiarities as compared to electronic systems.

A. Topological phases of matter

According to Bloch's theorem, the eigenstates of a quantum particle in a periodic potential are organized into energy bands separated by energy gaps. This band structure determines the metallic/insulating nature of different solid-state materials (Ashcroft and Mermin, 1976). Besides the energy dispersion of the bands, the

geometrical structure of the Bloch eigenstates in momentum space can also have an impact on the electronic properties of materials, as first discovered by [Karplus and Luttinger, 1954](#) and [Adams and Blount, 1959](#). This geometrical structure is reflected also in integer-valued global topological invariants associated to each band, as we see below. In spite of their seemingly abstract nature ([Simon, 1983](#)), nontrivial values of these topological invariants have observable consequences such as the quantized bulk conductance in the quantum Hall effect and in the emergence of topologically protected edge states located on the physical boundary of the system ([Bernevig and Hughes, 2013](#); [Hasan and Kane, 2010](#); [Qi and Zhang, 2011](#); [Volovik, 2009](#)).

This subsection is devoted to an introduction to the basic concepts of such *topological phases of matter*. Readers familiar with (electronic) topological insulators and related fields can skip this part and move on to [Sec.II.B](#).

1. Integer quantum Hall effect

The quantum Hall effect is historically the first phenomenon where momentum-space topology was recognized to lead to observable physical phenomena. The integer quantum Hall effect was discovered in a two-dimensional electron gas subject to a strong perpendicular magnetic field by [Klitzing et al., 1980](#), who observed a robust quantization of the Hall conductance in units of e^2/h , where e is the charge of an electron and h is Planck's constant. Soon after, the extremely robust quantization of the Hall conductance was related to the topology of bands in momentum space by Thouless, Kohmoto, Nightingale, and den Nijs (TKNN) in ([Thouless et al., 1982](#)).

In order to review this landmark result, we first need to introduce the basic geometrical and topological properties of eigenstates in momentum space, such as the (local) Berry connection and Berry curvature and the (global) Chern number, respectively. We consider a single-particle Hamiltonian $\hat{H}(\hat{\mathbf{r}}, \hat{\mathbf{p}})$ in generic dimension d , where $\hat{\mathbf{r}}$ and $\hat{\mathbf{p}}$ are, respectively, the position and momentum operators. We assume that the Hamiltonian obeys the spatial periodicity condition $\hat{H}(\hat{\mathbf{r}} + \mathbf{a}_i, \hat{\mathbf{p}}) = \hat{H}(\hat{\mathbf{r}}, \hat{\mathbf{p}})$, where $\{\mathbf{a}_i\}$ are a set of d lattice vectors. Thanks to the spatial periodicity, one can invoke Bloch's theorem to write the eigenstates as

$$\psi_{n,\mathbf{k}}(\mathbf{r}) = e^{i\mathbf{k}\cdot\mathbf{r}} u_{n,\mathbf{k}}(\mathbf{r}), \quad (1)$$

where n is the band index and \mathbf{k} is the crystal momentum defined within the first Brillouin zone. The Bloch state $u_{n,\mathbf{k}}(\mathbf{r})$ obeys the same periodicity as the original Hamiltonian $u_{n,\mathbf{k}}(\mathbf{r} + \mathbf{a}_i) = u_{n,\mathbf{k}}(\mathbf{r})$ and is an eigenstate of the Bloch Hamiltonian

$$\hat{H}_{\mathbf{k}} \equiv e^{-i\mathbf{k}\cdot\hat{\mathbf{r}}} \hat{H}(\hat{\mathbf{r}}, \hat{\mathbf{p}}) e^{i\mathbf{k}\cdot\hat{\mathbf{r}}}, \quad (2)$$

namely

$$\hat{H}_{\mathbf{k}} u_{n,\mathbf{k}}(\mathbf{r}) = E_n(\mathbf{k}) u_{n,\mathbf{k}}(\mathbf{r}), \quad (3)$$

where $E_n(\mathbf{k})$ is the energy dispersion of the n -th band ([Ashcroft and Mermin, 1976](#)).

The physics of an energy band is captured in part by its dispersion relation $E_n(\mathbf{k})$, but also by the *geometrical* properties of how its eigenstates $u_{n,\mathbf{k}}(\mathbf{r})$ vary as a function of \mathbf{k} ([Adams and Blount, 1959](#); [Karplus and Luttinger, 1954](#); [Nagaosa et al., 2010](#); [Resta, 1994, 2011](#)). This geometry of the eigenstates is encoded by the Berry phase ([Berry, 1984](#); [Hannay, 1985](#); [Pancharatnam, 1956](#)), which is defined in the following. Whereas the Berry phase can be defined for a very general parameter space, in our discussion of topological phases of matter we restrict ourselves to the case where the parameters are the crystal momentum $\mathbf{k} = (k_x, k_y, k_z)$ varying over the first Brillouin zone. Then, if one prepares a localized wavepacket from states in band n and makes it adiabatically move along a closed path in momentum space, it will acquire a dynamical phase, determined by the time-integral of its \mathbf{k} -dependent energy, but also a Berry phase ([Xiao et al., 2010](#))

$$\gamma = \oint \mathcal{A}_n(\mathbf{k}) \cdot d\mathbf{k}, \quad (4)$$

that is geometrically determined by an integral, over the same momentum-space path, of the Berry connection, defined as

$$\mathcal{A}_n(\mathbf{k}) \equiv i \langle u_{n,\mathbf{k}} | \nabla_{\mathbf{k}} | u_{n,\mathbf{k}} \rangle. \quad (5)$$

Note that the definition of the Bloch states via [Eq. \(3\)](#) does not specify the overall phase of $|u_{n,\mathbf{k}}\rangle$, so one can freely choose the phase of the Bloch states. Under a gauge transformation $|u_{n,\mathbf{k}}\rangle \rightarrow e^{i\chi(\mathbf{k})} |u_{n,\mathbf{k}}\rangle$, the Berry connection is not gauge invariant and transforms as $\mathcal{A}_n(\mathbf{k}) \rightarrow \mathcal{A}_n(\mathbf{k}) - \nabla_{\mathbf{k}}\chi(\mathbf{k})$. However, the single-valuedness of $e^{i\chi(\mathbf{k})}$ at the beginning and the end of the path imposes that the Berry phase [\(4\)](#) for a given closed path is gauge invariant modulo 2π . Additionally, from the gauge-dependent Berry connection $\mathcal{A}_n(\mathbf{k})$ one can construct a gauge invariant Berry curvature, which in three dimensions takes the following form:

$$\Omega_n(\mathbf{k}) = \nabla_{\mathbf{k}} \times \mathcal{A}_n(\mathbf{k}), \quad (6)$$

and which encodes the geometrical properties of the n -th band. In two dimensions, the Berry curvature has only one component:

$$\Omega_n(\mathbf{k}) = i (\langle \partial_{k_x} u_{n,\mathbf{k}} | \partial_{k_y} u_{n,\mathbf{k}} \rangle - \langle \partial_{k_y} u_{n,\mathbf{k}} | \partial_{k_x} u_{n,\mathbf{k}} \rangle). \quad (7)$$

Importantly, although the Berry curvature is a gauge invariant quantity that is continuously defined over the whole Brillouin zone, the phase of the Bloch states




FIG. 2 A schematic illustration of how the Brillouin zone is divided into two parts, S and S' . Thanks to the periodicity of the quasi-momentum, the two-dimensional Brillouin zone has a torus-like structure, in which top-bottom and left-right edges (purple and red) should be identified.

themselves cannot always be chosen to be continuous. Whether this is possible or not depends on the value of a topological invariant of the band, the Chern number, defined as the integral

$$C_n = \frac{1}{2\pi} \int_{\text{BZ}} d^2k \Omega_n(k_x, k_y), \quad (8)$$

over the whole first Brillouin zone. If one can define the phase of the Bloch state, and hence the Berry connection $\mathcal{A}_n(\mathbf{k})$, continuously over the whole Brillouin zone, a direct consequence of the definition $\Omega_n(\mathbf{k}) = \nabla_{\mathbf{k}} \times \mathcal{A}_n(\mathbf{k})$ and of Stokes' theorem is that the Chern number must necessarily be zero. Conversely, having a non-zero Chern number, implies that the Bloch state and hence the Berry connection $\mathcal{A}_n(\mathbf{k})$ cannot be continuously defined.

It is not difficult to see that the Chern number always takes an integer value (Kohmoto, 1985). To this purpose, we divide the integration domain of (8) into two regions S and S' as sketched in Fig. 2. Within S , we choose a continuous gauge for the Bloch state, giving the Berry connection $\mathcal{A}_n(\mathbf{k})$. Similarly, within S' , we choose a continuous gauge, which yields the Berry connection $\mathcal{A}'_n(\mathbf{k})$. Keeping in mind that the first Brillouin zone can be thought of as a torus, thanks to the periodicity of the quasi-momentum, we can use Stokes' theorem within S and S' and rewrite the Chern number in terms of the line integral along the common boundary $\partial S = -\partial S'$,

$$\begin{aligned} C_n &= \frac{1}{2\pi} \int_S d^2k \Omega_n(k_x, k_y) + \frac{1}{2\pi} \int_{S'} d^2k \Omega_n(k_x, k_y) \\ &= \frac{1}{2\pi} \oint_{\partial S} d\mathbf{r} \cdot \mathcal{A}_n(\mathbf{k}) - \frac{1}{2\pi} \oint_{\partial S} d\mathbf{r} \cdot \mathcal{A}'_n(\mathbf{k}) \\ &= \frac{1}{2\pi} (\gamma - \gamma'), \end{aligned} \quad (9)$$

where γ and γ' are the Berry phases along the contour ∂S calculated using $\mathcal{A}_n(\mathbf{k})$ and $\mathcal{A}'_n(\mathbf{k})$, respectively. As the Berry phases are calculated along the same path, they must be equal up to multiples of 2π . This in turn implies that the Chern number C_n must be an integer. Importantly, this integer-valued quantity has a profound topological origin (Avron *et al.*, 1983; Niu *et al.*, 1985; Simon, 1983), which indicates that its value must remain strictly constant under smooth perturbations that preserve the band-gaps separating the band n to neighboring

bands (Avron *et al.*, 1983). Fermionic systems in which the fermions completely fill Bloch bands with non-zero Chern numbers are generically termed *Chern insulators*.

Within linear response theory and ignoring interparticle interactions, one can show that the Hall conductance σ_{xy} of a two-dimensional insulator is (Thouless *et al.*, 1982),

$$\sigma_{xy} = -\frac{e^2}{h} \sum_n C_n, \quad (10)$$

where the Chern numbers are summed over the n occupied bands. Since the Chern numbers can take only integers, it follows that the Hall conductance is quantized in units of e^2/h . As we shall see shortly, in the simplest case of a uniform two dimensional electron gas under a strong magnetic field, the energy levels form flat Landau levels, and all the Landau levels have the same Chern number. Therefore the Hall conductance of the integer quantum Hall effect is proportional to the number of occupied Landau levels.

The quantization of the Hall conductance can also be related to the number of modes that propagate unidirectionally around the system, the so-called chiral edge modes. Indeed, each of such edge modes contributes $-e^2/h$ to the measured Hall conductance (Büttiker, 1988; Halperin, 1982; MacDonald and Středa, 1984). The existence of such current-carrying edge modes is also constrained by topology, in the sense that the sum of the Chern numbers associated with the occupied bulk bands is equal to the number of edge modes contributing to the edge current (Hatsugai, 1993a,b; Qi *et al.*, 2006). This relationship between a bulk topological invariant, such as the Chern number, and the number of localized edge modes is an example of the bulk-edge correspondence, i.e., a matching between the topological properties defined in the bulk of a material with its boundary phenomena (Bernevig and Hughes, 2013).

While the discussions above are based on single-particle energy bands in a perfect crystal, the definition of the Chern number can also be generalised to include the effects of interactions and disorder (Niu *et al.*, 1985). When the interparticle interactions become very strong, the Hall conductance can become quantized at *fractional values of e^2/h* (Tsui *et al.*, 1982). This is known as the “fractional quantum Hall effect” in which the quantum many-body ground state is strongly-correlated and topological. Remarkably, the excitations of such a fractional quantum Hall state can have a fractional charge and possibly even fractional statistics (Arovas *et al.*, 1984; Laughlin, 1983). Progress towards realizing analogue fractional quantum Hall states of light will be reviewed in Sec VII.B.

In the rest of this subsection, we proceed with a detailed discussion of a few important models for integer quantum Hall systems and Chern insulators. We shall

start by considering a two-dimensional electron gas under a strong and uniform magnetic field, which gives rise to Landau levels and to the integer quantum Hall effect (Prange *et al.*, 1989; Yoshioka, 2002). The second example is the Harper-Hofstadter model, that is a tight-binding lattice model in a uniform magnetic field (Azbel, 1964; Harper, 1955b; Hofstadter, 1976). The third one is the Haldane model (Haldane, 1988), which is the first example of a Chern insulator model with alternating magnetic flux patterns. We will then conclude by illustrating the bulk-edge correspondence on a simple Jackiw-Rabbi model (Jackiw and Rebbi, 1976).

Landau levels: The quantum Hall effect was originally found in a semiconductor hetero-junction where electrons are confined to move in a two-dimensional plane (Klitzing *et al.*, 1980). This system can be modeled, to a first approximation, as a two-dimensional electron gas in free space under a constant magnetic field. The single-particle Hamiltonian is

$$\hat{H} = \frac{(\hat{p}_x - eA_x(\hat{\mathbf{r}}))^2 + (\hat{p}_y - eA_y(\hat{\mathbf{r}}))^2}{2m}, \quad (11)$$

where $\mathbf{A}(\mathbf{r}) = (A_x(\mathbf{r}), A_y(\mathbf{r}), 0)$ is the magnetic vector potential, associated with the magnetic field $\mathbf{B} = \nabla_{\mathbf{r}} \times \mathbf{A}(\mathbf{r})$.

For a given magnetic field, different forms of the vector potential $\mathbf{A}(\mathbf{r})$ can be chosen. For our case of a constant magnetic field along the z -direction, $\mathbf{B} = (0, 0, B)$, the two most common choices are the Landau gauge, which keeps translational symmetry along one direction as $\mathbf{A}(\mathbf{r}) = (-yB, 0, 0)$ or $(0, xB, 0)$, and the symmetric gauge, $\mathbf{A}(\mathbf{r}) = (-yB/2, xB/2, 0)$, which keeps instead rotational symmetry. Physical observables such as the energy spectrum and the Hall conductance do not depend on the choice of gauge.

The single-particle energy spectrum of this system consists of equally spaced *Landau levels* of energy

$$E_n = \hbar\omega_c (n + 1/2), \quad (12)$$

where $\omega_c \equiv |e|B/m$ is the cyclotron frequency and the integer $n \geq 0$. For a large system, each Landau level is highly degenerate with a degeneracy equal to the number of unit magnetic fluxes $\phi_0 = h/|e|$ piercing the system.

Regarding each Landau level as an energy band, the system can be considered as a Chern insulator when the Fermi energy lies within an energy gap. The Chern number is one for any Landau level. Then, from the TKNN formula (10), the Hall conductance is thus proportional to the number of occupied Landau levels, which explains the basic phenomenology of the integer quantum Hall effect.

Harper-Hofstadter model: The next model we consider is the discrete lattice version of the Landau level problem, the Harper-Hofstadter model (Azbel, 1964; Harper, 1955b; Hofstadter, 1976). In tight-binding models, the




FIG. 3 (Left) Energy spectrum of the Harper-Hofstadter model, which is called the Hofstadter butterfly (Hofstadter, 1976). (Right) The colored Hofstadter butterfly in which the color of each band-gap indicates the topological invariant of the gap, given by the sum of the Chern numbers of all bands below. Warm colors indicate a positive topological invariant, whereas the cool colors indicate a negative topological invariant. The horizontal axes are the energies and the vertical axes are the flux α .

magnetic vector potential $\mathbf{A}(\mathbf{r})$ enters as a non-trivial phase of the hopping amplitude between neighboring sites, called the *Peierls phase*. In the simplest cases, the phase accumulated when hopping from a site at \mathbf{r}_1 to a site at \mathbf{r}_2 can be written in terms of the vector potential as

$$\Phi_{\mathbf{r}_1 \rightarrow \mathbf{r}_2} = \frac{e}{\hbar} \int_{\mathbf{r}_1}^{\mathbf{r}_2} \mathbf{A}(\mathbf{r}) \cdot d\mathbf{r}, \quad (13)$$

where the integral is taken along a straight line connecting the two points (Luttinger, 1951; Peierls, 1933).

Choosing for definiteness the Landau gauge along the y -direction, $\mathbf{A}(\mathbf{r}) = (0, Bx, 0)$, the Hamiltonian of the Harper-Hofstadter model on a square lattice is:

$$\hat{H} = -J \sum_{x,y} \left(\hat{a}_{x+a,y}^\dagger \hat{a}_{x,y} + e^{i2\pi\alpha x/a} \hat{a}_{x,y+a}^\dagger \hat{a}_{x,y} + \text{H.c.} \right), \quad (14)$$

where $\hat{a}_{x,y}$ is the annihilation operator of a particle at site (x, y) , J is the magnitude of the (isotropic) hopping amplitude and a is the lattice spacing. The intensity of the magnetic field in the system is quantified by the parameter α , obeying $\alpha\phi_0 = Ba^2$, which identifies the magnetic flux per plaquette of the lattice in units of the magnetic flux quantum ϕ_0 . The main distinction from the Landau level case discussed above is that in the Hofstadter model there are two competing lengthscales: the lattice spacing and the magnetic length. As a result, the electrons paths interfere to give the fractal energy spectrum as a function of α , which is widely known as the *Hofstadter butterfly* and which is plotted in the left panel of Fig. 3. The first experimental demonstration of the Hofstadter butterfly was performed in a microwave waveguide, exploiting the analogy between the transfer matrix governing the transmission of microwaves and the eigenvalue equation of the Harper-Hofstadter model (Kuhl and Stöckmann, 1998).

To get more insight into this spectrum, it is useful to concentrate on cases where α is a rational number,

$\alpha = p/q$ with p and q being co-prime integers. Because of the spatially-varying hopping phase, the Hamiltonian breaks the basic periodicity of the square lattice. Periodicity is, however, recovered if we consider a larger unit cell of $q \times 1$ plaquettes: this is called the *magnetic unit cell* (Dana *et al.*, 1985; Zak, 1964). As the number of bands in lattice models is equal to the number of lattice sites per (magnetic) unit cell, the Harper-Hofstadter model with $\alpha = p/q$ has q bands.

To find the geometrical and topological properties of the model, one can diagonalize the momentum-space Hamiltonian (Azbel, 1964; Harper, 1955b; Hofstadter, 1976)

$$\hat{H}_{\mathbf{k}} = -J \sum_{i=0}^{q-1} (\cos(k_y - 2\pi\alpha i) \hat{a}_i(\mathbf{k})^\dagger \hat{a}_i(\mathbf{k}) + e^{-ik_x} \hat{a}_{i+1}(\mathbf{k})^\dagger \hat{a}_i(\mathbf{k}) + \text{H.c.}), \quad (15)$$

where i , defined mod q , indicates the site within a magnetic unit cell, and the momentum \mathbf{k} is chosen within the magnetic Brillouin zone: $-\pi/q \leq k_x \leq \pi/q$ and $-\pi \leq k_y \leq \pi$. An explicit calculation shows that the Chern numbers of all isolated bands of the Harper-Hofstadter model are nonzero and can be found as solutions of a simple Diophantine equation (Thouless *et al.*, 1982). As shown in the right panel of Fig. 3, this model exhibits a very rich structure of positive and negative Chern numbers depending on the magnetic flux.

Haldane model: The Haldane model (Haldane, 1988) is the first model system that exhibits a non-zero quantized Hall conductance in a non-uniform magnetic field with a vanishing average flux per plaquette. This model demonstrated that, to obtain the quantum Hall effect, the essential feature required is, in fact, not a net magnetic field but the breaking of time-reversal symmetry. As the Haldane model consists of a honeycomb lattice with suitable hopping amplitudes, it is useful to start by briefly reviewing the physics of a tight-binding model on a honeycomb lattice, which is often used to describe electrons in graphene (Castro Neto *et al.*, 2009) and which has recently been widely implemented in photonics, as we will review in Sec. III.D.

The honeycomb lattice with nearest-neighbor hopping is one of the simplest examples of a system which exhibits *Dirac cones* in the band structure, namely linear crossings of the energy dispersion of two neighboring bands. A honeycomb lattice has two lattice sites per unit cell, which gives two bands. These are degenerate at two isolated and inequivalent points in the Brillouin zone, called Dirac points. Around the Dirac points, the effective Hamiltonian of the two bands takes the following form in the sublattice basis:

$$\hat{H}_D \approx \hbar v_D (q_x \sigma_x + q_y \sigma_y), \quad (16)$$

where $v_D \equiv 3t_1/2$ is called the Dirac velocity, with t_1 being the nearest-neighbor hopping amplitude, and (q_x, q_y)




FIG. 4 (a) A plaquette of the Haldane model. In addition to the usual nearest-neighbour hoppings, there are also complex next-nearest-neighbour hoppings. For the latter, hopping along the arrows carries a phase of ϕ , whereas the hopping opposite to the arrows carries the opposite phase of $-\phi$. (b) The phase diagram of the Haldane model as a function of the next-nearest-neighbor hopping phase ϕ and the sublattice energy difference $2M$. (c) Bulk band structure of the honeycomb lattice with nearest-neighbor hopping only. Conical touchings of bands are Dirac points. (d) Typical bulk band structure of the Haldane model in the presence of a band gap. (e) Typical band structure with the zig-zag edge for a gapless honeycomb lattice. Flat-band localized edge states (indicated in purple) connect between the Dirac cones, with one per edge. (f) Typical band structure with edges on both sides of the system when a topological gap opens. Red and blue lines indicate edge states on left and right edges, respectively.

is the momentum measured from a Dirac point. As a result, the dispersion around the Dirac point is linear, $E = \pm \hbar v_D \sqrt{q_x^2 + q_y^2}$, and referred to as the Dirac cone. A complete plot of the band dispersion is shown in Fig. 4(c).

In order to open an energy gap at Dirac points, one needs to add a term proportional to σ_z in \hat{H}_D . This can be achieved by either breaking time-reversal symmetry or inversion symmetry, which implies that as long as both time-reversal symmetry and inversion symmetry are preserved the gapless Dirac points are protected (Bernevig and Hughes, 2013).

The key novelty of the Haldane model is to add two more sets of terms to the nearest-neighbor honeycomb lattice model, which open energy gaps at the Dirac cones in complementary ways, by breaking inversion symmetry or time-reversal symmetry. The first set of terms

is a constant energy difference $2M$ between two sublattices, which breaks inversion symmetry. The second set of terms are next-nearest-neighbor hoppings with magnitude t_2 and complex hopping phases breaking time-reversal symmetry, i.e. hopping along the arrows in Fig. 4(a) carries a phase of ϕ , whereas the hopping along the opposite direction carries the opposite phase of $-\phi$. While adding the energy difference between sublattices opens a trivial gap, in the sense that the resulting bands are topologically trivial, adding complex next-nearest-neighbor hoppings results in opening a gap with topologically nontrivial bands. This different behavior is due to the breaking of time-reversal symmetry by the complex hopping phases, a necessary condition to obtain a nonzero Chern number (Bernevig and Hughes, 2013). The full topological phase diagram of the Haldane model as a function of the magnitude of the two gap-opening terms is summarized in Fig. 4(b), while an example of the energy band dispersion for a bulk system is shown in Fig. 4(d).

The nontrivial topology of the Haldane model can also be seen from the appearance of chiral propagating edge states. The dispersion of a finite slab of Haldane model is displayed in Fig. 4(e,f) for the gapless case and the gapped case with a topologically nontrivial gap, respectively. Along the y -direction the system is taken to be periodic, so the momentum k_y is a good quantum number. In the other x -direction, we have a large but finite size system with a sharp edge. While the edge states of the unperturbed honeycomb lattice are flat at the energy of the Dirac points (Fig. 4(e)), when a bulk topological gap opens, two new states appear, which traverse the energy gap in opposite directions along opposite edges of the system. Propagation of each of these states is therefore unidirectional and is protected by the fact that these edge states are spatially-separated, suppressing scattering processes from one edge state into the other.

The Haldane model, and its generalization, which is often termed the quantum anomalous Hall effect, has been realized in solid-state systems by Chang *et al.*, 2013, as well as in photonics (Rechtsman *et al.*, 2013b) and ultracold atomic gases (Jotzu *et al.*, 2014).

Bulk-edge correspondence: The relationship between a topologically nontrivial band structure and the presence of topologically protected edge states is a very general phenomenon known as the bulk-edge correspondence (Hatsugai, 1993a,b; Qi *et al.*, 2006). We now illustrate the bulk-edge correspondence through a simple model. Note that a solid mathematical formulation of this bulk-edge correspondence has been developed based on the index theorem (Callias, 1978; Chiu *et al.*, 2016).

As we have seen in the Haldane model, one can obtain topological bands by adding proper gap-opening terms, proportional to σ_z , to the gapless unperturbed Hamiltonian (16). Looking at the phase diagram of the Haldane model, Fig. 4(b), a topological phase transition can be in-




FIG. 5 A schematic illustration of how the interface of two regions with different topological numbers can host a localized state. The spatial dependence of the mass term $m(x)$ and the wavefunction residing at the interface ($x = 0$) are schematically plotted. Sketched above are the schematic dispersion for values of m at corresponding positions, showing that the bulk band-gap closes when $m(0) = 0$.

duced by changing M/t_2 or ϕ . In the Dirac Hamiltonian description, this topological phase transition corresponds to adding a σ_z term and changing the sign of its coefficient (Bernevig and Hughes, 2013; Haldane, 1988). We can therefore model the interface of two Haldane models with different Chern numbers by considering the following Jackiw-Rebbi Hamiltonian (Jackiw and Rebbi, 1976):

$$\hat{H} = \hbar v_D (q_x \sigma_x + q_y \sigma_y) + m(x) \sigma_z, \quad (17)$$

where the mass term $m(x)$ varies along x direction, obeying $m(x) < 0$ at $x < 0$, $m(0) = 0$, and $m(x) > 0$ at $x > 0$. The gap closes at $x = 0$, and the system is divided into two parts with $m < 0$ and $m > 0$, which are sketched in Fig. 5 for illustration and which have different Chern numbers. Writing the momentum operators in terms of a spatial derivative ($q_{x,y} = -i\nabla_{x,y}$), it is straightforward to see that the wavefunction of the form

$$\psi(x) \propto e^{ik_y y} \exp\left(-\frac{1}{\hbar v_D} \int_0^x m(x') dx'\right) \begin{pmatrix} 1 \\ i \end{pmatrix} \quad (18)$$

is an eigenstate of the Hamiltonian with the energy $\hbar v_D k_y$. This state $\psi(x)$ is localized around $x = 0$, and has a positive group velocity along the y -direction. As there is no other (normalizable) state around $x = 0$ with a negative group velocity, this state is chiral and robust against back-scattering. Analogously, edge states around a generic topologically nontrivial system can be understood as interface states between the system and the topologically trivial vacuum (Hasan and Kane, 2010).

2. Quantum spin-Hall system

In all the models that we have seen so far, time-reversal symmetry was explicitly broken through either the ap-

plied magnetic field or the complex hopping phase. When time-reversal symmetry is present, the Berry curvature obeys $\Omega_n(-\mathbf{k}) = -\Omega_n(\mathbf{k})$ for non-degenerate bands, implying that the Chern number, which is an integral of the Berry curvature over the Brillouin zone, is necessarily zero. A similar argument also holds for degenerate bands: no Chern bands can be found in two-dimensional systems preserving time-reversal symmetry (Bernevig and Hughes, 2013).

In 2005, new classes of two dimensional topological phases were proposed (Bernevig *et al.*, 2006; Bernevig and Zhang, 2006; Kane and Mele, 2005a,b). These models consist of two copies of a Chern insulator, one for each spin, where the magnetic field acting on two spins are opposite and hence the Chern number for spin up C_{up} is opposite to that for spin down $C_{\text{down}} = -C_{\text{up}}$. Since the magnetic fields for two spins are opposite, time-reversal-symmetry is preserved, and the sum of the Chern numbers for the two spins is zero. In this model, as long as there are no spin-flip processes, the two spin components are uncoupled and independently behave as Chern insulators with opposite Chern numbers $C_{\text{up,down}}$. As a result, there are the same number of edge states in the two spin states, but with opposite propagation direction; instead of chiral, the edge states are then called *helical*.

Even when the spin is not conserved, at least one pair of edge states survives and is topologically protected, as long as time-reversal symmetry holds. This feature is a consequence of Kramers' theorem, which holds in fermionic systems in the presence of time-reversal symmetry. The theorem tells us that if there is a state with energy E and momentum \mathbf{k} , there must exist another distinct state with the same energy but with the opposite momentum $-\mathbf{k}$. In particular, at time-reversal-symmetric momenta such as $\mathbf{k} = 0$, states should be doubly degenerate. As a consequence, when there is a pair of edge states with spin-up and spin-down crossing at $\mathbf{k} = 0$, the edge states cannot open a gap, and hence there are topologically-protected helical edge states. This is clearly different from a trivial insulator where there are no robust edge states traversing the gap. The topological invariant characterizing the system is given by $C_{\text{up}} \bmod 2$ ($= C_{\text{down}} \bmod 2$), which takes values of either 0 (trivial) or 1 (non-trivial), and thus is called the \mathbb{Z}_2 topological invariant. This \mathbb{Z}_2 topological invariant keeps the same value even when spin-mixing terms are added, provided time-reversal symmetry is maintained (Fu and Kane, 2006; Kane and Mele, 2005b; Roy, 2009; Sheng *et al.*, 2006). Such topological phases are called the quantum spin-Hall insulators or the \mathbb{Z}_2 topological insulators. The \mathbb{Z}_2 topological insulator has been experimentally realized in HgTe quantum wells (König *et al.*, 2007) following the theoretical proposal (Bernevig *et al.*, 2006). Shortly afterwards, \mathbb{Z}_2 topological insulators were found to exist also in three dimensions (Fu *et al.*, 2007; Moore and Balents, 2007; Qi *et al.*, 2008; Roy, 2009).

One may also envisage an analog of the quantum spin-Hall insulators for photons by using, for example, the polarization degree of freedom as pseudospins. However, the bosonic nature of photons forbids the existence of direct photonic analogs of the quantum spin-Hall insulators (De Nittis and Lein, 2017). For Kramers' theorem to hold, one needs that the time reversal operator \mathcal{T} be fermionic, which satisfies $\mathcal{T}^2 = -1$, while the bosonic time-reversal operator obeys $\mathcal{T}^2 = +1$.

However, if there is no coupling between pseudospins, i.e. if there is an extra symmetry in the system, then each pseudospin component can independently behave as a model with nonzero Chern number, and hence shows helical edge states in the presence of the bosonic time-reversal symmetry (Albert *et al.*, 2015; Hafezi *et al.*, 2011). Note however that topological edge states of such systems are not robust against terms coupling different pseudospin states, which would be the photonic analog of time-reversal-symmetry breaking magnetic impurities for electronic \mathbb{Z}_2 topological insulators. Photonic models showing analogs of quantum spin-Hall systems with no (or little) couplings between different pseudospin states will be reviewed in Sec.III.B.

3. Topological phases in other dimensions

We have so far reviewed the topological phases of matter in two dimensions with and without time-reversal symmetries. Generally speaking, in the presence of a given symmetry, one can consider topological phases which are protected as long as the symmetry is preserved, which lead to the concept of the *symmetry protected topological phases*. A complete classification of non-interacting fermionic topological phases in any dimension based on the time-reversal, particle-hole, and chiral symmetries is known in the literature (Chiu *et al.*, 2016; Kitaev *et al.*, 2009; Ryu *et al.*, 2010; Schnyder *et al.*, 2009; Teo and Kane, 2010).

As the topological band structure is a single-particle property and does not depend on the statistics of the particles, this classification, originally derived for fermionic systems, directly applies to bosonic systems as well, provided the Hamiltonian conserves the number of particles. The situation is in fact different when the number of particles are allowed to change, e.g. in Bogoliubov-de Gennes Hamiltonians of superconductors; in this case the bosonic and fermionic band structures are different. The fermionic case is included in the above-mentioned classification, while the bosonic case will be reviewed in Sec. VI.B of this article. We now focus on three specific examples of topological phases of matter in dimensions other than two: one-dimensional Hamiltonians with chiral symmetry, three-dimensional Weyl points, and higher-dimensional quantum Hall systems.

One-dimensional chiral Hamiltonian: One dimensional




FIG. 6 Schematic illustration of the Su-Schrieffer-Heeger model. Dashed squares indicate the unit cell of the lattice; each unit cell contains two lattice sites, one belonging to the A sublattice and other to the B sublattice. The lattice terminates on the left-hand side with a complete unit cell.

Hamiltonians with chiral symmetry can have topologically nontrivial phases characterized by an integer-valued *winding number*. In non-interacting tight-binding models, chiral symmetry is equivalent to having a bipartite lattice, i.e. a lattice that can be divided into two sublattices with hopping occurring only between different sublattices. When a discrete translational symmetry is present, the Hamiltonian in momentum space can be written in the following generic form (Ryu *et al.*, 2010):

$$\hat{H}_k = \begin{pmatrix} \mathbf{0} & Q(k)^\dagger \\ Q(k) & \mathbf{0} \end{pmatrix}, \quad (19)$$

where $Q(k)$ is an $n \times n$ matrix satisfying $Q(k) = Q(k + 2\pi/a)$, each unit cell consists of $2n$ sites, and a is again the lattice spacing.

When there is a gap at zero energy, namely $\det Q(k) \neq 0$ for any k , the topology of this Hamiltonian is characterized by the winding number defined through the phase of $\det Q(k) \equiv |\det Q(k)|e^{i\theta(k)}$ as (Kane and Lubensky, 2014; Zak, 1989)

$$\mathcal{W} = \frac{1}{2\pi} \int_0^{2\pi/a} dk \frac{d\theta(k)}{dk}. \quad (20)$$

The winding number tells us the number of times $\det Q(k)$ wraps around the origin when plotted in a complex plane as one varies k along the Brillouin zone. The bulk-boundary correspondence states that the number of zero energy edge modes on one side of the one dimensional chain is given by the absolute value of the winding number $|\mathcal{W}|$ (Delplace *et al.*, 2011; Ryu and Hatsugai, 2002). Such zero energy edge modes are topologically protected in the sense that they remain locked at zero energy even in the presence of chiral-symmetry-preserving perturbations provided the gap remains open.

The prototypical example of a one-dimensional chiral Hamiltonian with non-trivial topology is the Su-Schrieffer-Heeger (SSH) model (Su *et al.*, 1979), which is a chain with two alternating hopping strengths as sketched in Fig. 6. The system can be divided into two sublattices A and B, and the tight-binding Hamiltonian can be written as

$$\hat{H}_{\text{SSH}} = \sum_x \left(t\hat{b}_x^\dagger \hat{a}_x + t'\hat{a}_{x+1}^\dagger \hat{b}_x + \text{H.c.} \right), \quad (21)$$

where a_x and b_x are annihilation operators for A and B sublattice sites at position x . The intra- and inter-cell hoppings are described by the (real) hopping amplitudes t and t' , respectively. After a Fourier transformation, one sees that the momentum-space Hamiltonian has the form of Eq. (19) with $n = 1$ and $Q(k) = t + t'e^{ik}$. The system is gapped as long as $t \neq t'$, and the corresponding winding numbers are $\mathcal{W} = 0$ for $t > t'$ and $\mathcal{W} = 1$ for $t < t'$. Therefore, when the chain is terminated at one end with the final hopping of t , there exists a zero energy topological edge state if $t < t'$ (Ryu and Hatsugai, 2002).

One dimensional photonic structures with nontrivial topology will be discussed in Sec. IV.

Three-dimensional Weyl points: As briefly introduced above, two-dimensional band-structures can host Dirac cones, corresponding to gapless points around which bands disperse linearly with respect to the two quasi-momenta (16). In three dimensions, the analog of a Dirac point is a Weyl point (Armitage *et al.*, 2017; Lu *et al.*, 2013b; Wan *et al.*, 2011): a point degeneracy between two bands which have a linear dispersion in all three directions in momentum space at low energy, as described by the Weyl Hamiltonian:

$$H_W = \hbar v(q_x \sigma_x + q_y \sigma_y + q_z \sigma_z), \quad (22)$$

where $\mathbf{q} = (q_x, q_y, q_z)$ is the momentum measured relative to the degenerate point and v is the group velocity, taken here to be locally isotropic.

Close to a Weyl point, the resulting Berry curvature (6) is reminiscent of the magnetic field around a magnetic monopole, where the field can either point outward or inward towards the Weyl point. In analogy with magnetic monopoles, a quantized “charge” can be associated with a Weyl point; this is nothing other than a Chern number calculated by integrating the Berry curvature over a two-dimensional surface enclosing the Weyl point, generalising Eq. (8). It can also be shown that Weyl points only generate non-zero Berry curvature when either \mathcal{P} (parity) and/or \mathcal{T} (time-reversal symmetry) is broken. Consequently, to get Weyl points in a bandstructure, one can break \mathcal{P} , \mathcal{T} or both symmetries. The photonic realization of Weyl points and other three-dimensional gapless phases will be discussed in Sec. V.A.

Higher-dimensional Quantum Hall Systems: The two-dimensional integer quantum Hall effect, introduced above, is just the first in a family of related quantum Hall phenomena that can exist in higher dimensions (Chiu *et al.*, 2016). Here, we briefly review what happens in three or four spatial dimensions, highlighting key differences that emerge due to having odd/even numbers of spatial dimensions.

Firstly, in an odd number of spatial dimensions, the possible quantum Hall responses are closely related to those of a lower-dimensional system, as they are only associated with lower-dimensional topological invariants. In three dimensions, for example, a gapped band can

be labelled by a triad of Chern numbers, e.g. $[C^{(1)} \equiv (C_x^{(1)}, C_y^{(1)}, C_z^{(1)})]$, where each Chern number is calculated from Eq. (8) by integrating the Berry curvature over a 2D momentum plane orthogonal to the quasi-momentum direction indicated (Avron *et al.*, 1983). This triad of two-dimensional topological invariants leads to a linear three-dimensional quantum Hall effect, calculated by extending Eq. (10) to three spatial dimensions (Halperin, 1987; Kohmoto *et al.*, 1992; Montambaux and Kohmoto, 1990). In the simplest case, such a three-dimensional quantum Hall system could be constructed by stacking 2D quantum Hall systems along a third direction and weakly coupling the layers together (Balents and Fisher, 1996; Bernevig *et al.*, 2007; Chalker and Dohmen, 1995; Störmer *et al.*, 1986). In Sec. V.B, we shall discuss three-dimensional quantum Hall systems and other gapped 3D topological phases in the context of photonics.

Secondly, in even numbers of spatial dimensions, such as four dimensions, new types of quantum Hall effect can emerge that are intrinsically associated with that higher dimensionality (Meng, 2003; Prodan and Schulz-Baldes, 2016). These quantum Hall effects can be distinguished from their lower-dimensional cousins as they appear as a higher-order response to perturbing electromagnetic fields and because they are related to topological invariants that vanish in fewer spatial dimensions.

The four-dimensional quantum Hall effect, in particular, was first discussed by Fröhlich and Pedrini, 2000 and Zhang and Hu, 2001, and later played an important role in understanding time-reversal invariant topological insulators in two and three dimensions through a dimensional reduction procedure from four dimensions (Qi *et al.*, 2008). This quantum Hall effect consists of a non-linear quantized current response in one direction, when perturbative electric and magnetic fields are applied in the other three directions. We assume for simplicity that only one non-degenerate band is occupied and an electromagnetic gauge potential $\mathbf{A} = (A_0, A_1, A_2, A_3, A_4)$ is applied as a perturbation, where A_0 is the electric potential and (A_1, A_2, A_3, A_4) is the magnetic vector potential in four dimensions. The current in response to the applied electric field $E_\nu \equiv \partial_0 A_\nu - \partial_\nu A_0$ and the applied magnetic field $B_{\rho\sigma} \equiv \partial_\rho A_\sigma - \partial_\sigma A_\rho$ is then given by (Price *et al.*, 2015)

$$j^\mu = -\frac{e^2}{h} E_\nu \int_{\text{BZ}} \Omega^{\mu\nu} \frac{d^4 k}{(2\pi)^3} + \frac{e^3}{2h^2} \varepsilon^{\mu\nu\rho\sigma} E_\nu B_{\rho\sigma} C^{(2)}, \quad (23)$$

where $\mu, \nu, \rho,$ and σ run through spatial indices. The Berry curvature along μ - ν plane is defined as $\Omega^{\mu\nu} = \partial_{k_\mu} \mathcal{A}_\nu - \partial_{k_\nu} \mathcal{A}_\mu$ in terms of the usual Berry connection $\mathcal{A}_\mu = i \langle u_k | \partial_{k_\mu} | u_k \rangle$ and the integral is now over the four-dimensional Brillouin zone. Here, $\varepsilon^{\mu\nu\rho\sigma}$ is the 4D Levi-Civita symbol.

The second term in (23) vanishes in fewer than four spatial dimensions, and so this is the new quantum Hall effect that can emerge in a 4D system. It depends also on a four-dimensional topological invariant (Nakahara, 2003)

$$C^{(2)} = \frac{1}{32\pi^2} \int_{\text{BZ}} \varepsilon_{\alpha\beta\gamma\delta} \Omega^{\alpha\beta} \Omega^{\gamma\delta} d^4 k, \quad (24)$$

which is known as the second Chern number. In contrast, the Chern number (8) appearing in the two-dimensional quantum Hall effect is sometimes called the first Chern number. Note that both first and second Chern numbers can be extended to the case of degenerate bands by taking a trace of the integrand over the degenerate bands. The first term in (23) is a contribution to the current that is reminiscent of the two-dimensional quantum Hall response, where only two directions are involved, and is characterized by the first Chern number in the μ - ν plane; physically, this is similar to the quantum Hall physics of odd dimensions introduced above. Note that when the system possesses time-reversal symmetry, the first term of (23) vanishes and only the second term remains (Zhang and Hu, 2001). Experimental observations of the four-dimensional quantum Hall effect through topological pumping are discussed in Sec. IV.B, and proposals for directly observing the four-dimensional quantum Hall effect using synthetic dimensions are discussed in Sec. V.C.

4. Topological Pumping

Topological invariants, such as the first and second Chern numbers defined above, can also lead to a quantization of particle transport in systems which are ‘‘pumped’’ periodically and adiabatically in time. In this section, we introduce the concept of topological pumping by reviewing connections between the Archimedes screw pump and a topological pump in the semiclassical limit. We then discuss how a 1D topological pump can be related to the 2D quantum Hall effect, and how the topological framework developed so far leads to a simple lattice model for a topological pump.

Archimedes Screw pump: A pump is a device that moves fluids by mechanical action, i.e., it consumes energy to perform mechanical work by moving the fluid. One of the oldest pumps known to man is the so-called Archimedes’ screw pump, which is still in use today. In this device, a fluid is pumped by turning a screw-shaped surface inside a cylindrical shaft, see Fig. 7(a). As the screw-shaped surface is made to rotate around its axis, a volume of fluid is collected at one end of the device. It is then pushed along the tube by the rotating helicoid until it pours out at the other end of the device. Ideally, at each full turn of the pump, the collected volume is identical and the fluid is homogeneously transported a unit of




FIG. 7 (a) Schematic of an Archimedes screw pump, which mechanically transports fluid in the direction of the blue arrow as the helicoidal surface is rotated. In a quantum mechanical treatment, the screw pump can be approximated by a series of parabolic potentials, as indicated by the blue dashed lines. (b) Illustration of the quantum Hall effect for Landau levels; the spectrum is sketched in the Landau gauge on a cylinder, with open boundary conditions along x and periodic boundary conditions along y . Each state in a given Landau level is indexed by the transverse momentum k_y , setting the center of the state along x . As magnetic flux is threaded through the cylinder, an electric field is generated along y and there is a spectral flow of states along x , corresponding to the quantum Hall transport. At the system boundaries, the Landau levels bend up in energy, forming chiral-propagating edge states, as expected from the bulk-boundary correspondence. At the left-hand side, the threading of flux means empty states will move down in energy and be filled up from an external reservoir as they cross the Fermi level, μ . At the right-hand side, states will flow up in energy and particles will be ejected into a reservoir. Artefacts of these chiral edge states persist in a 1D topological pump as edge modes that cross the bulk energy-gap as the pump parameter is tuned.

distance along the device. Consequently, the screw pump is used as a variable rate feeder to deliver a measured rate or quantity of material in industrial processes.

Let us now adopt a quantum mechanical description of the screw pump. At any given time, the fluid is approximately confined within a series of parabolic potentials, see Fig. 7(a). Assuming that the fluid is noninteracting, it suffices to write the Hamiltonian for a single particle of mass m in the resulting chain of parabolic potentials

$$\hat{H} = \frac{\hat{p}_x^2}{2m} + V \sum_j (\hat{x} - x_j(t))^2 \theta[(x_{j-1}(t) + x_j(t))/2] \theta[-(x_{j+1}(t) + x_j(t))/2], \quad (25)$$

where \hat{x} and \hat{p}_x are the position operator and its conjugate, respectively. The parabolic potentials are characterised by time-dependent minima located at points $x_m(t)$, with a potential amplitude V , and equidistant separation $\Delta x = x_{m+1} - x_m$. The heavyside function $\theta[x]$ cuts the influence of neighboring sites on one another. Neglecting the coupling between neighboring wells, each well hosts the standard harmonic oscillator states of a single particle. Within a pump cycle, the minima can be parameterised as $x_m(t) = x_{m,0} + U(\varphi(t))\Delta x$, where

$x_{m,0}$ are some initial positions, $\varphi(t) = 2\pi t/T_p$ is the periodic pumping parameter, and the displacement satisfies $U(\varphi + 2\pi) = U(\varphi) + 1$. After a full period T_p of the pumping, each minimum moves by one site, and thus the Hamiltonian is invariant. Therefore, the eigenstates of the Hamiltonian are periodic as $\varphi \rightarrow \varphi + 2\pi$.

Turning on weak tunnel coupling between the localized states leads to one-dimensional Bloch bands that span the device, with time-dependent Bloch states $u_{n,k_x,\varphi}(x)$ and Bloch energies $\omega_{n,k_x,\varphi}$. In the semiclassical limit, the transport of particles by pumping can generally be captured using the semiclassical equations of motion for a wavepacket that is prepared with a well-defined center-of-mass position x and momentum k_x in a given instantaneous Bloch band n . Under adiabatic modulation of the pumping parameter, the wave packet remains in the instantaneous band and evolves with a velocity

$$\dot{x}_n = \frac{\partial \omega_{n,k_x,\varphi}}{\partial k_x} + \Omega_n \partial_t \varphi \quad (26)$$

resulting from the sum of the usual group velocity plus an anomalous velocity term. The latter is determined by the Berry curvature $\Omega_n(k_x, \varphi) = i(\langle \partial_\varphi u_n | \partial_{k_x} u_n \rangle - \langle \partial_{k_x} u_n | \partial_\varphi u_n \rangle)$ associated with the instantaneous Bloch states (Karplus and Luttinger, 1954; Xiao *et al.*, 2010).

Hence, changing the pump parameter φ induces an extra motion of the particle which, depending on the sign of Ω_n , can be either in the same or in the opposite direction as the motion of the lattice. The resulting displacement of the wavepacket after one pump cycle is obtained by integrating \dot{x}_n and can in principle be arbitrary. As this anomalous particle transport depends on the geometrical Berry curvature, it is often referred to as “geometrical pumping”. Experiments along these lines have been performed with cold atoms in (Lu *et al.*, 2016a) and in photonics in (Wimmer *et al.*, 2017).

For a filled or homogeneously populated bulk band n , however, the periodicity of the Bloch energy $\omega_{n,k_x,\varphi}$ in k_x around the Brillouin zone guarantees that the group velocity contribution integrates to zero. The displacement per cycle can then be related to the 2D topological first Chern number C of the pumping process:

$$C = \frac{1}{2\pi} \int_{\text{BZ}} \int_0^{2\pi} \Omega_n(k_x, \varphi) d\varphi dk_x \quad (27)$$

whose expression closely resembles Eq. (8) for the Chern number of a band. As the displacement is proportional to this topological invariant, it neither depends on the pumping speed, provided adiabaticity still holds, nor on the specific lattice parameters as long as band crossings do not occur. Hence, the transport is highly robust against perturbations such as interaction effects or disorder (Niu and Thouless, 1984). For the screw pump potential (25), one intuitively expects the fluid to move along

with the moving potential. This is in agreement with our effective Bloch-bands description, where all lowest bands of the system appear with $C = 1$ corresponding to the displacement by Δx per unit cycle. The situation is of course much richer in the wave mechanics case where the displacement can also be in the opposite direction depending on the sign of C .

There is a deep connection between the screw pump and the physics of the quantum Hall effect, as suggested by the quantization of the pumped particles per cycle in terms of the topological first Chern number. To explore this further, we start from the Landau level Hamiltonian introduced above in (11) in the Landau gauge, $\mathbf{A} = (0, xB, 0)$. Then the transverse momentum $\hbar k_y$ is a good quantum number, and, for a given state, the Hamiltonian reduces to that of a shifted 1D harmonic oscillator

$$\hat{H}_{\text{LL}} = \frac{\hat{p}_x^2}{2m_a} + \frac{1}{2}m\omega_c^2 \left(\hat{x} - \frac{\hbar k_y}{m\omega_c} \right)^2 \quad (28)$$

where ω_c is the cyclotron frequency introduced above, and $x_c = \hbar k_y / m\omega_c$ is the shifted center of the harmonic potential. Comparing this expression with Eq. (25), one can see that a state in a given potential minimum j in the 1D screw pump is analogous to a state with a given k_y in a 2D Landau level.

In the Landau gauge, we can imagine putting the system on a cylinder, i.e., applying periodic boundary conditions in the y -direction, and threading a magnetic flux through the cylinder. (Note that applying periodic boundary conditions puts k_y on a lattice of discretely allowed values given by the Born von-Karman boundary conditions, analogously to Eq. (25).) This threading of flux generates an electromotive electric field E_y in the y -direction, leading to a Hall response in the x -direction. An adiabatic threading of flux causes continuous evolution of eigenvalues known as the spectral flow (Phillips, 1996; Waterstraat, 2016). The number of states which cross the gap in the positive direction over the flux-threading cycle is equal to the Chern number of the spectral gap, σ_n . The central position x_c of each eigenstate shifts to that associated with the next allowed value of k_y , similar to how minima move in the screw pump, and in agreement with Laughlin's pumping argument (Laughlin, 1981). Indeed, Landau levels have first Chern numbers $|C| = 1$ in agreement with the intuition developed from the screw pump.

This analogy shows that the pump parameter φ of the 1D screw pump can be thought of as a threaded magnetic flux and hence a momentum in a perpendicular fictitious dimension; this correspondence can be exploited in a procedure called "dimensional extension" to derive the appropriate higher-dimensional model beginning from the lower-dimensional pump (Kraus *et al.*, 2012, 2013; Kraus and Zilberberg, 2012; Prodan and Schulz-Baldes, 2016; Qi *et al.*, 2008; Verbin *et al.*, 2013, 2015). Finally, as sketched in Fig. 7, the bulk-edge correspondence of the

quantum Hall effect can be used to explain how a topological pump functions with open boundary conditions and couples to particle reservoirs.

Thouless Pump & the Hofstadter Model: Turning the above analogy around, we can start with a 2D quantum Hall system, such as the Hofstadter model Eq. (14), and obtain the corresponding topological pump. For the Hofstadter model in the Landau gauge, we proceed by Fourier-transforming the model only in the y -direction to obtain

$$\hat{H} = -J \sum_{x, k_y} \left[\hat{a}_{x, k_y}^\dagger \hat{a}_{x+a, k_y} + h.c. + 2 \cos(2\pi\alpha x/a - k_y a) \hat{a}_{x, k_y}^\dagger \hat{a}_{x, k_y} \right]. \quad (29)$$

Then applying the procedure of "dimensional reduction", the momentum along y is now considered as an external parameter $\varphi = k_y$; this reduces the dimensionality of the Hamiltonian by one dimension by removing the sum over k_y and making the operators k_y -independent (Thouless, 1983). This 1D model Eq. (29) is then commonly known as the Harper model (Harper, 1955a) and will be discussed in detail in Sec.IV.B. This system can be adiabatically pumped by slowly changing the external parameter φ , and hence modulating the onsite energy periodically in time. For a filled band insulator, this is known as a Thouless pump (Thouless, 1983), as there is a quantization of particle transport over each pump cycle, as set by the sum over the first Chern numbers of the filled bands.

The main distinction from the Landau level case discussed above is that in the Hofstadter model there are two competing length scales: the lattice spacing and the magnetic length. As a result, the electrons paths interfere to give the fractal Hofstadter butterfly, with a resulting band structure composed of bands with positive and negative Chern numbers. Similarly, in the 1D Thouless pump, the on-site potential imposes a new length scale, and in the resulting band structure, we encounter bands that will pump against the direction in which the potential is moved. This is a purely wave-physics interference effect in contrast to the classical particle picture used to understand the screw pump.

In the lattice configuration realized in the ultracold atomic experiment (Lohse *et al.*, 2016; Nakajima *et al.*, 2016), the lowest energy band had a positive $C > 0$. Observing pumping in the opposite direction then required working with higher bands, which was achieved using the atomic gas in strongly non-equilibrium conditions. As we shall review in Sec. IV.B, photonic systems proved to be an ideal platform for realizing topological pumps in more complex and also higher dimensional geometries.

5. Floquet engineering: Topology through time-periodic modulations

After having introduced in the previous section the general concepts of topological bands, we now briefly review the manner by which topological band properties can be generated by subjecting a (static) system to an external time-periodic drive, an approach which is generally referred to as *Floquet engineering*, due to its direct relation to the Floquet theorem (Bukov *et al.*, 2015; Cayssol *et al.*, 2013; Eckardt, 2016; Goldman and Dalibard, 2014; Kitagawa *et al.*, 2010a, 2011; Lindner *et al.*, 2011; Oka and Aoki, 2009). We will describe how topological properties emerge in this general context, and will then briefly discuss how Floquet engineering can be exploited in photonics.

Let us first consider a generic quantum system described by the static Hamiltonian \hat{H}_0 . The aim of Floquet engineering is to modify the band structure of this Hamiltonian, effectively, by subjecting the system to a time-periodic modulation $\hat{V}(t+T) = \hat{V}(t)$, where $T = 2\pi/\Omega$ denotes the period of the drive. In the non-trivial case where $[\hat{H}_0, \hat{V}(t)] \neq 0$, the time-evolution operator $\hat{U}(t, t_0)$, which is associated with the total time-dependent Hamiltonian $\hat{H}(t) = \hat{H}_0 + \hat{V}(t)$, forms an intricate object that one can formally write as a time-ordered integral, $\hat{U}(t, t_0) = \mathcal{T} \exp\left(-\frac{i}{\hbar} \int_{t_0}^t dt' \hat{H}(t')\right)$. However, due to the time-periodicity inherent to the system, $\hat{H}(t+T) = \hat{H}(t)$, this time-evolution operator can be factorized, leading to the more suggestive form (Bukov *et al.*, 2015; Goldman and Dalibard, 2014; Kiss *et al.*, 1994; Rahav *et al.*, 2003)

$$\hat{U}(t, t_0) = e^{-i\hat{K}_{\text{kick}}(t)} e^{-i(t-t_0)\hat{H}_{\text{eff}}/\hbar} e^{i\hat{K}_{\text{kick}}(t_0)}, \quad (30)$$

where the operator \hat{H}_{eff} is time independent, and where $\hat{K}_{\text{kick}}(t+T) = \hat{K}_{\text{kick}}(t)$ has zero average over a period of the drive. The latter expression (30) indicates that the dynamics of the periodically-driven system is essentially ruled by an effective Hamiltonian, \hat{H}_{eff} , whose properties are potentially distinct from those associated with the initial static Hamiltonian \hat{H}_0 . In addition, the final “kick” $e^{-i\hat{K}_{\text{kick}}(t)}$ in Eq. (30) reflects the micro-motion, namely, the dynamics taking place within each period of the drive. Both the effective Hamiltonian \hat{H}_{eff} and the kick operator $\hat{K}_{\text{kick}}(t)$ result from a rich interplay between the static Hamiltonian \hat{H}_0 and the drive operator $\hat{V}(t)$; these two operators, and hence the time-evolution operator in Eq. (30), can be systematically computed using various perturbative treatments (Bukov *et al.*, 2015; Eckardt and Anisimovas, 2015; Goldman and Dalibard, 2014; Goldman *et al.*, 2015; Mikami *et al.*, 2016).

In traditional realizations, Floquet-engineering operates in the so-called “high-frequency” regime of the drive ($\Omega \rightarrow \infty$); physically, this corresponds to situations where the period T sets the shortest time scale in the sys-

tem (Kitagawa *et al.*, 2010a), and hence, where the micro-motion is typically irrelevant. In this regime of the drive, it is instructive to probe the dynamics stroboscopically, namely, by considering discrete observation times $t_N = NT$, where N is an arbitrary integer and $t_0 = 0$; up to a unitary (gauge) transformation, the long-time dynamics is then captured by the stroboscopic time-evolution operator [Eq. (30)]

$$\hat{U}(t_N) = e^{-it_N \hat{H}_{\text{eff}}/\hbar} = \left[e^{-iT \hat{H}_{\text{eff}}/\hbar} \right]^N = \left[\hat{U}(T) \right]^N. \quad (31)$$

Hence, in this framework, the relevant dynamics is governed by the Floquet operator $\hat{U}(T)$, or equally, by the effective Hamiltonian $\hat{H}_{\text{eff}} = (i\hbar/T) \log \hat{U}(T)$. As an important corollary, the topological properties of the system are then entirely dictated by the band structure of the effective Hamiltonian: for a proper choice of the drive protocol $[\hat{V}(t)]$, the effective Hamiltonian \hat{H}_{eff} can host topological properties, even when the underlying static system $[\hat{H}_0]$ is trivial. Consequently, in the “high-frequency” regime ($\Omega \rightarrow \infty$), the topological classification of periodically-driven systems is strictly equivalent to that of static systems (Cayssol *et al.*, 2013; Kitagawa *et al.*, 2010a; Lindner *et al.*, 2011): topological band theory (Hasan and Kane, 2010; Qi and Zhang, 2011) directly applies to the Bloch bands associated with the effective Hamiltonian \hat{H}_{eff} , i.e. the so-called “Floquet spectrum”.

A simple but important example of such driven-induced topological states is found when analyzing the properties of a particle hopping on a 2D circularly-shaken honeycomb lattice (Eckardt and Anisimovas, 2015; Jotzu *et al.*, 2014; Oka and Aoki, 2009; Plekhanov *et al.*, 2017; Rechtsman *et al.*, 2013b; Zheng and Zhai, 2014). In a frame moving with the shaken lattice, the drive takes the form of a time-periodic inertial force $\mathbf{F}(t)$, so that the time-dependent Hamiltonian can be written in the form

$$\hat{H}(t) = -J \sum_{\langle j,k \rangle} \hat{a}_j^\dagger \hat{a}_k - \sum_j \mathbf{F}(t) \cdot \mathbf{r}_j \hat{a}_j^\dagger \hat{a}_j, \quad (32)$$

where the first term describes hopping between nearest-neighbor sites of the honeycomb lattice, with hopping amplitude J , where $\mathbf{F}(t) = F [\cos(\Omega t) \mathbf{e}_x + \sin(\Omega t) \mathbf{e}_y]$ reflects the circular shaking, and where \mathbf{r}_j denotes the position of site j . Interestingly, we note that the time-dependent Hamiltonian in Eq. (32) is equivalent to that describing electrons in graphene when the latter is irradiated by a circularly-polarized light (Cayssol *et al.*, 2013; Oka and Aoki, 2009); in that case, the force $\mathbf{F}(t)$ is then directly related to the AC electric field of the radiation. The effective Hamiltonian \hat{H}_{eff} associated with the time-dependent Hamiltonian in Eq. (32) can be evaluated using the so-called $1/\Omega$ -expansion (Bukov *et al.*, 2015; Eckardt and Anisimovas, 2015; Goldman and Dalibard, 2014; Goldman *et al.*, 2015; Mikami *et al.*, 2016),

which yields

$$\hat{H}_{\text{eff}} \approx -J_{\text{eff}} \sum_{\langle j,k \rangle} \hat{a}_j^\dagger \hat{a}_k - J_{\text{eff}}^{\text{NNN}} \sum_{\langle\langle m,n \rangle\rangle} i^\zeta \hat{a}_m^\dagger \hat{a}_n, \quad (33)$$

where the first term describes the renormalization of the nearest-neighbor hopping term in Eq. (32), with $J_{\text{eff}} = J\mathcal{J}_0(Fd/\Omega)$, and where the second term corresponds to a next-nearest-neighbor hopping term, with effective hopping matrix elements $J_{\text{eff}}^{\text{NNN}} i^\zeta = \pm i(\sqrt{3}J^2/\Omega)\mathcal{J}_1^2(Fd/\Omega)$ whose sign depends on the orientation of the hopping event; here $\mathcal{J}_{0,1}$ denote Bessel functions of the first kind and d is the lattice spacing. Importantly, the effective Hamiltonian in Eq. (33) is equivalent to the Haldane model (Haldane, 1988) introduced in Sec.II.A.1.

In direct analogy with the original Haldane model, the spectrum associated with the effective Hamiltonian (33) displays two Bloch bands with non-zero Chern numbers and chiral edge states. In this Floquet-engineered context, we note that the time-reversal-symmetry breaking nature of the Chern bands naturally stem from the chirality of the circular drive. In condensed matter, a driven system exhibiting effective Bloch bands with non-zero Chern numbers is generally called a *Floquet Chern insulator*. Such a strategy was considered in various physical contexts, ranging from irradiated materials (Cayssol *et al.*, 2013; Lindner *et al.*, 2011; Oka and Aoki, 2009) to ultracold atoms in shaken optical lattices (Eckardt, 2016; Fläschner *et al.*, 2016; Jotzu *et al.*, 2014), but it was in fact pioneered in photonic experiments using femtosecond-laser-written lattices (Rechtsman *et al.*, 2013b) as reviewed in detail in Sec.III.A.2.

This Floquet-engineering strategy is not restricted to the Haldane model, but can indeed be extended to engineer other toy models of topological matter. For instance, the Harper-Hofstadter model (Hofstadter, 1976) discussed in Section II.A.1 [Eq. (14)] can be designed by subjecting a 2D square lattice to a proper time-periodic modulation; various theoretical proposals can be found in different physical contexts, including cold atoms (Creffield *et al.*, 2016; Kolovsky, 2011), ion-traps (Bermudez *et al.*, 2011) and photonics (Fang *et al.*, 2012b). As further discussed in Section III.A.5(b), these schemes build on the concept of resonant “modulation-assisted tunneling” (Goldman *et al.*, 2015), where a time-periodic modulation is set on resonance with respect to an intrinsic energy offset between neighboring sites. Cold-atom realizations of the Harper-Hofstadter model, using resonant moving optical potentials, were reported in Refs. (Aidelsburger *et al.*, 2013, 2014; Miyake *et al.*, 2013; Tai *et al.*, 2017a). More recently, a photonics implementation based on a circuit-QED architecture was described in Ref. (Roushan *et al.*, 2017b); see Fig. 30 and Section VII.B.

The simple topological characterization presented above for the high-frequency regime ($\Omega \rightarrow \infty$) breaks down when the period of the drive becomes comparable

to other times scales in the problem (e.g. when $\hbar\Omega$ becomes comparable to the bandwidth of the effective spectrum (Kitagawa *et al.*, 2010a)). Indeed, in that situation, the micro-motion becomes crucial and the topological classification based on \hat{H}_{eff} only must be revised (Carpentier *et al.*, 2015; Kitagawa *et al.*, 2010a; Nathan and Rudner, 2015; Rudner *et al.*, 2013). In particular, away from the high-frequency regime, topologically-protected edge modes are shown to exist even when the topological invariants (e.g. Chern numbers) associated with \hat{H}_{eff} are all trivial (Kitagawa *et al.*, 2010a; Rudner *et al.*, 2013). The discovery of such *anomalous Floquet topological phases* suggested that novel types of topological invariants had to be introduced in order to accurately recover the bulk-edge correspondence in this regime. Such topological invariants (winding numbers) were identified in Refs. (Asbóth, 2012; Bi *et al.*, 2017b; Carpentier *et al.*, 2015; Kitagawa *et al.*, 2010a; Nathan and Rudner, 2015; Rudner *et al.*, 2013; Yao *et al.*, 2017), and were indeed shown to depend on the complete time-evolution operator $\hat{U}(t, t_0)$; the crucial role played by the micro-motion in this topological characterization (Nathan and Rudner, 2015) indicates a shift of paradigm with respect to the standard topological classification of static systems.

As a final remark, we note that a fruitful approach to Floquet topological physics is offered by so-called *quantum walks* (Broome *et al.*, 2010; Kitagawa *et al.*, 2012, 2010b), where the time-evolution operator of a system is digitally built, by repeatedly applying a series of unitary operations $\hat{U}(T) = \hat{U}_M \hat{U}_{M-1} \dots \hat{U}_2 \hat{U}_1$; due to the T -periodicity of such quantum walks, their topological classification is equivalent to that of Floquet-engineered systems discussed above (Kitagawa *et al.*, 2012, 2010b).

As we shall see at multiple places in the course of this review, photonic systems have shown a great potential to implement Floquet techniques in different platforms. This has led to the experimental realization of intriguing Floquet phases (Bandres *et al.*, 2016; Bellec *et al.*, 2017; Noh *et al.*, 2017b; Rechtsman *et al.*, 2013b; Roushan *et al.*, 2017b), in particular anomalous Floquet topological states (Gao *et al.*, 2016a; Maczewsky *et al.*, 2017; Mukherjee *et al.*, 2017b) and topologically-protected states in quantum walks (Cardano *et al.*, 2017; Kitagawa *et al.*, 2012).

B. Features of photonic systems

Historically, the study of topological effects in quantum condensed matter systems originated from electric conduction experiments measuring the current vs. voltage characteristics of two-dimensional electron gases. In these systems, the basic constituents, the electrons, obey Fermi statistics; the topologically non-trivial states arise as the equilibrium state for sufficiently low temperature; and the electric conductivity is measured under weak or

moderate external fields that do not dramatically affect the underlying many-body state.

This equilibrium or quasi-equilibrium condition is shared by almost all condensed matter experiments, with a few remarkable exceptions such as Floquet topological insulators (Cayssol *et al.*, 2013; Inoue and Tanaka, 2010; Lindner *et al.*, 2011; Oka and Aoki, 2009) and light-induced superconductivity (Fausti *et al.*, 2011). In these systems, new phases of matter are induced by intentionally keeping the system far away from equilibrium by means of some incident electromagnetic radiation.

The situation is fundamentally different in photonic systems for at least two fundamental reasons: (i) the basic constituent – the photon, possibly dressed by some matter excitation into a polariton – obeys bosonic statistics; (ii) photons can reside in any realistic device only for a finite time and some external driving is needed to inject them into the system. In the next two subsections we shall review the consequences of these remarkable differences. In the last subsection we will review some basic concepts of nonlinear optics and we illustrate how a third-order $\chi^{(3)}$ nonlinearity can be seen as an effective binary interaction between photons, therefore opening the way towards many-body physics using gases of interacting photons (Carusotto and Ciuti, 2013).

1. Bosonic nature

As introduced above, the first key difference between quantum condensed matter systems, based on electron gases, and photonic systems is that the basic constituents of the latter are bosonic. In the previous Sec.II.A.2, we have seen how this difference can have an impact already at the single particle level as Kramer’s theorem, which underlies quantum spin-Hall physics, only holds in the presence of fermionic time-reversal operators satisfying $\mathcal{T}^2 = -1$.

At the many-body level, the difference between bosons and fermions is even more apparent as quantum statistics impose a specific symmetry on the many-body wavefunction under particle exchange: because of the Pauli exclusion principle, a non-interacting gas of fermions at low temperatures fills all states below the Fermi level with just one particle per state, while leaving all higher states empty. In the particular case of insulators, where the Fermi level lies within an energy gap, all valence (conduction) bands are filled (empty), so that integrals over the Brillouin zone naturally appear in the calculations. In contrast, a weakly interacting bosonic system at low temperatures consists of a Bose-Einstein condensate with a macroscopic population of particles accumulated in the single lowest-energy state (Huang, 1987; Pitaevskii and Stringari, 2016). As we shall see in the course of this section, the picture is made different in optical systems by the presence of losses and/or the propagating nature

of photons, so that the ground state of the system is typically a trivial vacuum state: generating and maintaining the photon gas in a state with interesting topological properties then requires injecting light from some external source.

2. Non-equilibrium nature

In any realistic optical device, photons are typically subject to a variety of processes that tend to reduce the number of photons present in the device. Depending on the geometry and the materials used, these processes can range from absorptive losses in the underlying material medium, which make the photons simply disappear, to radiative losses and/or a finite propagation time through the system, which result into the emission of light in the surrounding space as propagating radiation. As a consequence, some external pumping mechanism must be introduced to inject and sustain the light field in the device. Except for very specific cases (Hafezi *et al.*, 2015; Klaers *et al.*, 2010; Lebreuilly *et al.*, 2017; Rasmussen *et al.*, 2000; Silberberg *et al.*, 2009), the resulting state of the light field is thus quite distinct from a thermal equilibrium state.

Given this intrinsically non-equilibrium nature of photonic systems, the ways in which topological effects manifest, as well as the experimental probes and diagnostics that are available, can be completely different in photonic systems compared to condensed matter set-ups. For example, the light emitted by the device carries out detailed information on the field distribution and the photon statistics inside the device; depending on the specific set-up, this information can be extracted by imaging the emission in free space, as is typical for planar microcavities, and/or by collecting the emission with local probes such as antennas or waveguides. Different ways of injecting light into the system have also been experimentally used to highlight different properties of topological states.

We shall now proceed to review the main such pumping schemes, highlighting the key general features of each. The impact of the non-equilibrium condition on specific topological effects will be then be discussed throughout the following sections, e.g. on non-Hermitian topological photonics in Sec.VI, on the stabilization of many-body phases in Sec.VII.B, and on the measurement of topological invariants in Sec.VIII.C.

a. Coherent pumping In a typical coherent pumping scheme, the system is illuminated with an externally incident laser beam or by placing an antenna or an external waveguide in the system’s vicinity, so as to inject coherent light at specific spatial locations. Light propagation through the system is then monitored by collecting transmitted and/or scattered light with a second antenna or

a detector.

The conceptually simplest theoretical description of such an approach consists in solving Maxwell's equations, including the specific geometrical arrangement of dielectric and magnetic elements and suitable source terms to describe the emitting antenna. Since a complete analytical solution is typically beyond reach, a number of numerical methods have been developed for this task, ranging from the same plane wave expansions used to obtain the band structure, to finite element methods for the time-evolution (Joannopoulos *et al.*, 2011). Such techniques were for instance used in the theoretical calculations presented in (Wang *et al.*, 2009) and shown in Fig.1(b-c,e).

For non-bianisotropic materials with no magneto-electric coupling, one can eliminate the magnetic field and write Maxwell's equation for a field oscillating at frequency ω in the compact form (Joannopoulos *et al.*, 2011; Wang *et al.*, 2008)

$$\nabla_{\mathbf{r}} \times [\boldsymbol{\mu}^{-1}(\mathbf{r}) \nabla_{\mathbf{r}} \times \mathbf{E}(\mathbf{r})] = \omega^2 \boldsymbol{\epsilon}(\mathbf{r}) \mathbf{E}(\mathbf{r}). \quad (34)$$

where $\mathbf{E}(\mathbf{r})$ is the electric field, and $\boldsymbol{\mu}(\mathbf{r})$ and $\boldsymbol{\epsilon}(\mathbf{r})$ are the magnetic permeability and dielectric permittivity tensors, respectively. The gyro-magnetic photonic crystals used in the experiment (Wang *et al.*, 2009) and illustrated in Fig.1(a1-a2) were characterized by spatially periodic $\boldsymbol{\mu}(\mathbf{r})$ and $\boldsymbol{\epsilon}(\mathbf{r})$, as well as non-vanishing non-diagonal elements of $\boldsymbol{\mu}$, which quantify the strength of the gyro-magnetic effect. Emission by an antenna can be straightforwardly included in the model just by adding source terms on the RHS of (34).

In spatially periodic systems, a basis of solutions of (34) can be found which satisfies the Bloch theorem just like the Schrödinger equation for electrons in a periodic potential. In particular, these solutions are labelled by the crystal momentum \mathbf{k} and the band index n and the role of the Bloch wavefunction is played here by the electric field $\mathbf{E}_{n,\mathbf{k}}(\mathbf{r})$. For a given configuration, the specific form of the photonic Bloch bands and of the Bloch wavefunctions can be obtained with standard photonic techniques reviewed in (Joannopoulos *et al.*, 2011).

Given the different form of the wave equation (34) compared to the Schrödinger equation, calculation of the topological invariants for photonic crystals requires some specific work beyond the picture presented in Sec.II.A for electronic systems. While the operator acting on $\mathbf{E}(\mathbf{r})$ on the left hand side of (34) is still a Hermitian operator, the eigenvalue problem involves the dielectric permittivity $\boldsymbol{\epsilon}(\mathbf{r})$ multiplying on the right hand side. It is then useful to consider the modified scalar product,

$$\langle \mathbf{E}_1 | \mathbf{E}_2 \rangle = \int d^2\mathbf{r} \sum_{i,j} E_{1,i}^*(\mathbf{r}) \epsilon_{i,j}(\mathbf{r}) E_{2,j}(\mathbf{r}), \quad (35)$$

where $i, j = \{x, y, z\}$. The Berry connection introduced

in Eq. (5) then takes the form (Wang *et al.*, 2008)

$$\begin{aligned} \mathcal{A}_n(\mathbf{k}) &= i \langle \mathbf{E}_{n,\mathbf{k}} | \nabla_{\mathbf{k}} \mathbf{E}_{n,\mathbf{k}} \rangle = \\ &= i \int d^2\mathbf{r} \sum_{ij} E_{n,\mathbf{k},i}^*(\mathbf{r}) \epsilon_{ij}(\mathbf{r}) \nabla_{\mathbf{k}} E_{n,\mathbf{k},j}(\mathbf{r}) \end{aligned} \quad (36)$$

in terms of the (normalized) Bloch eigenfunctions $\mathbf{E}_{n,\mathbf{k}}(\mathbf{r})$ of the photon state of crystal momentum \mathbf{k} and band index n . Starting from the expression Eq. (36) for the Berry connection, the geometrical and topological invariants such as the Berry curvature, the Chern number and the bulk-boundary correspondence display the usual features as reviewed in Sec. II.A; in particular, as before, the Chern number can become nonzero only when the time-reversal symmetry is broken. Of course, equivalent geometrical and topological properties would be found if the Bloch wavefunction was written in terms of magnetic field after having eliminated the electric one.

While this *ab initio* technique can be applied in full generality to any photonic structure, a different strategy consists of developing simplified models that are able to capture the main physics, while, at the same time, providing some analytical insight as well as the possibility of extending to quantum optical features. The most celebrated such model, most suitable for discrete systems of coupled resonators, is inspired by the tight-binding picture of solid-state physics (Ashcroft and Mermin, 1976) and is naturally expressed in a quantum language. In the classical optics and photonics literature, it often goes under the name of coupled mode theory, as reviewed in (Haus and Huang, 1991).

The starting point is an expansion of the electromagnetic field

$$\mathbf{E}(\mathbf{r}) = \sum_j \mathbf{E}_j(\mathbf{r}) \hat{a}_j + \mathbf{E}_j^*(\mathbf{r}) \hat{a}_j^\dagger \quad (37)$$

onto a basis of localized quantized modes labelled (in the simplest case of single-mode resonators) by the site index j . The (suitably normalized) mode profiles $\mathbf{E}_j(\mathbf{r})$ are obtained as the eigenmodes of Maxwell's equations of eigenfrequency ω_j , and the quantum \hat{a}_j and \hat{a}_j^\dagger operators respectively destroy or create a photon in each resonator j and satisfy Bose statistics, that is $[\hat{a}_j, \hat{a}_{j'}] = 0$ and $[\hat{a}_j, \hat{a}_{j'}^\dagger] = \delta_{j,j'}$.

The corresponding Hamiltonian has the form of a collection of independent harmonic oscillators in which tunneling between neighboring sites $j' \rightarrow j$ can be straightforwardly included as hopping terms of amplitude $J_{j,j'}$,

$$H_{\text{res}} = \sum_j \hbar \omega_j \left[\hat{a}_j^\dagger \hat{a}_j + \frac{1}{2} \right] - \sum_{j,j'} J_{j,j'} \hat{a}_j^\dagger \hat{a}_{j'}. \quad (38)$$

Pioneering examples of the application of this tight-binding formalism are found in (Bayindir *et al.*, 2000; Yariv *et al.*, 1999). If the hopping amplitudes $J_{j,j'}$ can

be made complex, photons behave as if they are experiencing a synthetic magnetic vector potential. The site-dependence of the resonator frequency ω_j can model an external potential acting on photons.

The main difference between photonic systems and the usual solid-state ones is that photons can radiate away from the resonators into the surrounding empty space, e.g. through the non-perfectly reflecting cavity mirrors. At the level of the Hamiltonian Eq. (38), this requires the inclusion of a continuum of radiative modes \hat{A}_η , labelled by the index η and satisfying Bose commutation relations. These modes are linearly coupled to the localized resonator modes via terms of the form,

$$H = H_{\text{res}} + \int d\eta \hbar\omega_\eta \hat{A}_\eta^\dagger \hat{A}_\eta + \sum_j \int d\eta \left[\hbar g_{j,\eta} \hat{A}_\eta^\dagger \hat{a}_j + \text{h.c.} \right] \quad (39)$$

where ω_η is the frequency of a given radiative mode η and $g_{j,\eta}$ is the coupling between that radiative mode and the resonator mode j . As is discussed in full detail in quantum optics textbooks (Walls and Milburn, 2006), this Hamiltonian is the starting point of the so-called input-output formulation of the cavity field dynamics in terms of a quantum Langevin equation (Gardiner and Collett, 1985). Under the simplifying assumptions that the different resonators are coupled to independent continua of radiative modes with an approximately constant spectral weight within the frequency range of interest, one can write

$$i \frac{d\hat{a}_j}{dt} = \omega_j \hat{a}_j - \sum_{j'} J_{j,j'} \hat{a}_{j'} - \frac{i\gamma_j}{2} \hat{a}_j + \hat{A}_j^{\text{in}}(t). \quad (40)$$

where the radiative damping rate

$$\gamma_j = 2\pi |g_{j,\eta}|^2 \left| \frac{d\omega_\eta}{d\eta} \right|^{-1} \quad (41)$$

has to be evaluated for the resonant radiative mode such that $\omega_\eta = \omega_j$ and the (bosonic) input operators

$$\hat{A}_j^{\text{in}}(t) = - \int d\eta g_{j,\eta}^* \hat{A}_\eta \quad (42)$$

include the zero-point quantum noise as well as the incident radiation.

The model can of course be straightforwardly extended, e.g. to account for loss channels of non-radiative origin. More complex configurations may require including more field components on each site to describe multi-mode cavities; dissipative terms of different forms, e.g. a non-diagonal damping matrix $\gamma_{j,j'}$ to describe simultaneous coupling of several sites to the same continuum (Chen *et al.*, 1990; Cohen-Tannoudji *et al.*, 2008;

Ghulinyan *et al.*, 2014; Harris, 1989); and/or light amplification processes by population-inverted emitters (Gardiner and Zoller, 2004; Walls and Milburn, 2006).

In the most relevant case of a coherent incident field and a quadratic resonator Hamiltonian H_{res} , we can replace the operators with \mathbb{C} -number-valued expectation values α_j that evolve according to the ordinary differential equations,

$$i \frac{d\alpha_j}{dt} = \omega_j \alpha_j - \sum_{j'} J_{j,j'} \alpha_{j'} - \frac{i\gamma_j}{2} \alpha_j + F_j(t). \quad (43)$$

where the source term $F_j(t) = \langle \hat{A}_j^{\text{in}}(t) \rangle$ corresponds to the classical amplitude of the incident field. Techniques to efficiently evaluate the tight-binding parameters from classical transmission and reflection calculations are discussed, e.g., in (Hafezi *et al.*, 2011) for coupled ring resonator arrays. More details on other specific systems can be found, e.g. in (Bellec *et al.*, 2013a) for microwave resonators, in (Kruk *et al.*, 2017) for dielectric nanoparticles, in (Downing and Weick, 2017; Poddubny *et al.*, 2014) for plasmonic chains. Generalization to spatially continuous systems such as planar microcavities is reviewed in (Carusotto and Ciuti, 2013).

Apart from the last two terms describing driving and dissipation, this equation Eq. (43) has exactly the same form as the Schrödinger equation for non-interacting tight-binding electrons, where the field amplitude α_j plays the role of the (discrete) electron wavefunction. This formal equivalence between equations allows one to simulate the single-particle properties of tight-binding models using photonics.

Depending on the specific spatio-temporal shape of the coherent drive $F_j(t)$, the motion equation Eq. (43) can be used to describe various different phenomena such as the time-dependent response to a pulsed excitation or the non-equilibrium steady-state under a monochromatic excitation. As a general rule, a coherent drive selectively excites only those modes that are on or close to resonance with the pump frequency spectrum and that have a significant overlap with the pump profile.

For instance, when a monochromatic pump is shone on the bulk of the system, the injected light intensity dramatically depends on whether its frequency corresponds to an allowed energy band or to a band gap. On resonance with a band, a spatially extended and periodic pump can selectively excite Bloch states with specific k (Bardyn *et al.*, 2014), while a spatially localized pump generates an outward propagating field up to distances roughly proportional to the group velocity of the excited modes over the total photon decay rate (Ozawa and Carusotto, 2014). Within a forbidden gap, no propagating state is instead available and the spatial light intensity profile will show a sharp exponential decay, typically determined by the distance to the nearest band edge.

When the pump is focused on a system edge, pumping

in an energy band will result in light penetrating into the bulk, while pumping in a band gap will concentrate the excitation on edge states, if available. Of course, different modes can be selectively excited by playing with the spatial and/or polarization shape and symmetry of the pump spot.

In more complex geometries, the frequency selectivity of a coherent pump has been used to selectively excite specific localized modes, ranging from the Landau levels in the non-planar ring cavity of (Schine *et al.*, 2016) to complex Penrose tiling geometries (Vignolo *et al.*, 2016); an approach which may be extended to explore a variety of other interesting cases, including e.g. the relativistic Landau levels of strained honeycomb lattices and the momentum-space Landau levels that appear under a harmonic confinement, as theoretically studied in respectively (Salerno *et al.*, 2015) and (Berceanu *et al.*, 2016).

b. Incoherent pumping Photoluminescence experiments using incoherent pumping are a straightforward but powerful tool to visualize the energy distribution and the structure of the eigenstates of a system. In particular, this approach is routinely used in planar microcavity devices. Using this technique, states in a specific energy range can be isolated by spectrally resolving the emission, and then near- or far-field images recover the spatial profile or the k -space momentum distribution of modes. In the topological photonics context, this technique was used, e.g., to visualize the relativistic Dirac-like dispersion in honeycomb lattices (Jacqmin *et al.*, 2014) and the corresponding edge states (Milićević *et al.*, 2015; Milićević *et al.*, 2017).

Typical photoluminescence experiments are performed in a low pump power regime where the emission occurs spontaneously and is distributed fairly uniformly across all modes. Ramping up the pump power, experiments can enter a regime where bosonic stimulation and then mode-competition effects conspire to concentrate the emission into a reduced number of modes. For high enough pump power, above the so-called lasing threshold, stimulated emission exceeds losses and a new kind of steady-state is achieved: in this state, a strong and coherent light intensity is concentrated into a single mode, which absorbs all pump power and which has an emission line-width that is dramatically reduced (Gardiner and Zoller, 2004; Walls and Milburn, 2006).

As a rule of thumb, the lasing mode is typically selected by a largest gain condition; attention must however be paid to complex spatial mode deformation effects due to interplay of gain with the nonlinearity and the gain saturation (Türeci *et al.*, 2007), such as ballistic outward flows (Richard *et al.*, 2005; Wertz *et al.*, 2010; Wouters *et al.*, 2008) or solitonic-like self-bound modes (Jacqmin *et al.*, 2014; Tanese *et al.*, 2013). In the topological photonics context, in spite of these complications, ramping

the pump power above lasing threshold has been instrumental in resolving tiny spin-orbit coupling effects in hexagonal chains of micropillar resonators (Sala *et al.*, 2015; Zambon *et al.*, 2018). On-going advances in topological lasing will be outlined in the concluding section Sec.VIII.

c. Propagating geometries While both previous schemes are based on a driven-dissipative evolution of the light field, the conservative dynamics of light flowing through so-called "propagating geometries" has been exploited in a number of recent breakthrough experiments in topological photonics, starting from the realization of topological quantum walks in (Kitagawa *et al.*, 2012) to Floquet topological insulators in coupled waveguide systems in (Rechtsman *et al.*, 2013b) and Berry-phase-induced anomalous transport in (Wimmer *et al.*, 2017). Well before these advances, such propagating geometries had already been used to realize a variety of novel phenomena, e.g. spatial lattice solitons (Christodoulides *et al.*, 2003; Efremidis *et al.*, 2002; Fleischer *et al.*, 2003), two-dimensional Anderson localization (Lahini *et al.*, 2008; Schwartz *et al.*, 2007) and wave dynamics in quasicrystals (Freedman *et al.*, 2006, 2007; Levi *et al.*, 2011; Verbin *et al.*, 2013). All these experiments are based on purely classical properties of light and do not involve any quantum optical feature.

A convenient theoretical description of classical monochromatic light in these systems is based on paraxial diffraction theory (Boyd, 2008; Rosanov, 2002; Yariv, 1976). We start from classical Maxwell's wave equation for light propagating in a source-free, non-magnetic material of (spatially dependent) dielectric constant $\varepsilon = \varepsilon(x, y, z)$:

$$\nabla \times \nabla \times \mathbf{E} = \varepsilon \left(\frac{\omega}{c} \right)^2 \mathbf{E}, \quad (44)$$

where \mathbf{E} is the electric field; ω is the light frequency and c is the speed of light in vacuum. Using a standard vector identity and the fact that $\nabla \cdot (\varepsilon \mathbf{E}) = 0$, we arrive at:

$$-\nabla^2 \mathbf{E} = \varepsilon \left(\frac{\omega}{c} \right)^2 \mathbf{E} + \nabla \left(\mathbf{E} \cdot \frac{\nabla \varepsilon}{\varepsilon} \right). \quad (45)$$

Assuming that the dielectric constant $\varepsilon(x, y, z) = \varepsilon_0 + \Delta\varepsilon(x, y, z)$ displays relatively small variations from the background value $\varepsilon_0 = n_0^2$ and that the length scale of the spatial variation is large compared to the wavelength $\lambda_0 = 2\pi/k_0 = 2\pi c/(n_0\omega)$ in the medium, we can neglect the last term coupling the polarization and the orbital degrees of freedom and assume that the two polarizations evolve independently according to a scalar equation. In the experiment of (Rechtsman *et al.*, 2013b), ε varies by $\Delta\varepsilon \sim 10^{-3}$ on a length scale $\sim 10\lambda_0$.

We further assume that light is made to propagate along *paraxial* directions close to the axis of the waveguide.

uides (denoted as the positive z -direction). This guarantees that the wavevector component in the z -direction must be much greater than those in the x and y directions ($k_0 \simeq k_z \gg k_{x,y}$). This suggests to write the electric field in the carrier-envelope form

$$\mathbf{E}(x, y, z) = \hat{\mathbf{e}} \tilde{E}(x, y, z) \exp[ik_0 z] \quad (46)$$

where $\hat{\mathbf{e}}$ is the unit vector in the direction of polarization and \tilde{E} is a slowly varying function satisfying $|\nabla \tilde{E}| \ll |k_0 \tilde{E}|$. Plugging this expression into the propagation equation Eq. (45), we find that:

$$-\partial_z^2 \tilde{E} - 2ik_0 \partial_z \tilde{E} - \nabla_{\perp}^2 \tilde{E} + k_0^2 \tilde{E} = \varepsilon \left(\frac{\omega}{c}\right)^2 \tilde{E}, \quad (47)$$

where the ∇_{\perp} operator acts on the transverse (x, y)-plane. We now use the fact that \tilde{E} varies slowly in the z -direction to neglect the $\partial_z^2 \tilde{E}$ term from Eq. (47). Using for convenience the refractive index $n = \sqrt{\varepsilon} = n_0 + \Delta n \simeq n_0 + \Delta\varepsilon/(2n_0)$, we arrive at the paraxial equation for the diffraction of light through the structure:

$$i\partial_z \tilde{E} = -\frac{1}{2k_0} \nabla_{\perp}^2 \tilde{E} - \frac{k_0 \Delta n}{n_0} \tilde{E}. \quad (48)$$

For a strong enough confinement within the waveguides, the usual tight-binding approximation (Ashcroft and Mermin, 1976) can be performed on the paraxial equation Eq. (48), which leads to evolution equations for the field amplitude α_j in each waveguide of the form

$$i \frac{d\alpha_j}{dz} = k_j^z \alpha_j - J_{j,j'} \alpha_{j'}, \quad (49)$$

where the wavevector k_j^z of light propagating inside the waveguide j is determined by the background index n_0 as well as by the lateral confinement length, while the tunneling matrix $J_{j,j'}$ depends on the overlap of the evanescent tails of j, j' modes.

The formal similarity of the paraxial propagation equation Eq. (48) with the Schrödinger equation of quantum mechanics establishes a close analogy between the diffraction of classical light and the motion of a spinless massive quantum particle, where the diffraction along the xy plane sets the particle mass and the refractive index modulation gives the external potential. Note, however, that the role of the temporal coordinate is played here by the propagation direction z , whose spatial derivative replaces on the LHS of Eq. (48) the usual temporal t -derivative of the Schrödinger equation. As a result, the Floquet approach discussed in Sec.II.A.5 can be implemented in propagating geometries by spatially modulating system properties along the z direction, as experimentally pioneered in (Rechtsman *et al.*, 2013b) and reviewed here in Sec.III.A.2.

Going beyond the assumption of monochromaticity, the propagating light can have non-trivial temporal dynamics, such that the physical time t becomes like a third

spatial coordinate, in addition to the x, y transverse coordinates and the propagation equation for the slowly varying field $\tilde{E}(\mathbf{r}, t; z)$ must include an additional kinetic energy term with respect to the temporal direction,

$$i \frac{\partial \tilde{E}}{\partial z} = -\frac{1}{2k_0} \nabla_{\perp}^2 \tilde{E} - \frac{k_0 \Delta n}{n_0} \tilde{E} - \frac{1}{2m_t} \frac{\partial \tilde{E}}{\partial t^2} \quad (50)$$

whose mass-like coefficient $m_t^{-1} \equiv -d^2 k/d\omega^2$ (with $k(\omega) = \omega n(\omega)/c$) is proportional to the group velocity dispersion in the medium of frequency-dependent refractive index $n(\omega)$ (Boyd, 2008).

In order to go beyond the classical light case considered in the above equations, theoretical works (Lai and Haus, 1989a,b; Larré and Carusotto, 2015) have developed a full quantum theory to map the light propagation of quasi-monochromatic quantum light onto a standard many-body theory of interacting bosons. In the near future, this reformulation may be useful in studying the interplay of quantum fluctuations, strong nonlinearities, and topological effects.

As a consequence of the exchanged roles of the z coordinate and the physical time t , radiation that propagates through such a device does not provide a real-time monitoring of the field evolution, as is usual for the cavity systems discussed above, but instead only offers access to the field state at the endpoint $z = z_{fin}$ of the evolution. Correspondingly, extracting spectral information about eigenmodes requires the inclusion of additional elements, such as an extra one-dimensional chain of waveguides that can be used to selectively inject light at one side of the system, according to a k_z resonance condition (Noh *et al.*, 2017a). Analogously, the role of the incident field on the front interface of the device is only to set the initial condition of the evolution at $z = z_{in}$. The counterpart of these limitations is that such propagating geometries allow us to study time-dependent problems with conservative dynamics, extending in the long run even to quantum many-body physics (Larré and Carusotto, 2016; Polkovnikov *et al.*, 2011).

From an experimental point of view, focusing on works related to topological photonics, it is worth noting that a discrete version of the position-to-time mapping was exploited in (Schreiber *et al.*, 2012) to obtain a 2D analogue quantum walk by encoding two extra spatial variables in the arrival time of optical pulses. A related quantum walk was used in the experiment (Wimmer *et al.*, 2017) to reconstruct the geometrical Berry curvature of a photonic lattice model from anomalous transport features.

3. Basics of nonlinear optics

We conclude this section reviewing the main aspects of photonic systems with a brief summary of the key concepts of nonlinear optics. In view of the on-going developments towards the realization of strongly correlated




FIG. 8 QED Feynman diagram contributing to photon-photon scattering in vacuum via creation of a virtual electron-positron pair. Wavy lines represent photons and directed arrows represent electrons and positrons.

many-photon states in strongly nonlinear systems, as will be reviewed in Sec. VII.B, we will pay special attention to the reformulation of $\chi^{(3)}$ Kerr nonlinearities in terms of a binary interaction between photons.

The standard semi-classical description of nonlinear optical processes is based on Maxwell's equations, including nonlinear terms resulting from the nonlinear dependence of the dielectric polarization on the applied field (Boyd, 2008; Butcher and Cotter, 2008). In the most naive form, this reads

$$P = \chi^{(1)} E + \chi^{(2)} E^2 + \chi^{(3)} E^3 + \dots \quad (51)$$

where the linear electric susceptibility $\chi^{(1)}$ is responsible for the usual refractive index; the second-order susceptibility $\chi^{(2)}$ gives rise, e.g., to second harmonic generation, optical rectification, and parametric down-conversion processes; and the third-order susceptibility $\chi^{(3)}$ leads to four-wave mixing processes as well as an intensity-dependent refractive index. In the simplest form, this last effect can be reformulated as

$$n(\mathbf{r}) = n_0 + n_{nl} |\mathbf{E}(\mathbf{r})|^2, \quad (52)$$

where n_0 is refractive index in the linear regime, and the n_{nl} coefficient, proportional to the $\chi^{(3)}$ susceptibility, quantifies the refractive index dependence on the local light intensity.

From the point of view of photons as quantum mechanical particles, such an intensity-dependent refractive index can be reinterpreted in terms of binary interactions between photons, which are mediated by the nonlinear polarization of the underlying medium. This picture of interacting photons was pioneered in the calculation of the effective $\chi^{(3)}$ third-order nonlinear polarizability of the vacuum arising from the exchange of a virtual electron-positron pair (Heisenberg and Euler, 1936; Karplus and Neuman, 1951), as sketched in the Feynman diagram for photon-photon scattering shown in Fig. 8. Given the large mass m_e of electrons and positrons as

compared to the optical energies, the low-energy cross-section of such processes

$$\sigma \propto \alpha^4 \left(\frac{\hbar}{m_e c} \right)^2 \left[\frac{\hbar \omega}{m_e c^2} \right]^6 \quad (53)$$

is very low in vacuo: in the $\sigma \approx 10^{-68} \text{ m}^2$ range for 1 eV optical photons. This has made the experimental observation of this physics in vacuo extremely challenging. Given the $(\hbar \omega)^6$ dependence of the scattering cross section, the most natural strategy is to use high energy photons, e.g. in the γ -ray range. A first experimental observation of photon-photon scattering using the electromagnetic fields surrounding ultra-relativistic colliding ions has been recently reported in (Aaboud *et al.*, 2017).

As compared to the vacuum, condensed-matter media offer the much more accessible option of replacing electron-positron pairs of MeV-ranged mass m_e with electron-hole pairs of eV-ranged rest mass (set by the band gap of the material). According to (53) the corresponding reduction of the intermediate-state detuning provides a dramatic reinforcement of the cross section by ≈ 36 orders of magnitude. This corresponds to a significant value of the nonlinear $\chi^{(3)}$ polarizability in Eq. (51), which leads to many nonlinear optical phenomena, including the intensity-dependent refractive index of Eq. (52), two-photon absorption and parametric amplification/oscillation, etc. (Boyd, 2008; Butcher and Cotter, 2008).

Beyond the basic Feynman diagram sketched in Fig. 8, more complex configurations arising in specific materials may offer interesting advantages to experiment, e.g. when photons are dressed by optical transitions in discrete emitters (Noh and Angelakis, 2017) or excitons in solid-state materials (Ciuti *et al.*, 1998), when the collision process occurs via a long-lived biexcitonic bound state (Carusotto *et al.*, 2010; Takemura *et al.*, 2014; Wouters, 2007), or when the nonlinearity inherits the long-range character of the interactions between Rydberg states in solid-state (Kazimierczuk *et al.*, 2014) or gaseous media (Chang *et al.*, 2014; Saffman *et al.*, 2010).

In most media, the third-order $\chi^{(3)}$, introduced in Eq. (51), can however be viewed in the many-body language as the result of simple two-photon collisions. This alternative perspective shines new light on basic nonlinear optical phenomena from a novel angle and allows to take advantage of the artillery of many-body techniques originally developed in the context of condensed-matter and nuclear physics to predict new optical effects. In the last decade, the resulting concept of *quantum fluids of light* (Carusotto and Ciuti, 2013) has led to the experimental observation of a remarkable number of many-body effects in weakly interacting gases of photons, such as Bose-Einstein condensation, superfluidity and quantum hydrodynamics, etc.

Under the simplifying assumptions that the frequency-dependence of $\chi^{(3)}$ is negligible and that the rotating-

wave approximation can be performed, the real part of $\chi^{(3)}$ provides a interaction Hamiltonian

$$\hat{H}_{\text{int}} = \frac{g^{(3)}}{2} \int d^3\mathbf{r} \hat{E}^\dagger(\mathbf{r}) \hat{E}^\dagger(\mathbf{r}) \hat{E}(\mathbf{r}) \hat{E}(\mathbf{r}), \quad (54)$$

which is quartic in the electric field operators $\hat{E}(\mathbf{r})$ and where the amplitude $g^{(3)}$ is proportional to the real part of the $\chi^{(3)}$ nonlinearity. The imaginary part gives instead two-body losses from two-photon absorption. The assumption of a spatially- and temporally-local polarization response of the medium to the applied electric field medium that underlies (54) is valid in most common silicon- or silica-based materials used for topological photonics in the infrared and visible range (Hafezi *et al.*, 2013b; Rechtsman *et al.*, 2013b), as well as in Jaynes-Cummings-type lattices embedding strongly detuned emitters (Fitzpatrick *et al.*, 2017; Hoffman *et al.*, 2011). Nonetheless, one must not forget that several other experiments make use of thermal (Vocke *et al.*, 2015), photorefractive (Fleischer *et al.*, 2003; Jia and Fleischer, 2009), or Rydberg-mediated nonlinearities (Chang *et al.*, 2014; Saffman *et al.*, 2010), whose response may be slow in time and/or long-range in space due to heat and charge diffusion effects and to the inherently long-range nature of dipole interactions.

From a quantitative point of view, it is crucial to keep in mind that the optical nonlinearity of most commonly used materials results in very weak interactions between *single* photons, so that an accurate theoretical description is provided by a mean-field approach. Under this approximation, high-order averages are split into products of the mean-field, e.g.

$$\langle \hat{E}^\dagger(\mathbf{r}) \hat{E}(\mathbf{r}) \hat{E}(\mathbf{r}) \rangle \simeq \langle \hat{E}^\dagger(\mathbf{r}) \rangle \langle \hat{E}(\mathbf{r}) \rangle \langle \hat{E}(\mathbf{r}) \rangle, \quad (55)$$

and the Heisenberg equation of motion for the field expectation value $E(\mathbf{r}) = \langle \hat{E}(\mathbf{r}) \rangle$ recovers the classical Maxwell's equations including a nonlinear polarization term Eq. (51). Of course, the extremely small intensity of single photon nonlinear effects does not preclude that a huge number of photons can collectively have a dramatic impact on the macroscopic optical response to a strong light beam.

Within the mean-field approximation Eq. (55), an intensity-dependent refractive index can be included in the classical equations of motion Eq. (43) of the tight-binding formalism described in the previous subsection Sec.II.B.2, by simply adding to the RHS of the motion equation for α_j an additional term of the form

$$+ \omega_{nl} |\alpha_j|^2 \alpha_j, \quad (56)$$

where the nonlinearity parameter ω_{nl} is proportional (with an opposite sign) to the real part of the $\chi^{(3)}$ nonlinearity and, typically, inversely proportional to the spatial

volume of the optical mode under consideration (Carusotto and Ciuti, 2013). In propagating geometries, instead, an interaction term of the form

$$- \frac{k_0 n_{nl}}{n_0} |\tilde{E}(\mathbf{r})|^2 \tilde{E}(\mathbf{r}) \quad (57)$$

has to be added to the right-hand side of the paraxial propagation equation Eq. (48) for monochromatic light, which then takes the form of a Gross-Pitaevskii equation of dilute Bose-Einstein condensates (Pitaevskii and Stringari, 2016). In both cases, a non-vanishing imaginary part of $\chi^{(3)}$ and n_{nl} in (56-57) can be included to model saturable absorption, two-photon absorption or gain saturation effects.

Going beyond the mean-field regime and realizing strongly correlated photon states requires very special materials with extremely strong nonlinearities. Finding such materials is one of the most active research lines in modern nonlinear optics (Carusotto and Ciuti, 2013; Chang *et al.*, 2014; Roy *et al.*, 2017a). So far, most exciting results have been obtained using polaritons in gases of coherently driven atoms in a Rydberg-EIT configuration (Firstenberg *et al.*, 2013; Gorshkov *et al.*, 2011; Peyronel *et al.*, 2012) – the so-called Rydberg polaritons – or circuit-QED devices, where microwave cavity photons are strongly coupled to a superconducting qubit element (Gu *et al.*, 2017; Houck *et al.*, 2012; Schoelkopf and Girvin, 2008; You and Nori, 2011). Even though a complete and quantitative account of the complex features of these optical nonlinearities calls for a more sophisticated theoretical description of the interactions between Rydberg polaritons (Bienias *et al.*, 2014; Jachymski *et al.*, 2016) and of the Josephson dynamics in circuit-QED devices (Bourassa *et al.*, 2012), the simplest form (54) of the interaction Hamiltonian is typically sufficient to capture the main physics.

The quantum Langevin equation for the cavity field dynamics Eq. (40) is also straightforwardly extended to interacting regimes by adding a two-photon interaction term

$$H_{\text{nl}} = \sum_j \frac{\hbar \omega_{nl}}{2} \hat{a}_j^\dagger \hat{a}_j^\dagger \hat{a}_j \hat{a}_j, \quad (58)$$

to the resonator Hamiltonian H_{res} . Actual calculations are often simpler to perform by recasting the input-output formalism in terms of a master equation for the density matrix $\hat{\rho}$. For a coherent drive, the driving and dissipation terms have the form

$$\begin{aligned} \frac{d\hat{\rho}}{dt} = & -\frac{i}{\hbar} [\hat{H}_{\text{res}} + H_{\text{nl}} + \sum_j F_j(t) \hat{a}_j^\dagger + F_j^*(t) \hat{a}_j, \hat{\rho}] + \\ & + \sum_j \frac{\gamma_j}{2} [2\hat{a}_j \hat{\rho} \hat{a}_j^\dagger - \hat{a}_j^\dagger \hat{a}_j \hat{\rho} - \hat{\rho} \hat{a}_j^\dagger \hat{a}_j]. \end{aligned} \quad (59)$$

Generalization of this approach to incoherent pumps, as discussed in Sec.II.B.2, can be found in quantum optics

textbooks (Gardiner and Zoller, 2004; Walls and Milburn, 2006).

As introduced in Sec.II.B.2, a quantum description of light propagation in the propagating geometries crucially requires going beyond the monochromatic light assumption. Among many recent developments in this direction (Bienias *et al.*, 2014; Gorshkov *et al.*, 2011; Gullans *et al.*, 2016; Maghrebi *et al.*, 2015a; Moos *et al.*, 2015; Petrosyan *et al.*, 2011), a particularly promising theoretical approach is based on a quantum version of the $t-z$ mapping. As discussed in (Lai and Haus, 1989a,b; Larré and Carusotto, 2015), this reformulation leads to a model of interacting bosons, again with the physical roles of time t and propagation coordinate z exchanged.

III. TOPOLOGICAL PHOTONICS IN TWO DIMENSIONS

Having reviewed basic ideas of topological physics and of optical and photonic systems in the previous section, we are now in the position to dive into the exciting recent advances of topological photonics. Taking inspiration from the well-known classification of electronic topological insulators (Ryu *et al.*, 2010), the next sections will be organized according to the dimensionality and to the symmetry class to which each topological system belongs.

The present section will be focussed on two-dimensional systems, starting from the analogue quantum Hall systems that sparked the whole field of topological photonics. In the following sub-sections we will then move to quantum spin-Hall systems, anomalous Floquet insulators and, finally, gapless systems such as honeycomb lattices. For each class, we will present the principal material platforms that have been developed and the main topological effects that each system has been able to observe.

A. Analogue quantum Hall systems in photonics

In this first sub-section we will review two-dimensional photonic systems in which time-reversal symmetry is explicitly broken, so that the topology can be classified in terms of the integer-valued (first) Chern number. These systems can be considered as direct photonic analogues of integer quantum Hall states of a two dimensional electron gas in the presence of a strong out-of-plane magnetic field.

Our attention will be focussed on those optical platforms that have led to major experimental advances in the field, namely gyromagnetic photonic crystals and arrays of coupled waveguides, but we will briefly discuss also other platforms that have been theoretically proposed and are presently under experimental investigation. While most of the experiments focussed on the chiral edge modes and the resulting topologically pro-




FIG. 9 (Color online) Comparison of theoretical and experimental results for the setup shown in Fig. 1(a) for varying magnetic field. Panel (a): Theoretical topological gap map as a function of the static magnetic field applied. Each band gap is labeled by its gap Chern number, the sum of Chern numbers of the bulk bands below the gap. Panel (b): Calculations of the edge state dispersion for a few different values of the magnetic field. Panel (c): Experimental edge dispersions obtained by Fourier-transforming the edge-mode profiles containing both intensity and phase information. Panel (d): Experimental bulk transmission as a function of magnetic field and frequency, in agreement with the gap map in (a). Figure adapted from (Skirlo *et al.*, 2015).

tected one-way propagation, we will conclude by briefly reviewing theoretical work in the direction of measuring geometrical and topological properties of the bulk.

1. Gyro-magnetic photonic crystals

In a nutshell, photonic crystals (Joannopoulos *et al.*, 2011) consist of a spatially periodic arrangement of material elements giving spatially periodic dielectric permittivity $\epsilon_{ij}(\mathbf{r})$ and magnetic permeability $\mu_{ij}(\mathbf{r})$ tensors. In such a geometry, one can apply to photons the Bloch theorem originally developed in solid-state physics for electrons in crystalline solids (Ashcroft and Mermin, 1976): photon states organize themselves in allowed bands separated by forbidden gaps and are labelled by their wavevector defined within the first Brillouin zone of the periodic lattice.

Most of the early literature on photonic crystals focussed on the possibility of realizing a complete photonic band gap (John, 1987; Yablonovitch, 1987) that could, e.g., suppress spontaneous emission on embedded emitters and, in the presence of defects in the otherwise crystalline order, create strongly localized states. The study of such in-gap states gave rise to a number of exciting developments in view of photonic applications

such as high-Q photonic crystal cavities (Noda, 2016), high-performance, low-noise semiconductor lasers (Altug *et al.*, 2006; Painter *et al.*, 1999), and low-loss-waveguides insensitive to bends (Bayindir *et al.*, 2000; Lin *et al.*, 1998; Mekis *et al.*, 1996; Yariv *et al.*, 1999). On the other hand, propagating band states regained major attention when it was realized that photonic bands are not completely characterized by their energy dispersion, but also encode geometrical and topological features, which could lead to photonic analogues of the electronic quantum Hall effect (Haldane and Raghunathan, 2008).

The seminal proposal of Haldane and Raghunathan focussed on the case of a gyoelectric photonic crystal where a pair of Dirac cones are gapped under a static magnetic field which breaks the time-reversal symmetry (Haldane and Raghunathan, 2008; Raghunathan and Haldane, 2008). Soon after, this was followed by a realistic design of a gyromagnetic photonic crystal operating at microwave frequencies and displaying a Chern number of one (Chong *et al.*, 2008; Wang *et al.*, 2008). The basic formalism, such as how to define the Berry curvature of photonic bands, has been reviewed in Sec. II.B.2.

Experiments (Wang *et al.*, 2009) were performed using the material platform sketched in Fig.1(a), namely a periodic array of ferrite rods of vanadium-doped calciumiron-garnet (VCIG), a material that under a strong static magnetic field shows strong gyro-magnetic properties encoded in the non-diagonal matrix elements of the magnetic tensor μ_{ij} . The theoretical dispersion of photonic energy bands for such a system is displayed in Fig.1(e), which includes labels indicating the Chern number of the different bands.

In the experiment, the photonic crystal slab had of course a finite spatial size and was bounded by a metal wall on one side: since the reflecting gap of the surrounding metal has a topologically trivial nature, the bulk-boundary correspondence predicts that a topologically protected chiral edge mode appears within the bulk energy gaps of the photonic crystal, as indicated by the red line in the dispersion plot in Fig.1(e). Since time-reversal symmetry is broken by the external magnetic field, the number of edge states in each gap and their direction of propagation is determined by the sum of the Chern numbers of all lower-lying bands.

The main phenomenological consequence of such chiral edge states is that they support propagation in one direction only, so that back-scattering from defects and scatterers is completely suppressed independently of their nature and strength. Since no state is available at the same energy that propagates in the opposite direction, any wave incident on the defect can only circumnavigate it and then recover its original path along the edge of the system, at most accumulating some phase shift. This remarkable feature is apparent in the numerical simulation shown in Fig.1(b) and is in stark contrast with standard waveguides where generic defects are responsible for a

strong back-scattering of light and, therefore, a significantly suppressed transmission.

In contrast to the reciprocal behaviour of the bulk visible in Fig.1(c), the huge non-reciprocity of the transmission between a pair of antennas located on the edge shown in Fig.1(d), as well as its insensitivity to the presence of a metallic scatterer located in between them was the smoking gun of the non-trivial topology.

Very similar experiments were soon performed by other groups (Fu *et al.*, 2011a,b, 2010a,b; Li *et al.*, 2014a,c; Lian *et al.*, 2012a,b; Poo *et al.*, 2016, 2011, 2012; Yang *et al.*, 2013). The huge possibilities of these systems to engineer a variety of different band topologies were then explored. For instance, photonic bands with large Chern numbers were theoretically identified in (Skirlo *et al.*, 2014) by simultaneously gapping multiple pairs of Dirac cones. This prediction was confirmed by the experiment (Skirlo *et al.*, 2015), where the edge mode profiles were directly scanned and Fourier-transformed, so to observe chiral dispersions of Chern number ranging from 1 to 4, as illustrated in Fig. 9.

As further features of topological edge states, it was later shown that they can self-guide in air (Ao *et al.*, 2009; Li *et al.*, 2015c, 2014b; Liu *et al.*, 2012; Lu *et al.*, 2013a; Poo *et al.*, 2011), appear in coupled defect cavities (Fang *et al.*, 2011), have robust local density of states (Asatryan *et al.*, 2013, 2014), be modeled in time domain (Li *et al.*, 2013), self-collimate unidirectionally (Li *et al.*, 2015a), be realized in materials of Telle-gen magnetoelectric couplings (He *et al.*, 2016b; Jacobs *et al.*, 2015; Ochiai, 2015b; Sun *et al.*, 2017b), form bulk flat bands (Yang *et al.*, 2017b,c), and be immune to disorder in the bulk (Mansha and Chong, 2017; Xiao and Fan, 2017a). Remarkably, while it was previously known that one-way modes can exist on the surfaces of continuous magnetic media (Deng *et al.*, 2015; Gangaraj and Hanson, 2017; Hartstein *et al.*, 1973; Ochiai, 2015a; Shen *et al.*, 2015; Yu *et al.*, 2014, 2008; Zhang *et al.*, 2012), the topological origin of these modes was only recently unveiled (Silveirinha, 2015, 2016a,b). Topologically robust, zero-dimensional defect states localized on a dislocation in a two-dimensional topological photonic crystal were theoretically predicted and experimentally observed in (Li *et al.*, 2018).

From the application point of view, these one-way edge waveguides inspired novel device designs for tunable delays and phase shifts with unity transmission (Wang *et al.*, 2008), reflectionless waveguide bends and splitters (He *et al.*, 2010a,b; Liu *et al.*, 2010; Wang *et al.*, 2013), signal switches (Zang and Jiang, 2011), directional filters (Fu *et al.*, 2010b) and couplers (Wang *et al.*, 2011; Zhu and Jiang, 2011), broadband circulators (Qiu *et al.*, 2011; Zhang *et al.*, 2013), slow-light waveguides (Yang *et al.*, 2013), terahertz circuits (Bahari *et al.*, 2016), photonic pulling force (Wang *et al.*, 2015a), and other functions (Wu *et al.*, 2017b).

Whereas all these experiments were carried out using magneto-optic photonic crystals in the microwave domain, there is a strong push towards extending these ideas towards optical frequencies. In this domain, the magneto-optical effects are typically weaker by at least two to three orders of magnitude, but the material is continuously being improved (Onbasli *et al.*, 2016) and enhanced (Luo *et al.*, 2016). Even though the resulting topological band gap is correspondingly smaller than that in the microwave range, such a small bandwidth is still enough to provide topological features in narrow-band phenomena such as topological laser operation (Bahari *et al.*, 2017a). More details on these very recent developments are given in the outlook Sec. VIII.

2. Propagating geometries

In this section, we will discuss an alternative approach that does not make use of magneto-optical effects, thus enabling the realization of a photonic topological system in the optical frequency range: by using three-dimensional systems (in particular, arrays of optical waveguides (Szameit and Nolte, 2010)), it is possible to use one of the dimensions as a temporal coordinate and thus observe topological protection in the orthogonal plane (Rechtsman *et al.*, 2013b). For waveguide arrays, this means breaking inversion symmetry along the direction of propagation (z) in order to observe chiral edge states in the (x, y) -plane. This is akin to substituting *optical activity* (i.e., circular birefringence) for the Faraday effect required for the Haldane-Raghu mechanism. In this section, we start by introducing the fundamentals of waveguide arrays and how they may be described as a 2+1-dimensional system (two spatial dimensions, and one temporal) in which diffraction of optical wavepackets substitutes for temporal evolution of quantum mechanical particles. We then describe how such arrays may break z -reversal symmetry and realize topological protection through the Floquet-engineering approach introduced in Section II.A.5. The waveguide array geometry (they are also called ‘photonic lattices’) have been previously used across different physical systems to realize novel phenomena such as spatial lattice solitons (Christodoulides *et al.*, 2003; Efremidis *et al.*, 2002; Fleischer *et al.*, 2003), two-dimensional Anderson localization (Lahini *et al.*, 2008; Schwartz *et al.*, 2007), wave dynamics in quasicrystals (Freedman *et al.*, 2006, 2007; Levi *et al.*, 2011; Verbin *et al.*, 2013), among many others. There is also a proposal for obtaining a non-Abelian braiding phase for photons (Iadecola *et al.*, 2016) by guiding light through zero-energy mid-gap modes in a two-dimensional host lattice (though braiding of photons would not be of use for quantum computing in same spirit as the braiding of Majorana fermions).

A typical waveguide array geometry is depicted




FIG. 10 (a) Schematic diagram of honeycomb lattice waveguide array, with straight waveguides. Adapted from (Rechtsman *et al.*, 2013c). (b) Microscope image of input facet of a honeycomb lattice waveguide array. Adapted from (Rechtsman *et al.*, 2013a). (c) Schematic of an array of helical waveguides; each waveguide helix has the same radius, period, and phase (i.e., they are all moving in concert). Adapted from (Rechtsman *et al.*, 2013b).

schematically in Fig. 10(a). The waveguides are fabricated using the direct laser writing method, and their properties are highly dependent on the host material (Fig. 10(b) shows a microscope image of a transverse cross-section of the structure). Below, we will be describing experiments performed in fused silica glass (refractive index $n = 1.46$ at wavelength $\lambda = 633\text{nm}$). The specifics of the fabrication procedure are described in detail elsewhere (Szameit *et al.*, 2007; Szameit and Nolte, 2010). Typical parameters that describe waveguide properties are: refractive index increase from the background, $\Delta n \sim 1.0 \times 10^{-3}$; waveguide radii in the x and y directions are approximately $r_x = 2\mu\text{m}$ and $r_y = 5\mu\text{m}$, and their shapes may be described with a hypergaussian functional form:

$\Delta n(x, y) = \Delta n_0 \exp(-[(x/r_x)^2 + (y/r_y)^2]^\alpha)$, with the exponent $\alpha = 3$ in fused silica glass. In a typical waveguide array experiment, a beam of light is injected at the input facet of the array ($z=0$) and allowed to propagate through until it exits the array, at which point it is imaged onto a CCD camera. As discussed in Sec. II.B.2.c, it is described by the paraxial equation for the diffraction of light:

$$i\partial_z \tilde{E} = -\frac{1}{2k_0} \nabla_\perp^2 \tilde{E} - \frac{k_0 \Delta n}{n_1} \tilde{E}, \quad (60)$$

where \tilde{E} represents the envelope function of the electric field. Note that the paraxial equation takes the form of a Schrödinger equation, even though it describes the diffraction of classical light rather than the motion of a massive quantum particle. However, in the usual Schrödinger equation of quantum mechanics, the left side of the equation has a time derivative; here it is a derivative in z , the spatial coordinate in the propagation direction. Therefore, z takes the role of a temporal coordinate and the transverse (x, y) plane takes the role of an artificial two-dimensional material. The diffraction of static (CW) light therefore emulates the evolution of the wavefunction of a single quantum mechanical particle.

The structure shown schematically in Fig. 10(a) and in an experimental image in Fig. 10(b) is a honeycomb lattice of waveguides. Each waveguide is single-mode, meaning it can be thought of as a potential well with a single bound state. The waveguides are placed at a distance from one another such that the modes of neighboring waveguides can evanescently couple (i.e., ‘tunnel’) between neighbors (typical spacing $d = 15\mu\text{m}$). This results in a typical hopping parameter (a.k.a., coupling constant) between waveguides of $J \sim 1\text{cm}^{-1}$, but this can be tuned by changing the wavelength, waveguide refractive index, and/or spacing between the waveguides. The length of the sample (which corresponds to the amount of ‘time’ - i.e, propagation distance - that the optical wavefunction can propagate) is typically taken to be on the order $Z \sim 10\text{cm}$. Given that this particular array is a honeycomb lattice, the diffraction of photons therefore has a perfect correspondence with the motion of non-interacting electrons in graphene. Honeycomb waveguide arrays were first used to demonstrate optical Dirac physics via the observation of conical diffraction (Peleg *et al.*, 2007).

Since each waveguide acts as an ‘artificial atom’ in the analogy between waveguide arrays and two-dimensional materials, it is possible to employ the tight-binding approximation to Eq. (60), as described in Sec. II.B.2.c. In this approximation, the wavefunction \tilde{E} is expanded in a subspace composed of bound modes of each waveguide. Thus, we may write the paraxial equation as:

$$i\partial_z \alpha_m = - \sum_{\langle m, n \rangle} J_{mn} \alpha_n, \quad (61)$$

where α_m is the amplitude of the mode in waveguide m , J_{mn} is the hopping strength between waveguides m and n , and the summation is taken over neighboring waveguides (nearest, next-nearest, and so on, as necessary). Here we henceforth assume the tight-binding description due to its ubiquity across other experimental platforms (condensed matter, ultracold atoms, coupled resonators, among others). The bulk and edge band structures of the honeycomb lattice of Fig. 10(a) are shown in Fig. 4(c,e), assuming nearest-neighbor hopping only. For a discussion of edge band structure, see Sec. III.D. This is exactly the band structure of graphene (Neto *et al.*, 2009; Wallace, 1947) exhibiting Dirac cones - conical touchings between bands at the Brillouin zone corners. We note here this system may be explored beyond the tight-binding limit by performing photonic ‘ab-initio’ simulations by diagonalizing the full continuum Schrödinger equation (Eq. (60)). Most often, continuum simulations yield only minor quantitative corrections to tight-binding, but in special cases can reveal profound qualitative differences, including the presence of edge states in regions of the edge Brillouin zone where tight-binding predicts none (Plotnik *et al.*, 2014). A rigorous description of continuous topological systems can be found in a series of works by M. Weinstein and collaborators (Fefferman and Weinstein, 2012; Fefferman *et al.*, 2014; Fefferman and Weinstein, 2014; Lee-Thorp *et al.*, 2016).

A key requirement of realizing topologically protected chiral edge states is breaking time-reversal symmetry, as described above. Since z acts as a temporal coordinate in waveguide arrays, breaking z -reversal symmetry can allow for topologically-protected edge states in the transverse (x, y) plane. This is accomplished by using helical, instead of straight, waveguides in a honeycomb waveguide array, as depicted in Fig. 10(c). Similar helical waveguide arrays have been used to demonstrate dynamical localization (Crespi *et al.*, 2013). To describe the diffraction of light through the helical array, we move into a coordinate frame co-moving with the helices: $x \rightarrow x + R \cos \Omega z$, $y \rightarrow y + R \sin \Omega z$, $z \rightarrow z$. In the new coordinate system, the Laplacian remains unchanged, but the z -derivative transforms as:

$$\partial_z \rightarrow R\Omega (-\sin(\Omega z) \partial_x + \cos(\Omega z) \partial_y) + \partial_z \quad (62)$$

We now rewrite the z -derivative of Eq. (60) in this new coordinate system and find:

$$i\partial_z \tilde{E} = (i\nabla_\perp - \mathbf{A}(z))^2 \tilde{E} - \frac{k_0 \Delta n}{n_1} \tilde{E} + \frac{k_0}{2} R^2 \Omega^2 \tilde{E}, \quad (63)$$

where $\mathbf{A}(z) = k_0 R \Omega (-\sin \Omega z, \cos \Omega z)$ is the vector potential induced by the helical rotation, and the final term in Eq. (63) can be ignored because it is simply proportional to the identity. This vector potential \mathbf{A} corresponds to a circularly rotating electric field (note that it is curl-free, so the corresponding magnetic field is zero).

It can be incorporated into Eq. (61) simply by including the appropriate Peierls phase factors in the hopping, namely $J_{mn} \rightarrow J_{mn} \exp[-i\mathbf{A}(\mathbf{z}) \cdot \mathbf{r}_{mn}]$, where \mathbf{r}_{mn} is the vector that connects site m to site n . Therefore, the Schrödinger equation of Eq. (61) is time-periodic, and must be solved using the machinery of Floquet engineering; see Section II.A.5. This equation is perfectly analogous to the Schrödinger equation that describes the motion of electrons in graphene under the influence of circularly polarized classical light; see also Eq. (32) and paragraph below. The rotating field acts to break time-reversal symmetry (actually z -reversal symmetry, z being the temporal coordinate), without requiring the presence of a magnetic field (even a fictitious one). This Floquet system precisely maps to the Haldane model (Haldane, 1988) in the high-frequency driving limit; see Eq. (33) for a description of the effective Haldane-like Hamiltonian. The bulk and edge band structures for the Haldane model are shown in Fig. 4(d) and (f), respectively; these are qualitatively akin to those of the waveguide array described here. Comparing Fig. 4(c,e) with (d,f), it can be seen that the helicity acts to break the degeneracy at the Dirac points and open a bulk band gap. In the edge band structure (Fig. 4(f)), edge states are present (these are the two bands crossing the gap), with one localized to the top of the structure and the other localized to the bottom. They are part of a single chiral edge state that flows around the edges.

The topologically-protected chiral edge states can be directly observed experimentally. A series of such experiments were performed in Ref. (Rechtsman *et al.*, 2013b), observing perfect transmission around corners and past defects; we highlight one of these in Fig. 11. Here, light is injected in the top-left corner of a honeycomb lattice arranged in an equilateral triangle geometry, for a series of samples of increasing helix radius. The optical wavefunction travels clockwise around the structure, with increasing group velocity as a function of helix radius (i.e., gauge field strength) until about $R \sim 8\mu\text{m}$, whereupon the group velocity decreases and goes to zero at $R \sim 16\mu\text{m}$. This is quantitatively consistent with the lattice model, which predicts that the band gap opens, and then closes, as a function of increased helix radius (Rechtsman *et al.*, 2013b). The high background signal that is clearly observable at $R = 16\mu\text{m}$ arises from bending loss associated with the helicity of the waveguides (this corresponds to the phenomenon of ‘heating’ in condensed matter Floquet systems). Note that in each case, some bulk modes are excited (i.e., not all light is localized to the edge of the structure). This is attributable to the fact that the bulk modes have some support at the corner of the structure and are therefore excited along with the edge states when light is injected at the corner waveguide.

Since the advent of topological phenomena in the paraxial geometry described above, there has been sig-




FIG. 11 (a) Microscope image of input facet of waveguide array (Rechtsman *et al.*, 2013b). The yellow circle indicates the point of injection of input light. (b-f) Output facet of waveguide array after 10cm of propagation, for different devices with increasing helix radius. The edge states do not backscatter as they turn around the top-right corner, i.e., in (d). Their group velocity increases and then decreases with R , as predicted from the Floquet lattice model. Significant bending loss can be observed in the background for $R = 16\mu\text{m}$. Helix period is 1cm in all cases.

nificant progress in this direction. To highlight a few examples, waveguide array geometries have been used either theoretically or experimentally to demonstrate the optical Rashba effect (Plotnik *et al.*, 2016), the photonic anomalous Floquet topological insulator state (Bellec *et al.*, 2017; Maczewsky *et al.*, 2017; Mukherjee *et al.*, 2017b; Rudner *et al.*, 2013), topological transitions (Guglielmon *et al.*, 2017; Leykam *et al.*, 2016), chiral edge states in quasicrystals (in particular, Penrose tilings) (Bandres *et al.*, 2016), protected zero-dimensional cavity modes in two-dimensional lattices (Noh *et al.*, 2018) (protected modes two dimensions lower than host lattice), type-II Weyl points in three dimensions (Noh *et al.*, 2017b), and the topological Anderson insulator (Stützer *et al.*, 2018), a one-dimensional version of which was realized in an ultracold atomic gas (Meier *et al.*, 2018). Furthermore, Floquet topological insulators were realized in waveguide arrays fabricated with two-photon polymerization in photoresist materials; these were used to demonstrate protection against time-dependent defects (Jörg *et al.*, 2017). Due to the difficulty of breaking time-reversal symmetry in a planar geometry, the paraxial platform described here provides a rich methodology for photonic topological phenomena, including those that incorporate non-Hermiticity, nonlinearity and other effects that go beyond solid-state physics. Recently, a

state-recycling technique has been developed to significantly enhance the effective timescales over which dynamics take place within arrays of coupled optical waveguides (Mukherjee *et al.*, 2017a). This scheme consists in placing the photonic lattice into a cavity, which allows the optical state to be re-injected many times into the lattice. This approach also allows one to image real-time (stroboscopic) evolution in photonic lattices, by recording the state of the system after each round trip.

3. Optomechanics

There have been also interesting developments in breaking time-reversal-symmetry and implementing synthetic gauge fields in optomechanical systems. In general, the field of optomechanics deals with the coherent interaction between photons and acoustic phonons confined in a cavity or an array, which can be controlled at the single phonon level (Aspelmeyer *et al.*, 2014). This field has generated a lot of excitement due to its potential applications, ranging from sensing to quantum information processing.

In 2012, it was proposed that time-reversal symmetry could be broken in optomechanical resonators, and therefore, one could use them as an optical isolator and non-reciprocal phase shifter (Hafezi and Rabl, 2012). Specifically, a directional laser pump was used to select one circulation direction, such that, consequently, the manifestation of time-reversal breaking could be observed in the non-reciprocal optical response (Ruesink *et al.*, 2016; Shen *et al.*, 2016). One can also switch the role of phonon and photons and study non-reciprocal transport of phonons (Fleury *et al.*, 2014; Habraken *et al.*, 2012; Kim *et al.*, 2017; Stannigel *et al.*, 2012).

More recently, there has been an intriguing proposal for the implementation of synthetic gauge fields in optomechanical crystals (Schmidt *et al.*, 2015). Thanks to the uniformity in their fabrication, optomechanical crystals can form a 1D or 2D array of resonators, with a significant degree of controllability. Initial experimental demonstration of such an approach for a few sites has been recently reported in Ref. (Fang *et al.*, 2017). These advances could lead to the realization of various topological phases in optomechanical crystals (Schmidt *et al.*, 2015) and non-reciprocal baths and amplifiers (Metelmann and Clerk, 2015). Other optomechanical systems to implement synthetic gauge fields include quantum wells (Poshakinskiy and Poddubny, 2017), and superconducting circuits (Chu *et al.*, 2017; Clark *et al.*, 2017; Gustafsson *et al.*, 2014; O’Connell *et al.*, 2010).

4. Cavity- and Circuit-QED systems

Arrays of cavity- and circuit-QED devices can offer several advantages for exploring topological states of light, including highly controllable geometry and connectivity, as well as the possibility of reaching the regime of strong effective photon-photon interactions. Depending on the material platform and the frequency domain considered, these interactions can be obtained by coupling photons to a variety of emitters, e.g. atoms, chemical impurities in a solid-state material, artificial atoms such as quantum dots, or even superconducting circuits (Gu *et al.*, 2017). While progress in interacting topological photonics will be discussed in Sec. VII, we here introduce experimental and theoretical advances towards realizing non-interacting analogue quantum Hall states in cavity- and circuit-QED related systems.

One approach in this direction relies upon engineering arrays of tunnel-coupled re-entrant coaxial microwave cavities, as outlined theoretically in (Anderson *et al.*, 2016) and demonstrated experimentally in (Owens *et al.*, 2018). In this set-up, time-reversal symmetry is broken through the chiral on-site spatial wave-function of every fourth resonator, as depicted in Fig. 12. This chiral on-site wave-function is engineered through the coupling of a ferrimagnetic Yttrium-Iron-Garnet (YIG) crystal in a magnetic field to the near-degenerate P-modes of a triple-coax cavity. The resulting model corresponds to a Harper-Hofstadter model (Eq. 14) with an effective magnetic flux per plaquette of $\alpha = 1/4$. By using a real magnetic field to break time-reversal symmetry, this approach is akin to earlier work (Wang *et al.*, 2009) at room temperature, but allows to substantially suppress losses by focussing on those topological bands that are dark to the YIG crystals. Measuring and compensating disorder in this system relies upon Hamiltonian tomography techniques based upon 1- and 2- point network analysis of the lattice (Ma *et al.*, 2017). Forthcoming challenges include extensions of these ideas to higher dimensional topological circuits, and then to strongly interacting systems. For the latter, quality factors will need to be significantly higher as could likely be realised using superconducting cavities, and the system will need to be operated in a cryogenic environment where $\hbar\omega_0 \ll k_B T$. Specific issues related to the population of strongly correlated states will be addressed in Sec. VII.B.

A different strategy to induce an artificial gauge field in a coupled cavity array was proposed in (Cho *et al.*, 2008). The idea is to trap in each cavity a single atom with two ground states, that can be described as a lattice of effective spin-1/2 and can thus be mapped onto a lattice of impenetrable bosons. Hopping between neighboring sites is provided by the effective spin-exchange coupling that results from inter-cavity hopping of virtually-excited photons when atoms are optically dressed. The artificial magnetic field is imposed by making the opti-




FIG. 12 (a) Connectivity of a microwave cavity model engineered to realise a Harper-Hofstadter model (Eq. 14). Open circles are fundamental-mode resonators exhibiting an s -like onsite orbital. Blue→orange circles are chiral resonators with a single isolated $p_x + ip_y$ orbital, employed to induce an effective magnetic flux per plaquette of $\alpha = 1/4$. (b) A photograph of the lattice from overhead, with a single four-site magnetic unit cell highlighted in green. Fundamental-mode resonators consist of a single-post coaxial cavity; chiral resonators consist of a three-post cavity where time-reversal symmetry is broken using a Yttrium-Iron-Garnet sphere in a magnetic field. Tunneling between cavities is achieved via a slot cut between them. (c) A cut-out side-view of the same structure: now evident are aluminum screws threaded into the center-post of the fundamental-mode resonators, used to tune their frequencies to degeneracy. (d) Site-resolved measurement of the band-structure and edge-dispersion of the lattice. Frequencies on the vertical axis are in GHz. Figures taken from (Owens *et al.*, 2018).

cal dressing field spatially dependent. While this first proposal focused on the fractional quantum Hall regime, similar ideas have since been applied to non-interacting polaritons in a hybrid circuit QED system (Yang *et al.*, 2012), where a superconducting resonator at each lattice site is coupled to a nitrogen-vacancy center ensemble, whose internal states are dressed by spatially-dependent microwave sources.

Another proposal to realise a synthetic magnetic field in a circuit-QED architecture by means of passive circu-

lator elements was put forward in (Koch *et al.*, 2010): the circulators mediating the tunnel-coupling between resonators are designed to break time-reversal symmetry and can be implemented with simple superconducting circuits, such as three-junction Josephson rings. Extended to a lattice of resonators, this approach could be used to realise a Chern insulator, for example, in a Kagome geometry as also further studied in (Petrescu *et al.*, 2012). Along these lines, a general strategy to break time-reversal symmetry in a resonator lattice by exploiting the magneto-optical effect in the waveguide or resonator elements mediating inter-site coupling was discussed for photonic crystal resonator lattices in (Fang and Fan, 2013b)

5. Other proposals

Beyond current experiments, there have been many proposals for alternative routes to breaking time-reversal symmetry in photonics by exploiting light-matter coupling, dynamical modulation and/or novel lattice elements. As we now briefly review, these developments may lead to analogue quantum Hall effects in a variety of new systems, including microcavity polaritons and different realizations of resonator lattices.

a. Topolaritons Microcavity polaritons provide a particularly suitable photonic platform to address the physics of lattices with broken time-reversal symmetry. Polaritons are mixed light-matter quasiparticles arising from the strong coupling between photons and excitons – electron-hole bound pairs – confined in a semiconductor microcavity (Carusotto and Ciuti, 2013). While they are neutral particles, excitons possess a non-negligible magnetic moment arising from the spin of the electron and hole in the pair. Thus, polaritons, via their excitonic component, show significant Zeeman splittings (Δ_Z) when subject to an external magnetic field (Mirek *et al.*, 2017). In this situation, the lowest energy mode of polaritons confined in a single resonator splits into two states of different emission energy characterized by opposite circular polarizations.

This feature has been exploited in a number of theoretical works to propose a Chern insulator based on a polariton lattice in an external magnetic field (Bardyn *et al.*, 2015; Karzig *et al.*, 2015; Nalitov *et al.*, 2015; Yi and Karzig, 2016). The combination of the polariton Zeeman splitting and the TE-TM splitting (Δ_{TE-TM}) characteristic of the photonic part of polaritons (Kavokin *et al.*, 2005) results in the opening of a topological gap whenever band crossings are present in the spectrum of the lattice. Simultaneously, in finite size samples, protected chiral edge states emerge at the boundaries, with a chiral direction determined by the sign of the external magnetic

field. A prominent example of these *topolaritons* is a honeycomb lattice of semiconductor micropillars in a presence of an external magnetic field (Nalitov *et al.*, 2015). In this case, the external magnetic field is expected to open a gap at the Dirac cones with a magnitude given by Δ_Z and Δ_{TE-TM} ; the resulting bands acquire a Chern number of ∓ 2 or ± 1 depending on the ratio of these splittings to the nearest-neighbor hopping (Bleu *et al.*, 2016, 2017b). One of the most attractive features of polaritons is the possibility of combining these topological properties with significant Kerr nonlinearities; more discussion on interacting topological systems is given in Sec. VII. Recently, topolaritons have been experimentally realized in the context of topological lasers (Klemmt *et al.*, 2018), which are discussed in more details in Sec. VIII.

Besides exciton-polaritons in suitable semiconductor devices, it was recognized in (Jin *et al.*, 2016) that magneto-plasmons arising from the coupling of the electromagnetic field with electron-hole excitations in electronic quantum Hall systems also have nontrivial topological properties. In particular, since magneto-plasmons possess a particle-hole symmetry while breaking time-reversal symmetry, they are a unique example of class D 2D topological systems (Ryu *et al.*, 2010).

b. Dynamical modulation A flexible and powerful way to break time-reversal symmetry in both the microwave and optical domain is offered by the dynamical modulation of the properties of a resonator array. Typically, one either dynamically tunes the resonance frequencies of different cavities (Hayward *et al.*, 2012; Minkov and Savona, 2016), or hopping amplitudes (Fang *et al.*, 2012b). In this approach of *modulation assisted tunneling*, a large difference, $\Delta\omega$, in the on-site resonance frequencies between neighbouring tight-binding lattice sites initially suppresses particle tunneling. This tunneling can be restored by applying a suitable resonant time-dependent modulation at frequency $\Delta\omega$. The phase of the external modulation then appears in the phase of the tunneling amplitudes, simulating the effects of a gauge field on a charged particle. The idea was first introduced by (Jaksch and Zoller, 2003) and experimentally implemented in (Aidelsburger *et al.*, 2013; Miyake *et al.*, 2013) for ultracold gases. A related idea was proposed in a classical mechanical framework in (Salerno *et al.*, 2016).

For a suitable spatial-dependence of phases in a square geometry, the dynamical resonator array maps directly onto the Harper-Hofstadter model (Eq. 14). Although this mapping relies on the RWA, non-trivial topological features persist even if the inter-resonator coupling is ultra-strong and the RWA breaks down (Yuan and Fan, 2015b). Other choices of the modulation phases can be used to engineer spatially-inhomogeneous effective gauge fields (Fang and Fan, 2013a; Lin and Fan, 2014) or effective electric fields (Yuan and Fan, 2015a;

Yuan *et al.*, 2016b) that may also be useful in controlling light. Complex modulations were implemented in (Lumer *et al.*, 2018) to realize novel light guiding effects by a spatially dependent gauge field, as proposed in (Lin and Fan, 2014). Going up in geometrical complexity, non-reciprocal propagation based on interband photonic transitions induced in a waveguide by means of a running-wave-shaped, electrically-driven modulation of the refractive index was experimentally demonstrated in (Lira *et al.*, 2012). In general, note that dynamical modulation as well as magneto-optical effects could also be exploited to imprint a synthetic magnetic field for photons in a resonator-free implementation, based on a waveguide network (Lin and Fan, 2015).

The actual implementation of the general temporal modulation idea depends on the specific photonic system under consideration. In the infrared and optical domains, the main limitation is posed by the rate at which photonic structures can actually be modulated (e.g., by carrier injection or optomechanics), which is typically significantly lower than the optical frequency itself (GHz vs 100THz). A resonator lattice with hoppings on the order of the modulation frequency (or lower) is thus required, which in turn imposes stringent bounds on the quality factor of the resonators. In circuit-QED architectures, schemes to dynamically modulate the superconducting circuit elements (e.g. SQUIDs) used to couple the lattice of resonators together, were proposed in (Peropadre *et al.*, 2013; Wang *et al.*, 2015b, 2016d). As we shall see in more detail in Sec. VII.B, this strategy was employed for a pioneering demonstration of the interplay of magnetic and interaction effects in (Roushan *et al.*, 2017b). In propagating geometries based on waveguide arrays, the role of time and the propagation direction are exchanged (see Sec. III.A.2), and so a suitable “time-dependent” modulation may be realized by spatially varying the refractive index of the medium (Dubček *et al.*, 2015b; Longhi, 2013). Pioneering experiments along these lines were reported in Mukherjee *et al.*, 2015 and Rechtsman *et al.*, 2013b and are reviewed in Sec. III.A.2.

B. Analogue quantum spin Hall systems in photonics

The second main class of topological photonics systems in 2D are those which preserve time-reversal symmetry for photons and which are analogous to quantum spin Hall systems in condensed matter. In this class of systems, the one-way edge modes are only topologically protected if spin-changing scattering processes can be neglected. In the first six subsections we will provide an in-depth discussion of key systems that have been experimentally realised. The last subsection will briefly review some promising theoretical proposals.

1. Silicon ring resonator arrays

This first subsection is devoted to a review of the experiments in (Hafezi *et al.*, 2013b; Mittal *et al.*, 2014, 2016a) using arrays of silicon ring resonators. In these systems, a synthetic magnetic field for photons can be engineered by controlling the differential optical paths, when photons hops between neighboring sites in two opposite directions. A pseudospin-1/2 degree of freedom naturally arises as ring resonators support a pair of degenerate whispering gallery-like modes, propagating in opposite clock-wise and counter-clock-wise directions. As the device does not contain any real magnetic element, time-reversal symmetry imposes that the two spin states experience opposite synthetic magnetic fields, and therefore, the corresponding edge states have opposite chiralities, in direct analogy to quantum spin Hall systems.

We start by explaining how a non-symmetric phase hopping can be engineered between a pair of optical resonators. By arranging these pairs into a square lattice, one can synthesize a uniform magnetic field. Furthermore, we review the experimental progress in this platform, from imaging topological edge states, demonstration of their robustness in transport observables such as transmission and delay time, and finally measurement of the associated topological invariants. We also discuss the prospect of extension of these ideas in the quantum regime, in particular, in generation of photon pairs and quantum transport of two-photons.

Two-resonator case. As the first step, we show a non-zero hopping phase can be obtained for photons hopping back and forth between a pair of coupled site-resonators. As originally proposed in (Hafezi *et al.*, 2011), this can be obtained when two neighboring resonators are coupled through a third off-resonant link ring, as sketched in Fig. 13(a). We consider the length of resonators to be $m\lambda$, where $2m$ is an even integer and λ is the resonant wavelength. The off-resonant ring link length is set to $2m\lambda + 3\lambda/2$ to guarantee that it remains anti-resonant. The physics would be analogous for an odd integer, except for the sign of tunneling. Now, if the off-resonant ring link is vertically shifted by x between site resonators, as shown in Fig. 13(a), the accumulated optical phase during the hopping from one site to the other is asymmetrical, and we achieve the desired hopping phase between two sites, given by $\phi = 4\pi x/\lambda$, as shown in Fig. 13(b).

The formal derivation of this hopping phase is presented in (Hafezi *et al.*, 2011) using the input-output formalism (Gardiner and Collett, 1985), which, as discussed in Sec.II.B.2, is equivalent to coupled mode-theory in the non-interacting case. An alternative approach using transfer matrices is presented in the supplemental material of (Hafezi *et al.*, 2013b). We briefly, recapitulate the input-output formalism approach. Specifically, the boundary condition for the electric field is given by:




FIG. 13 Panel (a) Schematic sketch of the coupling mechanism between two site resonators via an off-resonant link ring. Panel (b) After integrating out the off-resonant link, its asymmetric location leads a non-zero hopping phase ϕ between site resonators. (c) A two-dimensional array of rings realizing the Harper-Hofstadter model for photons. Figure from (Hafezi *et al.*, 2011).

$\hat{E}_i^{\text{out}} = \hat{E}_i^{\text{in}} + \sqrt{2\kappa}\hat{a}_i$, where κ is the coupling efficiency and the site index can be left or right, $i = R, L$. Assuming no intrinsic loss in waveguides, the dynamics of each optical resonator is given by: $\dot{\hat{a}}_i = -\kappa\hat{a}_i - \sqrt{2\kappa}\hat{E}_i^{\text{in}}$. The input of each resonator is related to the output of the other resonator, by a propagation phase which depends on the optical length. For our set of parameters, $\hat{E}_{R(L)}^{\text{in}} = -i\hat{E}_{L(R)}^{\text{out}} \exp(\mp 2\pi i\phi)$. For counter-clockwise modes of the side resonators, as shown in Fig. 13(a), photons acquire a phase proportional to the upper or lower connecting optical lengths, when they hop from left to right and vice versa. If the link ring is positioned asymmetrically, this hopping phase can be different for forward/backward hopping. Specifically, by eliminating the fields in the link rings ($\hat{E}_R^{\text{in}}, \hat{E}_L^{\text{in}}, \hat{E}_R^{\text{out}}, \hat{E}_L^{\text{out}}$), the photon dynamics is given by $\hat{a}_{R(L)} = i\kappa \exp(\mp 2\pi i\phi)\hat{a}_{L(R)}$ which is equivalent to having a Hamiltonian

$$H = -\kappa\hat{a}_R^\dagger\hat{a}_L \exp(-2\pi i\phi) + \text{h.c.} \quad (64)$$

Note that since the overall length of the link resonators is fixed, to maintain the anti-resonance condition, the

acquired phases for forward/backward hops are not independent from each other. Specifically, they are equal in magnitude and opposite in sign. In the general case of an arbitrary off-resonant link ring, the non-zero hopping phases in the forward and backward directions remain opposite in sign to guarantee the hermiticity of the Hamiltonian. However, if the anti-resonancy condition is not met, the resonant frequency of the resonators gets shifted, see the discussion in the supplementary material of (Hafezi *et al.*, 2013b).

2D lattice: By arranging the resonator in a lattice structure, one can implement an effective uniform magnetic field in form of the Harper-Hofstadter model, which was discussed in Sec. II.A.1. In particular, we implement the Landau gauge, as in Eq. 14, to obtain the same phase α over each square plaquette, except that here x-y directions are swapped. The phase pattern can be obtained by arranging the positioning of the link rings according to the Landau gauge. Specifically, the link rings responsible for tunneling in the y-direction are symmetrically positioned to get zero hopping phase. In contrast, the link ring for tunneling in the x-direction are asymmetrically positioned, in an increasing order in y-direction, so giving rise to a $e^{i 2\pi\alpha y}$ hopping phase. As discussed in Sec. II.A.1, for a fixed magnetic field, which is set by the parameter α , any finite lattice displays bulk and edge bands, the former being organized in the celebrated Hofstadter butterfly, and the latter being located in the gaps of the bulk dispersion.

Experimental setup. The above model was implemented in (Hafezi *et al.*, 2013b) using standard silicon-on-insulator (SOI) technology working in the telecom range at $\lambda \simeq 1.55 \mu\text{m}$. As it is illustrated in Fig. 14, high-quality silicon ring resonators with a Q factor exceeding 10^4 were fabricated on top of an oxide substrate using deep-UV projection photolithography. The cross-section of the waveguides, which form the site resonators and off-resonant link rings, was designed to ensure single-mode propagation of the transverse electric (TE) mode. The evanescent coupling between the site resonators and the link rings was controlled by the thickness of the air gaps separating the elements. Due to surface roughness of waveguides, a fraction of the light in the resonators scatters orthogonally to the plane and can be captured by a microscope. In addition to transmission measurement through the input and output waveguides coupled to specific site resonators, imaging this scattered light on a CCD camera gives direct information on the spatial profile of the photonic modes. The good quality of the resonators guarantees that the spin-flip-like coupling between clock-wise and counter-clock-wise propagating modes are effectively negligible.

Topologically protected edge states When the system is excited via the input waveguide with a laser field at a frequency resonant with one of the edge modes, the photons are guided through the edge and exit the sys-




FIG. 14 Upper row (a): sketch of the experimental setup. Second row (b): spatial intensity profile showing light propagation along the edge of the system. Third row (c): spatial intensity profile showing routing of light along the edge and around a missing resonator on the edge. (d) SEM image of the system with a missing resonator, intentionally removed in the design. Adapted from Ref. (Hafezi *et al.*, 2013b)

tem from the output waveguide. Fig. 14(b) shows the light propagation clockwise along the edge of the system. The transverse width of the edge state was about one to two resonators, as observed both in experiment and simulation. As a direct manifestation of the topological protection of edge states, when a resonator is removed from the path of an edge state, the photons route around the missing resonator and then continue their path to the output coupler without being back-reflected [Fig. 14(c)].

Beside the spatial imaging the edge states and the qualitative study of their robustness, a more quantitative experimental demonstration of their robustness was reported in (Mittal *et al.*, 2014) using the structure described above and sketched in Fig. 14(a). Fig. 15 shows the transmission and delay spectra interferometrically measured at the output port for eight different 8×8 lattice devices. Overall, the transmission spectra of the different devices show significant fluctuations because of in-




FIG. 15 (a) Measured transmission and (b) delay-time spectra for eight different 8×8 lattice devices. Two regions with reduced variance in both the transmission and the delay-time are indicated by the red and green shading. A noisy region of propagation through bulk states is indicated by the blue shading. Lower panel: scaling of the transmission as a function of system size for 2D and 1D devices. Solid markers with error bars are the measured average and standard deviation (65% confidence band) values. Solid lines with shaded areas are the simulated average and standard deviation. Adapted from Ref.(Mittal *et al.*, 2014)

intrinsic fabrication variations in the dimensions of the site resonators and link rings. On closer inspection, one can see that the fluctuations in both the transmission and the delay-time are suppressed in the two regions indicated by the red and green shading in Fig. 15(a,b) corresponding to propagation via edge states in the counter-clock-wise and clock-wise direction along the short and long edges, respectively.

More insight on this physics can be seen in Fig. 15(c), which shows the measured average transmission and its standard deviation for a sample of 95 devices. The transmission through the topological edge states of a 2D lattice and the one through the (non-topological) band of an analogous 1D array (a so-called CROW) are plotted as a function of system size, i.e. the number of resonators travelled from input to output. In both cases, the transmission decays exponentially with system size. However, the decay rate is slower for the topological 2D system compared to the 1D system. The shaded regions are simulation results, using the experimentally estimated parameters, which agree with the experimental observation.

In order to differentiate the decay of transmission stemming from resonator losses from the one due to disorder – both resulting in exponential attenuation – the simulated result in the absence of disorder is presented as a dashed line: while losses affect both 2D and 1D systems in the same way, transport through topological edge states of a 2D system appears to be much less disturbed by disorder than the 1D counterpart.

As a further feature of topologically protected edge states, the work (Mittal *et al.*, 2014) experimentally demonstrated how transport in lossy edge states was unambiguously distinguished from tunneling through localized bulk states by considering the statistical distribution of the delay time during propagation. Specifically, the delay distribution for edge states is approximately Gaussian with a Gaussian width independent of system size, as typical of diffusive transport in one-dimensional systems (Cooper *et al.*, 2010). On the other hand, the distribution for bulk states is asymmetric with the most probable value being less than the average, as typical of transport governed by localization and earlier observed in one-dimensional systems in the microwave domain (Chabanov and Genack, 2001).

Invariant measurement The hallmark feature of topological physics is the presence of one-way propagating modes at the system boundary, whose chirality is a consequence of topological character of the bulk. Specifically, the bulk-boundary correspondence dictates that the number of chiral edge modes is completely determined by the bulk topological invariant, the Chern number. Following a proposal in Ref.(Hafezi, 2014), the winding number associated to a spectral flow of edge states (see Sec. II.A.4) was experimentally studied in (Mittal *et al.*, 2016a).

While the transverse conductance experiment usually performed in electronic systems is not applicable to photonic systems, the general spectral flow argument (Halperin, 1982; Laughlin, 1981) is in fact applicable also to this case. To model the spectral flow of a quantum Hall edge of winding number $k = 1$, one can consider a linear edge dispersion $E_p = vp$ where E_p is the energy, v is the group velocity, and p is the momentum along the edge. When a gauge flux (θ) is coupled to the edge, the momentum is replaced by the covariant momentum, i.e., $E_p = v(p - q\frac{\theta}{L})$, where L is the length of the edge and q is the charge of the edge excitations. For non-interacting photons, the (synthetic) charge can be set to $q = 1$ so that the corresponding vector potential is simply θ/L . For a finite system, quantization of momentum on the edge results in

$$E_n = \frac{2\pi v}{L} \left(n - \frac{\theta}{2\pi} \right), \quad (65)$$

where n is an integer. Therefore, the insertion of $\theta = 2\pi$ flux shifts $E_n \rightarrow E_{n-1}$, resulting in a spectral flow.




FIG. 16 Panel (a): sketch of the experimental device (left). The edge states considered in the experiment lie on the outer edge. The tunable gauge field coupled only to the edge states is introduced by fabricating heaters on link rings situated on the lattice edges. SEM image showing heaters fabricated on top of link rings (top-right) and schematic of the waveguide cross section showing the ring resonators, the metal heaters and the metal routing layer (bottom-right). Panel (b): measured transmission as a function of the coupled flux θ and the incident laser frequency ω . Insets: zoom-in of the edge state bands. Adapted from Ref. (Mittal *et al.*, 2016a)

To experimentally observe and measure this spectral flow, the synthetic gauge field system described above should be supplemented with an extra tunable gauge flux (Mittal *et al.*, 2016a). To couple a tunable gauge field to the edges, metallic heaters were fabricated above the link ring waveguides on the lattice edge, as shown in Fig. 16(a). These heaters use the thermo-optic effect to modify the accumulated phase of light propagating through the waveguides and hence result in a gauge flux.

Fig. 16(b) shows the measured transmission spectrum as a function of the coupled flux θ . Edge states of the outer edge and the bulk states are easily identifiable, as bright and dark regions, respectively. As the coupled flux θ increases, the energy of the clockwise edge states decreases, whereas the energy of counter-clockwise edge states increases. For a 2π increase in flux, the edge state resonances move by one resonance to replace the position once held by its neighbor. This flow indicates that the measured winding number is $k = +1.0(1)$ for the clockwise circulating edge states, and $k = -1.0(2)$ for

the counter-clockwise circulating edge states.

Specific theoretical proposals to directly detect the bulk topological invariants without using the edge physics have been put forward by many authors. A short review of the main ones can be found in Sec. VIII.C.

2. Topological RF Circuits

RF circuits are an excellent substrate for the study of topological band structures, and eventually strongly interacting topological phases of matter. Their connectivity can literally be wired in an arbitrary manner, with arbitrary numbers of connections per node and long-range connections, allowing a wide range of exotic band structures to be realized. Furthermore, the physical size of the components is macroscopically large compared to atoms, phonons, and optical photons, enabling easy access for site- and time-resolved measurements. Moreover, topological properties in linear lattice models can be observed at room temperature and in the presence of significant dissipation, despite the fact that there are many thermal photons present. This is a vivid demonstration that topological band structures are property of waves rather than of quantum mechanical interactions.

Two types of RF topological circuits have been experimentally demonstrated so far; the first type was discussed in Sec. III.A.4 and was used to realise a topological model with time-reversal symmetry breaking and non-zero Chern numbers. The second type of circuit, which will be the focus of this subsection, employed kHz frequency lumped element inductors and capacitors to implement a time-reversal invariant topological system (Ningyuan *et al.*, 2015). In this system, it was possible to observe many signature effects of a topological insulator, as well as several features that would be difficult or impossible to find in a solid-state material system.

In what follows, we briefly describe the experimental realization of a quantum spin-Hall version of the Harper-Hofstadter model (Goldman *et al.*, 2010) with a quarter flux per plaquette, that uses only inductors and capacitors; it was subsequently pointed out (Albert *et al.*, 2015) that a minimal circuit model could be realized at a flux per plaquette of $\frac{1}{3}$. As shown in Fig. 17, each lattice site in the experimental circuit consisted of two inductors (labeled ‘A’ and ‘B’), allowing for the representation of two pseudo-spin states. Tunneling between lattice sites was achieved by capacitive coupling, with the sign of the coupling reflected in which ends of the inductors are coupled to one-another. Finally, the spin-orbit coupling was implemented by changing, on a site-by-site basis, whether A was coupled to A, $-A$, B, or $-B$.

In more detail, a localized excitation on a single lattice site was represented by RF fields in a $\frac{1}{\sqrt{2}}(A \pm iB)$ superposition of the two inductors on the lattice site, where $+$ ($-$) corresponded to spin up (down). Under these




FIG. 17 (a) Schematic of the experimental circuit used to realize a time-reversal-invariant quantum spin Hall system. In this set-up, the periodic structure is composed of inductors and coupling capacitors (black) that are connected by wires (light and dark blue lines). Each lattice site consists of two inductors, labeled “A” and “B”, corresponding to right and left circularly polarized spins. As a photon hops around a single plaquette (indicated in orange) it accumulates a Berry phase of $\pi/2$; this is engineered by braiding the capacitive couplings, as detailed in (b) and indicated here in green. (b) The synthetic spin-orbit coupling is engineered through the structure of the coupling elements between lattice sites. Each of the four tunneling phases implemented (indicated by the left column of each set of figures) is induced by a particular coupling between inductors (middle column), as described in the corresponding rotation matrix (right column); the signs of the couplings are controlled by whether the ‘+’ end of one inductor is coupled to the ‘+’ or ‘-’ end of the adjacent inductor. (c) Photograph of the experimental circuit, in which the inductors (black cylinders) are coupled via the capacitors (blue). Inset: Zoom-in on a single plaquette that consists of four adjacent lattice sites. Figure taken from (Ningyuan *et al.*, 2015).

conditions, for a spin-up excitation, coupling $(A, B) \rightarrow (A, B)$ between adjacent lattice sites corresponded to implementing a tunneling phase of 0° , while connecting $(A, B) \rightarrow (-A, -B)$ corresponded to a tunneling phase of 180° , and $(A, B) \rightarrow \pm(B, -A)$ to a tunneling phase of $\pm 90^\circ$. Repeating $(0^\circ, 90^\circ, 180^\circ, -90^\circ)$ horizontal tunnel-couplers every four lattice sites, with all vertical tunnel couplers having a phase of 0° , resulted in a Harper-Hofstadter model with a quarter flux per plaquette. Spin-down excitations experienced the opposite Peierls phase, and hence the opposite effective flux per plaquette.

It is important to note that this approach transcends

the tight-binding regime: instead of a full LC resonator on each A- and B- site, only an inductor was included. When combined with coupling capacitors, this resulted in a Harper-Hofstadter spin-band-structure exhibiting non-zero spin Chern numbers identical to those observed in the analogous tight-binding model; the lack of on-site inductors changes the band widths and gaps, but, perhaps surprisingly, does *not* change the topology of the spin-bands. An equivalent construction swaps inductors and capacitors, sending $\omega \rightarrow \frac{\omega_0^2}{\omega}$ (with $\omega_0 \equiv \frac{1}{\sqrt{LC}}$), in much the same way that swapping inductors and capacitors in the (topologically trivial) 1D lumped-element transmission line converts the transmission line between left- and right-handed.

By forgoing on-site resonators and using *onsite* inductors with *coupling* capacitors, this circuit operates in what can be referred to as a “massless, left-handed” configuration. This requires fewer inductors, which are the primary source of loss and disorder, and provides band gaps of order ω_0 , making the system less susceptible to disorder than tight-binding approaches with an onsite resonator, where the band-gaps are all reduced by the ratio of the on-site capacitance to the coupling capacitance.

Because this approach exhibits tunneling energies comparable to the photon energy ω_0 , it can be operated with effective quality factors (resulting from inductor loss R) of order ~ 100 (characteristic of off-the-shelf room-temperature electronic components), with tunneling still observed over ~ 30 lattice sites within the photon $\frac{1}{e}$ lifetime; furthermore, a few percent disorder in the components, typical of off-the-shelf electronics, do not induce noticeable backscattering within the photon lifetime. Fig. 18a-b shows the dynamical evolution of a spin-mixed pulse injected on the edge, as it splits into spin-components which move with opposite chiralities.

Because this system is a *circuit*, its global connectivity is easily modified. Previous work has seen the exploration of Möbius topologies (Fig. 18 c-d), with only a single edge (Ningyuan *et al.*, 2015); by connecting the left and right edges with a twist and spin-flip (c), a Möbius strip geometry is realized, where spin-ful excitations propagate to the right along the top edge, undergo a spin flip, propagate to the left along the bottom edge, and repeat. Related prospects include for creation of conical defects to explore inter-Landau-level states (Biswas and Son, 2016).

A topoelectrical circuit (Lee *et al.*, 2018) displaying zero-dimensional topological corner midgap states, protected by the bulk spectral gap, reflection symmetries, and a spectral symmetry has also experimentally been realized in (Imhof *et al.*, 2018). A three-dimensional circuit displaying Weyl points was realized in (Lu *et al.*, 2018), providing access to the Weyl points and the Fermi arcs as well as to the Berry curvature distribution underlying the quantized chiral charge of the Weyl points. Finally,




FIG. 18 Time-Resolved transport dynamics of the edge modes in the topological circuit (Ningyuan *et al.*, 2015), shown in Fig. 17. (a)&(b) The spin-resolved time-evolution of sites around the edge of the system, following the initial excitation of an A inductor at the edge. This corresponds, in the pseudo-spin basis, to an initial localized excitation of $(\uparrow + \downarrow)/\sqrt{2}$ at a single lattice site. (a) When exciting a topological edge state (at 145 kHz), the \uparrow (red) and \downarrow (blue) signals propagate around the edge in opposite directions, demonstrating the expected spin-momentum locking. Despite the presence of disorder, two round-trips are visible (as sites 0 and 42 are equivalent). (b) When exciting a non-topological edge state (at 225 kHz), disorder immediately leads to backscattering. Inset: The grey arrow indicates the initially-excited lattice site, with the edge site numbering convention indicated by the red and blue arrows. (c) Illustration of quantum spin Hall edge states on a Möbius strip (Top Panel), with arrows indicating the edge propagation direction, and colors representing the spin states. Experimentally, this was realized by imposing the connectivity of a Möbius strip on the circuit (Bottom Panel), with the printed circuit board shown in orange and the additional external connections colored according to the spin. (d) Spin-resolved detection of edge-transport after the excitation of \uparrow ; the \uparrow (red) and \downarrow (blue) signals show the conversion from \uparrow to \downarrow as the excitation moves from one edge to the other, with three round-trips being visible. Figure taken from (Ningyuan *et al.*, 2015).

a microwave network was used in (Hu *et al.*, 2015) to measure a topological edge invariant.

3. Twisted Optical Resonators

Exploring Landau level physics with charge-neutral particles is a persistent goal of the synthetic matter and meta-material communities, both because learning to create “effective magnetic fields” for charge-neutral particles illuminates the meaning of a magnetic field, and because interacting topological matter in the continuum (fractional quantum Hall phases, for example) admits simpler theoretical description than lattice analogues (fractional Chern insulators).

Proposals to explore Landau-level physics with light rely upon coupling an optical field to a rotating atomic medium (Otterbach *et al.*, 2010) or phase plate (Longhi, 2015), to inject angular momentum. The connection to synthetic magnetic fields may be understood by realizing that sending light through a rotating medium induces an image rotation (Franke-Arnold *et al.*, 2011), thus turning the laboratory frame into a rotating frame. When viewed in a (constantly) rotating frame, massive particles experience fictitious Coriolis and centrifugal forces $2m\vec{\Omega} \times \vec{v}$, and $m\vec{\Omega} \times \vec{\Omega} \times \vec{r} = m|\Omega|^2\vec{r}_\perp$, respectively, where \vec{r} and \vec{v} are the particle position and velocity, $\vec{\Omega}$ is the angular velocity of the rotating frame, and m is the particle mass. The Coriolis force has the same form as the Lorentz force $q\vec{v} \times \vec{B}$, with the identification $q\vec{B} \equiv m\vec{\Omega}$. Thus, to investigate Landau level physics of light in the *lab* frame, it is only necessary to induce a continuous image rotation on the light, and to somehow compensate for the centrifugal force that this rotation applies to the light.

The combination of image rotation and centrifugal force cancellation was first realized in (Schine *et al.*, 2016), using techniques proposed in (Sommer and Simon, 2016) and depicted in Fig. 19. The image rotation is achieved by directing the light repeatedly through a non-planar path using a four-mirror optical resonator: akin to a pair of back-to-back dove prisms, this extremely low-loss construction is able to send the light through the non-planar path *thousands* of times, inducing an image rotation on each round-trip set by the non-planarity of the resonator.

To achieve the centrifugal-force cancellation, the mirrors that produce the image rotation are curved to repeatedly focus the light towards the resonator axis. The interplay of wave propagation, reflection off of curved surfaces, and image rotation results in a complex mode-structure for such a resonator. Qualitatively, wave-propagation in the paraxial limit is equivalent to evolution of a free massive particle, and thus imbues the photon with mass; reflection off of the curved mirror surface provides a radial impulse which is proportional to the distance from the resonator axis, providing harmonic trapping, while the resonator twist (non-planarity) induces the Lorentz and centrifugal forces.

This analogy may be sharpened through 2D ABCD matrices (Siegman, 1986), or more intuitively, by treat-




FIG. 19 Engineering Landau Levels for optical photons. Light in an optical resonator behaves as 2D particles in a harmonic trap. (a) When the ray trajectory is followed over many round-trips through a two-mirror resonator, its “hit-pattern” in the central plane of the resonator corresponds to the stroboscopic evolution of a classical massive trapped particle. The projection of the eigenmodes of the resonator onto the central plane of the resonator are Hermite-Gauss, akin to a quantum harmonic oscillator. (b) When a four-mirror resonator is twisted out of the plane, the ray trajectories undergo a round-trip rotation about the resonator axis, equivalent to Lorentz (Coriolis) and centrifugal forces; when the centrifugal force precisely cancels the harmonic confinement induced by the mirror curvature, the resulting Lorentz force produces Landau levels. (c) A realistic rendering of the four-mirror resonators employed in (Schine *et al.*, 2016), from where this figure is taken.

ing the repeated passage of the optical field through the resonator as a periodically driven system (Sommer and Simon, 2016), resulting in an effective “Floquet” Hamiltonian for the optical field in a particular plane of the resonator. The final result is manifolds of degenerate resonator eigen-modes with energies $\frac{E_{npq}}{\hbar} = \omega_{npq} = n\frac{2\pi c}{L} + p\omega_{cyc}$, where c is the speed of light in vacuum, L is the resonator round-trip length, and ω_{cyc} is the effective cyclotron frequency which determines the energy-gap between Landau levels. The n quantum number determines how many wavelengths fit within the resonator longitudinally, or equivalently which “Floquet” copy we are referring to. Then p is the Landau-level index, and q is the angular momentum index of the eigenstate within the Landau-level.

Stabilizing the Landau Levels against Astigmatism In practice, such resonators are sensitive to mirror astigmatism (due to off-axis incidence of the optical field on the mirrors), which results in different harmonic confinement along x - and y - axes, and a consequent destabilization of the Landau level. This destabilization may be understood in various ways: (1) astigmatism means that centrifugal force cannot be simultaneously canceled along both x - and y - axes, and because particles in magnetic fields move along equi-potentials, the residual confinement along a single axis guides the particles off to infinity; (2) when the astigmatism is optimally canceled, the residual confining potential takes the form $(x^2 - y^2) = r^2(e^{i2\theta} + e^{-i2\theta})$ — a potential that drives $\Delta l = \pm 2$ transitions within the (degenerate) manifold of states comprising the Landau level. This effect has also been observed in rotating atomic gases (Cooper, 2008).

To realize a degenerate Landau level, this astigmatism issue must be addressed. One approach is to engineer a Landau-level-like scenario which does not exhibit states whose angular momenta are separated by $2\hbar$; this is possible by further twisting the resonator, resulting in a situation where the new “lowest” manifold of degenerate states exhibits only every third unit of angular momentum $q = 0, 3, 6, \dots$. This new, demonstrably stable system corresponds to a Landau level on a cone with opening angle $\alpha = \arcsin(\frac{1}{3})$, which may be understood by realizing that the allowed values of q only support three-fold symmetric light-patterns; when a photon leaves a particular third, it re-enters that third from the opposite edge, but the dynamics are otherwise that of a planar Landau level. This is precisely how a charged particle behaves when constrained to the surface of a cone, in a B -field normal to the cone’s surface. This platform has enabled the first direct measurement of the mean orbital spin (of the LLL, see Fig 20), a topological quantum number which quantifies the coupling of density to manifold curvature through the Wen-Zee action (Wen and Zee, 1992).

Time-reversal symmetry breaking is particularly important in interacting systems, as collisions between particles in a quantum spin-Hall system can induce back-scattering which would otherwise be symmetry-protected. The twisted optical resonator does not break time-reversal symmetry, and so it must have a hidden spin degree of freedom which when flipped induces the photons to experience the opposite magnetic field. This turns out to arise from the order in which the photon traverses the mirrors of the running-wave twisted resonator. Forward- and backward- modes exhibit opposite synthetic magnetic fields. To break the degeneracy between them, it is sufficient to differentially couple to the polarization of the modes, which are opposite relative to a fixed axis (though they are the same relative to the direction of propagation). This Faraday-type coupling was recently demonstrated with an optically pumped ensemble in a weak magnetic field (Ningyuan *et al.*, 2017).

4. Intrinsic spin-orbit interactions for light

The fundamental properties of Maxwell’s equations lead to an “intrinsic” spin-orbit coupling for light, in contrast to “extrinsic” spin-orbit effects, engineered by the design of photonic materials as discussed in the previous sections. Intrinsic spin-orbit coupling plays an important role on length-scales comparable to the wavelength of light, and so has attracted attention across photonics, nano-optics and plasmonics (Bliokh *et al.*, 2015b). It can also lead to an analogue of the quantum spin Hall effect for light (Bliokh *et al.*, 2015a), as we briefly introduce.

In free space, a propagating plane-wave has two spin states, given by the left and right-handed circular polarizations, which have opposite helicities $\sigma = \pm 1$. The




FIG. 20 Translational Invariance of Landau Levels, and Topological Dynamics on a Conical Manifold. The astigmatism-robust Landau levels exhibit a three-fold symmetry, manifest in their spectrum by only every third angular momentum state being degenerate $q = 0, 3, 6, \dots$, or $q = 1, 4, 7, \dots$ or $q = 2, 5, 8, \dots$. Panels in the top, central, bottom row correspond to starting with $q = 0, 1, 2$, respectively. Away from the cone tip, the system supports translational invariance, as well as the possibility of generating states with arbitrary angular momentum. Left column: Injection of the two lowest Landau levels centered away from the cone tip: within each group of images, the lower subpanels are for the lowest Landau level, while the upper ones are for the first excited one. In each case, the resonator responds with three copies of the Landau level under consideration: far enough away from the cone tip (left subpanels) the copies are independent of one another. When they are closer (right subpanels), the form of the interference of the copies near the cone tip is sensitive to the initial $q = 0, 1, 2$, corresponding to flux-threading the tip itself. Right panels: The breaking of translational invariance near the cone-tip reflects the Wen-Zee coupling of the lowest Landau level to the geometric curvature of the cone tip; the excess density for the unthreaded cone reflects the mean-orbital spin of the integer quantum Hall state. Figure taken from (Schine *et al.*, 2016).

corresponding spin vector is $\mathbf{S} = \sigma \mathbf{k}/|\mathbf{k}|$ in units of \hbar , and so is aligned with the propagation vector \mathbf{k} . This is an intrinsic coupling between the orbital and spin degrees of freedom for light, underlying a wide-range of phenomena (Bliokh *et al.*, 2015b).

When light is strongly-confined transverse to its propagation direction, intrinsic spin-orbit coupling can lead to behaviour reminiscent of the electronic quantum spin Hall effect (Bliokh *et al.*, 2015a; Kavokin *et al.*, 2005; Leyder *et al.*, 2007; Mechelen and Jacob, 2016). In a nutshell, the idea is the following. As a general consequence of Gauss' law in free space, $\nabla \cdot \mathbf{E} = 0$, the transversality condition ($\mathbf{k} \cdot \mathbf{E} = 0$) implies that the electric field polarization depends directly on the wavevector \mathbf{k} . Since the wavevector \mathbf{k} in the evanescent tail of a confined optical mode has an imaginary component orthogonal to the surface, the polarization acquires a circular component corresponding to a non-vanishing transverse spin component, whose sign changes with the propagation direction (Bliokh *et al.*, 2014; Bliokh and Nori, 2012).

An analogy can thus be drawn between this so-called “spin-direction” (or “spin-momentum”) locking of confined optical modes and the spin-momentum locked edge states of a 2D quantum spin Hall system or surface states of a 3D topological insulator. However, a key difference is that the optical modes are bosonic and so not topologically-protected by time-reversal symmetry (Bliokh *et al.*, 2015a). Formally, this can be seen by noting that, although a non-zero topological Chern number can be found for each uncoupled helicity state in free space, these Chern numbers are even and so the \mathbb{Z}_2 topological invariant, introduced in Sec. II.A.2, vanishes.

Nevertheless, previous experiments had observed how spin-polarized emitters give rise to a spin-controlled unidirectional excitation of surface or guided modes in a wide-range of systems, including metal surfaces (Lee *et al.*, 2012; O'Connor *et al.*, 2014; Rodríguez-Fortuño *et al.*, 2013), optical nano-fibers (Mitsch *et al.*, 2014; Petersen *et al.*, 2014; Sayrin *et al.*, 2015) and waveguides (Le Feber *et al.*, 2015; Söllner *et al.*, 2015). These experiments also show that, despite the lack of topological protection, the spin-direction locking and this optical analogue of the quantum spin Hall effect is very robust to system details (Bliokh *et al.*, 2015b).

5. Bianisotropic Metamaterials

In this section, we review how a quantum spin Hall effect for photons can be realised in electromagnetic metamaterials, namely artificial composite materials containing sub-wavelength dielectric and/or metallic structures (Liu and Zhang, 2011). The key advantage of such materials is the great flexibility that they offer for engineering the effective dielectric permittivity ε , magnetic permeability μ , and bianisotropy or magneto-electric coupling χ that appear in the electric and magnetic response to long-wavelength fields,

$$\begin{pmatrix} \mathbf{D} \\ \mathbf{B} \end{pmatrix} = \begin{pmatrix} \varepsilon & i\chi \\ -i\chi^T & \mu \end{pmatrix} \begin{pmatrix} \mathbf{E} \\ \mathbf{H} \end{pmatrix}. \quad (66)$$

As first proposed in (Khanikaev *et al.*, 2013), a properly designed metamaterial structure can exhibit a quantum spin Hall effect for light, as experimentally evidenced by the presence of topologically robust spin-dependent edge states, analogous to the helical edge states of an electronic topological insulator.

Following the initial theoretical proposal, the first step in this direction is to construct photonic modes which mimic the electron spin-1/2 eigenstates. This can be achieved, for example, through an enforced matching of $\varepsilon = \mu$ in a metamaterial, ensuring that, in the absence of bianisotropy, the TE- and TM-polarized modes propagate along a 2D metamaterial slab with the same wavenumbers, restoring the $(\mathbf{E}, \mathbf{H}) \rightarrow (-\mathbf{H}, \mathbf{E})$ duality of the electromagnetic field in free space. For a given wavevector \mathbf{k} , linear combinations $\psi_{\mathbf{k}}^{\pm}$ of these degenerate TE/TM modes can be constructed with the special property that $\psi_{\mathbf{k}}^{+}$ can be transformed into $\psi_{-\mathbf{k}}^{-}$ by a suitable symmetry operation \hat{D} , similarly to how an electron's spin is flipped by time-reversal symmetry. Since the square of the transformation satisfies the usual $\hat{D}^2 = -1$ condition of electronic time-reversal, these states can be identified as a pair of photonic pseudo-spin-1/2 states and show symmetry-protected Kramers doublets for time-reversal symmetric momenta.

The second step required is then to engineer an appropriate bianisotropy or magneto-electric coupling χ (Serdyukov *et al.*, 2001) that generates a strong spin-orbit coupling between the pseudo-spin states, mimicking that found in a topological insulator. To first order, the effect of a finite χ can in fact be recast in reciprocal space as an explicit coupling between photon momentum and polarization, $\mathbf{D} = \epsilon\mathbf{E} + (ic/\omega)\chi\mu^{-1}\mathbf{k} \times \mathbf{E}$ and $\mathbf{B} = \mu\mathbf{H} + (ic/\omega)\chi^T\epsilon^{-1}\mathbf{k} \times \mathbf{H}$. While the bianisotropy χ of materials found in Nature, such as optically-active media containing chiral molecules, is typically small, a very large value can be obtained in metamaterial structures containing, for example, split-ring resonators (Li *et al.*, 2009; Marqués *et al.*, 2002; Pendry *et al.*, 1999; Shelby *et al.*, 2001).

This approach was exploited in (Khanikaev *et al.*, 2013) in the design of $\varepsilon = \mu$ -matched metamaterial rods arranged into a “meta-crystal” in the form of a hexagonal lattice, which had a significant value of the bianisotropy $\chi_{xy} = -\chi_{yx}$ terms. In such a hexagonal geometry, the photonic bands host doubly-degenerate Dirac points, which are gapped out by the bianisotropy χ . Around the gapped Dirac points, the effective low-energy model can then be mapped to the Kane-Mele model for the quantum spin Hall effect (see Section II.A.2), where the topological states are protected by the engineered symmetry \hat{D} of the electromagnetic field in the metamaterial.

Experimentally, the key signature of this photonic quantum spin Hall effect is the appearance of robust polarisation-dependent edge states. Such “spin”-




FIG. 21 (a) (Top) Displacing a metallic collar relative to the two metal plates of a waveguide introduces an effective bianisotropy, coupling the electric and magnetic fields. Shifting a collar from the “down” to “up” position reverses the sign of this term, and so changes the sign of the “mass” term m that opens a gap at the Dirac cone, once these rods are arranged into a hexagonal or triangular lattice. (Middle and Bottom) Photograph of the experiment in (Cheng *et al.*, 2016), in which collars in a triangular array can be moved up and down to create arbitrary and reconfigurable topological domain walls. (b) Configuration of the topological switch where in the red (blue) regions, the metallic collars are always up (down), and where in the green region, the collars are dynamically moved from down to up to change the location of the domain wall. (c) and (d) The time-resolved switching of transmission via the edge states through (c) Port 2 and (d) Port 3, with insets illustrating the power flow. All figures adapted from (Cheng *et al.*, 2016).

polarized transport has been observed for microwave photons in (Chen *et al.*, 2014) for a uniaxial hexagonal meta-crystal of non-resonant meta-atoms sandwiched between two metallic plates, where the effective bianisotropy arises from field inhomogeneities. The topological robustness of such edge modes was also further demonstrated in (Slobozhanyuk *et al.*, 2016b), who used near-field imaging in a square lattice of bianisotropic metamolecules to directly show the lack of backscattering around sharp corners.

Following works have shown that topological photonic states displaying the photonic analog of the quantum spin Hall effect can also be realised in an even more simple structure, as recently proposed (Ma *et al.*, 2015) and experimentally realized (Cheng *et al.*, 2016; Lai *et al.*, 2016; Xiao *et al.*, 2016a). In this “meta-waveguide” set-up, a parallel-plate metal waveguide is filled with a hexagonal or triangular lattice of metallic cylinders, which are connected at the top and bottom to the metal plates (see Figure 21 (a)). The geometry of the cylinders and plates is carefully optimized such that the spectrum contains an overlapping degenerate pair of Dirac cones for the TE

and TM modes, allowing for the introduction of photonic pseudo-spin states. The bianisotropy is then introduced by either allowing a finite air gap between a cylinder and one of the metal plates or by adding an asymmetrically-placed collar to the cylinder (Ma *et al.*, 2015). Moving such a metallic collar relative to the metal plates can be understood as changing the sign of the “mass” term that opens up the gaps at the Dirac cones (see Eq. 17 in Sec. II.A.1) allowing for the straightforward and reconfigurable engineering of arbitrary topological domain walls. As proposed in (Cheng *et al.*, 2016), this could form the basis of a topological switch, in which the movement of metallic collars is used to switch the propagation path from one port to another port (see Figure 21 (b)-(d)).

Very recently, these ideas have also been extended to all-dielectric bianisotropic metamaterials (Slobozhanyuk *et al.*, 2016a, 2017), which may offer advantages for reducing (Ohmic) losses, scaling up to optical frequencies, and increasing compatibility with on-chip integration. In these systems, arrays of dielectric disks are carefully designed such that the electric and magnetic dipole modes are degenerate and electromagnetic duality is restored (Slobozhanyuk *et al.*, 2016a). The bianisotropy term is then introduced by adding a raised circular notch on one of the flat faces of the disk; when the disks are arranged into a hexagonal or triangular meta-crystal, this gaps out the Dirac points, leading to a photonic topological insulator. This has been experimentally realised in the microwave regime for a 2D array of ceramic disks (Slobozhanyuk *et al.*, 2017). Such arrays could also be layered to make a 3D system, analogous to a “weak” 3D topological insulator (Slobozhanyuk *et al.*, 2017), as discussed further in Sec. V.B.

6. Photonic structures with crystalline symmetries

As we have just seen, the coupling of electric and magnetic fields in bianisotropic materials requires subwavelength structures that are asymmetric in the direction perpendicular to the plane of the topological metacrystal. Due to the small size of the features required for their fabrication, their implementation remains challenging for visible and near-infrared wavelengths. A different method to implement an analogue of the quantum spin Hall effect for photons in two-dimensions was proposed by L.-H. Wu and X. Hu (Wu and Hu, 2015). The configuration is based on subwavelength dielectric structures with inversion symmetry with respect to the 2D plane, which is in principle easier to implement experimentally.

The idea is to consider the lowest TM mode of a slab of cylindrical sub-wavelength dielectric rods surrounded by air and confined between two metallic plates (see Fig. 22(b)). The rods are arranged in a honeycomb geometry resulting in a photonic Dirac dispersion at K and K' points. If instead of taking the usual two-sites unit cell




FIG. 22 (a)-(c) Scheme of the lattice of hexagonal clusters of rods with dielectric constant ϵ_d . (d)-(f) Energy-momentum dispersion corresponding to expanded (d), centered-honeycomb (e), and contracted (f) clusters, showing in the latter case the opening of a topological gap. Panels (b), (d)-(f) from Wu and Hu, 2015.

of the honeycomb lattice, we consider a hexagonal cluster of rods as the unit cell, the additional band folding translates the Dirac points to the center of the Brillouin zone, resulting in a doubly degenerate Dirac crossing at the Γ point (see Fig. 22(e)). Solving Maxwell’s equations for the lowest TM mode of the slab shows that the in-plane magnetic field distributions of the doubly degenerate lower Dirac bands present a p_x, p_y character, while the upper Dirac bands contain states with $d_{xy}, d_{x^2-y^2}$ symmetry. Their symmetric and antisymmetric combinations, p_{\pm} and d_{\pm} , constitute a pseudospin basis (Fig. 22(e), red and blue lines, respectively).

When shifting the rods towards the center of the hexagonal clusters, the dispersion shows the opening of a trivial gap at the Dirac point, with the upper and lower bands preserving their symmetry (Fig. 22(a),(d)). If the rods are pushed away from the center of each cluster a gap opens again, but this time it is the result of a band inversion at the Dirac point, with the new states having d/p character for the lower/upper bands (Fig. 22(c),(f)) (Wu and Hu, 2015; Xu *et al.*, 2016). The bands possess now nonzero pseudospin Chern numbers, analogue to the case of an electronic \mathbb{Z}_2 topological insulator. If we now consider a ribbon of the topologically nontrivial photonic crystal, on a given edge, two bands of edge states with opposite group velocities traverse the gap around the Γ point. The wavefunctions of each edge band are associated to a specific pseudospin. A similar situation takes place at the interface between two photonic crystals with trivial/nontrivial arrangements.

We can relate these topological features to a photonic *pseudo* quantum spin Hall effect. Indeed, while Maxwell’s equation respect bosonic time reversal symmetry operations ($\mathcal{T}^2 = +1$), in the presence of the C_{6v} symmetry, we can construct an antiunitary opera-

tor $\tilde{\mathcal{T}}^2 = (\mathcal{K}\mathcal{T})^2 = -1$ in which \mathcal{K} respects the parity of p and d eigenstates with respect to $\pi/2$ and $\pi/4$ rotations, respectively (Wu and Hu, 2015). This pseudo time-reversal symmetry provides a Kramers doublet at the expense of keeping the $C_{6\nu}$ geometry of the lattice. In particular, the breaking of the crystalline order at the edge of the ribbon couples the two pseudospins and results in the opening of a very small gap at the crossing of the edge state bands at the time-reversal symmetric point at the Γ point.

The simplicity of this proposal has triggered its implementation in a wide variety of systems, ranging from acoustics (Yves *et al.*, 2017b) to phononics (Brendel *et al.*, 2017; He *et al.*, 2016a; Xia *et al.*, 2017). In optics it has been implemented for microwaves (Yang *et al.*, 2018; Yves *et al.*, 2017a), and for infrared photons in a photonic crystal slab (Anderson and Subramania, 2017; Barik *et al.*, 2018, 2016; Zhu *et al.*, 2018b) and metasurfaces (Gorlach *et al.*, 2018). In the experiments of Barik *et al.*, 2018 it was shown that the two pseudospins characterizing each interface state band correspond to opposite circular polarizations of the confined photons. A semiconductor quantum dot embedded in the photonic crystal was used to couple single photons to the topological interface states. In a different configuration, using coupled waveguides, it has been shown that the lattice configuration with the non-trivial gap can hold localised zero-dimensional defect states (Noh *et al.*, 2018). All these features open interesting perspectives in views of engineering novel whispering galling mode geometries (Siroki *et al.*, 2017) or exploring quantum chiral optics (Lodahl *et al.*, 2017).

7. Other proposals

Two alternative routes to simulating an artificial gauge field for a coupled optical cavity array have been proposed in (Umucalılar and Carusotto, 2011), by harnessing the polarization degree of freedom for light. In the first scheme, a nontrivial tunneling phase between lattice sites is generated by coupling together the orbital and polarization degrees of freedom, through, for example, a suitable embedding of either birefringent slabs or optically-active layers within an array of distributed Bragg reflector microcavities. In the second scheme, photons move in a single planar microcavity, where a suitable periodic lateral patterning of the cavity generates both a confining lattice potential but also a position-dependent polarisation mixing. As a result, the polarization of a photon then traces a closed loop in polarization-space when it evanescently tunnels between lattice wells, and so gains a geometrical Berry phase; this effect corresponds to a generalization for evanescent waves of how propagating photons can be imprinted with geometrical Pancharatnam (Pancharatnam, 1956) or Berry phases (Chiao *et al.*,

1988; Tomita and Chiao, 1986). This second scheme, in particular, offers potential for reaching the strongly-interacting photon regime (see Sec. VII.B), if the lateral patterning is scaled down to the micrometer scale, where the tighter confinement of light within the lattice wells will lead to greatly enhanced photon-photon interactions.

C. Anomalous Floquet topological insulators

The third main class of 2D topological photonic systems are the so-called anomalous Floquet topological insulators, where unusual topological properties can emerge due to periodically driving the system in time. In this subsection, we will set the focus on the two recent experiments of Refs. (Maczewsky *et al.*, 2017; Mukherjee *et al.*, 2017b), which realized such topological properties experimentally by designing suitable propagating waveguide arrays for photons.

As already discussed in Section II.A.5, topological band structures can be engineered through Floquet engineering, namely, by subjecting a system to a time-periodic modulation. In two dimensions, an emblematic example is provided by the “Floquet Chern insulator”, which can be realized by subjecting a honeycomb (graphene-like) lattice to a circular shaking (either produced by a mechanical modulation of the lattice (Jotzu *et al.*, 2014; Rechtsman *et al.*, 2013b), or by irradiation (Lindner *et al.*, 2011; Oka and Aoki, 2009)). In the high-frequency regime of the drive, i.e. when $\hbar\Omega$ is much larger than any other energy scale in the system, the dynamics is well captured by an effective Hamiltonian, whose band structure (the “Floquet” spectrum) exhibits Bloch bands with non-zero Chern numbers [Section II.A.5]. In this high-frequency regime, the topological characterization of the driven system reduces to applying the standard topological band theory to the effective Hamiltonian (i.e. to the Floquet band structure); in particular, the usual bulk-edge correspondence guarantees that a Floquet Chern insulator exhibits chiral edge modes at its boundaries. This approach was pioneered in photonics (Rechtsman *et al.*, 2013b), where a 2D honeycomb-shaped array of optical waveguides was circularly modulated along a third spatial direction playing the role of “time”.

This simple topological characterization breaks down when the bandwidth of a Floquet Chern insulator approaches $\hbar\Omega$ (i.e. when the system no longer operates in the high-frequency regime $\Omega \rightarrow \infty$). This can be understood in two ways. First, let us recall that the Floquet spectrum, which is associated with the Floquet operator $\hat{U}(T) = e^{-(i/\hbar)T\hat{H}_{\text{eff}}}$, is only uniquely defined within a Floquet-Brillouin zone $[-\hbar\Omega/2; \hbar\Omega/2]$; consequently, when the effective bandwidth approaches $\hbar\Omega$, gap-closing events are possible at the Floquet-Brillouin zone’s bound-




FIG. 23 (a) Sketch of the Rudner *et al.* toy model for anomalous Floquet topological phases (Rudner *et al.*, 2013), where hopping on a square lattice is activated in a cyclic and time-periodic manner. (b) Energy bands and sketch of the corresponding states, in the ideal case where the activated-hopping amplitude J is exactly set on resonance, $J = \hbar\Omega$, where Ω is the drive frequency. The dispersionless bulk band (blue) corresponds to cyclotron-localized states and is associated with a zero Chern number; dispersive branches (green and red) correspond to propagating edge states (skipped-cyclotron orbits). These edge states are topologically protected by a winding number that takes the full-time-evolution (including the micro-motion) into account (Nathan and Rudner, 2015). (c) The optical waveguide implementation of Ref. (Mukherjee *et al.*, 2017b), where the sequential activation of neighboring coupling was finely engineered. (d) Experimental evidence for the simultaneous existence of robust propagating edge states and (quasi) localized bulk states. Panels (a,b) adapted from Ref. (Rudner *et al.*, 2013) and (c,d) from Ref. (Mukherjee *et al.*, 2017b).

aries (Kitagawa *et al.*, 2010a). This can produce a cancellation of the Chern numbers associated with the Floquet bands, however, and this is a crucial observation, topological chiral edge states can still be present in the inner part of the spectrum. Such a co-existence of chiral edge modes with *seemingly trivial* Bloch bands, which is in apparent contradiction with the bulk-edge correspondence, is referred to as “anomalous Floquet topological phases” (Rudner *et al.*, 2013), as opposed to the standard Floquet Chern insulators discussed above. A second, and more fundamental, reason for this breakdown of the usual topological characterization stems from the fact that the micro-motion plays a crucial role as soon as the period of the drive T no longer sets the shortest time scale in the problem: in the “low-frequency” regime, the dynamics (including the topological nature of the system) cannot possibly be ruled by \hat{H}_{eff} only; see Section II.A.5. All together, this indicates that the “anomalous Floquet topological phase” cannot be accurately characterized by the Chern number related to the effective Hamiltonian,

but rather, by a distinct topological invariant: a “winding number”, which fully takes the micro-motion into account (Carpentier *et al.*, 2015; Kitagawa *et al.*, 2010a; Nathan and Rudner, 2015; Rudner *et al.*, 2013).

An instructive toy model leading to a dramatic instance of an anomalous Floquet topological phase was introduced in (Rudner *et al.*, 2013). In a certain parameter regime, the system simultaneously exhibits a *single completely flat* Bloch band (with a trivial Chern number) together with a topologically-protected chiral edge mode. As represented in Fig. 23(a), the driven system consists in a 2D square lattice, whose allowed hopping events are activated in a sequential and T -periodic manner: the time-evolution operator describes a four-step quantum walk. One should note that the circular nature of the sequence imprints a chirality to the system, similarly to the circular shaking leading to Floquet Chern insulators in graphene-like lattices (Jotzu *et al.*, 2014; Lindner *et al.*, 2011; Oka and Aoki, 2009; Rechtsman *et al.*, 2013b); such a chirality is crucial for the emergence of chiral

edge modes in the system. The effective (Floquet) band structure of this model is shown in Fig. 23(b), for the extreme case where the activated-hopping amplitude J is exactly set on resonance, $J = \hbar\Omega$. The resulting unique (two-degenerate) Bloch band is completely flat, which reflects the fact that the stroboscopic motion, i.e. the motion over each period of the drive T , is necessarily restricted to closed loops in any region of the bulk; however, as in the quantum Hall effect, such “orbits” reduce to skipped-orbits at the boundaries, hence resulting in a chiral edge mode [see Fig. 23(b)]. This simple picture is confirmed by the topological analysis of the model, which associates a non-trivial winding number to the Floquet Bloch band (Rudner *et al.*, 2013).

Recently, two independent experimental teams (Maczewsky *et al.*, 2017; Mukherjee *et al.*, 2017b) reported on the realization of this intriguing “Rudner-toy” model, using an array of laser-inscribed coupled waveguides using the technique described in (Szameit and Nolte, 2010). Both teams considered a 2D array of optical waveguides, defined in the $x - y$ plane, with a propagation direction z playing the role of time. As illustrated in Fig. 23(a), the model relies on a sequential activation of neighboring coupling within this 2D square lattice; in Refs. (Maczewsky *et al.*, 2017; Mukherjee *et al.*, 2017b), the experimentalists achieved this goal by spatially modulating the waveguides along the z direction, in such a way that different pairs of waveguides are locally moved together (to activate the coupling) and then apart (to switch it off); see Fig. 23(c) for a sketch of this protocol. These independent teams implemented two distinct configurations of the model, the activated couplings being homogeneous in Ref. (Maczewsky *et al.*, 2017), while these were chosen to be inhomogeneous in Ref. (Mukherjee *et al.*, 2017b); however, both experiments reached the aforementioned “anomalous” regime of the topological phase diagram, where chiral edge modes are uniquely determined by the non-trivial winding number of Refs. (Kitagawa *et al.*, 2010a; Nathan and Rudner, 2015; Rudner *et al.*, 2013). In Ref. (Mukherjee *et al.*, 2017b), the realization of the anomalous Floquet topological phase was demonstrated through a thoughtful study of the activated-coupling strength (which uniquely identified the realized “anomalous” phase within the topological phase diagram) combined with direct observations of the chiral edge states propagation and bulk localization [see Fig. 23(d)]; this analysis was further validated through numerical simulations based on the theoretical model. In Ref. (Maczewsky *et al.*, 2017), the “anomalous” phase was also signaled by demonstrating the dispersionless nature of the bulk (i.e. the existence of a flat band) and the chiral nature of the edge mode; this latter work also analyzed a topological transition from the “anomalous” topological phase to a trivial phase (characterized by the absence of edge mode) by designing

lattices with decreasing coupling strengths. These two experiments demonstrated the high tunability offered by laser-inscribed photonic crystals, in view of designing intriguing toy models and simulating exotic phases of matter.

Various other schemes have been proposed to reach the anomalous regime of Floquet topological systems, which can be applied to a variety of physical platforms (Kitagawa *et al.*, 2010a,b; Leykam *et al.*, 2016; Quelle *et al.*, 2017; Reichl and Mueller, 2014). In this broader context, robust localized states, associated with non-trivial winding numbers (Kitagawa *et al.*, 2010b), were first demonstrated in a photonic setup realizing a 1D quantum walk (Kitagawa *et al.*, 2012); a similar setup was recently explored in view of directly extracting winding numbers through Zak-phase measurements performed in the bulk (Cardano *et al.*, 2017). Besides, the winding number of 2D anomalous Floquet topological insulators was also measured in a microwave network, using a dimensional-reduction (topological pump) approach (Hu *et al.*, 2015). Finally, the existence of anomalous Floquet edge modes was also shown in a designer surface plasmon structure operating in the microwave regime (Gao *et al.*, 2016a).

D. Topology in gapless photonic systems

Another important class of topological systems are gapless photonic lattices with Dirac points. The primary example of this kind is the honeycomb lattice of coupled photonic resonators or waveguides. The Hamiltonian describing the dynamics of photons in these lattices is equivalent to that of p_z electrons in graphene, giving rise to two bands with linear crossings, at the Dirac points, as illustrated in Fig. 24(d) for a lattice of polariton resonators. In the absence of external fields, spin-orbit coupling or temporal modulation the system remains ungapped. Nevertheless, this type of lattice presents features that are topological in the sense that they can be related to certain topological invariants or geometrical properties of the system. These features include topological edge states, topological phase transitions, and the emergence of synthetic gauge fields when suitably deforming the lattice.

To analyze these topological properties, let us consider the hopping of the lowest photonic mode of each individual resonator to its nearest neighbor. In the tight-binding approximation with nearest neighbour hoppings, the Hamiltonian in momentum space is chiral and takes the form of Eq. 19 in the A, B sublattice basis depicted in Fig. 25(h) by green and blue dots. In this case, $Q(\mathbf{k})$ in Eq. 19 takes the form $Q(\mathbf{k}) = -t_1 e^{i\mathbf{k}\cdot\mathbf{R}_1} - t_2 e^{i\mathbf{k}\cdot\mathbf{R}_2} - t_3 e^{i\mathbf{k}\cdot\mathbf{R}_3} \equiv |Q(\mathbf{k})| e^{-i\theta(\mathbf{k})}$, where $t_{1,2,3}$ are the nearest neighbor hopping amplitudes and $\mathbf{R}_{1,2,3}$ are the vectors connecting a site to its three nearest neighbors (see Fig. 25(h)). For equal hoppings ($t_1 = t_2 = t_3 \equiv t$) the

two bands of eigenvalues of the honeycomb Hamiltonian are $\epsilon(\mathbf{k}) = \pm t|Q(\mathbf{k})|$ (Wallace, 1947), resulting in Dirac-like crossings at the K, K' points in the first Brillouin zone. The eigenfunctions are $|u_{k,\pm}\rangle = (1/\sqrt{2})(e^{-i\theta(\mathbf{k})}, \pm 1)^\dagger$.

The honeycomb Hamiltonian has been implemented in a number of photonic systems including microwave resonators (Bellec *et al.*, 2013a; Bittner *et al.*, 2010), photo-refractive crystals (Peleg *et al.*, 2007; Song *et al.*, 2015; Sun *et al.*, 2018), atomic vapour cells (Zhang *et al.*, 2018), coupled microlasers (Nixon *et al.*, 2013), polariton lattices (Jacqmin *et al.*, 2014; Kusudo *et al.*, 2013) and propagating waveguides (Plotnik *et al.*, 2014; Rechtsman *et al.*, 2013c). Some examples are shown in Fig. 24.

The first noticeable feature of the eigenstates is their pseudospin structure, with two components that reflect the underlying A, B interlaced triangular sublattices. The pseudospin structure results in particular scattering properties close to the Dirac cones: using a honeycomb lattice imprinted in a photo-refractive medium it has been shown that the conical diffraction characteristic of Dirac crossings (Peleg *et al.*, 2007) presents an orbital angular momentum $l = \pm 1$, depending on whether sublattice A or B is excited (Song *et al.*, 2015).

A second relevant feature of the eigenfunctions is their nontrivial Berry phase: if the eigenfunctions are transported adiabatically on a close loop in momentum space around one of the Dirac points, the eigenfunctions change sign (Castro Neto *et al.*, 2009). In other words, they get a Berry phase of $\pm\pi$, the sign being opposite for the K and K' points. This effect is present even when a gap is opened at the Dirac cones, which can be induced by introducing an on-site energy difference Δ between the A and B sublattices. In this case, a non-zero Berry curvature extends around the Dirac points. Therefore, if a wave packet is created close to one of the Dirac points and subject to acceleration, the Berry curvature results in an anomalous velocity whose sign depends on the Dirac point around which the wavepacket is created (Ozawa and Carusotto, 2014). An efficient way of accelerating the photon wavepacket is to design a lattice whose resonators continuously increase in size from one lattice site to the next: the decreasing photon confinement results in an on-site energy gradient. If the photon lifetime is long enough, the force can induce magnetic Bloch oscillations with a displacement perpendicular to the gradient and a direction determined by the Dirac point around which oscillations take place (Cominotti and Carusotto, 2013).

Photonic simulators can also be used to explore other properties of propagating wavepackets in a honeycomb lattice. For instance, the chiral symmetry of the honeycomb Hamiltonian is preserved in the presence of a potential step, resulting in phenomena like Veselago lensing (Cheianov *et al.*, 2007), Goos-Hänchen effect (Grosche *et al.*, 2016) or Klein tunneling (Dreisow *et al.*, 2012; Solnyshkov *et al.*, 2016b). Dissipation,

present for example in polariton lattices, does not significantly affect these phenomena (Ozawa *et al.*, 2017). Dirac cones can also be used to tailor the dispersion of photonic structures, even producing “epsilon-near-zero” materials (Huang *et al.*, 2011; Moitra *et al.*, 2013).

Edge states in Dirac systems - One of the characteristics of the honeycomb lattice is the existence of zero-energy edge states in finite size ribbons (Klein, 1994; Nakada *et al.*, 1996). These edge states are topological in the sense that they are related to the winding properties of the bulk Hamiltonian (Ryu and Hatsugai, 2002). To understand this bulk-edge relation, let us consider a graphene ribbon of finite size in the direction perpendicular to the edge (\perp), and infinite in the parallel direction (\parallel). By Fourier transforming the real-space Hamiltonian along this axis, we can reduce it to an effective 1D Hamiltonian for each value of k_{\parallel} (Castro Neto *et al.*, 2009; Delplace *et al.*, 2011). This effective 1D chiral Hamiltonian has the same form as the SSH Hamiltonian discussed in Sec. II.A.3, characterized by the complex function $Q(k_{\perp}, k_{\parallel}) = |Q(k_{\perp}, k_{\parallel})| e^{i\theta(k_{\perp}, k_{\parallel})}$. We can then apply the topological arguments discussed in Sec. II.A.3. The number of pairs of zero-energy edge states, appearing at the two edges of the ribbon, is thus determined by the winding of $\theta(k_{\perp}, k_{\parallel})$, Eq. 20, along the k_{\perp} direction over the first Brillouin zone (Delplace *et al.*, 2011; Mong and Shivamoggi, 2011; Ryu and Hatsugai, 2002):

$$\mathcal{W}(k_{\parallel}) = \frac{1}{2\pi} \int_{BZ} dk_{\perp} \frac{d\theta(k_{\perp}, k_{\parallel})}{dk_{\perp}}. \quad (67)$$

The information about the specific type of edge is contained in the choice of unit cell dimer and unit vectors when writing down the honeycomb Hamiltonian, such that they allow for the full reconstruction of the lattice (including the edges). Therefore, the information on the edges is reflected in the specific form of $\theta(k_{\perp}, k_{\parallel})$ when writing down Eq. 19 (Delplace *et al.*, 2011). Figure 25(b) and (e) show $\theta(k_{\perp}, k_{\parallel})$ for zigzag and bearded edges, respectively. The colored areas indicate the values of k_{\parallel} for which $\mathcal{W} = 1$, corresponding to the existence of edge states.

The direct access to the wavefunctions in photonic lattices has been employed to study the local properties of these edge states. Their existence has been evidenced experimentally in lattices of coupled waveguides (Plotnik *et al.*, 2014), microwave resonators (Bellec *et al.*, 2014; Bittner *et al.*, 2012) and polaritons (Milićević *et al.*, 2015), showing that edge states for zigzag and bearded terminations connect Dirac cones in complementary regions in momentum space (Fig. 25(b) and (e)), while armchair terminations do not possess any edge state ($\mathcal{W} = 0$ for any k_{\parallel}).

The topological arguments used to predict the existence of edge states in a honeycomb lattice can be extended to other Hamiltonians, for instance, when more




FIG. 24 (a) Image of a honeycomb lattice of microwave resonators with armchair, bearded and zigzag edges. (b) Measured density of states as a function of microwave frequency ν . The peak at the Dirac energy ν_d indicates the presence of zero energy states. From Bellec *et al.*, 2014. (c) Scanning electron microscope image of a honeycomb lattice of coupled polariton micropillars. (d) Photoluminescence spectrum measured at the center of the lattice showing Dirac crossings (left), and at the zigzag edge (right), showing a flat band of edge states (pointed by the arrow). From Milićević *et al.*, 2015. (e) Representation of a lattice of coupled waveguides and (f) measured momentum space distribution of the transmission through the zigzag and bearded edges. Red lines show the position of the Dirac points. From Plotnik *et al.*, 2014.

than one mode per site is involved. As long as the system possesses the chiral symmetry, they can be written in the form of Hamiltonian 19. With more than one mode per site, $Q(\mathbf{k})$ takes the form of a $n \times n$ matrix whose determinant can be written as $\det Q(k) \equiv |\det Q(k)|e^{i\theta(k)}$. The existence of pairs of zero energy edge states is again given by the winding of $\theta(\mathbf{k})$ along the momentum direction perpendicular to the edge (Kane and Lubensky, 2014). An example of this kind of chiral Hamiltonians is the p -orbital version of graphene, in which orbitals with p_x, p_y geometry are considered at each lattice site (Wu *et al.*, 2007). This 4×4 orbital Hamiltonian has been implemented in a polariton honeycomb lattice when considering the doubly degenerate first excited states of each coupled micropillar. The spectrum consists of four bands with Dirac crossings, and the presence of zero-energy edge states for different kinds of terminations is well accounted for by the analysis of the winding of $\theta(\mathbf{k})$ we have just presented (Milićević *et al.*, 2017).

Similar arguments can be applied to Dirac Hamiltonians without chiral symmetry, for instance with next-nearest neighbor hopping or with a staggered potential in the A, B sublattices, both effects giving rise to non-zero diagonal terms in Eq. 19. The presence of edge states can also be determined via winding arguments, but their energy is not necessarily zero (Mong and Shivamoggi, 2011).

Valley Hall edge states - Propagating edge states with weak topological protection can be engineered in lattices

with appropriate staggered potentials. The staggered potential between the A, B sublattices Δ breaks the inversion symmetry and opens a gap at the Dirac points. By integrating the Berry curvature around each Dirac point, we can define a *valley* Chern number whose sign is opposite for K and K' points (the total Chern number of the band still being zero). If the staggered potential is changed to $(-\Delta)$, the gap is still open but the signs of the *valley* Chern numbers switch between K and K' points. When joining two honeycomb semi-infinite ribbons subject to opposite staggered potentials Δ and $-\Delta$, it has been shown that interface states appear in two bands that traverse the gap (Bleu *et al.*, 2017c; Chen and Dong, 2016; Goldman *et al.*, 2016b; Ma and Shvets, 2016; Qiu *et al.*, 2017; Weinstein *et al.*, 2016; Zhang *et al.*, 2011). Indeed, if the K and K' valleys of the two lattices have opposite valley Chern numbers, the gap needs to close locally at those points at the interface between the two lattices (Ma and Shvets, 2016). This situation is restricted to very specific conditions, for instance, it applies to zigzag interfaces. For this configuration, the propagation of a wavepacket in an interface state with a given valley polarization is protected against any perturbation that does not mix the two valleys, i.e., bends of the interface of 120° , which preserve the zigzag character of the interface (Chen and Dong, 2016; Ma and Shvets, 2016; Wu *et al.*, 2017a). Interface states between photonic crystals which exhibit valley-Hall effect and quantum spin-Hall effect have also been experimentally demonstrated (Kang *et al.*, 2018).




FIG. 25 Stream plots of $\theta(k_{\perp}, k_{\parallel})$ in momentum space for zigzag, bearded and armchair edges, and for different values of the homogeneous strain t_1/t_2 ($t_2 = t_3$). The colored areas show the regions of k_{\parallel} in which $\mathcal{W} = 1$, corresponding to the presence of edge states. Black solid and dashed lines represent the 1D Brillouin zones in k_{\parallel} and k_{\perp} , respectively. From Bellec *et al.*, 2014. Red points show the position of the Dirac points. (h) Scheme of the honeycomb lattice and nearest neighbor hoppings.

For armchair interfaces, the breaking of translational symmetry in the direction perpendicular to the interface mixes the K and K' points of each lattice, and the valley Chern number is not well defined. Similarly, if the gap becomes too large ($|\Delta| \gtrsim t$) in zigzag interfaces, the Berry curvatures associated to both valleys overlap. The valley Chern numbers are not well defined anymore and a gap opens in the edge state bands (Noh *et al.*, 2017a).

Despite the fact that valley Hall edge states are not robust against arbitrary spatial disorder at the interface, they can be used to route photons in photonic structures (Noh *et al.*, 2017a; Wu *et al.*, 2017a), and to design delay cavities in Si photonics technologies (Ma and Shvets, 2016). A route to recover topological protection is to use chiral nonlinear excitations (i.e. vortices) as proposed recently for a polariton fluid in a honeycomb lattice (Bleu *et al.*, 2017a).

Effect of strain - The application of strain to an ungapped honeycomb lattice ($\Delta = 0$) strongly modifies its spectrum and eigenfunctions and, consequently, its topological properties. We can consider two main classes of strain: (i) homogeneous strain, in which hopping is different along different spatial directions ($t_1 \neq t_2 \neq t_3$), and (ii) inhomogeneous strain, in which hopping takes

different values at different positions ($t_i(\mathbf{r})$). The first case was theoretically studied by Montambaux and co-workers (Montambaux *et al.*, 2009). They predicted a topological phase transition occurring when one of the three nearest neighbor hopping amplitudes (t_1) is twice as large as the other two ($t_2 = t_3$). At the transition point ($t_1 = 2t_2$), the two non-equivalent Dirac cones merge and disappear resulting in the opening of a full gap. This transition was first experimentally observed in cold atoms (Tarruell *et al.*, 2012), but it is in photonic lattices where its effect on the existence of edge states has been studied (see Fig. 25). Experiments in lattices of microwave resonators and coupled waveguides (Bellec *et al.*, 2014; Rechtsman *et al.*, 2013a), have shown that above the transition point ($t_1 > 2t_2$), ribbons with zigzag terminations contain a flat energy band of edge states covering the whole momentum space, while for bearded boundaries, edge states disappear. For armchair terminations, edge states appear for any value of the unidirectional strain as long as the anisotropy axis is not parallel to the edge, as in Fig. 25(g),(i) (Bellec *et al.*, 2013b, 2014). The existence of edge states in homogeneously strained honeycomb lattices can also be predicted from topological arguments. The Hamiltonian including this kind of strain still possesses the chiral symmetry, and the number of zero energy edge states is governed by the winding of $\theta(\mathbf{k})$. Interestingly, homogeneous strain has been employed to synthesise critically tilted Dirac cones in the orbital p-bands of lattices of coupled micropillars (Milićević *et al.*, 2018).

The second kind of strain concerns the continuous variation of the hopping over the lattice. Originally studied in the context of electronic graphene, this kind of inhomogeneous strain has been shown to induce a gauge field (Kane and Mele, 1997). If the strain takes the specific trigonal shape $(u_r, u_{\theta}) = \beta r^2(\sin 3\theta, \cos 3\theta)$, where




FIG. 26 Tight binding calculation of the level structure of a photonic graphene ribbon as a function of k_y , perpendicular to the strain direction. The hopping strength along the narrow dimension of the ribbon increases linearly from one edge to the other, resulting in a gauge field for photons. The Dirac points, located at $k_y = 1.2/a$ with a being the nearest-neighbor distance, split into Landau levels (red lines). The lowest one, $n_L = 0$ at $E = 0$, is flat, while the others have a non-zero group velocity. From Salerno *et al.*, 2015.

u_r, u_θ are the real space displacements of the positions of the carbon atoms in polar coordinates, the modified hopping induces a gauge field for electrons at the Dirac cones corresponding to a homogeneous pseudo-magnetic field perpendicular to the graphene sheet (Guinea *et al.*, 2010). In other words, the Hamiltonian describing the time evolution of electrons in the strained lattice can be cast in the form of an unstrained honeycomb Hamiltonian subject to a homogeneous pseudo-magnetic field perpendicular to the graphene plane. Note, however, that this field does not break time-reversal symmetry: it has opposite signs at the two Dirac points. Therefore, edge states appearing in the gaps between consecutive Landau levels are associated to propagation on both directions along the edge (Gopalakrishnan *et al.*, 2012; Salerno *et al.*, 2017) and backscattering-protected transport is not expected.

Translated to the photonic realm, this configuration provides a very efficient way of inducing a pseudo-magnetic field acting on photons as if they were charged particles. This precise idea was implemented by Rechtsman *et al.*, 2013c in a lattice of coupled waveguides by varying continuously their separation (i.e., the nearest-neighbor photon hopping) following the above mentioned strain configuration. The value of the valley dependent pseudo-magnetic field acting on the propagating photons can take effective values of several thousands Tesla, much higher than strengths of real magnetic field currently realizable in the laboratory. The main consequence is the emergence of Landau levels in the vicinity of the Dirac points (Castro Neto *et al.*, 2009). Analogously to the effect of a real magnetic field in graphene, the energy of the photonic Landau levels n_L scales as $\pm\sqrt{n_L}$. This was observed in numerical tight-binding calculations and experimentally via the localization of a wavepacket on the edge of a strained lattice (Rechtsman *et al.*, 2013c), attesting both the presence of flat Landau bands and of states localized at the edge, emerging from the gauge field.

The trigonal strain discussed so far is not the only way of inducing a homogeneous magnetic field. Salerno *et al.*, 2015 showed that linear uniaxial strain along one of the crystallographic directions results in a homogeneous pseudomagnetic field similar to that emerging from trigonal strain. Figure 26 shows the emergence of Landau levels associated to pseudo-magnetic fields of opposite sign at the K, K' points. The most noticeable feature in this tight-binding calculation is that, except for $n_L = 0$, the Landau levels are not completely flat. The reason is that the position dependent hopping results in a local Dirac velocity that varies along the lattice (de Juan *et al.*, 2012).

The engineering of Landau levels in photonic structures is particularly interesting in the quest for confined lasing geometries. The possibility of introducing flat gapped bands in the bulk of a photonic lattice, with a

high density of states, could be used to fabricate low threshold on-chip lasers. Moreover, the combination of gauge fields with gain and losses provides exciting perspectives on the study of the parity anomaly and sublattice selective lasing (Schomerus and Halpern, 2013).

IV. TOPOLOGICAL PHOTONICS IN ONE DIMENSION

The previous section was devoted to a review of two-dimensional photonics systems where topological concepts were first investigated. In the present section, we now turn our attention to one-dimensional models in Sec. IV.A and then, in Sec. IV.B, to the topological pumping effects that such systems have been used to study.

A. Topology in 1D chiral systems

In one dimension, topological phases of matter cannot exist without imposing symmetries on the system (Kitaev *et al.*, 2009; Schnyder *et al.*, 2009). An important symmetry in one dimension, which can lead to topological phases, is chiral symmetry, for which the representative one-dimensional topological model is the Su-Schrieffer-Heeger (SSH) model as introduced in Sec. II.A.3.

The first experimental realization of the SSH model and its associated topological edge state in a photonics context was in a photonic superlattice (Malkova *et al.*, 2009). Since then, the SSH model or related models have been discussed and realized in photonic crystals (Keil *et al.*, 2013; Xiao *et al.*, 2014), electromagnetic metamaterials (Tan *et al.*, 2014; Yannopapas, 2014), plasmonic and dielectric nanoparticles (Kruk *et al.*, 2017; Ling *et al.*, 2015; Poddubny *et al.*, 2014; Sinev *et al.*, 2015; Slobozhanyuk *et al.*, 2015, 2016c), plasmonic waveguide arrays (Bleckmann *et al.*, 2017; Cherpakova *et al.*, 2018), polariton micropillars (Solnyshkov *et al.*, 2016a), and coupled optical waveguides (Naz *et al.*, 2017). Lasing in the edge state of the SSH model has recently been experimentally observed in Parto *et al.*, 2018; St-Jean *et al.*, 2017; and Zhao *et al.*, 2018; these experiments constitute the first realizations of topological lasers, i.e. lasers which make use of topological edge states as we discuss more in Sec. VIII.B. The interplay between the SSH model and the radiative loss of photons have also been discussed in photonic crystals (Poshakinskiy *et al.*, 2014; Schomerus, 2013), microwave cavity arrays (Poli *et al.*, 2015), and coupled waveguide arrays (Zeuner *et al.*, 2015). The SSH model of optical waveguides with periodic modulation and its edge states have been studied with Floquet theory (Zhu *et al.*, 2018a). The SSH model has also been realized with electric circuits (Lee *et al.*, 2018), and the Zak phase (Berry phase across the Brillouin zone) has been measured to show its topological

nature (Goren *et al.*, 2018). By properly adding loss, the PT symmetric version of the SSH model was realized in (Weimann *et al.*, 2017). A one-dimensional electric circuit has also been shown to exhibit quantized energy transport due to the topological nature of the system in the presence of dissipation (Rosenthal *et al.*, 2018). More details on non-Hermitian topological models are given in Sec. VI.A. There is also a proposal to realize the one-dimensional Jackiw-Rebbi model, introduced in Sec. II.A.1, in a driven slow-light setup (Angelakis *et al.*, 2014), where the zero-energy bound mode can be probed through the transmission spectrum.

Another strategy to realize 1D chiral Hamiltonians with non-trivial topology involves discrete-time quantum walks (Kitagawa, 2012; Kitagawa *et al.*, 2010b). As it was discussed in Sec. II.A.5, a discrete-time quantum walk consists of a repeated application of a set of operations represented by a unitary matrix \hat{U} . Topological properties of such unitary evolution can be understood analogously to Floquet topological phases by defining an effective Hamiltonian \hat{H}_{eff} through $\hat{U} = e^{-i\hat{H}_{\text{eff}}T/\hbar}$. If \hat{H}_{eff} has non-trivial topology, its effect can be detected through the discrete-time quantum walk. The great flexibility in choosing a set of operations to realize \hat{U} and hence \hat{H}_{eff} makes the discrete-time quantum walk a powerful platform to explore topological phases of matter.

A topologically-nontrivial discrete-time quantum walk was first experimentally realized by Kitagawa *et al.*, 2012 using single photons going through a series of polarization rotations. In the experiment, a topological bound state between the interface of regions with different winding numbers has been observed. Subsequently, Cardano *et al.*, 2017, 2016, 2015 have studied the topological invariant of a quantum walk in the orbital angular momentum space and detected the topological transition between different phases. Following the proposal (Tarasinski *et al.*, 2014), the topological invariant of a one-dimensional quantum walk was measured and its robustness to disorder was assessed, using a fibre loop architecture based on the time-multiplexing technique (Barkhofen *et al.*, 2017)

B. Topological pumps

Electric currents are usually generated by applying a voltage across a material, inducing longitudinal charge transport. Using Faraday's induction law, we can similarly generate electric currents via a time-dependent variation of a magnetic flux. In both cases, the longitudinal conductivity is determined by the microscopic details of the material and can take arbitrary values. As we have seen in Sec. II, the situation is completely different in topological systems, where currents can show quantization effects.

In particular in topological charge pumps (Thou-




FIG. 27 Experimental observation of adiabatic pumping via topologically protected boundary states in a photonic waveguide array. (a) An illustration of the adiabatically modulated photonic waveguide array, constructed by slowly varying the spacing between the waveguides along the propagation axis z . Consequently, the injected light experiences an adiabatically modulated Hamiltonian, $H_{\text{off}}(\phi(z))$, as it propagates across the sample. (b) The spectrum of the model Eq. (68) as a function of the phase ϕ for $t = 40/75$, $2t_{xy} = 0.6$, $b = (1 + \sqrt{5})/2$. In the experiment, a 21 sites lattice was used and ϕ was scanned between 0.35π and 1.75π , marked by arrows (and red dots). The insets depict the spatial density of a boundary eigenstate as a function of the position at three different stages of the evolution: At $\phi = 0.35\pi$, the eigenstate is localized on the right boundary. At $\phi = \pi$, it is delocalized across the system, while at $\phi = 1.75\pi$ the state is again localized, but on the left boundary. (c) Experimental results: Light was injected into the rightmost waveguide at $z = 0$ with $\phi = 0.35\pi$. (c) The measured intensity distributions as a function of the position are presented at different stages of the adiabatic evolution, i.e., different propagation distances. It is evident that during the adiabatic evolution, the light crosses the lattice from right to left and is finally concentrated on the left-most waveguides. All panels are taken from Ref. (Kraus *et al.*, 2012).

less, 1983), Faraday's induction law is encoded in an adiabatic cyclic variation of a Hamiltonian potential that mimics the magnetic-flux threading in a higher-dimensional topological Chern insulator. Consequently, the charge transport across the system per unit cycle

of the pump parameter turns out to be quantized in much the same way that quantized Hall conductance appears in the higher-dimensional static Chern insulator. This feature has drawn much attention for controlled low-current nanoscale device applications (Kouwenhoven *et al.*, 1991).

The first experiments towards the implementation of a topological charge pump were conducted in solid state systems with demonstrations of quantized charge transport using single electron pumps (Geerligs *et al.*, 1990; Kouwenhoven *et al.*, 1991; Pothier *et al.*, 1991, 1992). The quantization, however, of the pumped charge in these devices relied on simple Coulomb blockade rather than on topological concepts. Later attempts using open mesoscopic systems incorporated geometrical ideas to generate a quasi-adiabatic, non-quantized current that is proportional to the area enclosed in parameter space (Brouwer, 1998; Möttönen *et al.*, 2008; Spivak *et al.*, 1995; Switkes *et al.*, 1999; Zhou *et al.*, 1999). The challenge of realizing quantized topological pumping was only recently accomplished using cold atoms in optical superlattices (Lohse *et al.*, 2016; Nakajima *et al.*, 2016).

In this Section, we introduce realizations of topological pumps in photonic systems. Unitary photonic topological pumps have been realized using coupled waveguide arrays (Kraus *et al.*, 2012; Verbin *et al.*, 2015), whereas non-Hermitian pumps with exceptional points were realized using microwave cavities (Hu *et al.*, 2015). In both realizations, the experiments focussed on states localized at the system's boundary which were directly excited by the incident light. Similarly to the aforementioned cold-atom experiments (Lohse *et al.*, 2016; Nakajima *et al.*, 2016), theoretical proposals have addressed the possibility of studying quantized bulk pumping in photonic systems (Ke *et al.*, 2016; Mei *et al.*, 2015). Geometric pumping has been experimentally realized using a fiber loop architecture (Wimmer *et al.*, 2017).

In the following, we shall focus on the first photonic realizations of topological pumps using waveguide arrays (Kraus *et al.*, 2012). While it is technologically challenging to modulate the waveguide profile in a way to precisely realize the on-site modulation involved in Thouless' original topological pump model (29), the excellent control in the waveguide spacing achievable with femtosecond laser microfabrication technology (Szameit *et al.*, 2007; Szameit and Nolte, 2010) allowed for the emulation of an off-diagonal pump model where the inter-waveguide hopping amplitudes are slowly modified along the propagation axis z and the light evolves according to the Hamiltonian

$$\hat{H}_{\text{off}} = -J \sum_x \left[\left(1 + \frac{2J_{xy}}{J} \cos(2\pi\alpha x/a + \phi(z)) \right) \hat{a}_x^\dagger \hat{a}_{x+a} + \text{h.c.} \right]. \quad (68)$$

where J is the bare hopping amplitude from waveguide

to waveguide $n - 1$, $2J_{xy}$ is the amplitude of its z -dependent modulation (which is equivalent to time-dependent modulation in propagating geometries, cf. Section III.A.2), and α is a spatial modulation frequency, see Fig. 27(a) for an illustration.

The mapping between the 2D quantum Hall effect on a lattice and the 1D pump discussed in Sec II.A.4 and Eqs. (14) and (29) can be extended beyond the Harper-Hofstadter model (Kraus and Zilberberg, 2012). Performing dimensional extension on (68), we obtain a 2D tight-binding model where motion along x occurs via a standard nearest neighbor hopping in the x direction and motion along y only occurs via diagonal hoppings to next-nearest-neighbors (Hatsugai and Kohmoto, 1990)

$$\hat{H} = -J \sum_{x,y} \left(\hat{a}_{x+a,y}^\dagger \hat{a}_{x,y} + \frac{J_{xy}}{J} e^{i2\pi\alpha x/a} \hat{a}_{x+a,y+a}^\dagger \hat{a}_{x,y} + \text{H.c.} \right), \quad (69)$$

Each plaquette in the model is threaded by $2\pi\alpha$ flux as in the Harper-Hofstadter model, cf. Eq. (14).

The model (68) is commonly known as the off-diagonal Harper model (Jitomirskaya and Marx, 2012; Ketoja and Satija, 1997). Figure 27(b) depicts its spectrum as a function of ϕ . We observe a characteristic gapped structure with topological modes crossing the gaps as a function of ϕ . These modes are localized at the system's boundary when they are well within the energy gap, while they become spatially extended when they spectrally approach the bulk modes, see insets of Fig. 27(b).

In the experiment [Fig. 27(c)], light is injected via fiber coupling directly into the left end. The initial ϕ is chosen in a way to support a localized state on this boundary, so that the injected light can directly excite this state. The value of ϕ is then scanned along the propagation axis by correspondingly varying the inter-waveguide distances. Depending on the final value of ϕ , the spatial intensity distribution at the output will recover the spatial shape of the wavefunction of the topological state in different regimes, e.g. either extended over the bulk or even completely localized at the other boundary of the system. Such pumping through the boundary states of the pump highlights the existence of localized states on both ends of the system for suitable values of ϕ , in agreement with the pump's bulk-edge correspondence.

Moreover, it also illustrates the fact that the topological boundary mode on one end of the system can be connected to the state localized on the other end through semiadiabatic scanning of ϕ in sufficiently short systems. The latter is an interesting effect that is not implied by the bulk topology of the system: the bulk topology implies in fact that a quantized number of boundary states will cross the gap on each side of the sample as a function of ϕ , but does not necessarily imply that these states have to be directly connected.

Using the same technology and extending this idea further, a topological pump was realized for an off-diagonal Fibonacci chain. This realization relied on a mapping between quasiperiodic chains and topological pumps (Kraus *et al.*, 2012; Kraus and Zilberberg, 2012; Verbin *et al.*, 2015). Thus, using a two-parameter pumping, the Fibonacci chain was deformed into an off-diagonal Harper model, pumped as above, and deformed back into a Fibonacci chain (Verbin *et al.*, 2015). Furthermore, topological phase transitions between quasiperiodic chains with smooth boundaries were studied using photonic waveguide arrays (Verbin *et al.*, 2013). A study of the spectral flow of edge states across the energy gaps of a Fibonacci quasicrystal was reported in (Baboux *et al.*, 2017) by scanning a suitable structural parameter through many copies of a polariton lattice device.

Interestingly, simultaneous realizations of atomic and photonic 2D topological pumps were recently reported. Such pumps were shown to be directly mappable to a 4D quantum Hall system (Kraus *et al.*, 2013). While the atomic experiment (Lohse *et al.*, 2018) performed a direct mapping of the Berry curvature by looking at anomalous transport in the bulk of the system, the photonic experiment (Zilberberg *et al.*, 2018) studied the boundary states associated with a second Chern number response using similar methods to those discussed above.

V. TOPOLOGICAL PHOTONICS IN HIGHER DIMENSIONS

Having reviewed the photonic realizations of two- and one-dimensional topological models, we now briefly highlight very recent and on-going works in higher-dimensional topological systems. In Secs. V.A and V.B, we focus on the study of three-dimensional topological photonics, for which macroscopic photonic crystals and metamaterials operating in the microwave domain have provided the main experimental platform. Then, in Sec. V.C, we will discuss topological physics in even higher spatial dimensions, including perspectives in this direction opened up through the concept of “synthetic dimensions”.

A. Three-dimensional gapless phases

1. Weyl points and helicoid surface states

As briefly reviewed in Sec. II.A.3, three-dimensional band-structures can host Weyl points, corresponding to gapless points around which bands disperse linearly with respect to the three quasi-momenta (Eq. (22)). Each Weyl point can be associated with a topological Chern number, calculated by integrating the Berry curvature over a two-dimensional surface enclosing the Weyl point. It can be shown that Weyl points have Chern numbers




FIG. 28 (Color online) (a) Illustrations of Weyl points of opposite Chern numbers (C). (b) Surface dispersion near the projections of a pair of Weyl points with opposite Chern numbers, where the red and blue cones represent the bulk states projection at k_+ and k_- . The surface state is plotted using the Riemann sheet of $\text{Im}[\log[(k - k_+)/ (k - k_-)]]$ in the complex plane of k . The green surface arc is an iso-frequency contour. (c), (d), (e) are double-Weyl points of two bands, three bands and four bands. (f) A Dirac point consists of two Weyl points of opposite Chern numbers. Panel (b) taken from (Fang *et al.*, 2016a), and other panels adapted from (Zhang *et al.*, 2018).

of ± 1 [Fig. 28(a)], and require the breaking of either \mathcal{P} (parity) and/or \mathcal{T} (time-reversal) symmetry. If \mathcal{T} is broken, the minimum number of Weyl points in the band-structure is two, whereas, if only \mathcal{P} is broken, as is typically much easier to implement in experiments, then the minimum number of Weyl points is four. For a strong enough tilting of the Weyl cone, the group velocities of the two crossing bands can have the same sign along one direction, in which case one speaks of a Type-II Weyl point (Soluyanov *et al.*, 2015).

The topological character of Weyl points is reflected in the appearance of topologically-protected states on the surface of the three-dimensional system. These surface states are topologically equivalent to helicoid Riemann surfaces (Fang *et al.*, 2016a; Zhang *et al.*, 2018) defined with the two-dimensional surface Brillouin zone as the complex plane, shown in Fig. 28(b). A helicoid surface is a non-compact Riemann surface, which is unbounded in the frequency axis, corresponding to the gapless nature of the Weyl surface state. Locally around each Weyl cone, the surface states can be expressed as $\omega \propto \text{Im}[\log(k^C)]$, where C is the Chern number of the Weyl point. The bulk Weyl points project onto the surface Brillouin zone as poles ($C > 0$) and zeros ($C < 0$) of the multivalued helicoid surface sheets winding around these singularities. Their winding direction is determined by the sign of C , while the order of the pole or zero is given by $|C|$. As shown in Fig. 28(b), the isofrequency contours of the helicoid surface are always open arcs connecting the sur-

face projections of the positive and negative bulk Weyl points. These open surface arcs are known as “Fermi arcs” in Weyl semimetals.

Theoretically, Weyl points were first proposed to appear in double-gyroid photonic crystals, with a breaking of \mathcal{P} or \mathcal{T} (Lu *et al.*, 2013b). Since then, theoretical studies have shown that Weyl points could be realised in optical lattices (Dubček *et al.*, 2015a; Roy *et al.*, 2017b), photonic superlattices (Bravo-Abad *et al.*, 2015), magnetized plasmas (Gao *et al.*, 2016b), chiral metamaterials (Gao *et al.*, 2015; Liu *et al.*, 2017; Xiao *et al.*, 2016b), Floquet networks (Ochiai, 2016; Wang *et al.*, 2016a), chiral woodpile crystals (Chang *et al.*, 2017), magnetic tetrahedral crystals (Yang *et al.*, 2017d), and classical (Luo *et al.*, 2018) and quantum (Mei *et al.*, 2016) circuits. In an ideal Weyl system, all Weyl points would be frequency isolated and symmetry-related at the exact same frequency (Wang *et al.*, 2016c). We also note that ideal Weyl points move the classical free-space scattering laws from DC to the Weyl frequency by design (Zhou *et al.*, 2017a), and that, after including losses, Weyl points evolve into exceptional lines (Xu *et al.*, 2017).

Experimentally, Weyl points were demonstrated at microwave frequencies in a double-gyroid photonic crystal (Lu *et al.*, 2015), metallic photonic crystals with multi-Weyl points and surface transport (Chen *et al.*, 2016), photonic metamaterials with Type-II Weyl points and surface arcs (Yang *et al.*, 2017a), in lumped-element circuits (Lu *et al.*, 2018), and at optical frequencies in coupled waveguides with Type-II Weyl points and surface states (Noh *et al.*, 2017b). Synthetic Weyl points in the parameter space of 1D dielectric stacks were also observed in Wang *et al.*, 2017b, and ideal Weyl points have been found in a metallic design (Yang *et al.*, 2018). In this latter platform, the helicoid surface states of the four Weyl points were experimentally mapped out and were topologically equivalent to a Riemann sheet defined by the Jacobi elliptical function, analytical in the whole double-periodic surface Brillouin zone.

2. Multi-Weyl and Dirac points

A Weyl point, of non-zero Chern number, does not require any symmetry for protection, other than translational symmetry. With an increase of symmetry, multi-Weyl points can stabilize at high-symmetry momenta (Chang *et al.*, 2017; Chen *et al.*, 2016; Fang *et al.*, 2012a; Xu *et al.*, 2011). For example, double Weyl points (Zhang *et al.*, 2018) of Chern number of ± 2 can form between two bands as a quadratic Weyl point, between three bands as a spin-1 Weyl point, or between four bands as a charge-2 Dirac point, as shown in Fig. 28(c), (d) and (e). In the latter case, charge-2 refers to the Berry charge (Chern number) of 2, corresponding to the overlapping of Weyl points of the same Chern num-

ber. The double-Weyl surface states can be mapped, in the entire Brillouin zone, to the double-periodic Weierstrass elliptic functions: a type of Riemann surface with second-order poles and zeros (Zhang *et al.*, 2018).

More generally, a Dirac point in 3D refers to the overlapping of any two Weyl points of opposite Chern numbers, as shown, for example, in Fig. 28(f). Such 3D Dirac points were discussed in Lu *et al.*, 2016b, Slobozhanyuk *et al.*, 2016a, Wang *et al.*, 2017a, 2016b and Guo *et al.*, 2017. Since a 3D Dirac point has zero Chern number, it does not require the breaking of either \mathcal{P} or \mathcal{T} .

3. Nodal lines and surface

As well as the above point degeneracies (nodal points), line degeneracies are also important in 3D. Such nodal lines (Fang *et al.*, 2016b) can be protected by \mathcal{PT} symmetry with π Berry phase, same as the 2D Dirac cones. The nodal lines known so far can be classified into several families, namely nodal rings (Burkov *et al.*, 2011), nodal chains (Bzdusek *et al.*, 2016), nodal links (Yan *et al.*, 2017) and nodal knots (Bi *et al.*, 2017a). In photonics, a nodal ring was proposed in gyroid photonic crystals (Lu *et al.*, 2013b) and nodal chains were proposed in a FCC lattice (Kawakami and Hu, 2016) and discovered experimentally in a simple-cubic metallic photonic crystal (Yan *et al.*, 2018). Nodal lines can also exist in two-dimensional photonic crystals (Lin *et al.*, 2017), such as at the zone boundary of two dimensional lattices with glide reflection symmetry and \mathcal{T} . Nodal lines can also carry a Z_2 charge (Fang *et al.*, 2015).

Nodal surfaces can be protected by screw rotations and \mathcal{T} . It can even carry non-zero Chern numbers (Xiao and Fan, 2017b).

B. Three-dimensional gapped phases

Gapping topological degeneracies such as Weyl and Dirac points is the most effective way to obtain 3D band gaps supporting various topological interfacial states.

1. 3D Chern Insulator

As introduced in Sec. II.A.3, a three-dimensional quantum Hall system is characterised by a triad of first Chern numbers [$\mathbf{C}^{(1)} \equiv (C_x^{(1)}, C_y^{(1)}, C_z^{(1)})$], as indicated in Fig. 29(a). In such systems, the gapless surface states are unidirectional sheets, whose number and directionality equals the magnitude and sign of the Chern numbers for that surface (Koshino *et al.*, 2002). In the context of photonics, the elementary case of $\mathbf{C}^{(1)} = (0, 0, 1)$ was proposed by annihilating a single pair of Weyl points (a Dirac point), by supercell coupling, in the magnetic gyroid photonic crystals (Lu and Wang, 2016). Note that

as (first) Chern numbers vanish in the presence of time-reversal symmetry, the breaking of \mathcal{T} is required to realise such topological phases.

2. One-way fibers

One-way fiber modes can form along topological line-defects in 3D magnetic photonic crystals, illustrated in Fig. 29(b). This was proposed by [Bi and Wang, 2015](#) by Dirac mass engineering and designed in the gyroid photonic crystal by [Lu and Wang, 2016](#). The coupling of two Weyl points of opposite Chern numbers makes a 3D Dirac point, as introduced above. The resulting Dirac Hamiltonian

$$H_D = \hbar v(q_x \sigma_x + q_y \sigma_y + q_z \sigma_z \tau_z) + m \tau_+ + m^* \tau_-, \quad (70)$$

has a complex mass term m , where τ_z and $\tau_{\pm} \equiv (\tau_x \pm i\tau_y)/2$ are Pauli matrices acting on the valley degrees of freedom. In-plane winding of its argument $\text{Arg}[m]$ generates a vortex line in 3D supporting a zero mode at the vortex core, topologically protected by the second Chern number $C^{(2)}$ in the 4D parameter space (k_x, k_y, k_z, θ) , where θ is the winding angle of m .




FIG. 29 (Color online) List of 3D gapped phases in photonics, in which (a), (b), (c) require \mathcal{T} -breaking while (d) and (e) do not. (a) Analogue of the 3D Chern insulator labeled by three first Chern numbers. (b) One-way fiber of second Chern number. (c) Single-Dirac cone surface state with a \mathbb{Z}_2 invariant. (d) Two surface Dirac cones similar to those of weak 3D topological insulators. (e) Spatial-symmetry protected gapless quadratic touchings with a \mathbb{Z}_2 invariant.

In the photonic context, such a topological defect line inside an otherwise fully gapped gyroid photonic crystal can be obtained by means of a helical winding of the super-cell modulation coupling the two Weyl points. Depending on the spatial pitch and the handedness of the helical winding, one-way fiber modes of arbitrary $C^{(2)}$ can then be readily designed with arbitrary number of one-way modes.

This is in direct contrast with the one-way edge mode in 2D Chern crystals where high Chern numbers are difficult to obtain ([Skirlo *et al.*, 2015, 2014](#)). Another advantage of the one-way fiber design is that all one-way modes have almost identical group and phase velocities, due to the absence of sharp boundaries.

3. Single surface Dirac cone

A single-Dirac-cone surface state, the hallmark of 3D topological insulators ([Fu *et al.*, 2007](#)), can also be realized on the surface of magnetic photonic crystals, as shown by [Lu *et al.*, 2016b](#) and illustrated in Fig. 29(c). Instead of the Kramers' degeneracy of electrons due to \mathcal{T} , the double-degeneracy in photonics can be replaced by a crystalline symmetry — glide reflection. On the other hand, \mathcal{T} has to be broken to split the dispersions in all surface directions away from degeneracy.

The starting point to construct this phase is a pair of 3D Dirac points pinned at the high-symmetry points of the bulk Brillouin zone. By breaking \mathcal{T} using magnetic materials, the authors gapped the 3D Dirac points and obtained the gapless single-Dirac cone surface states. The topological invariant is \mathbb{Z}_2 ([Fu *et al.*, 2007](#); [Moore and Balents, 2007](#); [Roy, 2009](#)). When the glide reflection symmetry is broken uniformly on the surface, the single Dirac cone opens a frequency gap. Other than that, the surface state is robust against arbitrary random disorder, i.e., when the glide symmetry is preserved on average on the surface ([Lu *et al.*, 2016c](#)).

4. Non-magnetic designs

The above topological phases with 3D bulk gaps require \mathcal{T} -breaking for highly-robust interfacial states. However, the magnetic response is extremely weak towards optical frequencies, which has motivated the search for \mathcal{T} -invariant designs. One approach has been to search for non-magnetic designs that mimic weak 3D topological insulators, which can be viewed as stacks of 2D topological insulators. In photonics, this was first proposed for all-dielectric bianisotropic metamaterials [Slobozhanyuk *et al.*, 2016a](#), as introduced in Sec. III.B.5, and has been experimentally realised very recently in a 3D array of bianisotropic metallic split-ring resonators ([Yang *et al.*, 2018](#)).

A second approach to \mathcal{T} -invariant designs has been to search for systems, where the topological properties are protected instead by crystalline symmetries. The first proposal of such a topological crystalline insulator (Fu, 2011) (see Sec. III.B.6 for the discussion of such states in two dimensions) utilized C_4 rotation symmetry and exhibited double degeneracies at two surface momentum points. This defined a \mathbb{Z}_2 invariant for its surface states to connect gaplessly. At the degenerate point, the dispersion was quadratic due to the C_4 symmetry and \mathcal{T} . As illustrated in Fig. 29(e), this phase is realizable in photonic crystals (Alexandradinata *et al.*, 2014; Yannopapas, 2011). A concrete design for a tetragonal photonic crystal was also proposed in Ochiai, 2017.

C. Towards even higher dimensions

Topological phases of matter with spatial dimensions of four or higher can also be of experimental relevance in photonics. As already introduced in Sec. IV.B, a recent experiment has used topological pumping to probe a four-dimensional quantum Hall system (Zilberberg *et al.*, 2018); in this approach, some of the dimensions are replaced by externally tuned parameters, effectively freezing out the dynamics along these directions. An alternative approach, which could also offer access to the dynamics of particles moving in effectively spatial four dimensions, is based on so-called *synthetic dimensions*. In this, the key concept is to reinterpret internal degrees of freedom as spanning additional spatial dimensions, so that higher dimensional lattice models are simulated in lower dimensional systems. In this section, we first review the development of synthetic dimensions in general, before discussing progress in the exploration of four-dimensional topological systems with photons.

1. Synthetic dimensions

There are several different ways to make the effective spatial dimensionality of a lattice system larger than the physical dimensionality of the real space in which the lattice is located. One natural idea for this purpose is to increase the connectivity of the lattice, as proposed by Tsomokos *et al.*, 2010 for superconducting qubit circuits, by Jukić and Buljan, 2013 for photonic lattices, by Schwartz and Fischer, 2013 for multi-dimensional laser-mode lattices, and by Graß *et al.*, 2015 for trapped ions. Another strategy that can allow for even greater flexibility is to use the internal degrees of freedom, reinterpreting these as if they label different sites along an additional synthetic dimension in the system, as originally proposed by Boada *et al.*, 2012 in the context of ultracold atomic gases, and later extended by Celi *et al.*, 2014 to allow for complex hoppings along the synthetic

direction, and so to realise quantum Hall systems. Methods to create lattice structures more complex than just a square lattice were proposed in Anisimovas *et al.*, 2016; Boada *et al.*, 2015; and Suszalski and Zakrzewski, 2016.

The idea of synthetic dimensions was soon experimentally realized in the context of cold atoms by two groups (Mancini *et al.*, 2015; Stuhl *et al.*, 2015), in which a two dimensional ladder with a magnetic field was simulated using a one dimensional chain of atoms. Following experiments have then extended the synthetic dimension idea by using the different electronic states of atoms (Kolkowitz *et al.*, 2017; Livi *et al.*, 2016), and discrete states in momentum space (An *et al.*, 2017). Furthermore, there are theoretical proposals to use harmonic oscillator eigenstates (Price *et al.*, 2017) and orbital angular momentum states (Pelegrí *et al.*, 2017) as synthetic dimensions. Typically, the inter-particle interaction along the synthetic direction is very long ranged, resulting in a variety of interesting phenomena (Barbarino *et al.*, 2016; Bilitewski and Cooper, 2016; Calvanese Strinati *et al.*, 2017; Graß *et al.*, 2014; Jünemann *et al.*, 2017; Łącki *et al.*, 2016; Taddia *et al.*, 2017; Zeng *et al.*, 2015).

In photonics the first proposal for how to implement a synthetic dimension was made in Luo *et al.*, 2015, and extended later in Luo *et al.*, 2017 and Zhou *et al.*, 2017b, in which different orbital angular momentum states of light, coupled via spatial light modulators, were regarded as the synthetic dimension. This was followed by a proposal in optomechanics (Schmidt *et al.*, 2015), in which photon and phonon degrees of freedom were considered as two lattice sites along the synthetic dimension. Ozawa *et al.*, 2016b and Yuan *et al.*, 2016a have proposed to use different frequency modes of a multi-mode ring resonator, coupled via external modulation of refractive index, as a synthetic dimension. By modulating a resonator with multiple frequencies, models with any dimensions can also be simulated (Yuan *et al.*, 2017b). A synthetic frequency dimension could also be realized in a Raman medium, where the synthetic magnetic field is controlled by the alignment of the two Raman beams (Yuan *et al.*, 2017a). Instead of different frequency modes, the angular coordinate within a ring-resonator may be used as a synthetic dimension (Ozawa and Carusotto, 2017), in which the inter-photon interaction is local along the synthetic direction, in contrast to extremely long-ranged interactions in other proposals.

There have also been many theoretical proposals for the different physics that could be accessed with synthetic dimensions. In a single resonator with a synthetic dimension, it may be possible to study the edge state of the SSH model (Zhou *et al.*, 2017b) and Bloch oscillations along the synthetic direction (Yuan and Fan, 2016). In a one-dimensional array of optical cavities with one synthetic dimension, the effect of topological edge states of two-dimensional Chern insulators may be observed (Luo *et al.*, 2015). Such a topological edge mode can be use-

ful for high-efficiency frequency conversion if there is an edge along the synthetic direction made of frequency modes (Ozawa *et al.*, 2016b; Yuan *et al.*, 2016a) and for realizing an optical isolator if an edge is along the spatial direction (Ozawa *et al.*, 2016b). A two-dimensional array of resonators could also be augmented by one frequency dimension to realise photonic Weyl points (Lin *et al.*, 2016; Sun *et al.*, 2017a) or a weak 3D topological insulator (Lin *et al.*, 2018). In the long run, one may expect that the idea of synthetic dimensions could find applications in increasing the complexity of optical networks in photonic devices, also in connection with frequency-multiplexing (Saleh and Teich, 2007) and optical comb (Cai *et al.*, 2017; Schwartz and Fischer, 2013) techniques.

The experimental observation of topologically protected edge states in a photonic lattice with a synthetic dimension was very recently pioneered in (Lustig *et al.*, 2018). A two-dimensional array of single-mode waveguides was used, with the synthetic dimension being encoded in the modal space of coupled waveguides. This allowed for the generation of a synthetic magnetic field and, thus, of a topologically non-trivial two-dimensional model supporting chiral edge states.

2. Four-dimensional quantum Hall effect

As mentioned in Sec. IV.B, a recent photonic experiment has used topological pumping to probe the edge states of a four-dimensional quantum Hall system (Zilberberg *et al.*, 2018), based on the proposal of Kraus *et al.*, 2013. Concurrently with this experiment, the hallmarks of the quantized bulk response of the 4D quantum Hall effect (23), including the second Chern number, were measured through the topological pumping of a two-dimensional ultracold atomic system (Lohse *et al.*, 2018). By defining the second Chern number in a general parameter space, this topological invariant has also been shown to be experimentally relevant in helically modulated fibers (Lu and Wang, 2016), as discussed in Sec. V.B, where the angular coordinate in the cross section of a fiber acts as the fourth parameter, and in ultracold gases, where it was measured over a parameter space spanned by properties of two Raman lasers applied to the system (Sugawa *et al.*, 2016).

The first proposal for observing the full dynamics of a four-dimensional system was presented in Jukić and Buljan, 2013, based on using photonic lattices with high connectivity to study four-dimensional solitons. Using a synthetic dimension to directly observe the four-dimensional quantum Hall was then originally proposed in ultracold atomic gases (Price *et al.*, 2015) and soon after extended to photonics Ozawa *et al.*, 2016b. These proposals focused on the four-dimensional tight-binding

model (Kraus *et al.*, 2013):

$$\hat{H} = -J \sum_{\mathbf{r}} \left(\hat{a}_{\mathbf{r}+a\hat{e}_x}^\dagger \hat{a}_{\mathbf{r}} + \hat{a}_{\mathbf{r}+a\hat{e}_y}^\dagger \hat{a}_{\mathbf{r}} + e^{i2\pi\Phi_1 x/a} \hat{a}_{\mathbf{r}+a\hat{e}_z}^\dagger \hat{a}_{\mathbf{r}} + e^{i2\pi\Phi_2 y/a} \hat{a}_{\mathbf{r}+a\hat{e}_w}^\dagger \hat{a}_{\mathbf{r}} + \text{H.c.} \right), \quad (71)$$

where $a_{\mathbf{r}}$ is the annihilation operator of a particle at position specified by a four-dimensional vector $\mathbf{r} = (x, y, z, w)$ with w being the synthetic direction, and a being the lattice spacing. The fluxes Φ_1 and Φ_2 pierce the x - z plane and y - w plane, respectively. This is a generalization of the two-dimensional Harper-Hofstadter Hamiltonian to a 4D model with magnetic fields applied in two orthogonal planes, breaking time-reversal symmetry. However, unlike in 2D, time-reversal symmetry breaking is not the only route to observing a quantum Hall effect. Instead, 4D topological lattice models have also been proposed that exploit time-reversal-invariant spin-dependent gauge fields (Qi *et al.*, 2008), or lattice connectivity for spinless particles (Price, 2018).

Under the addition of weak electromagnetic perturbations, the filling of a 4D topological energy band would lead to a quantized nonlinear Hall current proportional to the second Chern number of the band(23). However, in contrast to the atomic case, in photonic systems with loss, this current j^μ is not a direct observable. Instead, it has been proposed to extract this topological response from the shift of the center of mass of the photonic steady-state intensity distribution under a monochromatic pump (Ozawa and Carusotto, 2014; Ozawa *et al.*, 2016b).

VI. GAIN AND LOSS IN TOPOLOGICAL PHOTONICS

In this section, we discuss the interplay of gain and loss with topology in photonics. We divide this subsection into two main parts; in the first, we discuss non-Hermitian topological models with gain and loss, while in the second, we focus on recent works concerning topology in Bogoliubov systems.

A. Non-Hermitian topological photonics

The study of topological physics with photons allows for the exploration of phenomena inaccessible in the context of condensed matter. A case in point is *non-Hermiticity* in the form of optical gain and loss. In photonics, gain and loss is much more common than in electrons in solids: gain media are the basis for lasers, and loss of photons is ubiquitous in every photonic device (loss is associated with absorption and surface roughness of a waveguide, for example).

There have thus far been a series of works delving into the interplay of non-Hermiticity and topology with a number of disparate aims (including a recent review on the topic (Alvarez *et al.*, 2018)). Inspired by a model proposed by Rudner and Levitov (Rudner and Levitov, 2009) Zeuner *et al.* (Zeuner *et al.*, 2015) used an optical waveguide array to demonstrate that the winding number of a one-dimensional topological system could be extracted from a non-Hermitian quantum walk. In that work, it was precisely the finite lifetime (induced by optical loss) of the ‘quantum walker’ that allowed for the observation of a topological transition. This amounts to the extraction of a topological number of a Hermitian system using non-Hermiticity, rather than exploring the topological invariants and edge states of non-Hermitian systems *per se*.

Another direction of non-Hermitian topological photonics is parity-time (\mathcal{PT}) symmetric (Makris *et al.*, 2008; Rüter *et al.*, 2010) topological systems. These are systems with balanced gain and loss such that the Hamiltonian commutes with the \mathcal{PT} operator (where \mathcal{P} represents parity and \mathcal{T} represents time reversal). It has been shown (Bender and Boettcher, 1998) that such systems may exhibit real eigenvalue spectra despite their non-Hermiticity; they have been the basis of a major research effort in photonics due to the possibility of overcoming parasitic loss and absorption in optical devices using gain. Thus, \mathcal{PT} -symmetric systems allow for the possibility of well-defined bands and gaps and are thus a natural place to start in studying non-Hermitian topological effects. That said, it was shown (Esaki *et al.*, 2011; Hu and Hughes, 2011) that a large class of systems that are \mathcal{PT} -symmetric in the bulk must have edge states that ‘break’ \mathcal{PT} ; namely, they have complex eigenvalues. Fortunately, under certain conditions, topological edge states with real eigenvalues can be found and have been demonstrated (Weimann *et al.*, 2017). Topological edge states in \mathcal{PT} -symmetric quantum walks have also been experimentally observed (Xiao *et al.*, 2017). Recent work has examined \mathcal{PT} -symmetric Hamiltonians in the context of topological superconductors (Kawabata *et al.*, 2018a).

A number of other unconventional phenomena arise when non-Hermiticity and topology are combined. One example is photonic ‘tachyon-like’ dispersion (Szameit *et al.*, 2011) that was demonstrated in the form of exceptional rings in photonic crystals (Zhen *et al.*, 2015) as well as ‘Fermi arc’-type states that connect between exceptional points in two-dimensional systems (Kozii and Fu, 2017; Malzard and Schomerus, 2018a; Zhou *et al.*, 2018); a related phenomenon was examined in the context of heavy-Fermion systems (Yoshida *et al.*, 2018). ‘Exceptional rings’ arise in three-dimensional topological systems exhibiting Weyl points (Szameit *et al.*, 2011; Xu *et al.*, 2017), though to date these have not been experimentally realized. The enhancement of topologi-

cal interface states in one-dimensional systems was proposed (Schomerus, 2013) and demonstrated in the microwave regime (Poli *et al.*, 2015) (along with a related work in optics (Pan *et al.*, 2018)); furthermore, it was shown that topological states absent when the system is Hermitian can be induced by adding losses (Malzard *et al.*, 2015). The interplay of non-Hermiticity and flat bands has been shown to result in a photonic analogue of Aharonov-Bohm caging (Leykam *et al.*, 2017b). Beyond photonic systems, the interplay between non-Hermiticity/dissipation and topology has been explored in a number of theoretical works in varying contexts (Bardyn *et al.*, 2013, 2012; Budich and Diehl, 2015; Budich *et al.*, 2015; Diehl *et al.*, 2011). Finally, topological edge state lasers (Bahari *et al.*, 2017b; Bandres *et al.*, 2018a; Harari *et al.*, 2018b), intrinsically non-Hermitian systems, have been observed recently - thus, it will become increasingly necessary to put on firm ground the complex interplay of non-Hermiticity and topological protection.

Despite this progress, the major challenge of non-Hermitian topological photonics remains the formulation of a general framework akin to that which exists for Hermitian systems. In particular, perhaps the key contemporary question is: what is the right topological invariant to consider for a given non-Hermitian Hamiltonian (Esaki *et al.*, 2011; Hu and Hughes, 2011), and what is its relevance to bulk-edge correspondence (some recent progress has been made in this direction (Kawabata *et al.*, 2018c; Lee, 2016; Leykam *et al.*, 2017a; Shen *et al.*, 2018; Weimann *et al.*, 2017; Yao *et al.*, 2018; Yao and Wang, 2018; Yin *et al.*, 2018))? In this spirit, there has been a recent effort to classify non-Hermitian topological systems (Gong *et al.*, 2018) in the same vein as Hermitian ones. This is certainly not a direct generalization however: it may be the case that the symmetries conventionally associated with the Hermitian case may not be well-suited to classify Hamiltonians in the non-Hermitian case. Specifically, recent work has shown that particle-hole and time-reversal symmetries are aspects of the same larger symmetry when Hermiticity is relaxed (Kawabata *et al.*, 2018b).

B. Emergent topology of Bogoliubov modes

Photons under a parametric driving can be described by a Hamiltonian with terms that do not conserve the number of photons. Such number non-conserving bosonic systems can have topological features which are qualitatively different from fermionic topological systems. To understand the origin of the number non-conserving terms, let us consider a photonic cavity whose resonant frequency is ω , and assume that the cavity is made of optically nonlinear material with a second-order nonlinear susceptibility $\chi^{(2)}$. When one pumps the system with fre-

quency 2ω , the nonlinearity converts the pumped photon into two photons with frequency ω in the cavity. Assuming that the pump beam is sufficiently strong and can be treated classically, the effective Hamiltonian describing the cavity takes the following form (Gerry and Knight, 2005):

$$\hat{H}_{\text{cavity}} = i\hbar\chi^{(2)} (\beta^* \hat{a}^2 - \beta \hat{a}^{\dagger 2}), \quad (72)$$

which does not conserve the number of photons, where \hat{a} is the annihilation operator of a photon in the cavity and a \mathbb{C} -number β characterizes the pumping field. Such a cavity can be aligned to form a periodic lattice. The second-quantized momentum-space Hamiltonian of the lattice system can be written in the following form:

$$\hat{H}_{\text{lattice}} = \frac{1}{2} \sum_{\mathbf{k}} \begin{pmatrix} \hat{\Psi}_{\mathbf{k}}^\dagger & \hat{\Psi}_{-\mathbf{k}} \end{pmatrix} H_{\mathbf{k}} \begin{pmatrix} \hat{\Psi}_{\mathbf{k}} \\ \hat{\Psi}_{-\mathbf{k}}^\dagger \end{pmatrix},$$

$$H_{\mathbf{k}} = \begin{pmatrix} A(\mathbf{k}) & B(\mathbf{k}) \\ B(-\mathbf{k})^* & A(-\mathbf{k})^t \end{pmatrix}, \quad (73)$$

where $\hat{\Psi}_{\mathbf{k}}$ is an N -component vector of annihilation operators with crystal momentum \mathbf{k} , and N is the number of lattice sites per unit cell. The N -by- N matrix $A(\mathbf{k})$ is Hermitian and $B(\mathbf{k})^t = B(-\mathbf{k})$. The terms due to $B(\mathbf{k})$ do not conserve the number of photons. At first glance, the Hamiltonian (73) is similar to the Bogoliubov-de Gennes Hamiltonian of superconducting electronic systems. In fact, the Hamiltonian (73) has particle-hole symmetry as in the fermionic Bogoliubov-de Gennes Hamiltonian, and the spectrum is symmetric with respect to the zero of the energy. However, the transformation needed to diagonalize the Hamiltonian is drastically different between bosons and fermions. The fermionic counterpart of $H_{\mathbf{k}}$ for the Bogoliubov-de Gennes Hamiltonian can be diagonalized by a unitary matrix to obtain eigenenergies of the systems which are guaranteed to be all real. On the other hand, in order to preserve the bosonic commutation relations, the bosonic Bogoliubov Hamiltonian $H_{\mathbf{k}}$ should be diagonalized by Bogoliubov transformations which are not a unitary matrix but a paraunitary matrix \hat{U} obeying $\hat{U}^\dagger (\sigma_z \otimes \mathbb{I}_N) \hat{U} = \sigma_z \otimes \mathbb{I}_N$. The associated eigenenergies can be complex. These differences imply that the standard wisdom on topological phases of matter known for fermions may not hold for bosonic Bogoliubov Hamiltonians. Because of the possibility of having complex eigenvalues, we also need to pay attention to the possibility of instability.

Bosonic Bogoliubov Hamiltonians appear not only in photonic systems. In fact, the topological properties of such Hamiltonians were first discussed in the context of magnons in ferromagnetic crystals (Shindou *et al.*, 2013a; Shindou and Ohe, 2014; Shindou *et al.*, 2013b), where analogs of the Chern insulators in bosonic Bogoliubov Hamiltonians were discussed. It was found that the relevant Berry connection of the n -th band of the Bogoliubov

Hamiltonian is

$$\mathcal{A}_n(\mathbf{k}) = i\langle u_{n,\mathbf{k}} | \sigma_z \nabla_{\mathbf{k}} | u_{n,\mathbf{k}} \rangle, \quad (74)$$

where $|u_{n,\mathbf{k}}\rangle$ is the Bloch state of the n -th band. Note the additional σ_z in the definition of the Berry connection. The Berry curvature is then defined as $\Omega_n = \nabla_{\mathbf{k}} \times \mathcal{A}_n(\mathbf{k})$. The Chern number calculated by integrating this Berry curvature over the Brillouin zone is guaranteed to be integer, and is related to the number of chiral edge modes.

Bogoliubov excitations are typically gapless at zero energy, but there can be gaps between higher energy bands. In exciton-polaritons, Bardyn *et al.*, 2016 and Bleu *et al.*, 2016 have analyzed the Bogoliubov modes of exciton-polariton condensates and proposed models which have topological edge states at the gaps with nonzero excitation energy. The topological edge states at higher energy gaps of Bogoliubov excitations were also discussed in ultracold atomic gases (Di Liberto *et al.*, 2016a; Engelhardt and Brandes, 2015; Furukawa and Ueda, 2015; Li *et al.*, 2015b). In a lattice of photonic cavities under parametric driving, Peano *et al.*, 2016a proposed a model which has a nonzero gap at zero energy. In order to have a stable system, the gap at zero energy cannot have an edge state, so the sum of the Chern numbers of bands at the negative energy is zero, but gaps between higher energy bands can have topological edge states.

A distinctive feature of the bosonic Bogoliubov Hamiltonian (73) is that the eigenenergies can become complex, hence triggering parametric instabilities (Shi *et al.*, 2017). Peano *et al.*, 2016b proposed a model where the topological edge states become unstable, even though all the bulk modes are stable. Such an unstable edge mode could be used as a traveling wave parametric amplifier. Instability caused by the topological edge modes was also analyzed in the context of ultracold atomic gases (Barnett, 2013; Engelhardt *et al.*, 2016; Galilo *et al.*, 2015). The interplay between the topology and the parametric instability has also been discussed in classical harmonic oscillators under periodic driving (Salerno *et al.*, 2016).

Finally, combined with strong optical nonlinearities (see Sec. VII.B), a p -wave version of parametric driving underlies the proposal in Bardyn and Imamoğlu, 2012 to obtain Majorana modes in a one-dimensional system of strongly interacting, fermionized photons.

VII. TOPOLOGICAL EFFECTS FOR INTERACTING PHOTONS

Most of the discussion of the previous sections concerned linear optical systems whose physics can be accurately described in terms of the standard Maxwell's equations including suitable linear dielectric and magnetic susceptibilities. In this regime, photons behave as independent particles. In this last section of the Review,

we focus our attention on the novel features that originate from the interplay of the topology with nonlinear optical effects.

Basing ourselves on the general introduction to the basic nonlinear optics concepts of Sec. II.B.3, the next two subsections Secs. VII.A and VII.B will summarize the main effects of an intensity-dependent refractive index in, respectively, the cases of weak and strong nonlinearity; in the former, a classical mean-field description based on Maxwell's equations with a nonlinear polarization term is accurate, while, in the latter, the physics is dominated by quantum optical effects due to the discreteness of the photon. We also note that Sec. VI.B provides a brief review of how parametric processes generated by a $\chi^{(2)}$ optical nonlinearity can give rise to a rich emergent topological structure for linear Bogoliubov modes.

A. Weak nonlinearities

For sufficiently weak values of the optical nonlinearities, one can legitimately perform the mean-field approximation of Eq. (55), in which the photons lose their particle character and collectively behave as a macroscopic wave, experiencing effective material properties that depend on the local amplitude of the light field according to the model of classical nonlinear polarization Eq. (51). While some parametric processes generated by a $\chi^{(2)}$ optical nonlinearity in the topological photonics context were reviewed in Sec. VI.B, in this subsection we shall focus on the case of an intensity-dependent refractive index Eq. (52): As we shall review in the following, theoretical works have anticipated that the modification of the refractive index induced by the nonlinearity may have dramatic observable consequences such as modifying the effective topology experienced by the wave.

A first and most natural question of nonlinear topological physics was to understand how solitons (Eisenberg *et al.*, 1998; Fleischer *et al.*, 2003; Segev *et al.*, 1992) or vortices (Kivshar and Agrawal, 2003) are affected by the underlying geometry and topology of the band. This physics has attracted great interest in many fields such as ultracold atomic gases, where relativistic solitons and vortices in honeycomb geometries have been studied in (Haddad and Carr, 2011, 2015; Haddad *et al.*, 2015).

Focussing on optical systems, intense research has been devoted to solitons in either the bulk or the edges of the Floquet photonic topological insulators of (Rechtsman *et al.*, 2013b), that were introduced in Sec. III.A.2. The first work in this direction (Lumer *et al.*, 2013b) highlighted different families of long-lived, self-localized wave packets residing in the bulk of the system. Depending on their size, the wavepackets may rotate in opposite directions, either following or opposite to the global Floquet modulation of the lattice. Most interestingly, the

current profile of a rotationally-symmetric six-site wide wavepacket can be understood as an edge state residing on the inner boundary of a self-induced effective hole due to the nonlinearity.

The study of the effect of nonlinearity on topological edge states has been pioneered by (Ablowitz *et al.*, 2014, 2015, 2013), where the linear edge states of the Floquet bands of the experiment (Rechtsman *et al.*, 2013b) were classified as a function of the Floquet modulation parameters and their nonlinear evolution recast in terms of an effective one-dimensional nonlinear Schrödinger-like equation, possibly including higher-order derivative terms. Based on this equation, unidirectionally propagating edge soliton states have been identified: the topological robustness of linear edge states to backscattering translates into an enhanced robustness of edge solitons against higher-order terms. Following works have then analyzed the topological robustness of edge solitons traveling around sharp corners (Ablowitz and Ma, 2015) and developed a general methodology to understand the tight-binding approximation in the context of nonlinear Floquet systems (Ablowitz and Cole, 2017).

Dynamical modulational instabilities of edge states under the effect of nonlinearity and their eventual break-up into a train of solitons has been explored by several authors. In (Lumer *et al.*, 2016), nonlinear extended edge states of the Floquet system of (Rechtsman *et al.*, 2013b) have been shown to be always unstable independently of the sign of their linear dispersion and the actual strength of the nonlinearity. This modulational instability eventually leads to the break up of the extended wave into soliton-like localized wavepackets. Depending on the strength of the nonlinearity, such solitons can extend over many sites along the edge or localize to a single site. For polariton honeycomb lattices, the modulational instability of edge states and the consequent appearance of long-lived quasi-soliton edge states was studied in (Kartashov and Skryabin, 2016), while, for kagome-shaped polariton lattices, topological edge solitons were studied in (Gulevich *et al.*, 2017). This latter work also highlighted the wide tunability of the edge soliton group velocity from positive to negative values as well as the robustness of topological edge solitons upon inter-soliton collisions.

The idea of nonlinear effects inducing transitions between states with different symmetries was pioneered in (Lumer *et al.*, 2013a) with a theoretical study of nonlinearity-induced transitions between \mathcal{PT} -broken and \mathcal{PT} -symmetric states in a non-Hermitian system and, then, in (Katan *et al.*, 2016) with a theoretical study of the effect of long-range nonlinearities on topological transport. Along these lines, a novel kind of topological solitons were investigated in (Leykam and Chong, 2016): nonlinear effects locally induce a topological transition in an otherwise topologically trivial lattice and solitons naturally arise as the edge states at the topological interface. Possible applications of such nonlinearly-induced

topological transition to optical isolation were explored in different geometries in (Zhou *et al.*, 2017c). A related study in a nonlinear but conservative one-dimensional SSH model was reported in (Hadad *et al.*, 2016).

B. Strong nonlinearities

When nonlinearities are large, the discrete nature of the photons constituting the field starts being important and one has to resort to a fully quantum description. Correspondingly, the physics of these systems is qualitatively different, as they are expected to support strongly correlated states of light that closely resemble their electronic counterparts, e.g. fractional quantum Hall liquids (Carusotto and Ciuti, 2013).

The simplest example of a quantum nonlinear effect is the so-called *photon blockade* phenomenon (Imamoglu *et al.*, 1997), that occurs in single-mode nonlinear cavities when the single photon nonlinearity ω_{nl} (i.e. the frequency shift experienced by the mode for a single photon occupation, as introduced in Eq. (58)) exceeds the damping rate γ of the cavity mode. For an incident beam on resonance with the empty cavity mode, a first photon can freely enter the cavity, but a second one will find the effective resonance shifted by ω_{nl} and can not enter until the first has left. In analogy with the Coulomb blockade of electronics, one can think of the first photon blocking the entrance of the second: hence the term *photon blockade*.

In the last decade or so, photon blockade has been observed in a variety of cavity configurations using different optically nonlinear elements in (Birnbaum *et al.*, 2005; Faraon *et al.*, 2008; Lang *et al.*, 2011; Reinhard *et al.*, 2012). In relation to topological photonics, the most promising platforms to combine photon blockade with synthetic gauge fields and/or non-trivial band topologies are the non-planar cavities containing coherently dressed atomic gases in a Rydberg-EIT configuration and circuit-QED devices embedding strongly nonlinear superconducting elements, as recently pioneered in (Jia *et al.*, 2017, 2018) and (Roushan *et al.*, 2017b), respectively.

The extension of photon blockade to many-cavity configurations so as to obtain complex strongly correlated many-photon states started attracting the interest of researchers in the mid-2000's with several proposals for how to realize Mott-insulator states of light (Angelakis *et al.*, 2007; Greentree *et al.*, 2006; Hartmann *et al.*, 2006). Along these lines, the first proposal of a quantum Hall effect for light appeared in (Cho *et al.*, 2008). While all these pioneering works made the quite strong assumption of a quasi-equilibrium photon gas, which is able to equilibrate and/or be adiabatically manipulated before disappearing due to losses, specific studies of the consequences of the intrinsically driven-dissipative nature of photon systems appeared just a few years later (Caru-

sotto *et al.*, 2009; Gerace *et al.*, 2009).

A first theoretical study of the interplay of strong interactions with a synthetic gauge field in a driven-dissipative context appeared in (Nunnenkamp *et al.*, 2011), where strongly-correlated states of photons in few-sites lattices were highlighted, together with the signatures of such states in the transmission properties of a device. Soon after, a proposal to generate fractional quantum Hall states of light as a driven-dissipative steady state of a lossy Bose-Hubbard model with non-trivial hopping phases under a coherent pump was reported in (Umucalilar and Carusotto, 2012). The selective excitation of the desired many-body state can be obtained via the same multiphoton frequency selection mechanism first introduced in (Carusotto *et al.*, 2009) for Tonks-Girardeau gases of light. Signatures of the strongly-correlated nature of quantum Hall states are then anticipated to transfer to the quantum statistical properties of the emitted light. While the coherent pumping scheme considered in Carusotto *et al.*, 2009 and further investigated in Hafezi *et al.*, 2013a is promising for the generation of few-photon quantum Hall states, its performance does not scale up favourably to larger numbers of photons: the frequency selection mechanism loses efficiency as many-photon peaks get spectrally closer, and the effective matrix element of the many-photon transition to the quantum Hall state may be quickly decreasing.

Soon afterwards, the coherent pumping scheme was extended to single cylindrical cavity geometries in (Umucalilar and Carusotto, 2013). In analogy with related research in rotating atomic gases (Cooper, 2008), one can take advantage of the formal similarity between the magnetic Lorentz force and the Coriolis one to study quantum Hall physics in the rotating fluids of light that are generated by a Laguerre-Gauss shaped coherent drive. Serious difficulties of this scheme were quickly pointed out (Grusdt *et al.*, 2013): any deviation from the perfect rotational symmetry of the cavity would result in a quick spin-down of the rotating photon gas, while the spectral detuning between Landau levels prevents the use of narrow-band nonlinear elements such as Rydberg-EIT atoms.

These difficulties have been solved by the twisted optical resonators used in the experiments of (Schine *et al.*, 2016) reviewed in Sec.III.B.3. Replacing the mechanical rotation of the fluid of light with a synthetic magnetic field recovers the degeneracy between states in the lowest Landau level and, at the same time, introduces a sizable detuning between states of opposite angular momentum, which prevents the cloud from spinning-down. Crucial experimental steps towards the generation of quantum Hall fluids of light in these systems have been recently reported, including the observation of Rydberg cavity-polaritons in running wave resonators (Ningyuan *et al.*, 2016), the use of optically pumped atomic samples to break time-reversal and thus suppressing back-scattering




FIG. 30 Upper right panel: sketch of the circulation dynamics of a single- (left) and two-photon (right) state superimposed on the circuit-QED system under consideration. Upper left panel: time-evolution of the excitation probability in the three $Q_{1,2,3}$ qubits for a single-photon state at magnetic flux $\Phi_B = -\pi/2$. Lower (b) panel: the same for a two-photon case. Figure from (Roushan *et al.*, 2017b)

processes that would otherwise effectively reverse the synthetic gauge field (Jia *et al.*, 2018), and the use of the ultra-strong photon-photon interactions of Rydberg EIT to get photon blockade in the Gaussian-shaped fundamental mode of a cavity (Jia *et al.*, 2017).

In the meantime, further theoretical work has anticipated an exotic phase diagram resulting from the short-distance saturation of the Rydberg-Rydberg interaction (Grusdt and Fleischhauer, 2013). On the other hand, since these experiments are exploring quantum Hall physics on the surface of a cone, direct measurement of the properties of anyonic excitations appears to be possible through the density distribution in the vicinity of the cone tip, which will reflect the central charge of the topological fluid (Can *et al.*, 2016). Additional theoretical work on the many-body aspects of this challenge was presented in (Dutta and Mueller, 2017; Sommer *et al.*, 2015).

Along related directions, fractional quantum Hall states in Jaynes-Cummings-Hubbard lattices were theoretically explored in (Hayward *et al.*, 2012). The possibility of realizing topological models based on the spin dynamics of Rydberg polaritons confined in a widely-spaced microcavity array was explored in (Maghrebi *et al.*, 2015b); here, hopping between sites does not occur by photon tunneling between neighboring cavities, but rather by dipolar interactions which exchange the spin state of neighboring dark polaritons.

In parallel to these advances with macroscopic optical cavities embedding atoms, the first experimental studies of magnetic effects in a strongly interacting gas of photons have been reported in (Roushan *et al.*, 2017b) using a circuit-QED architecture with a closed loop three-site geometry. The synthetic magnetic fields is obtained fol-

lowing the theoretical proposal in (Fang *et al.*, 2012b), where the hopping phase is determined by the oscillation phase of the temporal modulation needed to compensate the frequency mismatch of neighboring sites. This technique falls within the class of Floquet techniques discussed in Sec.II.A.5 and was pioneered in the cold atom context in (Aidelsburger *et al.*, 2013; Miyake *et al.*, 2013).

As is illustrated in Fig.30, a directional circulation of photons is the signature of broken time-reversal symmetry. Whereas non-interacting photons would independently rotate as single photons in a direction fixed by the synthetic magnetic field, the effect of strong interactions is manifested in the opposite circulation direction of photon vacancies. A similar inversion of the rotation direction as a consequence of strong interactions was very recently reported also in the cold-atom context in (Tai *et al.*, 2017b).

Thanks to the relatively long lifetime of photons, the experiment (Roushan *et al.*, 2017b) could be performed by initializing the system in a suitable one- or two-photon Fock state and then following its dynamics in the absence of any pump during the evolution: such an approach is able to directly probe the coherent quantum dynamics, but it is strongly limited by the decay rate of the full many-photon state, that typically scales proportionally to the photon number (Milman *et al.*, 2000). As a further important result of (Roushan *et al.*, 2017b), the adiabatic generation of a few-body interacting ground state for arbitrary synthetic magnetic flux Φ_B was also reported by slowly ramping up Φ_B to the desired final value. In (Roushan *et al.*, 2017a), this same platform was used in the presence of a quasi-periodic potential for photons to observe the Hofstadter butterfly of one-photon states and, then, localization effects in the two-photon sector.

This experimental result is all the more promising as several adiabatic protocols to create strongly correlated macroscopic topological states have been theoretically investigated, based on either the melting of Mott-insulator states (Cho *et al.*, 2008) or by a sequence of flux-insertion and then quasi-hole refilling processes (Grusdt *et al.*, 2014; Ivanov *et al.*, 2018; Letscher *et al.*, 2015). An interesting proposal to manipulate quantum states of light by means of a generalized Thouless pumping in strongly nonlinear arrays was put forward in (Tangpanitanon *et al.*, 2016). One has however to keep in mind that all these adiabatic schemes typically require that the process must be completed in a time-scale shorter than the lifetime of the quantum many-body state of interest, a condition that may become extremely demanding for macroscopic photon fluids.

An alternative approach to dynamically stabilize topological many-body states of light against losses without any active intervention from external observers was started in (Kapit *et al.*, 2014). In a circuit-QED context, this can be achieved using, e.g., the frequency-selective

parametric emission from a shadow lattice: refilling of holes occurs at a fast rate as long as it spectrally coincides with the emission bandwidth, while the injection of extra photons on top of the topological state is suppressed via a generalized blockade phenomenon due to the many-body energy gap. In a related proposal (Hafezi *et al.*, 2015), parametric coupling to a thermal bath was proposed to generate a tunable chemical potential for effectively thermalized light. A conceptually similar idea based on population-inverted two-level systems as frequency-dependent light emitters was investigated in (Biella *et al.*, 2017; Lebreuilly *et al.*, 2016) in the context of Mott insulator states of light. The significant advantages of replacing the Lorentzian emission spectrum with a more sophisticated square spectrum were pointed out in (Lebreuilly *et al.*, 2017). The study of frequency-dependent incoherent pumping schemes applied to the generation of quantum Hall states in the twisted optical resonators of (Schine *et al.*, 2016) was discussed in (Umucalilar and Carusotto, 2017).

In the long run, the application of photonic systems as useful platforms for topological quantum computation crucially requires strongly correlated topological states that support excitations with non-Abelian braiding statistics (Nayak *et al.*, 2008). A promising candidate for this purpose are the so-called Pfaffian states. Originally predicted in the context of the quantum Hall effect of electrons (Moore and Read, 1991), they can be obtained as the ground state in the presence of suitably engineered three-body interactions. A proposal to put this strategy into practice in a circuit-QED context is presented in (Hafezi *et al.*, 2014).

Before concluding, it is worth mentioning a completely different approach to the quantum dynamics of systems of few interacting particles (Kramer and Khomeriki, 2011; Longhi, 2011): in the simplest formulation, the idea is to map the $x_{1,2}$ spatial coordinates of two quantum particles onto the x, y spatial coordinates of a single particle moving in two dimensions. The two-body interactions are then modeled by letting the $x_1 = x_2$ line of sites have slightly different linear optical properties. As the dynamics remains at a fully single-particle level, it can be simulated in any of linear optical devices discussed in the first sections of this review.

Along these lines, physically two-dimensional arrays of waveguides were used in (Mukherjee *et al.*, 2016) to show evidence of tunneling processes for two-particle bound states in regimes where single-particle tunneling is instead strongly suppressed. The experiment (Schreiber *et al.*, 2012) demonstrated encoding of the spatial coordinates of a two-dimensional quantum walk into the arrival time of a single optical pulse, with a potential generalization to higher-dimensional configurations.

Very recently, the application of this mapping to study the interplay of interactions and topology in the two-particle dynamics of a one-dimensional SSH model was

simultaneously proposed in the theoretical works (Di Liberto *et al.*, 2016b) and (Gorlach and Poddubny, 2017). An extension of this work exploiting driven-dissipative effects in resonator arrays was proposed in (Gorlach *et al.*, 2018). While these schemes are naturally transferred to the fermionic sector of the two-particle dynamics by means of the suitably anti-symmetric initialization, a proposal to study supersymmetry (SUSY) effects in a light-matter context appeared in (Tomka *et al.*, 2015).

VIII. CONCLUSION AND PERSPECTIVES

In the previous sections we have seen how topological photonics has grown from its first proposal (Haldane and Raghu, 2008; Raghu and Haldane, 2008) into a wide and mature field of research with a number of active exciting directions. In this final section of the review, we aim at summarizing those future developments that are promising for the next future.

A. Optical isolation and robust transport

Topological photonics may have short- and mid-term technological impact. The most straightforward applications involves the use of topologically protected unidirectional edge states as robust optical delay lines or optically isolating elements, as originally discussed in (Hafezi *et al.*, 2011). While a firm experimental evidence of robust unidirectional propagation is nowadays available in a number of systems, practical application of these ideas into actual devices is still a subject of active investigation.

In particular, the stringent conditions that an optical isolator device must fulfill to be of practical utility were discussed in (Jalas *et al.*, 2013). Different strategies to match the requirements are presently being actively explored using either magnetic elements (Bahari *et al.*, 2017a; Solnyshkov *et al.*, 2018), optical nonlinearities (Khanikaev and Alù, 2015; Shi *et al.*, 2015), or externally-modulated elements (Fang *et al.*, 2017; Hua *et al.*, 2016; Yu and Fan, 2009).

Another exciting direction in view of applications in quantum information processing is to extend these results from classical light fields to quantum optical ones and link with on-going developments in chiral quantum optics (Lodahl *et al.*, 2017). A first step in this direction is to show that the dynamics of externally-generated entangled photon pairs inherits the topological protection of single-photon states, as theoretically studied in (Mittal *et al.*, 2016b; Rechtsman *et al.*, 2016). The emission and robust propagation of single photon states into chiral edge states was experimentally demonstrated in (Barik *et al.*, 2018). A proposal to generate quantum states of light by coupling an array of two-level emitters to a

chiral edge state was reported in (Ringel *et al.*, 2014): interesting correlations between photons originate when an externally generated few photon wavepacket scatter off the strongly nonlinear emitters.

B. Quantum emitters and topological laser

One of the most active directions of development to date is the study of the interplay of topology with light emitters and with optical gain, which is expected to offer novel features to be exploited in light sources, amplifiers and laser devices. Strong motivations supported this study in view of optoelectronic and photonics applications, to improve the performance of topological devices compared to their trivial counterparts.

A detailed study of parametric amplification on the chiral edge states of a two-dimensional Harper-Hofstadter model in the presence of parametric downconversion or spontaneous four-wave mixing emission processes was reported in (Peano *et al.*, 2016b). With a suitably chosen pump frequency, amplification can be restricted to the edge states, while bulk modes remain quickly damped as in the passive system. The unidirectional nature of the edge state guarantees that amplification is not only quantum-noise-limited as in standard parametric amplifiers, but also non-reciprocal and almost perfectly insensitive to disorder. Interesting consequences of the topology on the zero-point quantum fluctuations and on the emission of entangled photon pairs are also pointed out.

The robustness of the generated entangled photon pairs against disorder was specifically studied in (Mittal and Hafezi, 2017). In particular, it was shown how the generation of entangled photons using spontaneous four-wave mixing into topological states can outperform their topologically-trivial counterparts.

1. Topological lasers: theory

A strong activity is presently being devoted to the theoretical and experimental study of laser oscillation in topological systems, the so-called *topological lasers*. For one-dimensional systems, theoretical explorations of laser operation in topological states have appeared in (Pilozzi and Conti, 2016), which focused on a 1D Aubry-André-Harper bichromatic photonic crystal, and in (Malzard and Schomerus, 2018b), which studied a 1D SSH chain of resonators, showing that the topological mode selection can persist even in the presence of nonlinearities.

In two dimensions, topological lasing has been shown to lead to a highly efficient laser operation that remains monomode even well above the threshold and is robust against disorder (Harari *et al.*, 2018a). Simulations of the nonlinear wave equation with on-site saturable gain

terms of the form

$$\frac{i}{2} \frac{P_j}{1 + |\alpha_j|^2/n_{\text{sat}}} \alpha_j \quad (75)$$

included into the lattice model of Eq. (43) without coherent pumping term ($F_j = 0$) were performed for both trivial and topological lattice models with a pump profile P_j concentrated on the edge. Gain saturation at high power is modeled by the saturation density n_{sat} . In the topologically trivial case, laser operation is not able to exhaust all available gain, because many other modes can easily go above threshold for increasing power, leading to a complex multimode operation. This problem is particularly serious in the presence of disorder, which further suppress the mode competition effect by spatially localizing modes.

As it was shown in (Harari *et al.*, 2018a), all these problems turn out to be no longer relevant in topological lattices: the single lasing mode is extended around the whole system perimeter, maintaining a unidirectional flow and a spatially very uniform intensity profile even in the presence of disorder and for high pumping levels well above the threshold. In laser terms, this means a robust monomode operation with a high slope efficiency.

Further theoretical work (Seclí, 2017) has shown a strong dependence of the laser threshold on the topology of the amplifying region: as a consequence of the distinction between the convective and absolute instabilities, a higher lasing threshold was found when the amplifying region does not surround the whole system.

2. Topological lasers: experiments

From the experimental point of view, the study of the interplay of lasing with geometrical and topological features was initiated in (Sala *et al.*, 2015). In this experiment, the linewidth narrowing effect associated to laser operation was exploited to spectrally resolve the effect of the spin-orbit coupling terms in a hexagonal chain of pillar microcavities. The first examples of lasing operation in a topological non-trivial system were reported in one-dimensional chains of pillar microcavities (St-Jean *et al.*, 2017) or ring resonators (Parto *et al.*, 2018; Zhao *et al.*, 2018). In both cases, for suitable pump geometries, a single-mode laser emission occurred into the edge states of the chain.

Very recently, lasing in the topological edge states of a time-reversal-breaking two dimensional photonic crystal embedding magnetic YIG elements was reported in (Bahari *et al.*, 2017a). This work made use of a photonic crystal structure where gain is provided by quantum well emitters and time-reversal symmetry is broken by bonding the photonic crystal to a magnetic YIG material. In spite of the small width of the magneto-optically induced

band gap, and the comparatively large linewidth of the bulk modes, lasing of modes on the edge was observed.

Soon after, another experiment (Bandres *et al.*, 2018b) reported topological laser operation in a topological ring-resonator array (as discussed in Sec.III.B.1) embedding, in addition, quantum well emitters that provide optical gain. This setup does not require magnetic elements. By selectively pumping the edge resonators, a highly efficient single-mode emission into the topologically protected edge state was obtained even for gain values high above threshold. The performances of the novel device and the robustness against disorder were benchmarked with an extensive comparison to a topologically trivial laser device. A technique to break the spin-like symmetry between the clock- and counter-clockwise modes of the rings by adding S-bend elements into the resonators was also demonstrated, further reinforcing the unidirectional properties of the emission.

A third experiment reported lasing in chiral edge modes using a hexagonal lattice of microcavity polariton resonators (Klembt *et al.*, 2018). The scheme relies on the sensitivity of exciton polaritons to an external magnetic field to break time-reversal symmetry, in the configuration described in Sec. III.A.5.

C. Measurement of bulk topological and geometrical properties

Topological photonics are also opening new perspectives in the study of topological effects of wide interest for quantum condensed-matter physics. As we have reviewed, a variety of new, possibly high-dimensional lattice configurations are becoming available thanks to the advances in photonic fabrication and manipulation. Furthermore, thanks to the high flexibility of the optical excitation and diagnostic schemes, an intense study is being devoted to the observable consequences of the geometrical and topological concepts: while the topologically protected chiral edge states have been the smoking gun of a non-trivial topology starting from the pioneering work (Wang *et al.*, 2009), the geometrical quantities characterizing the bulk bands are nowadays the subject of active study.

From early studies of transport in electronic systems (Xiao *et al.*, 2010), it is well known that the Berry curvature enters the semiclassical equations of motion for electrons as a sort of momentum space magnetic field. The corresponding Lorentz-like force in reciprocal space provides an anomalous velocity term which is responsible, e.g., for the anomalous and integer (Thouless *et al.*, 1982) quantum Hall effects. In the topological photonics context, this idea underlies the experimental reconstruction in (Wimmer *et al.*, 2017) of the k -space distribution of the bulk Berry curvature from the anomalous velocity of a wavepacket performing Bloch oscillations under an ex-

ternal force as theoretically proposed in (Cominotti and Carusotto, 2013; Dudarev *et al.*, 2004; Price and Cooper, 2012).

Application of the anomalous velocity idea to the coherently pumped systems discussed in Sec.II.B.2 was theoretically proposed in (Ozawa and Carusotto, 2014). In addition to putting forward driven-dissipative versions of the anomalous Hall effects, specific pumping schemes able to equally distribute photons among the different momentum states were identified. In analogy to the classical theory of the integer quantum Hall effect in electronic systems (Thouless *et al.*, 1982; Xiao *et al.*, 2010) and to recent experiments with ultracold atoms (Aidelsburger *et al.*, 2014), the spatial displacement of the center of mass of the light intensity distribution then provides an information on the Chern number of the band. Later works (Ozawa *et al.*, 2016b; Price *et al.*, 2015, 2016) have generalized this idea to the second Chern number of four-dimensional lattice models and to the measurement of other geometrical quantities such as the Fubini-Study metric tensor (Ozawa, 2018). On a similar basis, the concept of mean chiral displacement has been introduced to measure the winding of a chiral Hamiltonian from real space measurements (Cardano *et al.*, 2016; Maffei *et al.*, 2018; Meier *et al.*, 2018; Mondragon-Shem *et al.*, 2014). One of the most attractive aspects of this proposal is the possibility of measuring topological invariants in the presence of disorder, which breaks translational symmetry. An alternative scheme to extract the Chern number from the steady-state field amplitude of a small photonic lattice with twisted boundary conditions was discussed in (Bardyn *et al.*, 2014). It has also been suggested to calculate the full quantum geometric tensor, including the Berry curvature and the Fubini-Study metric, from direct measurements of the photon wave-function in radiative photonic systems (Bleu *et al.*, 2018). Furthermore, dissipation has been used in the context of non-Hermitian systems to probe topological invariants in 1D photonic systems (Rudner and Levitov, 2009; Zeuner *et al.*, 2015), as discussed in Sec. VI.A.

Building on top of the anomalous velocity concept, recent works (Bliokh and Bliokh, 2005; Price *et al.*, 2014) have undertaken the challenge of upgrading the semiclassical equations of motion to a full quantum mechanical theory of quantum particles in the presence of a non-vanishing momentum-space Berry curvature. As an example of application of this theory, the eigenstates of a topologically non-trivial lattice model subject to an additional external harmonic potential can be physically understood as momentum-space Landau levels on the torus-shaped first Brillouin zone, with a degeneracy set by the Chern number of the band (Price *et al.*, 2014). A proposal to investigate this physics in a driven-dissipative array of optical cavities with site-dependent resonance frequencies appeared in (Berceanu *et al.*, 2016).

Further on-going work in this direction is investigat-

ing how the momentum-space magnetic field may lead to momentum-space analogs of the quantum Hall effects (Claassen *et al.*, 2015; Ozawa *et al.*, 2015, 2016a). In these last works, the minima of the k -space dispersion play the role of lattice sites, the harmonic trapping provides the momentum space analog of the kinetic energy, and the Berry curvature of the band plays the role of the magnetic field. Periodic boundary conditions are automatically inherited from the topology of the first Brillouin zone and the phase twist can be adjusted via the position of the harmonic trap minimum within the unit cell.

D. Topological quantum computing

A most exciting long term perspective is to use topological photonics devices as a platform for novel quantum information storage and processing tasks that exploit topological effects to protect their operation from external disturbances. A crucial requirement for such *topological quantum computing* with light appears to be the availability of strongly nonlinear elements to generate and manipulate strongly correlated states of light. The key experimental issues that researchers are facing along this route have been discussed in Sec.VII, together with the promising results that were reported in the last few years. In the following of this section, we wish to highlight a possible strategy along which this research may develop in the next future.

In the presence of suitably tailored optical nonlinearities, the fluid of light can form quantum states, e.g. Pfaffian ones (Hafezi *et al.*, 2014), that are anticipated to display a manifold of topologically degenerate ground states protected by a finite energy gap and elementary excitations with non-Abelian anyonic statistics. In systems with such properties (Nayak *et al.*, 2008), quantum information can be encoded in the ground state manifold of states and the unitary transforms corresponding to quantum logical operations can be performed by braiding quasi-holes around each other. With respect to standard quantum information protocols, quantum computing based on these topological states of matter has the advantage that the states within the topologically degenerate manifold can not be coupled nor mixed with each other by local disturbances, at least as long as their amplitude is not able to cross the energy gap.

To date, several condensed matter systems such as quantum Hall states of two-dimensional electrons under strong magnetic fields (Tong, 2016) or Majorana fermions in suitable superconductor-based solid-state nanostructures (Elliott and Franz, 2015) are being seriously considered to this purpose, but to the best of our knowledge no experimental evidence is yet available of anyonic braiding statistics. First proposals taking advantage of the peculiarities of the optical systems to observe any-

onic statistics have been recently put forward (Dutta and Mueller, 2017; Umucalılar and Carusotto, 2013), but a key question that remains open is the degree of robustness of the topologically encoded quantum information against the typical dissipative processes of optical systems. On the other hand, using an all-optical platform will be extremely favourable in view of integration of the quantum processing unit into an optical communication network.

ACKNOWLEDGMENTS

T.O. was supported by the EU-FET Proactive grant AQuS (Project No. 640800), the ERC Starting Grant TopoCold, and the Interdisciplinary Theoretical and Mathematical Sciences Program (iTHEMS) at RIKEN.

H.M.P. received funding from the Royal Society and from the European Unions Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 656093: “SynOptic”.

A.A. was supported by the ERC grant HoneyPol, the EU-FET Proactive grant AQuS (Project No. 640800), the French National Research Agency (ANR) project Quantum Fluids of Light (ANR-16-CE30-0021) and the program Labex CEMPI (ANR-11-LABX-0007), the CPER Photonics for Society P4S, and the Métropole Européenne de Lille.

N.G. was supported by the FRS-FNRS (Belgium) and by the ERC Starting Grant TopoCold.

M.H. acknowledges Sunil Mittal, and was supported by AFOSR MURI Grant No. FA95501610323, the Sloan Foundation, and the Physics Frontier Center at the Joint Quantum Institute.

L.L. was supported by the National key R&D Program of China under Grant No. 2017YFA0303800, 2016YFA0302400 and supported by NSFC under Project No.11721404.

M.C.R. acknowledges the National Science Foundation under award numbers ECCS-1509546 and DMS-1620422, the David and Lucile Packard Foundation, the Charles E. Kaufman Foundation, a supporting organization of the Pittsburgh Foundation, and the Alfred P. Sloan Foundation under fellowship number FG-2016-6418.

D.S. and J.S. were supported by the University of Chicago Materials Research Science and Engineering Center, which is funded by the National Science Foundation under Award No. DMR-1420709. This work was supported by ARO Grant No. W911NF-15-1-0397. D.S. acknowledges support from the David and Lucile Packard Foundation. This work was supported by DOE grant DE-SC0010267 and AFOSR grant FA9550-16-1-0323.

O.Z. was supported by the Swiss National Science Foundation (SNSF).

I.C. acknowledges funding from Provincia Autonoma di Trento, partly through the SiQuero project (“On Sili-

con Chip Quantum Optics for Quantum Computing and Secure Communications”), from ERC through the QGBE grant and from the EU-FET Proactive grant AQU-S, Project No.640800 and EU-FET-Open grant MIR-BOSE Project No.737017.

REFERENCES

- Aaboud, Morad, *et al.* (ATLAS) (2017), “Evidence for light-by-light scattering in heavy-ion collisions with the ATLAS detector at the LHC,” *Nature Phys.* **13** (9), 852–858.
- Ablowitz, Mark J, and Justin T. Cole (2017), “Tight-binding methods for general longitudinally driven photonic lattices: Edge states and solitons,” *Phys. Rev. A* **96**, 043868.
- Ablowitz, Mark J, Christopher W. Curtis, and Yi-Ping Ma (2014), “Linear and nonlinear traveling edge waves in optical honeycomb lattices,” *Phys. Rev. A* **90**, 023813.
- Ablowitz, Mark J, Christopher W Curtis, and Yi-Ping Ma (2015), “Adiabatic dynamics of edge waves in photonic graphene,” *2D Materials* **2** (2), 024003.
- Ablowitz, Mark J, Christopher W. Curtis, and Yi Zhu (2013), “Localized nonlinear edge states in honeycomb lattices,” *Phys. Rev. A* **88**, 013850.
- Ablowitz, Mark J, and Yi-Ping Ma (2015), “Strong transmission and reflection of edge modes in bounded photonic graphene,” *Opt. Lett.* **40** (20), 4635–4638.
- Adams, EN, and E.I. Blount (1959), “Energy bands in the presence of an external force fieldii: Anomalous velocities,” *Journal of Physics and Chemistry of Solids* **10** (4), 286 – 303.
- Aidelsburger, M, M. Atala, M. Lohse, J. T. Barreiro, B. Paredes, and I. Bloch (2013), “Realization of the Hofstadter Hamiltonian with ultracold atoms in optical lattices,” *Phys. Rev. Lett.* **111**, 185301.
- Aidelsburger, M, M. Lohse, C. Schweizer, M. Atala, J. T. Barreiro, S. Nascimbène, N. R. Cooper, I. Bloch, and N. Goldman (2014), “Measuring the Chern number of Hofstadter bands with ultracold bosonic atoms,” *Nat. Phys.* **11**, 3171.
- Albert, Victor V, Leonid I Glazman, and Liang Jiang (2015), “Topological properties of linear circuit lattices,” *Physical Review Letters* **114** (17), 173902.
- Alexandradinata, A, Chen Fang, Matthew J Gilbert, and B Andrei Bernevig (2014), “Spin-orbit-free topological insulators without time-reversal symmetry,” *Physical Review Letters* **113** (11), 116403.
- Altug, Hatice, Dirk Englund, and Jelena Vučković (2006), “Ultrafast photonic crystal nanocavity laser,” *Nature Physics* **2** (7), 484–488.
- Alvarez, VM, JE Vargas, M Berdakin, and LEF Torres (2018), “Topological states of non-hermitian systems,” arXiv preprint arXiv:1805.08200.
- An, Fangzhao Alex, Eric J. Meier, and Bryce Gadow (2017), “Direct observation of chiral currents and magnetic reflection in atomic flux lattices,” *Science Advances* **3** (4), 10.1126/sciadv.1602685, <http://advances.sciencemag.org/content/3/4/e1602685.full.pdf>.
- Anderson, Brandon M, Ruichao Ma, Clai Owens, David I Schuster, and Jonathan Simon (2016), “Engineering topological many-body materials in microwave cavity arrays,” *Physical Review X* **6** (4), 041043.
- Anderson, P Duke, and Ganapathi Subramania (2017), “Unidirectional edge states in topological honeycomb-lattice membrane photonic crystals,” *Optics Express* **25** (19), 23293–23301.
- Angelakis, D G, P. Das, and C. Noh (2014), “Probing the topological properties of the Jackiw-Rebbi model with light,” *Scientific Reports* **4**, 6110.
- Angelakis, Dimitris G, Marcelo Franca Santos, and Sougato Bose (2007), “Photon-blockade-induced Mott transitions and xy spin models in coupled cavity arrays,” *Phys. Rev. A* **76** (3), 10.1103/PhysRevA.76.031805.
- Anisimovas, E, M. Račiūnas, C. Sträter, A. Eckardt, I. B. Spielman, and G. Juzeliūnas (2016), “Semisynthetic zigzag optical lattice for ultracold bosons,” *Phys. Rev. A* **94**, 063632.
- Ao, Xianyu, Zhifang Lin, and CT Chan (2009), “One-way edge mode in a magneto-optical honeycomb photonic crystal,” *Physical Review B* **80** (3), 033105.
- Armitage, NP, EJ Mele, and Ashvin Vishwanath (2017), “Weyl and Dirac semimetals in three dimensional solids,” arXiv preprint arXiv:1705.01111.
- Arovas, Daniel, J. R. Schrieffer, and Frank Wilczek (1984), “Fractional statistics and the quantum Hall effect,” *Phys. Rev. Lett.* **53**, 722–723.
- Arrayás, M, D Bouwmeester, and JL Trueba (2017), “Knots in electromagnetism,” *Physics Reports* **667**, 1 – 61.
- Asatryan, Ara A, Lindsay C Botten, Kejie Fang, Shanhui Fan, and Ross C McPhedran (2013), “Local density of states of chiral Hall edge states in gyrotropic photonic clusters,” *Physical Review B* **88** (3), 035127.
- Asatryan, Ara A, Lindsay C Botten, Kejie Fang, Shanhui Fan, and Ross C McPhedran (2014), “Two-dimensional greens tensor for gyrotropic clusters composed of circular cylinders,” *JOSA A* **31** (10), 2294–2303.
- Asbóth, J K (2012), “Symmetries, topological phases, and bound states in the one-dimensional quantum walk,” *Phys. Rev. B* **86**, 195414.
- Ashcroft, Neil W, and N. David Mermin (1976), *Solid state physics* (Brooks Cole).
- Aspelmeyer, Markus, Tobias J Kippenberg, and Florian Marquardt (2014), “Cavity optomechanics,” *Reviews of Modern Physics* **86** (4), 1391.
- Avron, J E, R. Seiler, and B. Simon (1983), “Homotopy and quantization in condensed matter physics,” *Phys. Rev. Lett.* **51**, 51–53.
- Azbel, Mark Ya (1964), “Energy spectrum of a conduction electron in a magnetic field,” *Sov. Phys. JETP* **19** (3), 634–645.
- Baboux, Florent, Eli Levy, Aristide Lemaître, Carmen Gómez, Elisabeth Galopin, Luc Le Gratiet, Isabelle Sagnes, Alberto Amo, Jacqueline Bloch, and Eric Akkermans (2017), “Measuring topological invariants from generalized edge states in polaritonic quasicrystals,” *Phys. Rev. B* **95**, 161114.
- Bahari, Babak, Abdoulaye Ndao, Felipe Vallini, Abdelkrim El Amili, Yeshaiahu Fainman, and Boubacar Kanté (2017a), “Nonreciprocal lasing in topological cavities of arbitrary geometries,” *Science* **358** (6363), 636–640.
- Bahari, Babak, Abdoulaye Ndao, Felipe Vallini, Abdelkrim El Amili, Yeshaiahu Fainman, and Boubacar Kanté (2017b), “Nonreciprocal lasing in topological cavities of arbitrary geometries,” *Science* **358** (6363), 636–640.
- Bahari, Babak, Ricardo Tellez-Limon, and Boubacar Kanté (2016), “Topological terahertz circuits using semiconductors,” *Applied Physics Letters* **109** (14), 143501.
- Balents, Leon, and Matthew P. A. Fisher (1996), “Chiral

- surface states in the bulk quantum hall effect,” *Phys. Rev. Lett.* **76**, 2782–2785.
- Bandres, Miguel A, Mikael C. Rechtsman, and Mordechai Segev (2016), “Topological photonic quasicrystals: Fractal topological spectrum and protected transport,” *Phys. Rev. X* **6**, 011016.
- Bandres, Miguel A, Steffen Wittek, Gal Harari, Midya Parto, Jinhan Ren, Mordechai Segev, Demetrios N Christodoulides, and Mercedeh Khajavikhan (2018a), “Topological insulator laser: Experiments,” *Science* **359** (6381), eaar4005.
- Bandres, Miguel A, Steffen Wittek, Gal Harari, Midya Parto, Jinhan Ren, Mordechai Segev, Demetrios N. Christodoulides, and Mercedeh Khajavikhan (2018b), “Topological insulator laser: Experiments,” *Science* [10.1126/science.aar4005](https://doi.org/10.1126/science.aar4005).
- Barbarino, Simone, Luca Taddia, Davide Rossini, Leonardo Mazza, and Rosario Fazio (2016), “Synthetic gauge fields in synthetic dimensions: interactions and chiral edge modes,” *New Journal of Physics* **18** (3), 035010.
- Bardyn, C-E, M A Baranov, C V Kraus, E Rico, A mamolu, P Zoller, and S Diehl (2013), “Topology by dissipation,” *New Journal of Physics* **15** (8), 085001.
- Bardyn, C-E, M. A. Baranov, E. Rico, A. İmamoglu, P. Zoller, and S. Diehl (2012), “Majorana modes in driven-dissipative atomic superfluids with a zero chern number,” *Phys. Rev. Lett.* **109**, 130402.
- Bardyn, C-E, and A. İmamoglu (2012), “Majorana-like modes of light in a one-dimensional array of nonlinear cavities,” *Phys. Rev. Lett.* **109**, 253606.
- Bardyn, Charles-Edouard, Sebastian D Huber, and Oded Zeitlinger (2014), “Measuring topological invariants in small photonic lattices,” *New Journal of Physics* **16** (12), 123013.
- Bardyn, Charles-Edouard, Torsten Karzig, Gil Refael, and Timothy C. H. Liew (2015), “Topological polaritons and excitons in garden-variety systems,” *Phys. Rev. B* **91**, 161413.
- Bardyn, Charles-Edouard, Torsten Karzig, Gil Refael, and Timothy C. H. Liew (2016), “Chiral bogoliubov excitations in nonlinear bosonic systems,” *Phys. Rev. B* **93**, 020502.
- Barik, Sabyasachi, Aziz Karasahin, Christopher Flower, Tao Cai, Hirokazu Miyake, Wade DeGottardi, Mohammad Hafezi, and Edo Waks (2018), “A topological quantum optics interface,” *Science* **359** (6376), 666–668.
- Barik, Sabyasachi, Hirokazu Miyake, Wade DeGottardi, Edo Waks, and Mohammad Hafezi (2016), “Two-dimensionally confined topological edge states in photonic crystals,” *New Journal of Physics* **18** (11), 113013.
- Barkhofen, Sonja, Thomas Nitsche, Fabian Elster, Lennart Lorz, Aurél Gábris, Igor Jex, and Christine Silberhorn (2017), “Measuring topological invariants in disordered discrete-time quantum walks,” *Phys. Rev. A* **96**, 033846.
- Barnett, Ryan (2013), “Edge-state instabilities of bosons in a topological band,” *Phys. Rev. A* **88**, 063631.
- Bayindir, Mehmet, B Temelkuran, and E Ozbay (2000), “Tight-binding description of the coupled defect modes in three-dimensional photonic crystals,” *Physical Review Letters* **84** (10), 2140.
- Bellec, M, C. Michel, H. Zhang, S. Tzortzakis, and P. Delplace (2017), “Non-diffracting states in one-dimensional floquet photonic topological insulators,” *EPL* **119**, 14003.
- Bellec, Matthieu, Ulrich Kuhl, Gilles Montambaux, and Fabrice Mortessagne (2013a), “Tight-binding couplings in microwave artificial graphene,” *Phys. Rev. B* **88**, 115437.
- Bellec, Matthieu, Ulrich Kuhl, Gilles Montambaux, and Fabrice Mortessagne (2013b), “Topological Transition of Dirac Points in a Microwave Experiment,” *Physical Review Letters* **110** (3), 033902.
- Bellec, Matthieu, Ulrich Kuhl, Gilles Montambaux, and Fabrice Mortessagne (2014), “Manipulation of edge states in microwave artificial graphene,” *New Journal of Physics* **16** (11), 113023.
- Bender, Carl M, and Stefan Boettcher (1998), “Real spectra in non-hermitian hamiltonians having p t symmetry,” *Physical Review Letters* **80** (24), 5243.
- Berceanu, Andrei C, Hannah M. Price, Tomoki Ozawa, and Iacopo Carusotto (2016), “Momentum-space landau levels in driven-dissipative cavity arrays,” *Phys. Rev. A* **93**, 013827.
- Bermudez, Alejandro, Tobias Schaetz, and Diego Porras (2011), “Synthetic gauge fields for vibrational excitations of trapped ions,” *Phys. Rev. Lett.* **107**, 150501.
- Bernevig, B Andrei, and Taylor L. Hughes (2013), *Topological insulators and topological superconductors* (Princeton University Press).
- Bernevig, B Andrei, Taylor L. Hughes, Srinivas Raghu, and Daniel P. Arovas (2007), “Theory of the three-dimensional quantum hall effect in graphite,” *Phys. Rev. Lett.* **99**, 146804.
- Bernevig, B Andrei, Taylor L. Hughes, and Shou-Cheng Zhang (2006), “Quantum spin hall effect and topological phase transition in hgte quantum wells,” *Science* **314** (5806), 1757–1761.
- Bernevig, B Andrei, and Shou-Cheng Zhang (2006), “Quantum spin hall effect,” *Phys. Rev. Lett.* **96**, 106802.
- Berry, Michael V (1984), “Quantal phase factors accompanying adiabatic changes,” in *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, Vol. 392 (The Royal Society) pp. 45–57.
- Bi, Ren, and Zhong Wang (2015), “Unidirectional transport in electronic and photonic weyl materials by dirac mass engineering,” *Physical Review B* **92** (24), 241109.
- Bi, Ren, Zhongbo Yan, Ling Lu, and Zhong Wang (2017a), “Nodal-knot semimetals,” arXiv preprint arXiv:1704.06849.
- Bi, Ren, Zhongbo Yan, Ling Lu, and Zhong Wang (2017b), “Topological defects in floquet systems: Anomalous chiral modes and topological invariant,” *Phys. Rev. B* **95**, 161115.
- Biella, Alberto, Florent Storme, José Lebreuilly, Davide Rossini, Rosario Fazio, Iacopo Carusotto, and Cristiano Ciuti (2017), “Phase diagram of incoherently driven strongly correlated photonic lattices,” *Phys. Rev. A* **96**, 023839.
- Bienias, P, S. Choi, O. Firstenberg, M. F. Maghrebi, M. Gullans, M. D. Lukin, A. V. Gorshkov, and H. P. Büchler (2014), “Scattering resonances and bound states for strongly interacting rydberg polaritons,” *Phys. Rev. A* **90**, 053804.
- Bilitewski, Thomas, and Nigel R. Cooper (2016), “Synthetic dimensions in the strong-coupling limit: Supersolids and pair superfluids,” *Phys. Rev. A* **94**, 023630.
- Birnbaum, KM, A Boca, R Miller, AD Boozer, TE Northup, and HJ Kimble (2005), “Photon blockade in an optical cavity with one trapped atom,” *Nature* **436** (7047), 87–90.
- Biswas, Rudro R, and Dam Thanh Son (2016), “Fractional charge and inter-landau-level states at points of singular curvature,” *Proceedings of the National Academy of Sci-*

- ences **113** (31), 8636–8641.
- Bittner, S, B. Dietz, M. Miski-Oglu, P. Oria Iriarte, A. Richter, and F. Schäfer (2010), “Observation of a dirac point in microwave experiments with a photonic crystal modeling graphene,” *Phys. Rev. B* **82**, 014301.
- Bittner, S, B. Dietz, M. Miski-Oglu, and A. Richter (2012), “Extremal transmission through a microwave photonic crystal and the observation of edge states in a rectangular Dirac billiard,” *Physical Review B* **85** (6), 064301.
- Bleckmann, Felix, Zlata Cherpakova, Stefan Linden, and Andrea Alberti (2017), “Spectral imaging of topological edge states in plasmonic waveguide arrays,” *Phys. Rev. B* **96**, 045417.
- Bleu, O, G. Malpuech, and D. D. Solnyshkov (2017a), “ z_2 topological insulator analog for vortices in an interacting bosonic quantum fluid,” arXiv:1709.01830 [1709.01830](#).
- Bleu, O, D. D. Solnyshkov, and G. Malpuech (2016), “Interacting quantum fluid in a polariton chern insulator,” *Phys. Rev. B* **93**, 085438.
- Bleu, O, D. D. Solnyshkov, and G. Malpuech (2017b), “Photonic versus electronic quantum anomalous hall effect,” *Phys. Rev. B* **95**, 115415.
- Bleu, O, D D Solnyshkov, and G Malpuech (2017c), “Quantum valley Hall effect and perfect valley filter based on photonic analogs of transitional metal dichalcogenides,” *Phys. Rev. B* **95** (23), 235431.
- Bleu, O, D. D. Solnyshkov, and G. Malpuech (2018), “Measuring the quantum geometric tensor in two-dimensional photonic and exciton-polariton systems,” *Phys. Rev. B* **97**, 195422.
- Bliokh, K Yu, and Yu. P. Bliokh (2005), “Spin gauge fields: From berry phase to topological spin transport and hall effects,” *Annals of Physics* **319** (1), 13 – 47.
- Bliokh, Konstantin Y, Aleksandr Y Bekshaev, and Franco Nori (2014), “Extraordinary momentum and spin in evanescent waves,” *Nature communications* **5**, 3300.
- Bliokh, Konstantin Y, and Franco Nori (2012), “Transverse spin of a surface polariton,” *Phys. Rev. A* **85**, 061801.
- Bliokh, Konstantin Y, Daria Smirnova, and Franco Nori (2015a), “Quantum spin hall effect of light,” *Science* **348** (6242), 1448–1451.
- Bliokh, KY, FJ Rodríguez-Fortuño, Franco Nori, and Anatoly V Zayats (2015b), “Spin-orbit interactions of light,” *Nature Photonics* **9** (12), 796–808.
- Boada, O, A. Celi, J. I. Latorre, and M. Lewenstein (2012), “Quantum simulation of an extra dimension,” *Phys. Rev. Lett.* **108**, 133001.
- Boada, Octavi, Alessio Celi, Javier Rodríguez-Laguna, José I Latorre, and Maciej Lewenstein (2015), “Quantum simulation of non-trivial topology,” *New Journal of Physics* **17** (4), 045007.
- Bourassa, J, F. Beaudoin, Jay M. Gambetta, and A. Blais (2012), “Josephson-junction-embedded transmission-line resonators: From kerr medium to in-line transmon,” *Phys. Rev. A* **86**, 013814.
- Boyd, R W (2008), *Nonlinear Optics* (Academic Press).
- Bravo-Abad, Jorge, Ling Lu, Liang Fu, Hrvoje Buljan, and Marin Soljačić (2015), “Weyl points in photonic-crystal superlattices,” *2D Materials* **2** (3), 034013.
- Brendel, Christian, Vittorio Peano, Oskar Painter, and Florian Marquardt (2017), “Snowflake Topological Insulator for Sound Waves,” arXiv:1701.06330 [arXiv:1701.06330](#).
- Broome, M A, A. Fedrizzi, B. P. Lanyon, I. Kassal, A. Aspuru-Guzik, and A. G. White (2010), “Discrete single-photon quantum walks with tunable decoherence,” *Phys. Rev. Lett.* **104**, 153602.
- Brouwer, P W (1998), “Scattering approach to parametric pumping,” *Phys. Rev. B* **58**, R10135–R10138.
- Budich, Jan Carl, and Sebastian Diehl (2015), “Topology of density matrices,” *Phys. Rev. B* **91**, 165140.
- Budich, Jan Carl, Peter Zoller, and Sebastian Diehl (2015), “Dissipative preparation of chern insulators,” *Phys. Rev. A* **91**, 042117.
- Bukov, Marin, Luca D’Alessio, and Anatoli Polkovnikov (2015), “Universal high-frequency behavior of periodically driven systems: from dynamical stabilization to floquet engineering,” *Advances in Physics* **64** (2), 139–226.
- Burkov, AA, MD Hook, and Leon Balents (2011), “Topological nodal semimetals,” *Physical Review B* **84** (23), 235126.
- Butcher, P N, and D. Cotter (2008), *The elements of nonlinear optics*, Cambridge Studies in Modern Optics (Cambridge University Press).
- Büttiker, M (1988), “Absence of backscattering in the quantum hall effect in multiprobe conductors,” *Phys. Rev. B* **38**, 9375–9389.
- Bzdušek, Tomáš, Quansheng Wu, Andreas Rüegg, Manfred Sigrist, and Alexey A Soluyanov (2016), “Nodal-chain metals,” *Nature* **538**, 75–78.
- Cai, Y, J Roslund, G Ferrini, F Arzani, X Xu, C Fabre, and N Treps (2017), “Multimode entanglement in reconfigurable graph states using optical frequency combs,” *Nature communications* **8**, 15645.
- Callias, Constantine (1978), “Axial anomalies and index theorems on open spaces,” *Communications in Mathematical Physics* **62** (3), 213–234.
- Calvanese Strinati, Marcello, Eyal Cornfeld, Davide Rossini, Simone Barbarino, Marcello Dalmonte, Rosario Fazio, Eran Sela, and Leonardo Mazza (2017), “Laughlin-like states in bosonic and fermionic atomic synthetic ladders,” *Phys. Rev. X* **7**, 021033.
- Can, T, YH Chiu, M Laskin, and P Wiegmann (2016), “Emergent conformal symmetry and geometric transport properties of quantum hall states on singular surfaces,” *Physical Review Letters* **117** (26), 266803.
- Cardano, Filippo, Alessio D’Errico, Alexandre Dauphin, Maria Maffei, Bruno Piccirillo, Corrado de Lisio, Giulio De Filippis, Vittorio Cataudella, Enrico Santamato, Lorenzo Marrucci, *et al.* (2017), “Detection of zak phases and topological invariants in a chiral quantum walk of twisted photons,” *Nature Communications* **8**, 15516.
- Cardano, Filippo, Maria Maffei, Francesco Massa, Bruno Piccirillo, Corrado De Lisio, Giulio De Filippis, Vittorio Cataudella, Enrico Santamato, and Lorenzo Marrucci (2016), “Statistical moments of quantum-walk dynamics reveal topological quantum transitions,” *Nature communications* **7**.
- Cardano, Filippo, Francesco Massa, Hammam Qassim, Ebrahim Karimi, Sergei Slussarenko, Domenico Paparo, Corrado de Lisio, Fabio Sciarrino, Enrico Santamato, Robert W Boyd, *et al.* (2015), “Quantum walks and wavepacket dynamics on a lattice with twisted photons,” *Science advances* **1** (2), e1500087.
- Carpentier, David, Pierre Delplace, Michel Fruchart, and Krzysztof Gawedzki (2015), “Topological Index for Periodically Driven Time-Reversal Invariant 2D Systems,” *Phys. Rev. Lett.* **114** (10), 106806.
- Carusotto, I, D. Gerace, H. E. Tureci, S. De Liberato, C. Ciuti, and A. Imamoglu (2009), “Fermionized photons

- in an array of driven dissipative nonlinear cavities,” *Phys. Rev. Lett.* **103**, 033601.
- Carusotto, I, T. Volz, and A. Imamoglu (2010), “Feshbach blockade: Single-photon nonlinear optics using resonantly enhanced cavity polariton scattering from biexciton states,” *EPL* **90** (3), 37001.
- Carusotto, Iacopo, and Cristiano Ciuti (2013), “Quantum fluids of light,” *Rev. Mod. Phys.* **85**, 299–366.
- Castro Neto, A H, F Guinea, N M R Peres, K S Novoselov, and A K Geim (2009), “The electronic properties of graphene,” *Rev. Mod. Phys.* **81** (1), 109–162.
- Cayssol, J, B. Dora, F. Simon, and R. Moessner (2013), “Floquet topological insulators,” *Phys. Status Solidi Rapid Res. Lett.* **7**, 101–108.
- Celi, A, P. Massignan, J. Ruseckas, N. Goldman, I. B. Spielman, G. Juzeliūnas, and M. Lewenstein (2014), “Synthetic gauge fields in synthetic dimensions,” *Phys. Rev. Lett.* **112**, 043001.
- Jukić, D, and H. Buljan (2013), “Four-dimensional photonic lattices and discrete tesseract solitons,” *Phys. Rev. A* **87**, 013814.
- Chabanov, A A, and A. Genack (2001), “Statistics of Dynamics of Localized Waves,” *Phys. Rev. Lett.* **87** (23), 233903.
- Chalker, J T, and A. Dohmen (1995), “Three-dimensional disordered conductors in a strong magnetic field: Surface states and quantum hall plateaus,” *Phys. Rev. Lett.* **75**, 4496–4499.
- Chang, Cui-Zu, Jinsong Zhang, Xiao Feng, Jie Shen, Zuocheng Zhang, Minghua Guo, Kang Li, Yunbo Ou, Pang Wei, Li-Li Wang, *et al.* (2013), “Experimental observation of the quantum anomalous hall effect in a magnetic topological insulator,” *Science* **340** (6129), 167–170.
- Chang, Darrick E, Vladan Vuletić, and Mikhail D Lukin (2014), “Quantum nonlinear optics [mdash] photon by photon,” *Nature Photonics* **8** (9), 685–694.
- Chang, Ming-Li, Meng Xiao, Wen-Jie Chen, and Che Ting Chan (2017), “Multiple weyl points and the sign change of their topological charges in woodpile photonic crystals,” *Physical Review B* **95** (12), 125136.
- Cheianov, Vadim V, Vladimir Fal’ko, and B L Altshuler (2007), “The focusing of electron flow and a Veselago lens in graphene p-n junctions.” *Science (New York, N.Y.)* **315** (5816), 1252–5.
- Chen, Ce, Yi-Yian Yin, and DS Elliott (1990), “Interference between optical transitions,” *Physical review letters* **64** (5), 507.
- Chen, Wen-Jie, Shao-Ji Jiang, Xiao-Dong Chen, Baocheng Zhu, Lei Zhou, Jian-Wen Dong, and Che Ting Chan (2014), “Experimental realization of photonic topological insulator in a uniaxial metacrystal waveguide,” *Nature communications* **5**, 5782.
- Chen, Wen-Jie, Meng Xiao, and Che Ting Chan (2016), “Photonic crystals possessing multiple weyl points and the experimental observation of robust surface states,” *Nature communications* **7**, 13038.
- Chen, Xiao-Dong, and Jian-Wen Dong (2016), “Valley-protected backscattering suppression in silicon photonic graphene,” *arXiv preprint arXiv:1602.03352* .
- Cheng, Xiaojun, Camille Jouvaud, Xiang Ni, S Hossein Mousavi, Azriel Z Genack, and Alexander B Khanikaev (2016), “Robust reconfigurable electromagnetic pathways within a photonic topological insulator,” *Nature materials* **15** (5), 542–548.
- Cherpakova, Z, C. Jörg, C. Dauer, F. Letscher, M. Fleischer, S. Eggert, S. Linden, and G. von Freymann (2018), “Breakdown of topological protection under local periodic driving,” *ArXiv e-prints arXiv:1807.02321 [physics.optics]*.
- Chiao, R Y, A. Antaramian, K. M. Ganga, H. Jiao, S. R. Wilkinson, and H. Nathel (1988), “Observation of a topological phase by means of a nonplanar mach-zehnder interferometer,” *Phys. Rev. Lett.* **60**, 1214–1217.
- Chiu, Ching-Kai, Jeffrey C. Y. Teo, Andreas P. Schnyder, and Shinsei Ryu (2016), “Classification of topological quantum matter with symmetries,” *Rev. Mod. Phys.* **88**, 035005.
- Cho, Jaeyoon, Dimitris G. Angelakis, and Sougato Bose (2008), “Fractional quantum hall state in coupled cavities,” *Phys. Rev. Lett.* **101**, 246809.
- Chong, Y D, Xiao-Gang Wen, and Marin Soljačić (2008), “Effective theory of quadratic degeneracies,” *Phys. Rev. B* **77**, 235125.
- Christodoulides, Demetrios N, Falk Lederer, and Yaron Silberberg (2003), “Discretizing light behaviour in linear and nonlinear waveguide lattices,” *Nature* **424** (6950), 817–823.
- Chu, Yiwen, Prashanta Kharel, William H Renninger, Luke D Burkhardt, Luigi Frunzio, Peter T Rakich, and Robert J Schoelkopf (2017), “Quantum acoustics with superconducting qubits,” *Science* , eaao1511.
- Ciuti, C, V Savona, C Piermarocchi, A Quattropani, and P Schwendimann (1998), “Role of the exchange of carriers in elastic exciton-exciton scattering in quantum wells,” *Phys. Rev. B* **58** (12), 7926–7933.
- Claassen, Martin, Ching Hua Lee, Ronny Thomale, Xiao-Liang Qi, and Thomas P Devereaux (2015), “Position-momentum duality and fractional quantum hall effect in chern insulators,” *Physical review letters* **114** (23), 236802.
- Clark, Jeremy B, Florent Lecocq, Raymond W Simmonds, José Aumentado, and John D Teufel (2017), “Sideband cooling beyond the quantum backaction limit with squeezed light,” *Nature* **541** (7636), 191–195.
- Cohen-Tannoudji, Claude, Jacques Dupont-Roc, and Gilbert Grynberg (2008), *AtomPhoton Interactions* (Wiley-VCH Verlag GmbH).
- Cominotti, Marco, and Iacopo Carusotto (2013), “Berry curvature effects in the bloch oscillations of a quantum particle under a strong (synthetic) magnetic field,” *EPL (Europhysics Letters)* **103** (1), 10001.
- Cooper, Michael L, Greeshma Gupta, Mark A. Schneider, William M. J. Green, Solomon Assefa, Fengnian Xia, Yurii A. Vlasov, and Shayan Mookherjea (2010), “Statistics of light transport in 235-ring silicon coupled-resonator optical waveguides.” *Opt. Express* **18** (25), 26505–26516.
- Cooper, Nigel R (2008), “Rapidly rotating atomic gases,” *Advances in Physics* **57** (6), 539–616.
- Creffield, C E, G Pieplow, F Sols, and N Goldman (2016), “Realization of uniform synthetic magnetic fields by periodically shaking an optical square lattice,” *New Journal of Physics* **18** (9), 093013.
- Crespi, Andrea, Giacomo Corrielli, Giuseppe Della Valle, Roberto Osellame, and Stefano Longhi (2013), “Dynamic band collapse in photonic graphene,” *New Journal of Physics* **15** (1), 013012.
- Dalibard, Jean, Fabrice Gerbier, Gediminas Juzeliūnas, and Patrik Öhberg (2011), “Colloquium: Artificial gauge potentials for neutral atoms,” *Reviews of Modern Physics* **83** (4), 1523.
- Dana, Itzhack, Yosi Avron, and J Zak (1985), “Quantised

- hall conductance in a perfect crystal,” *Journal of Physics C: Solid State Physics* **18** (22), L679.
- De Nittis, Giuseppe, and Max Lein (2017), “Symmetry classification of topological photonic crystals,” [arXiv preprint arXiv:1710.08104](https://arxiv.org/abs/1710.08104).
- Delplace, P, D. Ullmo, and G. Montambaux (2011), “Zak phase and the existence of edge states in graphene,” *Physical Review B* **84** (19), 195452.
- Deng, Xiaohua, Lujun Hong, Xiaodong Zheng, and Linfang Shen (2015), “One-way regular electromagnetic mode immune to backscattering,” *Applied optics* **54** (14), 4608–4612.
- Dennis, Mark R, Kevin O’Holleran, and Miles J Padgett (2009), “Singular optics: optical vortices and polarization singularities,” *Progress in Optics* **53**, 293–363.
- Di Liberto, M, A. Hemmerich, and C. Morais Smith (2016a), “Topological varma superfluid in optical lattices,” *Phys. Rev. Lett.* **117**, 163001.
- Di Liberto, M, A Recati, I Carusotto, and C Menotti (2016b), “Two-body physics in the su-schrieffer-heeger model,” *Physical Review A* **94** (6), 062704.
- Diehl, Sebastian, Enrique Rico, Mikhail A Baranov, and Peter Zoller (2011), “Topology by dissipation in atomic quantum wires,” *Nature Physics* **7** (12), 971–977.
- Downing, Charles A, and Guillaume Weick (2017), “Topological collective plasmons in bipartite chains of metallic nanoparticles,” *Physical Review B* **95** (12), 125426.
- Dreisow, F, R Keil, A Tünnermann, S Nolte, S Longhi, and A Szameit (2012), “Klein tunneling of light in waveguide superlattices,” *EPL* **97** (1), 10008.
- Dubček, Tena, Colin J Kennedy, Ling Lu, Wolfgang Ketterle, Marin Soljačić, and Hrvoje Buljan (2015a), “Weyl points in three-dimensional optical lattices: synthetic magnetic monopoles in momentum space,” *Physical review letters* **114** (22), 225301.
- Dubček, Tena, Karlo Lelas, Dario Jukić, Robert Pezer, Marin Soljačić, and Hrvoje Buljan (2015b), “The harper-hofstadter hamiltonian and conical diffraction in photonic lattices with grating assisted tunneling,” *New Journal of Physics* **17** (12), 125002.
- Dudarev, Artem M, Roberto B Diener, Iacopo Carusotto, and Qian Niu (2004), “Spin-orbit coupling and berry phase with ultracold atoms in 2d optical lattices,” *Physical review letters* **92** (15), 153005.
- Dutta, Shovan, and Erich J Mueller (2017), “Coherent generation of photonic fractional quantum hall states in a cavity and the search for anyonic quasiparticles,” [arXiv preprint arXiv:1711.08059](https://arxiv.org/abs/1711.08059).
- Eckardt, A, and E. Anisimovas (2015), “High-frequency approximation for periodically driven quantum systems from a floquet-space perspective,” *New J. Phys.* **17**, 093039.
- Eckardt, André (2016), “Atomic quantum gases in periodically driven optical lattices,” [arXiv:1606.08041](https://arxiv.org/abs/1606.08041).
- Efremidis, Nikos K, Suzanne Sears, Demetrios N Christodoulides, Jason W Fleischer, and Mordechai Segev (2002), “Discrete solitons in photorefractive optically induced photonic lattices,” *Physical Review E* **66** (4), 046602.
- Eisenberg, HS, Ya Silberberg, R Morandotti, AR Boyd, and JS Aitchison (1998), “Discrete spatial optical solitons in waveguide arrays,” *Physical Review Letters* **81** (16), 3383.
- Elliott, Steven R, and Marcel Franz (2015), “Colloquium,” *Rev. Mod. Phys.* **87**, 137–163.
- Engelhardt, G, M. Benito, G. Platero, and T. Brandes (2016), “Topological instabilities in ac-driven bosonic systems,” *Phys. Rev. Lett.* **117**, 045302.
- Engelhardt, G, and T. Brandes (2015), “Topological bogoliubov excitations in inversion-symmetric systems of interacting bosons,” *Phys. Rev. A* **91**, 053621.
- Esaki, Kenta, Masatoshi Sato, Kazuki Hasebe, and Mahito Kohmoto (2011), “Edge states and topological phases in non-hermitian systems,” *Physical Review B* **84** (20), 205128.
- Fang, Chen, Yige Chen, Hae-Young Kee, and Liang Fu (2015), “Topological nodal line semimetals with and without spin-orbital coupling,” *Physical Review B* **92** (8), 081201.
- Fang, Chen, Ling Lu, Junwei Liu, and Liang Fu (2016a), “Topological semimetals with helicoid surface states,” *Nature Physics* **12**, 936–941.
- Fang, Chen, Hongming Weng, Xi Dai, and Zhong Fang (2016b), “Topological nodal line semimetals,” *Chinese Physics B* **25** (11), 117106.
- Fang, Kejie, and Shanhui Fan (2013a), “Controlling the flow of light using the inhomogeneous effective gauge field that emerges from dynamic modulation,” *Phys. Rev. Lett.* **111**, 203901.
- Fang, Kejie, and Shanhui Fan (2013b), “Effective magnetic field for photons based on the magneto-optical effect,” *Phys. Rev. A* **88**, 043847.
- Fang, Kejie, Jie Luo, Anja Metelmann, Matthew H. Matheny, Florian Marquardt, Aashish A. Clerk, and Oskar Painter (2017), “Generalized non-reciprocity in an optomechanical circuit via synthetic magnetism and reservoir engineering,” *Nat Phys* **13** (5), 465–471.
- Fang, Kejie, Zongfu Yu, and Shanhui Fan (2011), “Microscopic theory of photonic one-way edge mode,” *Phys Rev B* **84**, 075477.
- Fang, Kejie, Zongfu Yu, and Shanhui Fan (2012a), “Photonic aharonov-bohm effect based on dynamic modulation,” *Phys. Rev. Lett.* **108**, 153901.
- Fang, Kejie, Zongfu Yu, and Shanhui Fan (2012b), “Realizing effective magnetic field for photons by controlling the phase of dynamic modulation,” *Nature photonics* **6** (11), 782–787.
- Faraon, Andrei, Ilya Fushman, Dirk Englund, Nick Stoltz, Pierre Petroff, and Jelena Vuckovic (2008), “Coherent generation of non-classical light on a chip via photon-induced tunnelling and blockade,” *Nature Phys.* **4** (11), 859–863.
- Fausti, Daniele, RI Tobey, Nicky Dean, Stefan Kaiser, A Dienst, Matthias C Hoffmann, S Pyon, T Takayama, H Takagi, and Andrea Cavalleri (2011), “Light-induced superconductivity in a stripe-ordered cuprate,” *science* **331** (6014), 189–191.
- Fefferman, Charles, and Michael Weinstein (2012), “Honeycomb lattice potentials and dirac points,” *Journal of the American Mathematical Society* **25** (4), 1169–1220.
- Fefferman, Charles L, James P Lee-Thorp, and Michael I Weinstein (2014), “Topologically protected states in one-dimensional continuous systems and dirac points,” *Proceedings of the National Academy of Sciences* **111** (24), 8759–8763.
- Fefferman, Charles L, and Michael I Weinstein (2014), “Wave packets in honeycomb structures and two-dimensional dirac equations,” *Communications in Mathematical Physics* **326** (1), 251–286.
- Firstenberg, Ofer, Thibault Peyronel, Qi-Yu Liang, Alexey V Gorshkov, Mikhail D Lukin, and Vladan Vuletić (2013), “Attractive photons in a quantum nonlinear medium,” *Nature* **502**, 71–75.

- ture **502** (7469), 71–75.
- Fitzpatrick, Mattias, Neereja M. Sundaresan, Andy C. Y. Li, Jens Koch, and Andrew A. Houck (2017), “Observation of a dissipative phase transition in a one-dimensional circuit qed lattice,” *Phys. Rev. X* **7**, 011016.
- Fläschner, N., D. Vogel, M. Tarnowski, B. S. Rem, D.-S. Lühmann, M. Heyl, J. C. Budich, L. Mathey, K. Senstock, and C. Weitenberg (2016), “Observation of a dynamical topological phase transition,” ArXiv e-prints [arXiv:1608.05616](https://arxiv.org/abs/1608.05616) [[cond-mat.quant-gas](https://arxiv.org/abs/1608.05616)].
- Fleischer, Jason W, Mordechai Segev, Nikolaos K Efremidis, and Demetrios N Christodoulides (2003), “Observation of two-dimensional discrete solitons in optically induced nonlinear photonic lattices,” *Nature* **422** (6928), 147–150.
- Fleury, Romain, Dimitrios Sounas, Michael R Haberman, and Andrea Alù (2015), “Nonreciprocal acoustics,” *Acoustics Today* **11** (EPFL-ARTICLE-223074), 14–21.
- Fleury, Romain, Dimitrios L Sounas, Caleb F Sieck, Michael R Haberman, and Andrea Alù (2014), “Sound isolation and giant linear nonreciprocity in a compact acoustic circulator,” *Science* **343** (6170), 516–519.
- Franke-Arnold, Sonja, Graham Gibson, Robert W Boyd, and Miles J Padgett (2011), “Rotary photon drag enhanced by a slow-light medium,” *Science* **333** (6038), 65–67.
- Freedman, Barak, Guy Bartal, Mordechai Segev, Ron Lifshitz, Demetrios N Christodoulides, and Jason W Fleischer (2006), “Wave and defect dynamics in nonlinear photonic quasicrystals,” *Nature* **440** (7088), 1166–1169.
- Freedman, Barak, Ron Lifshitz, Jason W Fleischer, and Mordechai Segev (2007), “Phason dynamics in nonlinear photonic quasicrystals,” *Nature materials* **6** (10), 776–781.
- Fröhlich, J, and B. Pedrini (2000), “New applications of the chiral anomaly,” in *Mathematical Physics 2000*, edited by A. Fokas, A. Grigoryan, T. Kibble, and B. Zegarlinski (Imperial College Press, London, United Kingdom) pp. 9–47.
- Fu, Jin-Xin, Jin Lian, Rong-Juan Liu, Lin Gan, and Zhi-Yuan Li (2011a), “Unidirectional channel-drop filter by one-way gyromagnetic photonic crystal waveguides,” *Applied Physics Letters* **98** (21), 211104–211104.
- Fu, Jin-Xin, Rong-Juan Liu, Lin Gan, and Zhi-Yuan Li (2011b), “Control and blockage of edge modes in magneto-optical photonic crystals,” *EPL (Europhysics Letters)* **93** (2), 24001.
- Fu, Jin-Xin, Rong-Juan Liu, and Zhi-Yuan Li (2010a), “Experimental demonstration of tunable gyromagnetic photonic crystals controlled by dc magnetic fields,” *EPL (Europhysics Letters)* **89** (6), 64003.
- Fu, Jin-Xin, Rong-Juan Liu, and Zhi-Yuan Li (2010b), “Robust one-way modes in gyromagnetic photonic crystal waveguides with different interfaces,” *Applied Physics Letters* **97** (4), 041112–041112.
- Fu, Liang (2011), “Topological crystalline insulators,” *Phys. Rev. Lett.* **106**, 106802.
- Fu, Liang, and C. L. Kane (2006), “Time reversal polarization and a Z_2 adiabatic spin pump,” *Phys. Rev. B* **74**, 195312.
- Fu, Liang, C. L. Kane, and E. J. Mele (2007), “Topological insulators in three dimensions,” *Phys. Rev. Lett.* **98**, 106803.
- Furukawa, Shunsuke, and Masahito Ueda (2015), “Excitation band topology and edge matter waves in bose–einstein condensates in optical lattices,” *New Journal of Physics* **17** (11), 115014.
- Galilo, Bogdan, Derek K. K. Lee, and Ryan Barnett (2015), “Selective population of edge states in a 2d topological band system,” *Phys. Rev. Lett.* **115**, 245302.
- Gangaraj, Seyyed Ali Hassani, and George W Hanson (2017), “Topologically protected unidirectional surface states in biased ferrites: duality and application to directional couplers,” *IEEE Antennas and Wireless Propagation Letters* **16**, 449–452.
- Gao, Fei, Zhen Gao, Xihang Shi, Zhaoju Yang, Xiao Lin, Hongyi Xu, John D Joannopoulos, Marin Soljačić, Hongsheng Chen, Ling Lu, *et al.* (2016a), “Probing topological protection using a designer surface plasmon structure,” *Nature communications* **7**, 11619.
- Gao, Wenlong, Mark Lawrence, Biao Yang, Fu Liu, Fengzhou Fang, Benjamin Béri, Jensen Li, and Shuang Zhang (2015), “Topological photonic phase in chiral hyperbolic metamaterials,” *Physical review letters* **114** (3), 037402.
- Gao, Wenlong, Biao Yang, Mark Lawrence, Fengzhou Fang, Benjamin Béri, and Shuang Zhang (2016b), “Photonic weyl degeneracies in magnetized plasma,” *Nature communications* **7**, 12435.
- Gardiner, C W, and M. J. Collett (1985), “Input and output in damped quantum systems: Quantum stochastic differential equations and the master equation,” *Phys. Rev. A* **31**, 3761–3774.
- Gardiner, C W, and P. Zoller (2004), *Quantum Noise* (Springer Verlag, Berlin).
- Gbur, Gregory J (2016), *Singular Optics* (CRC Press).
- Geerligs, L J, V. F. Anderegg, P. A. M. Holweg, J. E. Mooij, H. Pothier, D. Esteve, C. Urbina, and M. H. Devoret (1990), “Frequency-locked turnstile device for single electrons,” *Phys. Rev. Lett.* **64**, 2691–2694.
- Gerace, Dario, Hakan E. Türeci, Atac Imamoglu, Vittorio Giovannetti, and Rosario Fazio (2009), “The quantum-optical josephson interferometer,” *Nat. Phys.* **5** (4), 281–284.
- Gerry, Christopher, and Peter Knight (2005), *Introductory quantum optics* (Cambridge university press).
- Ghulinyan, Mher, Fernando Ramiro Manzano, Nikola Prtljaga, Martino Bernard, Lorenzo Pavesi, Georg Pucker, and Iacopo Carusotto (2014), “Intermode reactive coupling induced by waveguide-resonator interaction,” *Physical Review A* **90** (5), 053811.
- Goldman, N, JC Budich, and P Zoller (2016a), “Topological quantum matter with ultracold gases in optical lattices,” *Nature Physics* **12** (7), 639–645.
- Goldman, N, and J. Dalibard (2014), “Periodically driven quantum systems: Effective hamiltonians and engineered gauge fields,” *Phys. Rev. X* **4**, 031027.
- Goldman, N, J. Dalibard, M. Aidelsburger, and N. R. Cooper (2015), “Periodically driven quantum matter: The case of resonant modulations,” *Phys. Rev. A* **91**, 033632.
- Goldman, N, G. Jotzu, M. Messer, F. Görg, R. Desbuquois, and T. Esslinger (2016b), “Creating topological interfaces and detecting chiral edge modes in a two-dimensional optical lattice,” *Phys. Rev. A* **94**, 043611.
- Goldman, N, I. Satija, P. Nikolic, A. Bermudez, M. A. Martin-Delgado, M. Lewenstein, and I. B. Spielman (2010), “Realistic time-reversal invariant topological insulators with neutral atoms,” *Phys. Rev. Lett.* **105**, 255302.
- Goldman, Nathan, G Juzeliūnas, P Öhberg, and Ian B Spielman (2014), “Light-induced gauge fields for ultracold atoms,” *Reports on Progress in Physics* **77** (12), 126401.
- Gong, Zongping, Yuto Ashida, Kohei Kawabata, Kazuaki Takasan, Sho Higashikawa, and Masahito Ueda (2018),

- “Topological phases of non-hermitian systems,” arXiv preprint arXiv:1802.07964.
- Gopalakrishnan, Sarang, Pouyan Ghaemi, and Shinsei Ryu (2012), “Non-Abelian $SU(2)$ gauge fields through density wave order and strain in graphene,” *Physical Review B* **86** (8), 081403.
- Goren, Tal, Kirill Plekhanov, Félicien Appas, and Karyn Le Hur (2018), “Topological zak phase in strongly coupled lc circuits,” *Phys. Rev. B* **97**, 041106.
- Gorlach, Maxim, Marco Di Liberto, and ... (2018)), “...” .
- Gorlach, Maxim A, Xiang Ni, Daria A Smirnova, Dmitry Korobkin, Dmitry Zhirihin, Alexey P Slobozhanyuk, Pavel A Belov, Andrea Alù, and Alexander B Khanikaev (2018), “Far-field probing of leaky topological states in all-dielectric metasurfaces,” *Nat. Commun.* **9** (1), 909.
- Gorlach, Maxim A, and Alexander N Poddubny (2017), “Interaction-induced two-photon edge states in an extended hubbard model realized in a cavity array,” *Physical Review A* **95** (3), 033831.
- Gorshkov, Alexey V, Johannes Otterbach, Michael Fleischhauer, Thomas Pohl, and Mikhail D. Lukin (2011), “Photon-photon interactions via rydberg blockade,” *Phys. Rev. Lett.* **107**, 133602.
- Graß, Tobias, Alessio Celi, and Maciej Lewenstein (2014), “Quantum magnetism of ultracold atoms with a dynamical pseudospin degree of freedom,” *Phys. Rev. A* **90**, 043628.
- Graß, Tobias, Christine Muschik, Alessio Celi, Ravindra W. Chhajlany, and Maciej Lewenstein (2015), “Synthetic magnetic fluxes and topological order in one-dimensional spin systems,” *Phys. Rev. A* **91**, 063612.
- Greentree, Andrew D, Charles Tahan, Jared H. Cole, and Lloyd C. L. Hollenberg (2006), “Quantum phase transitions of light,” *Nature Phys.* **2** (12), 856–861.
- Grosche, Simon, Alexander Szameit, and Marco Ornigotti (2016), “Spatial Goos-Hänchen shift in photonic graphene,” *Physical Review A* **94** (6), 063831.
- Grusdt, Fabian, and Michael Fleischhauer (2013), “Fractional quantum hall physics with ultracold rydberg gases in artificial gauge fields,” *Physical Review A* **87** (4), 043628.
- Grusdt, Fabian, Fabian Letscher, Mohammad Hafezi, and Michael Fleischhauer (2014), “Topological growing of Laughlin states in synthetic gauge fields,” *Physical Review Letters* **113** (15), 155301.
- Grusdt, Fabian, Johannes Otterbach, and Michael Fleischhauer (2013), personal communication.
- Gu, Xiu, Anton Frisk Kockum, Adam Miranowicz, Yu-xi Liu, and Franco Nori (2017), “Microwave photonics with superconducting quantum circuits,” *Physics Reports* **718-719**, 1–102.
- Guglielmon, Jonathan, Sheng Huang, Kevin P Chen, and Mikael C Rechtsman (2017), “Prediction and realization of a photonic topological phase transition,” in *CLEO: QELS/Fundamental Science* (Optical Society of America) pp. FM2G–6.
- Guinea, F, M. I. Katsnelson, and A. K. Geim (2010), “Energy gaps and a zero-field quantum Hall effect in graphene by strain engineering,” *Nature Physics* **6** (1), 30–33.
- Gulevich, Dmitry R, Dmitry Yudin, Dmitry V Skryabin, Ivan V Iorsh, and Ivan A Shelykh (2017), “Exploring nonlinear topological states of matter with exciton-polaritons: Edge solitons in Kagome lattice,” *Scientific Reports* **7** (1), 1780.
- Gullans, M J, J. D. Thompson, Y. Wang, Q.-Y. Liang, V. Vuletić, M. D. Lukin, and A. V. Gorshkov (2016), “Effective field theory for rydberg polaritons,” *Phys. Rev. Lett.* **117**, 113601.
- Guo, Qinghua, Biao Yang, Lingbo Xia, Wenlong Gao, Hongchao Liu, Jing Chen, Yuanjiang Xiang, and Shuang Zhang (2017), “Three dimensional photonic dirac points in metamaterials,” arXiv preprint arXiv:1706.08173.
- Gustafsson, Martin V, Thomas Aref, Anton Frisk Kockum, Maria K Ekström, Göran Johansson, and Per Delsing (2014), “Propagating phonons coupled to an artificial atom,” *Science* **346** (6206), 207–211.
- Habraken, SJM, K Stannigel, Mikhail D Lukin, P Zoller, and P Rabl (2012), “Continuous mode cooling and phonon routers for phononic quantum networks,” *New Journal of Physics* **14** (11), 115004.
- Hadad, Yakir, Alexander B. Khanikaev, and Andrea Alù (2016), “Self-induced topological transitions and edge states supported by nonlinear staggered potentials,” *Phys. Rev. B* **93**, 155112.
- Haddad, LH, and LD Carr (2011), “Relativistic linear stability equations for the nonlinear dirac equation in bose-einstein condensates,” *EPL (Europhysics Letters)* **94** (5), 56002.
- Haddad, LH, and Lincoln D Carr (2015), “The nonlinear dirac equation in bose-einstein condensates: vortex solutions and spectra in a weak harmonic trap,” *New Journal of Physics* **17** (11), 113011.
- Haddad, LH, CM Weaver, and Lincoln D Carr (2015), “The nonlinear dirac equation in bose-einstein condensates: I. relativistic solitons in armchair nanoribbon optical lattices,” *New Journal of Physics* **17** (6), 063033.
- Hafezi, M, P Adhikari, and JM Taylor (2015), “Chemical potential for light by parametric coupling,” *Physical Review B* **92** (17), 174305.
- Hafezi, Mohammad (2014), “Measuring topological invariants in photonic systems,” *Phys. Rev. Lett.* **112**, 210405.
- Hafezi, Mohammad, Prabin Adhikari, and Jacob M Taylor (2014), “Engineering three-body interaction and pfaffian states in circuit qed systems,” *Physical Review B* **90** (6), 060503.
- Hafezi, Mohammad, Eugene A Demler, Mikhail D Lukin, and Jacob M Taylor (2011), “Robust optical delay lines with topological protection,” *Nature Physics* **7** (11), 907–912.
- Hafezi, Mohammad, Mikhail D Lukin, and Jacob M Taylor (2013a), “Non-equilibrium fractional quantum hall state of light,” *New Journal of Physics* **15** (6), 063001.
- Hafezi, Mohammad, S Mittal, J Fan, A Migdall, and JM Taylor (2013b), “Imaging topological edge states in silicon photonics,” *Nature Photonics* **7** (12), 1001–1005.
- Hafezi, Mohammad, and Peter Rabl (2012), “Optomechanically induced non-reciprocity in microring resonators,” *Optics express* **20** (7), 7672–7684.
- Haldane, F D M (1988), “Model for a quantum hall effect without landau levels: Condensed-matter realization of the “parity anomaly”,” *Phys. Rev. Lett.* **61**, 2015–2018.
- Haldane, F D M, and S. Raghu (2008), “Possible realization of directional optical waveguides in photonic crystals with broken time-reversal symmetry,” *Phys. Rev. Lett.* **100**, 013904.
- Halperin, Bertrand I (1982), “Quantized Hall Conductance, Current Carrying Edge States and Extended States in 2D Disordered Potential,” *Phys. Rev. B* **25** (4), 2185–2190.
- Halperin, Bertrand I (1987), “Possible states for a three-dimensional electron gas in a strong magnetic field,” *Japanese Journal of Applied Physics* **26** (S3-3), 1913.

- Hannay, J H (1985), “Angle variable holonomy in adiabatic excursion of an integrable hamiltonian,” *Journal of Physics A: Mathematical and General* **18** (2), 221.
- Harari, Gal, Miguel A. Bandres, Yaakov Lumer, Mikael C. Rechtsman, Y. D. Chong, Mercedeh Khajavikhan, Demetrios N. Christodoulides, and Mordechai Segev (2018a), “Topological insulator laser: Theory,” *Science* [10.1126/science.aar4003](https://doi.org/10.1126/science.aar4003).
- Harari, Gal, Miguel A Bandres, Yaakov Lumer, Mikael C Rechtsman, YD Chong, Mercedeh Khajavikhan, Demetrios N Christodoulides, and Mordechai Segev (2018b), “Topological insulator laser: Theory,” *Science* **359** (6381), eaar4003.
- Harper, P G (1955a), “Scaling analysis of quasiperiodic systems: Generalized harper model,” *Proc. Phys. Soc. London A* **68**, 874.
- Harper, P G (1955b), “Single band motion of conduction electrons in a uniform magnetic field,” *Proceedings of the Physical Society. Section A* **68** (10), 874.
- Harris, S E (1989), “Lasers without inversion: Interference of lifetime-broadened resonances,” *Phys. Rev. Lett.* **62**, 1033–1036.
- Hartmann, Michael J, Fernando G. S. L. Brandão, and Martin B. Plenio (2006), “Strongly interacting polaritons in coupled arrays of cavities,” *Nature Phys.* **2** (12), 849–855.
- Hartstein, A, E Burstein, AA Maradudin, R Brewer, and RF Wallis (1973), “Surface polaritons on semi-infinite gyromagnetic media,” *Journal of Physics C: Solid State Physics* **6** (7), 1266.
- Hasan, M Z, and C. L. Kane (2010), “Colloquium,” *Rev. Mod. Phys.* **82**, 3045–3067.
- Hatsugai, Yasuhiro (1993a), “Chern number and edge states in the integer quantum hall effect,” *Phys. Rev. Lett.* **71**, 3697–3700.
- Hatsugai, Yasuhiro (1993b), “Edge states in the integer quantum hall effect and the riemann surface of the bloch function,” *Phys. Rev. B* **48**, 11851–11862.
- Hatsugai, Yasuhiro, and Mahito Kohmoto (1990), “Energy spectrum and the quantum hall effect on the square lattice with next-nearest-neighbor hopping,” *Physical review B* **42** (13), 8282.
- Haus, Hermann A, and Weiping Huang (1991), “Coupled-mode theory,” *Proceedings of the IEEE* **79** (10), 1505–1518.
- Hayward, Andrew L C, Andrew M. Martin, and Andrew D. Greentree (2012), “Fractional quantum hall physics in jaynes-cummings-hubbard lattices,” *Phys. Rev. Lett.* **108**, 223602.
- He, Cheng, Xiao-Lin Chen, Ming-Hui Lu, Xue-Feng Li, Wei-Wei Wan, Xiao-Shi Qian, Ruo-Cheng Yin, and Yan-Feng Chen (2010a), “Left-handed and right-handed one-way edge modes in a gyromagnetic photonic crystal,” *Journal of Applied Physics* **107** (12), 123117.
- He, Cheng, Xiao-Lin Chen, Ming-Hui Lu, Xue-Feng Li, Wei-Wei Wan, Xiao-Shi Qian, Ruo-Cheng Yin, and Yan-Feng Chen (2010b), “Tunable one-way cross-waveguide splitter based on gyromagnetic photonic crystal,” *Applied Physics Letters* **96** (11), 111111.
- He, Cheng, Xu Ni, Hao Ge, Xiao-Chen Sun, Yan-Bin Chen, Ming-Hui Lu, Xiao-Ping Liu, and Yan-Feng Chen (2016a), “Acoustic topological insulator and robust one-way sound transport,” *Nature Physics* **12** (12), 1124–1129.
- He, Cheng, Xiao-Chen Sun, Xiao-Ping Liu, Ming-Hui Lu, Yulin Chen, Liang Feng, and Yan-Feng Chen (2016b), “Photonic topological insulator with broken time-reversal symmetry,” *Proceedings of the National Academy of Sciences* **113** (18), 4924–4928.
- Heisenberg, W, and H Euler (1936), “Folgerungen aus der diracschen theorie des positrons,” *Zeitschrift für Physik A Hadrons and Nuclei* **98** (11), 714–732.
- Hoffman, A J, S. J. Srinivasan, S. Schmidt, L. Spietz, J. Aumentado, H. E. Türeci, and A. A. Houck (2011), “Dispersive photon blockade in a superconducting circuit,” *Phys. Rev. Lett.* **107**, 053602.
- Hofstadter, Douglas R (1976), “Energy levels and wave functions of bloch electrons in rational and irrational magnetic fields,” *Phys. Rev. B* **14**, 2239–2249.
- Houck, Andrew A, Hakan E Türeci, and Jens Koch (2012), “On-chip quantum simulation with superconducting circuits,” *Nature Physics* **8** (4), 292–299.
- Hsu, Chia Wei, Bo Zhen, A Douglas Stone, John D Joannopoulos, and Marin Soljačić (2016), “Bound states in the continuum,” *Nature Reviews Materials* **1**, 16048.
- Hu, Wenchao, Jason C. Pillay, Kan Wu, Michael Pasek, Perry Ping Shum, and Y. D. Chong (2015), “Measurement of a topological edge invariant in a microwave network,” *Phys. Rev. X* **5**, 011012.
- Hu, Yi Chen, and Taylor L Hughes (2011), “Absence of topological insulator phases in non-hermitian p t-symmetric hamiltonians,” *Physical Review B* **84** (15), 153101.
- Hua, Shiyue, Jianming Wen, Xiaoshun Jiang, Qian Hua, Liang Jiang, and Min Xiao (2016), “Demonstration of a chip-based optical isolator with parametric amplification,” *Nature communications* **7**, 13657.
- Huang, K (1987), *Statistical Mechanics* (Wiley).
- Huang, Xueqin, Yun Lai, Zhi Hong Hang, Huihuo Zheng, and CT Chan (2011), “Dirac cones induced by accidental degeneracy in photonic crystals and zero-refractive-index materials,” *Nature materials* **10** (8), 582–586.
- Huber, Sebastian D (2016), “Topological mechanics,” *Nature Physics* **12** (7), 621–623.
- Iadecola, Thomas, Thomas Schuster, and Claudio Chamon (2016), “Non-abelian braiding of light,” *Phys. Rev. Lett.* **117**, 073901.
- Imamoglu, A, H Schmidt, G Woods, and M Deutsch (1997), “Strongly interacting photons in a nonlinear cavity,” *Phys. Rev. Lett.* **79** (8), 1467–1470.
- Imhof, Stefan, Christian Berger, Florian Bayer, Johannes Brehm, Laurens W Molenkamp, Tobias Kiessling, Frank Schindler, Ching Hua Lee, Martin Greiter, Titus Neupert, et al. (2018), “Topoelectrical-circuit realization of topological corner modes,” *Nature Physics* **14** (9), 925.
- Inoue, Jun-ichi, and Akihiro Tanaka (2010), “Photoinduced transition between conventional and topological insulators in two-dimensional electronic systems,” *Phys. Rev. Lett.* **105**, 017401.
- Ivanov, Peter A, Fabian Letscher, Jonathan Simon, and Michael Fleischhauer (2018), “Adiabatic flux insertion and growing of laughlin states of cavity rydberg polaritons,” arXiv preprint arXiv:1803.04156.
- Jachymski, Krzysztof, Przemysław Bienias, and Hans Peter Büchler (2016), “Three-body interaction of rydberg slow-light polaritons,” *Physical review letters* **117** (5), 053601.
- Jackiw, R, and C. Rebbi (1976), “Solitons with fermion number 1/2,” *Phys. Rev. D* **13**, 3398–3409.
- Jacobs, Daniel A, Andrey E Miroshnichenko, Yuri S Kivshar, and Alexander B Khanikaev (2015), “Photonic topological chern insulators based on tellegen metacrystals,” *New Journal of Physics* **17** (12), 125015.

- Jacqmin, T, I. Carusotto, I. Sagnes, M. Abbarchi, D. D. Solnyshkov, G. Malpuech, E. Galopin, A. Lemaitre, J. Bloch, and A. Amo (2014), “Direct observation of dirac cones and a flatband in a honeycomb lattice for polaritons,” *Phys. Rev. Lett.* **112**, 116402.
- Jaksch, D, and P Zoller (2003), “Creation of effective magnetic fields in optical lattices: the hofstadter butterfly for cold neutral atoms,” *New Journal of Physics* **5** (1), 56.
- Jalas, Dirk, Alexander Petrov, Manfred Eich, Wolfgang Freude, Shanhui Fan, Zongfu Yu, Roel Baets, Miloš Popović, Andrea Melloni, John D Joannopoulos, *et al.* (2013), “What is [mdash] and what is not [mdash] an optical isolator,” *Nature Photonics* **7** (8), 579–582.
- Jia, Ningyuan, Nathan Schine, Alexandros Georgakopoulos, Albert Ryou, Ariel Sommer, and Jonathan Simon (2017), “A strongly interacting polaritonic quantum dot,” arXiv preprint arXiv:1705.07475.
- Jia, Ningyuan, Nathan Schine, Alexandros Georgakopoulos, Albert Ryou, Ariel Sommer, and Jonathan Simon (2018), “Photons and polaritons in a broken-time-reversal nonplanar resonator,” *Phys. Rev. A* **97**, 013802.
- Jia, Shu, and Jason W Fleischer (2009), “Nonlinear light propagation in rotating waveguide arrays,” *Physical Review A* **79** (4), 041804.
- Jin, Dafei, Ling Lu, Zhong Wang, Chen Fang, John D Joannopoulos, Marin Soljačić, Liang Fu, and Nicholas X Fang (2016), “Topological magnetoplasmon,” *Nature communications* **7**, 13486.
- Jitomirskaya, Svetlana, and CA Marx (2012), “Analytic quasi-periodic cocycles with singularities and the lyapunov exponent of extended harpers model,” *Communications in mathematical physics* **316** (1), 237–267.
- Joannopoulos, John D, Steven G Johnson, Joshua N Winn, and Robert D Meade (2011), *Photonic crystals: molding the flow of light* (Princeton university press).
- John, Sajeew (1987), “Strong localization of photons in certain disordered dielectric superlattices,” *Phys. Rev. Lett.* **58**, 2486–2489.
- Jörg, Christina, Fabian Letscher, Michael Fleischhauer, and Georg von Freymann (2017), “Dynamic defects in photonic floquet topological insulators,” arXiv:1704.00533.
- Jotzu, G, M. Messer, R. Desbuquois, M. Lebrat, T. Uehlinger, D. Greif, and T. Esslinger (2014), “Experimental realization of the topological haldane model with ultracold fermions,” *Nature* **515**, 237.
- de Juan, Fernando, Mauricio Sturla, and María A. H. Vozmediano (2012), “Space Dependent Fermi Velocity in Strained Graphene,” *Physical Review Letters* **108** (22), 227205.
- Jünemann, J, A. Piga, S.-J. Ran, M. Lewenstein, M. Rizzi, and A. Bermudez (2017), “Exploring interacting topological insulators with ultracold atoms: The synthetic creutz-hubbard model,” *Phys. Rev. X* **7**, 031057.
- Kane, C L, and E. J. Mele (1997), “Size, Shape, and Low Energy Electronic Structure of Carbon Nanotubes,” *Physical Review Letters* **78** (10), 1932–1935.
- Kane, C L, and E. J. Mele (2005a), “Quantum spin hall effect in graphene,” *Phys. Rev. Lett.* **95**, 226801.
- Kane, C L, and E. J. Mele (2005b), “ Z_2 topological order and the quantum spin hall effect,” *Phys. Rev. Lett.* **95**, 146802.
- Kane, CL, and TC Lubensky (2014), “Topological boundary modes in isostatic lattices,” *Nature Physics* **10**, 39–45.
- Kang, Yuhao, Xiang Ni, Xiaojun Cheng, Alexander B Khanikaev, and Azriel Z Genack (2018), “Pseudo-spin-valley coupled edge states in a photonic topological insulator,” *Nature communications* **9** (1), 3029.
- Kapit, Eliot, Mohammad Hafezi, and Steven H. Simon (2014), “Induced self-stabilization in fractional quantum hall states of light,” *Phys. Rev. X* **4**, 031039.
- Karplus, Robert, and J. M. Luttinger (1954), “Hall effect in ferromagnetics,” *Phys. Rev.* **95**, 1154–1160.
- Karplus, Robert, and Maurice Neuman (1951), “The scattering of light by light,” *Phys. Rev.* **83**, 776–784.
- Kartashov, Yaroslav V, and Dmitry V Skryabin (2016), “Modulational instability and solitary waves in polariton topological insulators,” *Optica* **3** (11), 1228–1236.
- Karzig, Torsten, Charles-Edouard Bardyn, Netanel H. Lindner, and Gil Refael (2015), “Topological polaritons,” *Phys. Rev. X* **5**, 031001.
- Katan, Yaniv Tenenbaum, Rivka Bekenstein, Miguel A Bandres, Yaakov Lumer, Plotnik Yonatan, and Mordechai Segev (2016), “Induction of topological transport by long ranged nonlinearity,” in *CLEO: QELS-Fundamental Science* (Optical Society of America) pp. FM3A–6.
- Kavokin, Alexey, Guillaume Malpuech, and Mikhail Glazov (2005), “Optical Spin Hall Effect,” *Phys. Rev. Lett.* **95** (13), 136601.
- Kawabata, Kohei, Yuto Ashida, Hosho Katsura, and Masahito Ueda (2018a), “Parity-time-symmetric topological superconductor,” arXiv preprint arXiv:1801.00499.
- Kawabata, Kohei, Sho Higashikawa, Zongping Gong, Yuto Ashida, and Masahito Ueda (2018b), “Topological unification of time-reversal and particle-hole symmetries in non-hermitian physics,” arXiv preprint arXiv:1804.04676.
- Kawabata, Kohei, Ken Shiozaki, and Masahito Ueda (2018c), “Non-hermitian chern insulator,” arXiv preprint arXiv:1805.09632.
- Kawakami, Takuto, and Xiao Hu (2016), “Symmetry-guaranteed and accidental nodal-line semimetals in fcc lattice,” arXiv preprint arXiv:1611.07342.
- Kazimierczuk, Tomasz, Dietmar Fröhlich, Stefan Scheel, Heinrich Stolz, and Manfred Bayer (2014), “Giant rydberg excitons in the copper oxide Cu_2O ,” *Nature* **514** (7522), 343–347.
- Ke, Yongguan, Xizhou Qin, Feng Mei, Honghua Zhong, Yuri S Kivshar, and Chaohong Lee (2016), “Topological phase transitions and thouless pumping of light in photonic waveguide arrays,” *Laser & Photonics Reviews* **10** (6), 995–1001.
- Keil, Robert, Julia M Zeuner, Felix Dreisow, Matthias Heinrich, Andreas Tünnermann, Stefan Nolte, and Alexander Szameit (2013), “The random mass dirac model and long-range correlations on an integrated optical platform,” *Nature communications* **4**, 1368.
- Ketoja, Jukka A, and Indubala I Satija (1997), “The re-entrant phase diagram of the generalized harper equation,” *Journal of Physics: Condensed Matter* **9** (5), 1123.
- Khanikaev, Alexander B, and Andrea Alù (2015), “Optical isolators: nonlinear dynamic reciprocity,” *Nature Photonics* **9** (6), 359–361.
- Khanikaev, Alexander B, S Hossein Mousavi, Wang-Kong Tse, Mehdi Kargarian, Allan H MacDonald, and Gennady Shvets (2013), “Photonic topological insulators,” *Nature materials* **12** (3), 233–239.
- Khanikaev, Alexander B, and Gennady Shvets (2017), “Two-dimensional topological photonics,” *Nature Photonics* **11** (12), 763.
- Kim, Seunghwi, Xunngong Xu, Jacob M. Taylor, and Gaurav Bahl (2017), “Dynamically induced robust phonon trans-

- port and chiral cooling in an optomechanical system,” *Nature Communications* **8** (1), 205.
- Kiss, T, J. Janszky, and P. Adam (1994), “Time evolution of harmonic oscillators with time-dependent parameters: A step-function approximation,” *Phys. Rev. A* **49**, 4935–4942.
- Kitaev, Alexei, Vladimir Lebedev, and Mikhail Feigelman (2009), “Periodic table for topological insulators and superconductors,” *AIP Conference Proceedings*, **1134**, 22–30.
- Kitagawa, T, E. Berg, M. Rudner, and E. Demler (2010a), “Topological characterization of periodically-driven quantum systems,” *Phys. Rev. B* **82**, 235114.
- Kitagawa, Takuya (2012), “Topological phenomena in quantum walks: elementary introduction to the physics of topological phases,” *Quantum Information Processing*, 1–42.
- Kitagawa, Takuya, Matthew A Broome, Alessandro Fedrizzi, Mark S Rudner, Erez Berg, Ivan Kassal, Alán Aspuru-Guzik, Eugene Demler, and Andrew G White (2012), “Observation of topologically protected bound states in photonic quantum walks,” *Nature communications* **3**, 882.
- Kitagawa, Takuya, Takashi Oka, Arne Brataas, Liang Fu, and Eugene Demler (2011), “Transport properties of nonequilibrium systems under the application of light: Photoinduced quantum hall insulators without landau levels,” *Phys. Rev. B* **84**, 235108.
- Kitagawa, Takuya, Mark S. Rudner, Erez Berg, and Eugene Demler (2010b), “Exploring topological phases with quantum walks,” *Phys. Rev. A* **82**, 033429.
- Kivshar, Yuri S, and Govind Agrawal (2003), *Optical solitons: from fibers to photonic crystals* (Academic press).
- Klaers, Jan, Julian Schmitt, Frank Vewinger, and Martin Weitz (2010), “Bose-einstein condensation of photons in an optical microcavity,” *NATURE* **468** (7323), 545–548.
- Klein, DJ (1994), “Graphitic polymer strips with edge states,” *Chemical Physics Letters* **217** (3), 261–265.
- Klembt, S, T. H. Harder, O. A. Egorov, K. Winkler, R. Ge, M. A. Bandres, M. Emmerling, L. Worschech, T. C. H. Liew, M. Segev, C. Schneider, and S. Höfling (2018), “Exciton-polariton topological insulator,” [arXiv:1808.03179](https://arxiv.org/abs/1808.03179) [arXiv:1808.03179](https://arxiv.org/abs/1808.03179).
- Klitzing, K v, G. Dorda, and M. Pepper (1980), “New method for high-accuracy determination of the fine-structure constant based on quantized hall resistance,” *Phys. Rev. Lett.* **45**, 494–497.
- von Klitzing, Klaus (1986), “The quantized hall effect,” *Rev. Mod. Phys.* **58**, 519–531.
- Koch, Jens, Andrew A. Houck, Karyn Le Hur, and S. M. Girvin (2010), “Time-reversal-symmetry breaking in circuit-qed-based photon lattices,” *Phys. Rev. A* **82**, 043811.
- Kohmoto, Mahito (1985), “Topological invariant and the quantization of the hall conductance,” *Annals of Physics* **160** (2), 343 – 354.
- Kohmoto, Mahito, Bertrand I. Halperin, and Yong-Shi Wu (1992), “Diophantine equation for the three-dimensional quantum hall effect,” *Phys. Rev. B* **45**, 13488–13493.
- Kolkowitz, S, SL Bromley, T Bothwell, ML Wall, GE Marti, AP Koller, X Zhang, AM Rey, and J Ye (2017), “Spin-orbit-coupled fermions in an optical lattice clock,” *Nature* **542** (7639), 66–70.
- Kolovsky, A R (2011), “Creating artificial magnetic fields for cold atoms by photon-assisted tunneling,” *EPL* **93**, 20003.
- König, Markus, Steffen Wiedmann, Christoph Brüne, Andreas Roth, Hartmut Buhmann, Laurens W. Molenkamp, Xiao-Liang Qi, and Shou-Cheng Zhang (2007), “Quantum spin hall insulator state in hgte quantum wells,” *Science* **318** (5851), 766–770.
- Koshino, Mikito, Hideo Aoki, and Bertrand I. Halperin (2002), “Wrapping current versus bulk integer quantum hall effect in three dimensions,” *Phys. Rev. B* **66**, 081301.
- Kouwenhoven, L P, A. T. Johnson, N. C. van der Vaart, C. J. P. M. Harmans, and C. T. Foxon (1991), “Quantized current in a quantum-dot turnstile using oscillating tunnel barriers,” *Phys. Rev. Lett.* **67**, 1626–1629.
- Kozii, Vladyslav, and Liang Fu (2017), “Non-hermitian topological theory of finite-lifetime quasiparticles: Prediction of bulk fermi arc due to exceptional point,” [arXiv preprint arXiv:1708.05841](https://arxiv.org/abs/1708.05841).
- Kraus, Yaacov E, Yoav Lahini, Zohar Ringel, Mor Verbin, and Oded Zilberberg (2012), “Topological states and adiabatic pumping in quasicrystals,” *Phys. Rev. Lett.* **109**, 106402.
- Kraus, Yaacov E, Zohar Ringel, and Oded Zilberberg (2013), “Four-dimensional quantum hall effect in a two-dimensional quasicrystal,” *Phys. Rev. Lett.* **111**, 226401.
- Kraus, Yaacov E, and Oded Zilberberg (2012), “Topological equivalence between the fibonacci quasicrystal and the harper model,” *Phys. Rev. Lett.* **109**, 116404.
- Krimer, Dmitry O, and Ramaz Khomeriki (2011), “Realization of discrete quantum billiards in a two-dimensional optical lattice,” *Phys. Rev. A* **84**, 041807.
- Kruk, Sergey, Alexey Slobozhanyuk, Denitza Denkova, Alexander Poddubny, Ivan Kravchenko, Andrey Miroschnichenko, Dragomir Neshev, and Yuri Kivshar (2017), “Edge states and topological phase transitions in chains of dielectric nanoparticles,” *Small* **13** (11), 1603190.
- Kuhl, U, and H.-J. Stöckmann (1998), “Microwave realization of the hofstadter butterfly,” *Phys. Rev. Lett.* **80**, 3232–3235.
- Kusudo, Kenichiro, Na Young Kim, Andreas Löffler, Sven Höfling, Alfred Forchel, and Yoshihisa Yamamoto (2013), “Stochastic formation of polariton condensates in two degenerate orbital states,” *Physical Review B* **87** (21), 214503.
- Lahini, Yoav, Assaf Avidan, Francesca Pozzi, Marc Sorel, Roberto Morandotti, Demetrios N Christodoulides, and Yaron Silberberg (2008), “Anderson localization and nonlinearity in one-dimensional disordered photonic lattices,” *Physical Review Letters* **100** (1), 013906.
- Lai, Kueifu, Tsuhsuang Ma, Xiao Bo, Steven Anlage, and Gennady Shvets (2016), “Experimental realization of a reflections-free compact delay line based on a photonic topological insulator,” *Scientific reports* **6**, 28453.
- Lai, Y, and H. A. Haus (1989a), “Quantum theory of solitons in optical fibers. i. time-dependent hartree approximation,” *Phys. Rev. A* **40**, 844–853.
- Lai, Y, and H. A. Haus (1989b), “Quantum theory of solitons in optical fibers. ii. exact solution,” *Phys. Rev. A* **40**, 854–866.
- Lang, C, D. Bozyigit, C. Eichler, L. Steffen, J. M. Fink, A. A. Abdumalikov, Jr., M. Baur, S. Filipp, M. P. da Silva, A. Blais, and A. Wallraff (2011), “Observation of resonant photon blockade at microwave frequencies using correlation function measurements,” *Phys. Rev. Lett.* **106** (24), 10.1103/PhysRevLett.106.243601.
- Larré, Pierre-Élie, and Iacopo Carusotto (2015), “Propagation of a quantum fluid of light in a cavityless nonlinear optical medium: General theory and response to quantum

- quenches,” *Physical Review A* **92** (4), 043802.
- Larré, Pierre-Élie, and Iacopo Carusotto (2016), “Prethermalization in a quenched one-dimensional quantum fluid of light. intrinsic limits to the coherent propagation of a light beam in a nonlinear optical fiber,” *European Physical Journal D* **70**, 45.
- Laughlin, R B (1981), “Quantized hall conductivity in two dimensions,” *Phys. Rev. B* **23**, 5632–5633.
- Laughlin, R B (1983), “Anomalous quantum hall effect: An incompressible quantum fluid with fractionally charged excitations,” *Phys. Rev. Lett.* **50**, 1395–1398.
- Łącki, Mateusz, Hannes Pichler, Antoine Sterdyniak, Andreas Lyras, Vassilis E. Lembessis, Omar Al-Dossary, Jan Carl Budich, and Peter Zoller (2016), “Quantum hall physics with cold atoms in cylindrical optical lattices,” *Phys. Rev. A* **93**, 013604.
- Le Feber, B, N Rotenberg, and L Kuipers (2015), “Nanophotonic control of circular dipole emission,” *Nature communications* **6**, 6695.
- Lebreuilly, José, Alberto Biella, Florent Storme, Davide Rossini, Rosario Fazio, Cristiano Ciuti, and Iacopo Carusotto (2017), “Stabilizing strongly correlated photon fluids with non-markovian reservoirs,” *Phys. Rev. A* **96**, 033828.
- Lebreuilly, José, Michiel Wouters, and Iacopo Carusotto (2016), “Towards strongly correlated photons in arrays of dissipative nonlinear cavities under a frequency-dependent incoherent pumping,” *Comptes Rendus Physique* **17** (8), 836–860.
- Lee, Ching Hua, Stefan Imhof, Christian Berger, Florian Bayer, Johannes Brehm, Laurens W Molenkamp, Tobias Kiessling, and Ronny Thomale (2018), “Topoelectrical circuits,” *Communications Physics* **1** (1), 39.
- Lee, Seung-Yeol, Il-Min Lee, Junghyun Park, Sewoong Oh, Wooyoung Lee, Kyoung-Youm Kim, and ByoungHo Lee (2012), “Role of magnetic induction currents in nanoslit excitation of surface plasmon polaritons,” *Phys. Rev. Lett.* **108**, 213907.
- Lee, Tony E (2016), “Anomalous edge state in a non-hermitian lattice,” *Physical review letters* **116** (13), 133903.
- Lee-Thorp, James P, Iva Vukićević, Xinan Xu, Jinghui Yang, Charles L Fefferman, Chee Wei Wong, and Michael I Weinstein (2016), “Photonic realization of topologically protected bound states in domain-wall waveguide arrays,” *Physical Review A* **93** (3), 033822.
- Letscher, Fabian, Fabian Grusdt, and Michael Fleischhauer (2015), “Growing quantum states with topological order,” *Physical Review B* **91** (18), 184302.
- Levi, Liad, Mikael Rechtsman, Barak Freedman, Tal Schwartz, Ofer Manela, and Mordechai Segev (2011), “Disorder-enhanced transport in photonic quasicrystals,” *Science* **332** (6037), 1541–1544.
- Leyder, C, M. Romanelli, J. Ph. Karr, E. Giacobino, T. C. H. Liew, M. M. Glazov, A. V. Kavokin, G. Malpuech, and A. Bramati (2007), “Observation of the optical spin Hall effect,” *Nature Physics* **3** (9), 628–631.
- Leykam, Daniel, Konstantin Y Bliokh, Chunli Huang, YD Chong, and Franco Nori (2017a), “Edge modes, degeneracies, and topological numbers in non-hermitian systems,” *Physical Review Letters* **118** (4), 040401.
- Leykam, Daniel, and Y. D. Chong (2016), “Edge solitons in nonlinear-photonic topological insulators,” *Phys. Rev. Lett.* **117**, 143901.
- Leykam, Daniel, Sergej Flach, and YD Chong (2017b), “Flat bands in lattices with non-hermitian coupling,” arXiv:1704.00896.
- Leykam, Daniel, M. C. Rechtsman, and Y. D. Chong (2016), “Anomalous topological phases and unpaired dirac cones in photonic floquet topological insulators,” *Phys. Rev. Lett.* **117**, 013902.
- Li, Fei-Fei, Hai-Xiao Wang, Zhan Xiong, Qun Lou, Ping Chen, Rui-Xin Wu, Yin Poo, Jian-Hua Jiang, and Sajeed John (2018), “Topological light-trapping on a dislocation,” *Nature Communications* **9** (1), 2462.
- Li, Qing-Bo, Zhen Li, and Rui-xin Wu (2015a), “Bending self-collimated one-way light by using gyromagnetic photonic crystals,” *Applied Physics Letters* **107** (24), 241907.
- Li, Qing-Bo, Rui-Xin Wu, Yan Yang, and Hui-Ling Sun (2013), “Modeling 2d gyromagnetic photonic crystals by modified fdtd method,” *Chinese Physics Letters* **30** (7), 074208.
- Li, Yun, Pinaki Sengupta, George G. Batrouni, Christian Miniatura, and Benoît Grémaud (2015b), “Berry curvature of interacting bosons in a honeycomb lattice,” *Phys. Rev. A* **92**, 043605.
- Li, Zhaofeng, Koray Aydin, and Ekmel Ozbay (2009), “Determination of the effective constitutive parameters of bianisotropic metamaterials from reflection and transmission coefficients,” *Phys. Rev. E* **79**, 026610.
- Li, Zhen, Rui-xin Wu, Qing-bo Li, and Yin Poo (2015c), “Realization of self-guided unidirectional waveguides by a chain of gyromagnetic rods,” *Applied optics* **54** (6), 1267–1272.
- Li, Zhen, Rui-xin Wu, Yin Poo, Qing-bo Li, Rong-juan Liu, and Zhi-yuan Li (2014a), “An experimental study on the bandwidth and tunability of msp-based one-way transmission,” *Applied Physics A* **117** (2), 451–454.
- Li, Zhen, Rui-xin Wu, Yin Poo, Zhi-fang Lin, and Qing-Bo Li (2014b), “Fusing electromagnetic one-way edge states to achieve broadband unidirectional wave transmission,” *Journal of Optics* **16** (12), 125004.
- Li, Zhi-Yuan, Rong-Juan Liu, Lin Gan, Jin-Xin Fu, and Jin Lian (2014c), “Nonreciprocal electromagnetic devices in gyromagnetic photonic crystals,” *International Journal of Modern Physics B* **28** (02), 1441010.
- Lian, Jin, Jin-Xin Fu, Lin Gan, and Zhi-Yuan Li (2012a), “Experimental realization of a magnetically tunable cavity in a gyromagnetic photonic crystal,” *Chinese Physics Letters* **29** (7), 074208.
- Lian, Jin, Jin-Xin Fu, Lin Gan, and Zhi-Yuan Li (2012b), “Robust and disorder-immune magnetically tunable one-way waveguides in a gyromagnetic photonic crystal,” *Physical Review B* **85** (12), 125108.
- Lin, Jun Yu, Nai Chao Hu, You Jian Chen, Ching Hua Lee, and Xiao Zhang (2017), “Line nodes, dirac points, and lifshitz transition in two-dimensional nonsymmorphic photonic crystals,” *Phys. Rev. B* **96**, 075438.
- Lin, Q, X.-Q. Sun, M. Xiao, S.-C. Zhang, and S. Fan (2018), “Constructing three-dimensional photonic topological insulator using two-dimensional ring resonator lattice with a synthetic frequency dimension,” ArXiv e-prints arXiv:1802.02597 [physics.optics].
- Lin, Qian, and Shanhui Fan (2014), “Light guiding by effective gauge field for photons,” *Phys. Rev. X* **4**, 031031.
- Lin, Qian, and Shanhui Fan (2015), “Resonator-free realization of effective magnetic field for photons,” *New Journal of Physics* **17** (7), 075008.
- Lin, Qian, Meng Xiao, Luqi Yuan, and Shanhui Fan (2016),

- “Photonic weyl point in a two-dimensional resonator lattice with a synthetic frequency dimension,” *Nature communications* **7**, 13731.
- Lin, Shawn-Yu, Edmund Chow, Vince Hietala, Pierre R Villeneuve, and JD Joannopoulos (1998), “Experimental demonstration of guiding and bending of electromagnetic waves in a photonic crystal,” *Science* **282** (5387), 274–276.
- Lindner, N H, G. Refael, and V. Galitski (2011), “Floquet topological insulator in semiconductor quantum wells,” *Nat. Phys.* **7**, 490–495.
- Ling, CW, Meng Xiao, CT Chan, SF Yu, and Kin Hung Fung (2015), “Topological edge plasmon modes between diatomic chains of plasmonic nanoparticles,” *Optics express* **23** (3), 2021–2031.
- Lira, Hugo, Zongfu Yu, Shanhui Fan, and Michal Lipson (2012), “Electrically driven nonreciprocity induced by interband photonic transition on a silicon chip,” *Phys. Rev. Lett.* **109**, 033901.
- Liu, Changxu, Wenlong Gao, Biao Yang, and Shuang Zhang (2017), “Disorder-induced topological state transition in photonic metamaterials,” *Phys. Rev. Lett.* **119**, 183901.
- Liu, Kexin, Linfang Shen, and Sailing He (2012), “One-way edge mode in a gyromagnetic photonic crystal slab,” *Optics Letters* **37** (19), 4110–4112.
- Liu, Shiyang, Wanli Lu, Zhifang Lin, and ST Chui (2010), “Magnetically controllable unidirectional electromagnetic waveguiding devices designed with metamaterials,” *Applied Physics Letters* **97** (20), 201113.
- Liu, Yongmin, and Xiang Zhang (2011), “Metamaterials: a new frontier of science and technology,” *Chemical Society Reviews* **40** (5), 2494–2507.
- Livi, L F, G. Cappelini, M. Diem, L. Franchi, C. Clivati, M. Frittelli, F. Levi, D. Calonico, J. Catani, M. Inguscio, and L. Fallani (2016), “Synthetic dimensions and spin-orbit coupling with an optical clock transition,” *Phys. Rev. Lett.* **117**, 220401.
- Lodahl, Peter, Sahand Mahmoodian, Søren Stobbe, Arno Rauschenbeutel, Philipp Schneeweiss, Jürgen Volz, Hannes Pichler, and Peter Zoller (2017), “Chiral quantum optics,” *Nature* **541** (7638), 473–480.
- Lohse, Michael, Christian Schweizer, Hannah M. Price, Oded Zilberberg, and Immanuel Bloch (2018), “Exploring 4d quantum hall physics with a 2d topological charge pump,” *Nature* **553**, 55.
- Lohse, Michael, Christian Schweizer, Oded Zilberberg, Monika Aidelsburger, and Immanuel Bloch (2016), “A thouless quantum pump with ultracold bosonic atoms in an optical superlattice,” *Nature Physics* **12** (4), 350.
- Longhi, Stefano (2011), “Photonic bloch oscillations of correlated particles,” *Opt. Lett.* **36** (16), 3248–3250.
- Longhi, Stefano (2013), “Effective magnetic fields for photons in waveguide and coupled resonator lattices,” *Opt. Lett.* **38** (18), 3570–3573.
- Longhi, Stefano (2015), “Synthetic gauge fields for light beams in optical resonators,” *Optics Letters* **40** (13), 2941–2944.
- Lu, H-I, M. Schemmer, L. M. Aycock, D. Genkina, S. Sugawa, and I. B. Spielman (2016a), “Geometrical pumping with a bose-einstein condensate,” *Phys. Rev. Lett.* **116**, 200402.
- Lu, Jie, Linfang Shen, Xiaohua Deng, Xiaoler Li, and Xi-aodong Zheng (2013a), “Impact of photonic crystal boundary shape on the existence of one-way edge mode,” *Applied optics* **52** (21), 5216–5220.
- Lu, Ling, Chen Fang, Liang Fu, Steven G Johnson, John D Joannopoulos, and Marin Soljačić (2016b), “Symmetry-protected topological photonic crystal in three dimensions,” *Nature Physics* **12** (4), 337–340.
- Lu, Ling, Liang Fu, John D Joannopoulos, and Marin Soljačić (2013b), “Weyl points and line nodes in gyroid photonic crystals,” *Nature photonics* **7** (4), 294–299.
- Lu, Ling, John D Joannopoulos, and Marin Soljačić (2014), “Topological photonics,” *Nature Photonics* **8** (11), 821–829.
- Lu, Ling, John D Joannopoulos, and Marin Soljačić (2016c), “Topological states in photonic systems,” *Nature Physics* **12** (7), 626–629.
- Lu, Ling, Zhiyu Wang, Dexin Ye, Lixin Ran, Liang Fu, John D Joannopoulos, and Marin Soljačić (2015), “Experimental observation of weyl points,” *Science* **349** (6248), 622–624.
- Lu, Ling, and Zhong Wang (2016), “Topological one-way fiber of second chern number,” *arXiv preprint arXiv:1611.01998* .
- Lu, Yuehui, Ningyuan Jia, Lin Su, Clai Owens, Gediminas Juzeliūnas, David I Schuster, and Jonathan Simon (2018), “Probing the berry curvature and fermi arcs of a weyl circuit,” *arXiv preprint arXiv:1807.05243*.
- Lumer, Yaakov, Miguel A Bandres, Matthias Heinrich, Lukas Maczewsky, Hanan Herzog-Sheinfux, Alexander Szameit, and Mordechai Segev (2018), “Observation of light guiding by artificial gauge fields,” *arXiv preprint arXiv:1808.10207* .
- Lumer, Yaakov, Yonatan Plotnik, Mikael C Rechtsman, and Mordechai Segev (2013a), “Nonlinearly induced p t transition in photonic systems,” *Physical review letters* **111** (26), 263901.
- Lumer, Yaakov, Yonatan Plotnik, Mikael C Rechtsman, and Mordechai Segev (2013b), “Self-localized states in photonic topological insulators,” *Physical review letters* **111** (24), 243905.
- Lumer, Yaakov, Mikael C. Rechtsman, Yonatan Plotnik, and Mordechai Segev (2016), “Instability of bosonic topological edge states in the presence of interactions,” *Phys. Rev. A* **94**, 021801.
- Luo, Kaifa, Rui Yu, and Hongming Weng (2018), “Topological nodal states in circuit lattice,” *arXiv preprint arXiv:1801.05581* .
- Luo, Xi-Wang, Xingxiang Zhou, Chuan-Feng Li, Jin-Shi Xu, Guang-Can Guo, and Zheng-Wei Zhou (2015), “Quantum simulation of 2d topological physics in a 1d array of optical cavities,” *Nature communications* **6**, 7704.
- Luo, Xi-Wang, Xingxiang Zhou, Jin-Shi Xu, Chuan-Feng Li, Guang-Can Guo, Chuanwei Zhang, and Zheng-Wei Zhou (2017), “Synthetic-lattice enabled all-optical devices based on orbital angular momentum of light,” *Nature Communications* **8**, ncomms16097.
- Luo, Xiaoguang, Ming Zhou, Jingfeng Liu, Teng Qiu, and Zongfu Yu (2016), “Magneto-optical metamaterials with extraordinarily strong magneto-optical effect,” *Applied Physics Letters* **108** (13), 131104.
- Lustig, Eran, Steffen Weimann, Yonatan Plotnik, Yaakov Lumer, Miguel A Bandres, Alexander Szameit, and Mordechai Segev (2018), “Photonic topological insulator in synthetic dimensions,” *arXiv preprint arXiv:1807.01983*.
- Luttinger, J M (1951), “The effect of a magnetic field on electrons in a periodic potential,” *Phys. Rev.* **84**, 814–817.
- Ma, Ruichao, Clai Owens, Aman LaChapelle, David I. Schuster, and Jonathan Simon (2017), “Hamiltonian tomogra-

- phy of photonic lattices,” *Phys. Rev. A* **95**, 062120.
- Ma, Tzuhsuan, Alexander B. Khanikaev, S. Hossein Mousavi, and Gennady Shvets (2015), “Guiding electromagnetic waves around sharp corners: Topologically protected photonic transport in metawaveguides,” *Phys. Rev. Lett.* **114**, 127401.
- Ma, Tzuhsuan, and Gennady Shvets (2016), “All-Si valley-Hall photonic topological insulator,” *New Journal of Physics* **18** (2), 025012.
- MacDonald, A H, and P. Středa (1984), “Quantized hall effect and edge currents,” *Phys. Rev. B* **29**, 1616–1619.
- Maczewsky, Lukas J, Julia M Zeuner, Stefan Nolte, and Alexander Szameit (2017), “Observation of photonic anomalous floquet topological insulators,” *Nature communications* **8**, 13756.
- Maffei, Maria, Alexandre Dauphin, Filippo Cardano, Maciej Lewenstein, and Pietro Massignan (2018), “Topological characterization of chiral models through their long time dynamics,” *New Journal of Physics* **20** (1), 013023.
- Maghrebi, M F, M. J. Gullans, P. Bienias, S. Choi, I. Martin, O. Firstenberg, M. D. Lukin, H. P. Büchler, and A. V. Gorshkov (2015a), “Coulomb bound states of strongly interacting photons,” *Phys. Rev. Lett.* **115**, 123601.
- Maghrebi, Mohammad F, Norman Y. Yao, Mohammad Hafezi, Thomas Pohl, Ofer Firstenberg, and Alexey V. Gorshkov (2015b), “Fractional quantum hall states of rydberg polaritons,” *Phys. Rev. A* **91**, 033838.
- Makris, Konstantinos G, Ramy El-Ganainy, Demetrios N Christodoulides, and Ziad H Musslimani (2008), “Beam dynamics in p t symmetric optical lattices,” *Physical Review Letters* **100** (10), 103904.
- Malkova, Natalia, Ivan Hromada, Xiaosheng Wang, Garnett Bryant, and Zhigang Chen (2009), “Observation of optical shockley-like surface states in photonic superlattices,” *Optics letters* **34** (11), 1633–1635.
- Malzard, Simon, Charles Poli, and Henning Schomerus (2015), “Topologically protected defect states in open photonic systems with non-hermitian charge-conjugation and parity-time symmetry,” *Physical review letters* **115** (20), 200402.
- Malzard, Simon, and Henning Schomerus (2018a), “Bulk and edge-state arcs in non-hermitian coupled-resonator arrays,” arXiv preprint arXiv:1805.08161.
- Malzard, Simon, and Henning Schomerus (2018b), “Nonlinear mode competition and symmetry-protected power oscillations in topological lasers,” *New Journal of Physics* **20**, 063044.
- Mancini, M, G. Pagano, G. Cappellini, L. Livi, M. Rider, J. Catani, C. Sias, P. Zoller, M. Inguscio, M. Dalmonte, and L. Fallani (2015), “Observation of chiral edge states with neutral fermions in synthetic hall ribbons,” *Science* **349** (6255), 1510–1513.
- Mansha, Shampy, and YD Chong (2017), “Robust edge states in amorphous gyromagnetic photonic lattices,” arXiv preprint arXiv:1707.02687.
- Marqués, Ricardo, Francisco Medina, and Rachid Rafii-El-Idrissi (2002), “Role of bianisotropy in negative permeability and left-handed metamaterials,” *Phys. Rev. B* **65**, 144440.
- Mechelen, Todd Van, and Zubin Jacob (2016), “Universal spin-momentum locking of evanescent waves,” *Optica* **3** (2), 118–126.
- Mei, Feng, Zheng-Yuan Xue, Dan-Wei Zhang, Lin Tian, Chaohong Lee, and Shi-Liang Zhu (2016), “Witnessing topological weyl semimetal phase in a minimal circuit-qed lattice,” *Quantum Science and Technology* **1** (1), 015006.
- Mei, Feng, Jia-Bin You, Wei Nie, Rosario Fazio, Shi-Liang Zhu, and L. C. Kwek (2015), “Simulation and detection of photonic chern insulators in a one-dimensional circuit-qed lattice,” *Phys. Rev. A* **92**, 041805.
- Meier, Eric J, Fangzhao Alex An, Alexandre Dauphin, Maria Maffei, Pietro Massignan, Taylor L Hughes, and Bryce Gadway (2018), “Observation of the topological anderson insulator in disordered atomic wires,” arXiv preprint arXiv:1802.02109 .
- Mekis, Attila, JC Chen, I Kurland, Shanhui Fan, Pierre R Villeneuve, and JD Joannopoulos (1996), “High transmission through sharp bends in photonic crystal waveguides,” *Physical Review Letters* **77** (18), 3787.
- Meng, Guowu (2003), “Geometric construction of the quantum hall effect in all even dimensions,” *Journal of Physics A: Mathematical and General* **36** (36), 9415.
- Metelmann, A, and A. A. Clerk (2015), “Nonreciprocal photon transmission and amplification via reservoir engineering,” *Phys. Rev. X* **5**, 021025.
- Mikami, T, S. Kitamura, K. Yasuda, N. Tsuji, T. Oka, and H. Aoki (2016), “Brillouin-wigner theory for high-frequency expansion in periodically driven systems: Application to floquet topological insulators,” *Phys. Rev. B* **93**, 144307.
- Milićević, M, G Montambaux, T Ozawa, I Sagnes, A Lemaître, L Le Gratiet, A Harouri, J Bloch, and A Amo (2018), “Tilted and type-iii dirac cones emerging from flat bands in photonic orbital graphene,” arXiv preprint arXiv:1807.08650 .
- Milićević, M, T Ozawa, P Andreakou, I Carusotto, T Jacqmin, E Galopin, A Lemaître, L Le Gratiet, I Sagnes, J Bloch, *et al.* (2015), “Edge states in polariton honeycomb lattices,” *2D Materials* **2** (3), 034012.
- Milićević, M, T. Ozawa, G. Montambaux, I. Carusotto, E. Galopin, A. Lemaître, L. Le Gratiet, I. Sagnes, J. Bloch, and A. Amo (2017), “Orbital Edge States in a Photonic Honeycomb Lattice,” *Physical Review Letters* **118** (10), 107403.
- Milman, Pérola, Yvan Castin, and Luiz Davidovich (2000), “Decoherence as phase diffusion,” *Phys. Rev. A* **61**, 063803.
- Minkov, Momchil, and Vincenzo Savona (2016), “Haldane quantum hall effect for light in a dynamically modulated array of resonators,” *Optica* **3** (2), 200–206.
- Mirek, R, M. Król, K. Lekenta, J.-G. Rousset, M. Nawrocki, M. Kulczykowski, M. Matuszewski, J. Szczytko, W. Pascuski, and B. Piętko (2017), “Angular dependence of giant Zeeman effect for semimagnetic cavity polaritons,” *Physical Review B* **95** (8), 085429.
- Mitsch, R, C Sayrin, B Albrecht, P Schneeweiss, and A Rauschenbeutel (2014), “Quantum state-controlled directional spontaneous emission of photons into a nanophotonic waveguide,” *Nature communications* **5**, 5713.
- Mittal, S, J. Fan, S. Faez, A. Migdall, J. M. Taylor, and M. Hafezi (2014), “Topologically robust transport of photons in a synthetic gauge field,” *Phys. Rev. Lett.* **113**, 087403.
- Mittal, Sunil, Sriram Ganeshan, Jingyun Fan, Abolhassan Vaezi, and Mohammad Hafezi (2016a), “Measurement of topological invariants in a 2d photonic system,” *Nature Photonics* **10** (3), 180.
- Mittal, Sunil, and Mohammad Hafezi (2017), “Topologically robust generation of correlated photon pairs,” arXiv preprint arXiv:1709.09984.
- Mittal, Sunil, Venkata Vikram Orre, and Mohammad Hafezi

- (2016b), “Topologically robust transport of entangled photons in a 2d photonic system,” *Opt. Express* **24** (14), 15631–15641.
- Miyake, H, G. A. Siviloglou, C. J. Kennedy, W. C. Burton, and W. Ketterle (2013), “Realizing the harper hamiltonian with laser-assisted tunneling in optical lattices,” *Phys. Rev. Lett.* **111**, 185302.
- Moitra, Parikshit, Yuanmu Yang, Zachary Anderson, Ivan I Kravchenko, Dayl P Briggs, and Jason Valentine (2013), “Realization of an all-dielectric zero-index optical metamaterial,” *Nature Photonics* **7** (10), 791–795.
- Mondragon-Shem, Ian, Taylor L Hughes, Juntao Song, and Emil Prodan (2014), “Topological Criticality in the Chiral-Symmetric AIII Class at Strong Disorder,” *Phys. Rev. Lett.* **113** (4), 46802.
- Mong, Roger S K, and Vasudha Shivamoggi (2011), “Edge states and the bulk-boundary correspondence in Dirac Hamiltonians,” *Physical Review B* **83** (12), 125109.
- Montambaux, G, and M. Kohmoto (1990), “Quantized hall effect in three dimensions,” *Phys. Rev. B* **41**, 11417–11421.
- Montambaux, G, F. Piéchon, J.-N. Fuchs, and M. O. Goerbig (2009), “Merging of Dirac points in a two-dimensional crystal,” *Physical Review B* **80** (15), 153412.
- Moore, Gregory, and Nicholas Read (1991), “Nonabelions in the fractional quantum hall effect,” *Nuclear Physics B* **360** (2), 362 – 396.
- Moore, J E, and L. Balents (2007), “Topological invariants of time-reversal-invariant band structures,” *Phys. Rev. B* **75**, 121306.
- Moos, Matthias, Michael Höning, Razmik Unanyan, and Michael Fleischhauer (2015), “Many-body physics of rydberg dark-state polaritons in the strongly interacting regime,” *Phys. Rev. A* **92**, 053846.
- Möttönen, Mikko, Juha J. Vartiainen, and Jukka P. Pekola (2008), “Experimental determination of the berry phase in a superconducting charge pump,” *Phys. Rev. Lett.* **100**, 177201.
- Mukherjee, Seababrata, Harikumar K Chandrasekharan, Patrik Öhberg, Nathan Goldman, and Robert R Thomson (2017a), “State-recycling and time-resolved imaging in topological photonic lattices,” [arXiv preprint arXiv:1712.08145](https://arxiv.org/abs/1712.08145).
- Mukherjee, Seababrata, Alexander Spracklen, Debadipta Choudhury, Nathan Goldman, Patrik Öhberg, Erika Andersson, and Robert R Thomson (2015), “Modulation-assisted tunneling in laser-fabricated photonic wannier-stark ladders,” *New Journal of Physics* **17** (11), 115002.
- Mukherjee, Seababrata, Alexander Spracklen, Manuel Valiente, Erika Andersson, Patrik Öhberg, Nathan Goldman, and Robert R Thomson (2017b), “Experimental observation of anomalous topological edge modes in a slowly driven photonic lattice,” *Nature communications* **8**, 13918.
- Mukherjee, Seababrata, Manuel Valiente, Nathan Goldman, Alexander Spracklen, Erika Andersson, Patrik Öhberg, and Robert R. Thomson (2016), “Observation of pair tunneling and coherent destruction of tunneling in arrays of optical waveguides,” *Phys. Rev. A* **94**, 053853.
- Nagaosa, Naoto, Jairo Sinova, Shigeki Onoda, A. H. MacDonald, and N. P. Ong (2010), “Anomalous hall effect,” *Rev. Mod. Phys.* **82**, 1539–1592.
- Nakada, Kyoko, Mitsutaka Fujita, Gene Dresselhaus, and Mildred Dresselhaus (1996), “Edge state in graphene ribbons: Nanometer size effect and edge shape dependence,” *Physical Review B* **54** (24), 17954–17961.
- Nakahara, Miki (2003), *Geometry, Topology and Physics, Second Edition* (CRC Press).
- Nakajima, Shuta, Takafumi Tomita, Shintaro Taie, Tomohiro Ichinose, Hideki Ozawa, Lei Wang, Matthias Troyer, and Yoshiro Takahashi (2016), “Topological Thouless pumping of ultracold fermions,” *Nature Physics* **12** (4), 296.
- Nalitov, A V, D. D. Solnyshkov, and G. Malpuech (2015), “Polariton \mathbb{Z} topological insulator,” *Phys. Rev. Lett.* **114**, 116401.
- Nathan, Frederik, and Mark S Rudner (2015), “Topological singularities and the general classification of Floquet Bloch systems,” *New J. Phys.* **17** (12), 1–22.
- Nayak, Chetan, Steven H. Simon, Ady Stern, Michael Freedman, and Sankar Das Sarma (2008), “Non-abelian anyons and topological quantum computation,” *Rev. Mod. Phys.* **80**, 1083–1159.
- Naz, Ehsan Saei Ghareh, Ion Cosma Fulga, Libo Ma, Oliver G Schmidt, and Jeroen van den Brink (2017), “Topological phase transition in a stretchable photonic crystal,” [arXiv preprint arXiv:1712.01678](https://arxiv.org/abs/1712.01678).
- Neto, AH Castro, F Guinea, Nuno MR Peres, Kostya S Novoselov, and Andre K Geim (2009), “The electronic properties of graphene,” *Reviews of modern physics* **81** (1), 109.
- Ningyuan, Jia, Alexandros Georgakopoulos, Albert Ryou, Nathan Schine, Ariel Sommer, and Jonathan Simon (2016), “Observation and characterization of cavity rydberg polaritons,” *Physical Review A* **93** (4), 041802.
- Ningyuan, Jia, Clai Owens, Ariel Sommer, David Schuster, and Jonathan Simon (2015), “Time-and site-resolved dynamics in a topological circuit,” *Physical Review X* **5** (2), 021031.
- Ningyuan, Jia, Nathan Schine, Alexandros Georgakopoulos, Albert Ryou, Ariel Sommer, and Jonathan Simon (2017), “Photons and polaritons in a time-reversal-broken non-planar resonator,” [arXiv preprint arXiv:1709.00021](https://arxiv.org/abs/1709.00021).
- Niu, Qian, D. J. Thouless, and Yong-Shi Wu (1985), “Quantized hall conductance as a topological invariant,” *Phys. Rev. B* **31**, 3372–3377.
- Niu, Qian, and DJ Thouless (1984), “Quantised adiabatic charge transport in the presence of substrate disorder and many-body interaction,” *Journal of Physics A: Mathematical and General* **17** (12), 2453.
- Nixon, Micha, Eitan Ronen, Asher A. Friesem, and Nir Davidson (2013), “Observing Geometric Frustration with Thousands of Coupled Lasers,” *Physical Review Letters* **110** (18), 184102.
- Noda, Susumu (2016), “Photonic-crystal cavities,” in *Optical Fiber Communication Conference* (Optical Society of America) p. Th1K.1.
- Noh, Changsuk, and Dimitris G Angelakis (2017), “Quantum simulations and many-body physics with light,” *Reports on Progress in Physics* **80** (1), 016401.
- Noh, Jiho, Wladimir A Benalcazar, Sheng Huang, Matthew J Collins, Kevin P Chen, Taylor L Hughes, and Mikael C Rechtsman (2018), “Topological protection of photonic mid-gap defect modes,” *Nature Photonics* , 1.
- Noh, Jiho, Sheng Huang, Kevin Chen, and Mikael C. Rechtsman (2017a), “Observation of Photonic Topological Valley-Hall Edge States,” [arXiv:1706.00059](https://arxiv.org/abs/1706.00059).
- Noh, Jiho, Sheng Huang, Daniel Leykam, YD Chong, Kevin P Chen, and Mikael C Rechtsman (2017b), “Experimental observation of optical weyl points and fermi arc-like surface states,” *Nature Physics* **13** (6), 611–617.

- Nunnenkamp, A, Jens Koch, and S M Girvin (2011), “Synthetic gauge fields and homodyne transmission in jaynes-cummings lattices,” *New Journal of Physics* **13** (9), 095008.
- Ochiai, Tetsuyuki (2015a), “Non-reciprocity and topology in optics: one-way road for light via surface magnon polariton,” *Science and technology of advanced materials* **16** (1), 014401.
- Ochiai, Tetsuyuki (2015b), “Time-reversal-violating photonic topological insulators with helical edge states,” *Journal of the Physical Society of Japan* **84** (5), 054401.
- Ochiai, Tetsuyuki (2016), “Floquet–weyl and floquet-topological-insulator phases in a stacked two-dimensional ring-network lattice,” *Journal of Physics: Condensed Matter* **28** (42), 425501.
- Ochiai, Tetsuyuki (2017), “Gapless surface states originating from accidentally degenerate quadratic band touching in a three-dimensional tetragonal photonic crystal,” *Phys. Rev. A* **96**, 043842.
- O’Connell, Aaron D, Max Hofheinz, Markus Ansmann, Radoslaw C Bialczak, Mike Lenander, Erik Lucero, Matthew Neeley, Daniel Sank, H Wang, M Weides, *et al.* (2010), “Quantum ground state and single-phonon control of a mechanical resonator,” *Nature* **464** (7289), 697–703.
- O’Connor, D, P Ginzburg, FJ Rodríguez-Fortuño, GA Wurtz, and AV Zayats (2014), “Spin-orbit coupling in surface plasmon scattering by nanostructures,” *Nature communications* **5**, 5327.
- Oka, Takashi, and Hideo Aoki (2009), “Photovoltaic hall effect in graphene,” *Phys. Rev. B* **79**, 081406.
- Onbasli, Mehmet C, Lukáš Beran, Martin Zahradník, Miroslav Kučera, Roman Antoš, Jan Mistrík, Gerald F Dionne, Martin Veis, and Caroline A Ross (2016), “Optical and magneto-optical behavior of cerium yttrium iron garnet thin films at wavelengths of 200–1770 nm,” *Scientific Reports* **6**, 23640.
- Otterbach, J, J Ruseckas, RG Unanyan, G Juzeliūnas, and M Fleischhauer (2010), “Effective magnetic fields for stationary light,” *Physical review letters* **104** (3), 033903.
- Owens, Clai, Aman LaChapelle, Brendan Saxberg, Brandon M. Anderson, Ruichao Ma, Jonathan Simon, and David I. Schuster (2018), “Quarter-flux hofstadter lattice in a qubit-compatible microwave cavity array,” *Phys. Rev. A* **97**, 013818.
- Ozawa, Tomoki (2018), “Steady-state hall response and quantum geometry of driven-dissipative lattices,” *Phys. Rev. B* **97**, 041108(R).
- Ozawa, Tomoki, Alberto Amo, Jacqueline Bloch, and Iacopo Carusotto (2017), “Klein tunneling in driven-dissipative photonic graphene,” *Phys. Rev. A* **96**, 013813.
- Ozawa, Tomoki, and Iacopo Carusotto (2014), “Anomalous and quantum hall effects in lossy photonic lattices,” *Phys. Rev. Lett.* **112**, 133902.
- Ozawa, Tomoki, and Iacopo Carusotto (2017), “Synthetic dimensions with magnetic fields and local interactions in photonic lattices,” *Phys. Rev. Lett.* **118**, 013601.
- Ozawa, Tomoki, Hannah M Price, and Iacopo Carusotto (2015), “Momentum-space harper-hofstadter model,” *Physical Review A* **92** (2), 023609.
- Ozawa, Tomoki, Hannah M Price, and Iacopo Carusotto (2016a), “Quantum hall effect in momentum space,” *Physical Review B* **93** (19), 195201.
- Ozawa, Tomoki, Hannah M. Price, Nathan Goldman, Oded Zilberberg, and Iacopo Carusotto (2016b), “Synthetic dimensions in integrated photonics: From optical isolation to four-dimensional quantum hall physics,” *Phys. Rev. A* **93**, 043827.
- Painter, Oskar, RK Lee, Axel Scherer, A Yariv, JD O’Brien, PD Dapkus, and I Kim (1999), “Two-dimensional photonic band-gap defect mode laser,” *Science* **284** (5421), 1819–1821.
- Pan, Mingsen, Han Zhao, Pei Miao, Stefano Longhi, and Liang Feng (2018), “Photonic zero mode in a non-hermitian photonic lattice,” *Nature communications* **9** (1), 1308.
- Pancharatnam, S (1956), “Generalized theory of interference, and its applications,” *Proceedings of the Indian Academy of Sciences - Section A* **44** (5), 247–262.
- Parto, Midya, Steffen Wittek, Hossein Hodaei, Gal Harari, Miguel A Bandres, Jinhan Ren, Mikael C Rechtsman, Mordechai Segev, Demetrios N Christodoulides, and Mercedeh Khajavikhan (2018), “Edge-Mode Lasing in 1D Topological Active Arrays,” *Physical Review Letters* **120** (11), 113901.
- Peano, Vittorio, Martin Houde, Christian Brendel, Florian Marquardt, and Aashish A Clerk (2016a), “Topological phase transitions and chiral inelastic transport induced by the squeezing of light,” *Nature communications* **7**, 10779.
- Peano, Vittorio, Martin Houde, Florian Marquardt, and Aashish A. Clerk (2016b), “Topological quantum fluctuations and traveling wave amplifiers,” *Phys. Rev. X* **6**, 041026.
- Peierls, R (1933), “Zur theorie des diamagnetismus von leitungselektronen,” *Z. Phys.* **80**, 763–791.
- Peleg, Or, Guy Bartal, Barak Freedman, Ofer Manela, Mordechai Segev, and Demetrios N Christodoulides (2007), “Conical Diffraction and Gap Solitons in Honeycomb Photonic Lattices,” *Phys. Rev. Lett.* **98** (10), 103901.
- Pelegri, G, J. Polo, A. Turpin, M. Lewenstein, J. Mompart, and V. Ahufinger (2017), “Single-atom edgelike states via quantum interference,” *Phys. Rev. A* **95**, 013614.
- Pendry, John B, A J. Holden, DJ Robbins, and WJ Stewart (1999), “Magnetism from conductors and enhanced nonlinear phenomena,” *IEEE transactions on microwave theory and techniques* **47** (11), 2075–2084.
- Peropadre, Borja, David Zueco, Friedrich Wulschner, Frank Deppe, Achim Marx, Rudolf Gross, and Juan José García-Ripoll (2013), “Tunable coupling engineering between superconducting resonators: From sidebands to effective gauge fields,” *Phys. Rev. B* **87**, 134504.
- Petersen, Jan, Jürgen Volz, and Arno Rauschenbeutel (2014), “Chiral nanophotonic waveguide interface based on spin-orbit interaction of light,” *Science* **346** (6205), 67–71.
- Petrescu, Alexandru, Andrew A. Houck, and Karyn Le Hur (2012), “Anomalous hall effects of light and chiral edge modes on the kagomé lattice,” *Phys. Rev. A* **86**, 053804.
- Petrosyan, David, Johannes Otterbach, and Michael Fleischhauer (2011), “Electromagnetically induced transparency with rydberg atoms,” *Phys. Rev. Lett.* **107**, 213601.
- Peyronel, Thibault, Ofer Firstenberg, Qi-Yu Liang, Sebastian Hofferberth, Alexey V Gorshkov, Thomas Pohl, Mikhail D Lukin, and Vladan Vuletić (2012), “Quantum nonlinear optics with single photons enabled by strongly interacting atoms,” *Nature* **488** (7409), 57–60.
- Phillips, John (1996), “Self-adjoint fredholm operators and spectral flow,” *Canadian Mathematical Bulletin* **39** (4), 460–467.
- Pilozzi, Laura, and Claudio Conti (2016), “Topological lasing in resonant photonic structures,” *Phys. Rev. B* **93**, 195317.

- Pitaevskii, L P, and S. Stringari (2016), *Bose Einstein condensation and superfluidity* (Clarendon Press, Oxford).
- Plekhanov, Kirill, Guillaume Roux, and Karyn Le Hur (2017), “Floquet engineering of haldane chern insulators and chiral bosonic phase transitions,” *Physical Review B* **95** (4), 045102.
- Plotnik, Y, MA Bandres, S Stützer, Y Lumer, MC Rechtsman, A Szameit, and M Segev (2016), “Analogue of rashba pseudo-spin-orbit coupling in photonic lattices by gauge field engineering,” *Physical Review B* **94** (2), 020301.
- Plotnik, Yonatan, Mikael C Rechtsman, Daohong Song, Matthias Heinrich, Julia M Zeuner, Stefan Nolte, Yaakov Lumer, Natalia Malkova, Jingjun Xu, Alexander Szameit, *et al.* (2014), “Observation of unconventional edge states in photonic graphene,” *Nature materials* **13** (1), 57–62.
- Poddubny, Alexander, Andrey Miroshnichenko, Alexey Slobozhanyuk, and Yuri Kivshar (2014), “Topological majorana states in zigzag chains of plasmonic nanoparticles,” *ACS Photonics* **1** (2), 101–105.
- Poli, Charles, Matthieu Bellec, Ulrich Kuhl, Fabrice Mortesagne, and Henning Schomerus (2015), “Selective enhancement of topologically induced interface states in a dielectric resonator chain,” *Nature communications* **6**, 6710.
- Polkovnikov, Anatoli, Krishnendu Sengupta, Alessandro Silva, and Mukund Vengalattore (2011), “Colloquium,” *Rev. Mod. Phys.* **83**, 863–883.
- Poo, Yin, Cheng He, Chao Xiao, Ming-Hui Lu, Rui-Xin Wu, and Yan-Feng Chen (2016), “Manipulating one-way space wave and its refraction by time-reversal and parity symmetry breaking,” *Scientific reports* **6**, 29380.
- Poo, Yin, Rui-xin Wu, Zhifang Lin, Yan Yang, and CT Chan (2011), “Experimental realization of self-guiding unidirectional electromagnetic edge states,” *Physical Review Letters* **106** (9), 93903.
- Poo, Yin, Rui-xin Wu, Shiyang Liu, Yan Yang, Zhifang Lin, and ST Chui (2012), “Experimental demonstration of surface morphology independent electromagnetic chiral edge states originated from magnetic plasmon resonance,” *Applied Physics Letters* **101** (8), 081912.
- Poshakinskiy, A V, and A. N. Poddubny (2017), “Phonononic crystals with a synthetic magnetic field for an acoustic diode,” *Phys. Rev. Lett.* **118**, 156801.
- Poshakinskiy, A V, A. N. Poddubny, L. Piloizzi, and E. L. Ivchenko (2014), “Radiative topological states in resonant photonic crystals,” *Phys. Rev. Lett.* **112**, 107403.
- Pothier, H, P. Lafarge, P.F. Orfila, C. Urbina, D. Esteve, and M.H. Devoret (1991), “Single electron pump fabricated with ultrasmall normal tunnel junctions,” *Physica B: Condensed Matter* **169** (1), 573 – 574.
- Pothier, H, P. Lafarge, C. Urbina, D. Esteve, and M. H. Devoret (1992), “Single-electron pump based on charging effects,” *EPL (Europhysics Letters)* **17** (3), 249.
- Prange, Richard E, Steven M. Girvin, Marvin E Cage, Kv Klitzing, AM Chang, F Duncan, M Haldane, RB Laughlin, AMM Pruisken, and DJ Thouless (1989), *The quantum Hall effect* (Springer).
- Price, H M (2018), “Four-dimensional topological lattices without gauge fields,” ArXiv e-prints [arXiv:1806.05263](https://arxiv.org/abs/1806.05263) [[cond-mat.quant-gas](https://arxiv.org/archive/cond-mat)].
- Price, H M, O. Zilberberg, T. Ozawa, I. Carusotto, and N. Goldman (2015), “Four-dimensional quantum hall effect with ultracold atoms,” *Phys. Rev. Lett.* **115**, 195303.
- Price, Hannah M, Tomoki Ozawa, and Iacopo Carusotto (2014), “Quantum mechanics with a momentum-space artificial magnetic field,” *Phys. Rev. Lett.* **113**, 190403.
- Price, Hannah M, Tomoki Ozawa, and Nathan Goldman (2017), “Synthetic dimensions for cold atoms from shaking a harmonic trap,” *Phys. Rev. A* **95**, 023607.
- Price, Hannah M, Oded Zilberberg, Tomoki Ozawa, Iacopo Carusotto, and Nathan Goldman (2016), “Measurement of chern numbers through center-of-mass responses,” *Physical Review B* **93** (24), 245113.
- Price, HM, and NR Cooper (2012), “Mapping the berry curvature from semiclassical dynamics in optical lattices,” *Physical Review A* **85** (3), 033620.
- Prodan, Emil, and Hermann Schulz-Baldes (2016), “Bulk and boundary invariants for complex topological insulators,” K.
- Qi, Xiao-Liang, Taylor L. Hughes, and Shou-Cheng Zhang (2008), “Topological field theory of time-reversal invariant insulators,” *Phys. Rev. B* **78**, 195424.
- Qi, Xiao-Liang, Yong-Shi Wu, and Shou-Cheng Zhang (2006), “General theorem relating the bulk topological number to edge states in two-dimensional insulators,” *Phys. Rev. B* **74**, 045125.
- Qi, Xiao-Liang, and Shou-Cheng Zhang (2011), “Topological insulators and superconductors,” *Rev. Mod. Phys.* **83**, 1057–1110.
- Qiu, Pingping, Rui Liang, Weibin Qiu, Houbo Chen, Junbo Ren, Zhili Lin, Jia-Xian Wang, Qiang Kan, and Jiao-Qing Pan (2017), “Topologically protected edge states in graphene plasmonic crystals,” *Optics Express* **25** (19), 22587–22594.
- Qiu, W, Z Wang, and M Soljačić (2011), “Broadband circulators based on directional coupling of one-way waveguides.” *Optics express* **19** (22), 22248.
- Quelle, A, C. Weitenberg, K. Sengstock, and C. Morais Smith (2017), “Driving protocol for a Floquet topological phase without static counterpart,” ArXiv e-prints [arXiv:1704.00306](https://arxiv.org/abs/1704.00306) [[cond-mat.quant-gas](https://arxiv.org/archive/cond-mat)].
- Raghu, S, and F. D. M. Haldane (2008), “Analogues of quantum-hall-effect edge states in photonic crystals,” *Phys. Rev. A* **78**, 033834.
- Rahav, Saar, Ido Gilary, and Shmuel Fishman (2003), “Effective hamiltonians for periodically driven systems,” *Phys. Rev. A* **68**, 013820.
- Rasmussen, KØ, T Cretegny, PG Kevrekidis, and Niels Grønbech-Jensen (2000), “Statistical mechanics of a discrete nonlinear system,” *Physical review letters* **84** (17), 3740.
- Rechtsman, Mikael C, Yaakov Lumer, Yonatan Plotnik, Armando Perez-Leija, Alexander Szameit, and Mordechai Segev (2016), “Topological protection of photonic path entanglement,” *Optica* **3** (9), 925–930.
- Rechtsman, Mikael C, Yonatan Plotnik, Julia M. Zeuner, Daohong Song, Zhigang Chen, Alexander Szameit, and Mordechai Segev (2013a), “Topological creation and destruction of edge states in photonic graphene,” *Phys. Rev. Lett.* **111**, 103901.
- Rechtsman, Mikael C, Julia M Zeuner, Yonatan Plotnik, Yaakov Lumer, Daniel Podolsky, Felix Dreisow, Stefan Nolte, Mordechai Segev, and Alexander Szameit (2013b), “Photonic floquet topological insulators,” *Nature* **496** (7444), 196–200.
- Rechtsman, Mikael C, Julia M Zeuner, Andreas Tünnermann, Stefan Nolte, Mordechai Segev, and Alexander Szameit (2013c), “Strain-induced pseudomagnetic field and photonic landau levels in dielectric structures,” *Nature Photonics* **7** (2), 153–158.

- Reichl, Matthew D, and Erich J. Mueller (2014), “Floquet edge states with ultracold atoms,” *Phys. Rev. A* **89**, 063628.
- Reinhard, Andreas, Thomas Volz, Martin Winger, Antonio Badolato, Kevin J Hennessy, Evelyn L Hu, and Ataç Imamoglu (2012), “Strongly correlated photons on a chip,” *Nature Photonics* **6** (2), 93–96.
- Resta, Raffaele (1994), “Macroscopic polarization in crystalline dielectrics: the geometric phase approach,” *Rev. Mod. Phys.* **66**, 899–915.
- Resta, Raffaele (2011), “The insulating state of matter: a geometrical theory,” *The European Physical Journal B-Condensed Matter and Complex Systems* **79** (2), 121–137.
- Richard, M, J Kasprzak, R Romestain, R Andre, and LS Dang (2005), “Spontaneous coherent phase transition of polaritons in cde microcavities,” *Phys. Rev. Lett.* **94** (18), 10.1103/PhysRevLett.94.187401.
- Ringel, Matous, Mikhail Pletyukhov, and Vladimir Gritsev (2014), “Topologically protected strongly correlated states of photons,” *New Journal of Physics* **16** (11), 113030.
- Rodríguez-Fortuño, Francisco J, Giuseppe Marino, Pavel Ginzburg, Daniel O’Connor, Alejandro Martínez, Gregory A. Wurtz, and Anatoly V. Zayats (2013), “Near-field interference for the unidirectional excitation of electromagnetic guided modes,” *Science* **340** (6130), 328–330.
- Rosanov, N N (2002), *Spatial hysteresis and optical patterns* (Springer).
- Rosenthal, Eric I, Nicole K. Ehrlich, Mark S. Rudner, Andrew P. Higginbotham, and K. W. Lehnert (2018), “Topological phase transition measured in a dissipative metamaterial,” *Phys. Rev. B* **97**, 220301.
- Roushan, P, C Neill, J Tangpanitanon, VM Bastidas, A Megrant, R Barends, Y Chen, Z Chen, B Chiaro, A Dunsworth, *et al.* (2017a), “Spectroscopic signatures of localization with interacting photons in superconducting qubits,” *Science* **358** (6367), 1175–1179.
- Roushan, Pedram, Charles Neill, Anthony Megrant, Yu Chen, Ryan Babbush, Rami Barends, Brooks Campbell, Zijun Chen, Ben Chiaro, Andrew Dunsworth, *et al.* (2017b), “Chiral ground-state currents of interacting photons in a synthetic magnetic field,” *Nature Physics* **13** (2), 146.
- Roy, Dibyendu, C. M. Wilson, and Ofer Firstenberg (2017a), “Colloquium,” *Rev. Mod. Phys.* **89**, 021001.
- Roy, Rahul (2009), “Topological phases and the quantum spin hall effect in three dimensions,” *Phys. Rev. B* **79**, 195322.
- Roy, Sthitadhi, Michael Kolodrubetz, Nathan Goldman, and Adolfo G Grushin (2017b), “Tunable axial gauge fields in engineered weyl semimetals: Semiclassical analysis and optical lattice implementations,” [arXiv preprint arXiv:1708.08469](https://arxiv.org/abs/1708.08469).
- Rudner, Mark S, Netanel H Lindner, Erez Berg, and Michael Levin (2013), “Anomalous Edge States and the Bulk-Edge Correspondence for Periodically Driven Two-Dimensional Systems,” *Phys. Rev. X* **3** (3), 031005.
- Rudner, MS, and LS Levitov (2009), “Topological transition in a non-hermitian quantum walk,” *Physical review letters* **102** (6), 065703.
- Ruesink, Freek, Mohammad-Ali Miri, Andrea Alù, and Ewold Verhagen (2016), “Nonreciprocity and magnetic-free isolation based on optomechanical interactions,” *Nature Communications* **7**, 13662.
- Rüter, Christian E, Konstantinos G Makris, Ramy El-Ganainy, Demetrios N Christodoulides, Mordechai Segev, and Detlef Kip (2010), “Observation of parity–time symmetry in optics,” *Nature physics* **6** (3), 192–195.
- Ryu, Shinsei, and Yasuhiro Hatsugai (2002), “Topological origin of zero-energy edge states in particle-hole symmetric systems,” *Phys. Rev. Lett.* **89**, 077002.
- Ryu, Shinsei, Andreas P Schnyder, Akira Furusaki, and Andreas WW Ludwig (2010), “Topological insulators and superconductors: tenfold way and dimensional hierarchy,” *New Journal of Physics* **12**, 065010.
- Saffman, Mark, Thad G Walker, and Klaus Mølmer (2010), “Quantum information with rydberg atoms,” *Reviews of Modern Physics* **82** (3), 2313.
- Sala, VG, DD Solnyshkov, I Carusotto, T Jacqmin, A Lemaître, H Terças, A Nalitov, M Abbarchi, E Galopin, I Sagnes, *et al.* (2015), “Spin-orbit coupling for photons and polaritons in microstructures,” *Physical Review X* **5** (1), 011034.
- Saleh, BEA, and M.C. Teich (2007), *Fundamentals of Photonics*, Wiley Series in Pure and Applied Optics (Wiley).
- Salerno, Grazia, Tomoki Ozawa, Hannah M Price, and Iacopo Carusotto (2015), “How to directly observe landau levels in driven-dissipative strained honeycomb lattices,” *2D Materials* **2** (3), 034015.
- Salerno, Grazia, Tomoki Ozawa, Hannah M. Price, and Iacopo Carusotto (2016), “Floquet topological system based on frequency-modulated classical coupled harmonic oscillators,” *Phys. Rev. B* **93**, 085105.
- Salerno, Grazia, Tomoki Ozawa, Hannah M. Price, and Iacopo Carusotto (2017), “Propagating edge states in strained honeycomb lattices,” *Phys. Rev. B* **95**, 245418.
- Sayrin, Clément, Christian Junge, Rudolf Mitsch, Bernhard Albrecht, Danny O’Shea, Philipp Schneeweiss, Jürgen Volz, and Arno Rauschenbeutel (2015), “Nanophotonic optical isolator controlled by the internal state of cold atoms,” *Phys. Rev. X* **5**, 041036.
- Schine, Nathan, Albert Ryou, Andrey Gromov, Ariel Sommer, and Jonathan Simon (2016), “Synthetic landau levels for photons,” *Nature* **534**, 671–675.
- Schmidt, M, S. Kessler, V. Peano, O. Painter, and F. Marquardt (2015), “Optomechanical creation of magnetic fields for photons on a lattice,” *Optica* **2** (7), 635–641.
- Schnyder, Andreas P, Shinsei Ryu, Akira Furusaki, Andreas WW Ludwig, Vladimir Lebedev, and Mikhail Feigelman (2009), “Classification of topological insulators and superconductors,” *AIP Conference Proceedings*, **1134** (1), 10–21.
- Schoelkopf, RJ, and SM Girvin (2008), “Wiring up quantum systems,” *Nature* **451** (7179), 664–669.
- Schomerus, Henning (2013), “Topologically protected midgap states in complex photonic lattices,” *Optics letters* **38** (11), 1912–1914.
- Schomerus, Henning, and Nicole Yunger Halpern (2013), “Parity Anomaly and Landau-Level Lasing in Strained Photonic Honeycomb Lattices,” *Physical Review Letters* **110** (1), 013903.
- Schreiber, Andreas, Aurél Gábris, Peter P. Rohde, Kaisa Laiho, Martin Štefaňák, Václav Potoček, Craig Hamilton, Igor Jex, and Christine Silberhorn (2012), “A 2d quantum walk simulation of two-particle dynamics,” *Science* **336** (6077), 55–58, <http://science.sciencemag.org/content/336/6077/55.full.pdf>.
- Schwartz, Alon, and Baruch Fischer (2013), “Laser mode hyper-combs,” *Optics express* **21** (5), 6196–6204.
- Schwartz, Tal, Guy Bartal, Shmuel Fishman, and Mordechai Segev (2007), “Transport and anderson localization in

- disordered two-dimensional photonic lattices,” *Nature* **446** (7131), 52–55.
- Seclí, Matteo (2017), *Edge State Lasing in a 2D Topological Photonic System*, Master’s thesis (University of Trento).
- Segev, Mordechai, Bruno Crosignani, Amnon Yariv, and Baruch Fischer (1992), “Spatial solitons in photorefractive media,” *Physical Review Letters* **68** (7), 923.
- Serdyukov, A, Igor Semchenko, S Tertyakov, and A Sihvola (2001), *Electromagnetics of bi-anisotropic materials-Theory and Application*, Vol. 11 (Gordon and Breach Science Publishers).
- Shelby, R A, D. R. Smith, and S. Schultz (2001), “Experimental verification of a negative index of refraction,” *Science* **292** (5514), 77–79, <http://science.sciencemag.org/content/292/5514/77.full.pdf>.
- Shen, Huitao, Bo Zhen, and Liang Fu (2018), “Topological band theory for non-hermitian hamiltonians,” *Physical review letters* **120** (14), 146402.
- Shen, Linfang, Yun You, Zhuoyuan Wang, and Xiaohua Deng (2015), “Backscattering-immune one-way surface magnetoplasmons at terahertz frequencies,” *Optics express* **23** (2), 950–962.
- Shen, Zhen, Yan-Lei Zhang, Yuan Chen, Chang-Ling Zou, Yun-Feng Xiao, Xu-Bo Zou, Fang-Wen Sun, Guang-Can Guo, and Chun-Hua Dong (2016), “Experimental realization of optomechanically induced non-reciprocity,” *Nature Photonics* **10**, 657–661.
- Sheng, D N, Z. Y. Weng, L. Sheng, and F. D. M. Haldane (2006), “Quantum spin-hall effect and topologically invariant chern numbers,” *Phys. Rev. Lett.* **97**, 036808.
- Shi, T, HJ Kimble, and JI Cirac (2017), “Topological phenomena in classical optical networks,” *Proceedings of the National Academy of Sciences* **114**, E8967–E8976.
- Shi, Yu, Zongfu Yu, and Shanhui Fan (2015), “Limitations of nonlinear optical isolators due to dynamic reciprocity,” *Nature photonics* **9** (6), 388–392.
- Shindou, Ryuichi, Ryo Matsumoto, Shuichi Murakami, and Jun-ichiro Ohe (2013a), “Topological chiral magnonic edge mode in a magnonic crystal,” *Phys. Rev. B* **87**, 174427.
- Shindou, Ryuichi, and Jun-ichiro Ohe (2014), “Magnetostatic wave analog of integer quantum hall state in patterned magnetic films,” *Phys. Rev. B* **89**, 054412.
- Shindou, Ryuichi, Jun-ichiro Ohe, Ryo Matsumoto, Shuichi Murakami, and Eiji Saitoh (2013b), “Chiral spin-wave edge modes in dipolar magnetic thin films,” *Phys. Rev. B* **87**, 174402.
- Siegman, Anthony E (1986), *Lasers* (University Science Books).
- Silberberg, Yaron, Yoav Lahini, Yaron Bromberg, Eran Small, and Roberto Morandotti (2009), “Universal correlations in a nonlinear periodic 1d system,” *Physical review letters* **102** (23), 233904.
- Silveirinha, Mário G (2015), “Chern invariants for continuous media,” *Physical Review B* **92** (12), 125153.
- Silveirinha, Mário G (2016a), “Bulk-edge correspondence for topological photonic continua,” *Physical Review B* **94** (20), 205105.
- Silveirinha, Mário G (2016b), “ Z_2 topological index for continuous photonic materials,” *Physical Review B* **93** (7), 075110.
- Simon, Barry (1983), “Holonomy, the quantum adiabatic theorem, and berry’s phase,” *Phys. Rev. Lett.* **51**, 2167–2170.
- Sinev, Ivan S, Ivan S Mukhin, Alexey P Slobozhanyuk, Alexander N Poddubny, Andrey E Miroshnichenko, Anton K Samusev, and Yuri S Kivshar (2015), “Mapping plasmonic topological states at the nanoscale,” *Nanoscale* **7** (28), 11904–11908.
- Siroki, Gleb, Paloma A. Huidobro, and Vincenzo Giannini (2017), “Topological photonics: From crystals to particles,” *Physical Review B* **96** (4), 041408.
- Skirlo, Scott A, Ling Lu, Yuichi Igarashi, Qinghui Yan, John Joannopoulos, and Marin Soljačić (2015), “Experimental observation of large chern numbers in photonic crystals,” *Phys. Rev. Lett.* **115**, 253901.
- Skirlo, Scott A, Ling Lu, and Marin Soljačić (2014), “Multi-mode one-way waveguides of large chern numbers,” *Physical review letters* **113** (11), 113904.
- Slobozhanyuk, Alexey, S Hossein Mousavi, Xiang Ni, Daria Smirnova, Yuri S Kivshar, and Alexander B Khanikaev (2016a), “Three-dimensional all-dielectric photonic topological insulator,” *Nature Photonics* **11**, 130–136.
- Slobozhanyuk, Alexey, Alena V Shchelokova, Xiang Ni, S Hossein Mousavi, Daria A Smirnova, Pavel A Belov, Andrea Alù, Yuri S Kivshar, and Alexander B Khanikaev (2017), “Near-field imaging of spin-locked edge states in all-dielectric topological metasurfaces,” *arXiv preprint arXiv:1705.07841*.
- Slobozhanyuk, Alexey P, Alexander B Khanikaev, Dmitry S Filonov, Daria A Smirnova, Andrey E Miroshnichenko, and Yuri S Kivshar (2016b), “Experimental demonstration of topological effects in bianisotropic metamaterials,” *Scientific reports* **6**, 22270.
- Slobozhanyuk, Alexey P, Alexander N. Poddubny, Andrey E. Miroshnichenko, Pavel A. Belov, and Yuri S. Kivshar (2015), “Subwavelength topological edge states in optically resonant dielectric structures,” *Phys. Rev. Lett.* **114**, 123901.
- Slobozhanyuk, Alexey P, Alexander N Poddubny, Ivan S Sinev, Anton K Samusev, Ye Feng Yu, Arseniy I Kuznetsov, Andrey E Miroshnichenko, and Yuri S Kivshar (2016c), “Enhanced photonic spin hall effect with subwavelength topological edge states,” *Laser & Photonics Reviews* **10** (4), 656–664.
- Söllner, Immo, Sahand Mahmoodian, Sofie Lindskov Hansen, Leonardo Midolo, Alisa Javadi, Gabija Kiršanskė, Tommaso Pregolato, Haitham El-Ella, Eun Hye Lee, Jin Dong Song, *et al.* (2015), “Deterministic photon-emitter coupling in chiral photonic circuits,” *Nature nanotechnology* **10** (9), 775–778.
- Solnyshkov, D D, O Bleu, and G Malpuech (2018), “Topological optical isolator based on polariton graphene,” *Applied Physics Letters* **112** (3), 31106.
- Solnyshkov, D D, A. V. Nalitov, and G. Malpuech (2016a), “Kibble-zurek mechanism in topologically nontrivial zigzag chains of polariton micropillars,” *Phys. Rev. Lett.* **116**, 046402.
- Solnyshkov, Dmitry, Anton Nalitov, Berihu Teklu, Louis Franck, and Guillaume Malpuech (2016b), “Spin-dependent Klein tunneling in polariton graphene with photonic spin-orbit interaction,” *Physical Review B* **93** (8), 085404.
- Soluyanov, Alexey A, Dominik Gresch, Zhijun Wang, Quan-Sheng Wu, Matthias Troyer, Xi Dai, and B Andrei Bernevig (2015), “Type-ii weyl semimetals,” *Nature* **527** (7579), 495–498.
- Sommer, A, H. P. Büchler, and J. Simon (2015), “Quantum Crystals and Laughlin Droplets of Cavity Rydberg Polaritons,” *ArXiv e-prints arXiv:1506.00341 [cond-mat.quant-*

- gas].
- Sommer, Ariel, and Jonathan Simon (2016), “Engineering photonic floquet hamiltonians through fabry-pérot resonators,” *New Journal of Physics* **18** (3), 035008.
- Song, Daohong, Vassilis Paltoglou, Sheng Liu, Yi Zhu, Daniel Gallardo, Liqin Tang, Jingjun Xu, Mark Ablowitz, Nikolaos K Efremidis, and Zhigang Chen (2015), “Unveiling pseudospin and angular momentum in photonic graphene.” *Nat. Commun.* **6**, 6272.
- Spivak, B, F. Zhou, and M. T. Beal Monod (1995), “Mesoscopic mechanisms of the photovoltaic effect and microwave absorption in granular metals,” *Phys. Rev. B* **51**, 13226–13230.
- St-Jean, P, V Goblot, E Galopin, A Lemaître, T Ozawa, L Le Gratiet, I Sagnes, J Bloch, and A Amo (2017), “Lasing in topological edge states of a one-dimensional lattice,” *Nature Photonics* **11** (10), 651.
- Stannigel, K, Peter Komar, SJM Habraken, SD Bennett, Mikhail D Lukin, P Zoller, and P Rabl (2012), “Optomechanical quantum information processing with photons and phonons,” *Physical Review Letters* **109** (1), 013603.
- Störmer, HL, JP Eisenstein, AC Gossard, W Wiegmann, and K Baldwin (1986), “Quantization of the hall effect in an anisotropic three-dimensional electronic system,” *Physical review letters* **56** (1), 85.
- Stuhl, B K, H.-I. Lu, L. M. Aycock, D. Genkina, and I. B. Spielman (2015), “Visualizing edge states with an atomic bose gas in the quantum hall regime,” *Science* **349** (6255), 1514–1518.
- Stützer, Simon, Yonatan Plotnik, Yaakov Lumer, Paraj Titum, Netanel H Lindner, Mordechai Segev, Mikael C Rechtsman, and Alexander Szameit (2018), “Photonic topological anderson insulators,” *Nature* **560** (7719), 461.
- Su, W P, J. R. Schrieffer, and A. J. Heeger (1979), “Solitons in polyacetylene,” *Phys. Rev. Lett.* **42**, 1698–1701.
- Sugawa, Seiji, Francisco Salces-Carcoba, Abigail R Perry, Yuchen Yue, and Ian B Spielman (2016), “Observation of a non-abelian yang monopole: From new chern numbers to a topological transition,” [arXiv preprint arXiv:1610.06228](https://arxiv.org/abs/1610.06228).
- Sun, Bo Ye, Xi Wang Luo, Ming Gong, Guang Can Guo, and Zheng Wei Zhou (2017a), “Weyl semimetal phases and implementation in degenerate optical cavities,” *Phys. Rev. A* **96**, 013857.
- Sun, Xiao-Chen, Cheng He, Xiao-Ping Liu, Ming-Hui Lu, Shi-Ning Zhu, and Yan-Feng Chen (2017b), “Two-dimensional topological photonic systems,” *Progress in Quantum Electronics* **55**, 52–73.
- Sun, Yong, Daniel Leykam, Stephen Nenni, Daohong Song, Hong Chen, Y. D. Chong, and Zhigang Chen (2018), “Observation of valley landau-zener-bloch oscillations and pseudospin imbalance in photonic graphene,” *Phys. Rev. Lett.* **121**, 033904.
- Suszalski, Dominik, and Jakub Zakrzewski (2016), “Different lattice geometries with a synthetic dimension,” *Phys. Rev. A* **94**, 033602.
- Switkes, M, C. M. Marcus, K. Campman, and A. C. Gossard (1999), “An adiabatic quantum electron pump,” *Science* **283** (5409), 1905–1908, <http://science.sciencemag.org/content/283/5409/1905.full.pdf>.
- Szameit, Alexander, Felix Dreisow, Thomas Pertsch, Stefan Nolte, and Andreas Tünnermann (2007), “Control of directional evanescent coupling in fs laser written waveguides,” *Optics express* **15** (4), 1579–1587.
- Szameit, Alexander, and Stefan Nolte (2010), “Discrete optics in femtosecond-laser-written photonic structures,” *Journal of Physics B: Atomic, Molecular and Optical Physics* **43** (16), 163001.
- Szameit, Alexander, Mikael C Rechtsman, Omri Bahat-Treidel, and Mordechai Segev (2011), “P t-symmetry in honeycomb photonic lattices,” *Physical Review A* **84** (2), 021806.
- Taddia, Luca, Eyal Cornfeld, Davide Rossini, Leonardo Mazza, Eran Sela, and Rosario Fazio (2017), “Topological fractional pumping with alkaline-earth-like atoms in synthetic lattices,” *Phys. Rev. Lett.* **118**, 230402.
- Tai, M Eric, Alexander Lukin, Matthew Rispoli, Robert Schittko, Tim Menke, Dan Borgnia, Philipp M Preiss, Fabian Grusdt, Adam M Kaufman, and Markus Greiner (2017a), “Microscopy of the interacting harper-hofstadter model in the two-body limit,” *Nature* **546** (7659), 519.
- Tai, M Eric, Alexander Lukin, Matthew Rispoli, Robert Schittko, Tim Menke, Dan Borgnia, Philipp M Preiss, Fabian Grusdt, Adam M Kaufman, and Markus Greiner (2017b), “Microscopy of the interacting harper-hofstadter model in the two-body limit,” *Nature* **546** (7659), 519.
- Takemura, Naotomo, Stéphane Trebaol, Michiel Wouters, Marcia T Portella-Oberli, and Benoît Deveaud (2014), “Polaritonic feshbach resonance,” *Nature Physics* **10** (7), 500–504.
- Tan, Wei, Yong Sun, Hong Chen, and Shun-Qing Shen (2014), “Photonic simulation of topological excitations in metamaterials,” *Scientific reports* **4**, 3842.
- Tanese, D, H Flayac, D Solnyshkov, A Amo, A Lemaître, E Galopin, R Braive, P Senellart, I Sagnes, G Malpuech, *et al.* (2013), “Polariton condensation in solitonic gap states in a one-dimensional periodic potential,” *Nature communications* **4**, 1749.
- Tangpanitanon, Jirawat, Victor M. Bastidas, Sarah Al-Assam, Pedram Roushan, Dieter Jaksch, and Dimitris G. Angelakis (2016), “Topological pumping of photons in nonlinear resonator arrays,” *Phys. Rev. Lett.* **117**, 213603.
- Tarasinski, B, J. K. Asbóth, and J. P. Dahlhaus (2014), “Scattering theory of topological phases in discrete-time quantum walks,” *Phys. Rev. A* **89**, 042327.
- Tarruell, Leticia, Daniel Greif, Thomas Uehlinger, Gregor Jotzu, and Tilman Esslinger (2012), “Creating, moving and merging Dirac points with a Fermi gas in a tunable honeycomb lattice,” *Nature* **483** (7389), 302–305.
- Teo, Jeffrey C Y, and C. L. Kane (2010), “Topological defects and gapless modes in insulators and superconductors,” *Phys. Rev. B* **82**, 115120.
- Thouless, D J (1983), “Quantization of particle transport,” *Phys. Rev. B* **27**, 6083–6087.
- Thouless, D J, M. Kohmoto, M. P. Nightingale, and M. den Nijs (1982), “Quantized hall conductance in a two-dimensional periodic potential,” *Phys. Rev. Lett.* **49**, 405–408.
- Tomita, Akira, and Raymond Y. Chiao (1986), “Observation of berry’s topological phase by use of an optical fiber,” *Phys. Rev. Lett.* **57**, 937–940.
- Tomka, Michael, Mikhail Pletyukhov, and Vladimir Gritsev (2015), “Supersymmetry in quantum optics and in spin-orbit coupled systems,” *Scientific reports* **5**, 13097.
- Tong, D (2016), “Lectures on the quantum hall effect,” *ArXiv e-prints arXiv:1606.06687 [hep-th]*.
- Tsomokos, Dimitris I, Sahel Ashhab, and Franco Nori (2010), “Using superconducting qubit circuits to engineer exotic

- lattice systems,” *Phys. Rev. A* **82**, 052311.
- Tsui, D C, H. L. Stormer, and A. C. Gossard (1982), “Two-dimensional magnetotransport in the extreme quantum limit,” *Phys. Rev. Lett.* **48**, 1559–1562.
- Türeci, Hakan E, A. Douglas Stone, and Li Ge (2007), “Theory of the spatial structure of nonlinear lasing modes,” *Phys. Rev. A* **76**, 013813.
- Umucalilar, R O, and I. Carusotto (2011), “Artificial gauge field for photons in coupled cavity arrays,” *Phys. Rev. A* **84**, 043804.
- Umucalilar, R O, and I. Carusotto (2012), “Fractional quantum hall states of photons in an array of dissipative coupled cavities,” *Phys. Rev. Lett.* **108**, 206809.
- Umucalilar, R O, and I. Carusotto (2017), “Generation and spectroscopic signatures of a fractional quantum hall liquid of photons in an incoherently pumped optical cavity,” ArXiv e-prints [arXiv:1708.05441](https://arxiv.org/abs/1708.05441) [[cond-mat.quant-gas](https://arxiv.org/abs/1708.05441)].
- Umucalilar, RO, and I Carusotto (2013), “Many-body braiding phases in a rotating strongly correlated photon gas,” *Physics Letters A* **377** (34), 2074–2078.
- Verbin, Mor, Oded Zilberberg, Yaacov E. Kraus, Yoav Lahini, and Yaron Silberberg (2013), “Observation of topological phase transitions in photonic quasicrystals,” *Phys. Rev. Lett.* **110**, 076403.
- Verbin, Mor, Oded Zilberberg, Yoav Lahini, Yaacov E. Kraus, and Yaron Silberberg (2015), “Topological pumping over a photonic fibonacci quasicrystal,” *Phys. Rev. B* **91**, 064201.
- Vignolo, Patrizia, Matthieu Bellec, Julian Böhm, Abdoulaye Camara, Jean-Marc Gambaudo, Ulrich Kuhl, and Fabrice Mortessagne (2016), “Energy landscape in a penrose tiling,” *Physical Review B* **93** (7), 075141.
- Vocke, David, Thomas Roger, Francesco Marino, Ewan M Wright, Iacopo Carusotto, Matteo Clerici, and Daniele Faccio (2015), “Experimental characterization of nonlocal photon fluids,” *Optica* **2** (5), 484–490.
- Volovik, G E (2009), *The Universe in a Helium Droplet* (Oxford university press).
- Wallace, P R (1947), “The band theory of graphite,” *Phys. Rev.* **71**, 622–634.
- Walls, D F, and G. Milburn (2006), *Quantum Optics* (Springer Verlag, Berlin).
- Wan, Xiangang, Ari M Turner, Ashvin Vishwanath, and Sergey Y Savrasov (2011), “Topological semimetal and fermi-arc surface states in the electronic structure of pyrochlore iridates,” *Physical Review B* **83** (20), 205101.
- Wang, Danlu, Chengwei Qiu, Peter T Rakich, and Zheng Wang (2015a), “Guide-wave photonic pulling force using one-way photonic chiral edge states,” in *CLEO: QELS_Fundamental Science* (Optical Society of America) pp. FM2D–7.
- Wang, Hai-Xiao, Yige Chen, Zhi Hong Hang, Hae-Young Kee, and Jian-Hua Jiang (2017a), “Type-II Dirac photons,” arXiv preprint [arXiv:1703.09899](https://arxiv.org/abs/1703.09899).
- Wang, Hailong, Longwen Zhou, and YD Chong (2016a), “Floquet weyl phases in a three-dimensional network model,” *Physical Review B* **93** (14), 144114.
- Wang, HaiXiao, Lin Xu, HuanYang Chen, and Jian-Hua Jiang (2016b), “Three-dimensional photonic Dirac points stabilized by point group symmetry,” *Phys. Rev. B* **93**, 235155.
- Wang, Luyang, Shao-Kai Jian, and Hong Yao (2016c), “Topological photonic crystal with equifrequency weyl points,” *Physical Review A* **93** (6), 061801.
- Wang, Qiang, Meng Xiao, Hui Liu, Shining Zhu, and CT Chan (2017b), “Optical interface states protected by synthetic weyl points,” *Physical Review X* **7** (3), 031032.
- Wang, Yan-Pu, Wei Wang, Zheng-Yuan Xue, Wan-Li Yang, Yong Hu, and Ying Wu (2015b), “Realizing and characterizing chiral photon flow in a circuit quantum electrodynamics necklace,” *Scientific reports* **5**, 8352.
- Wang, Yan-pu, Wan-li Yang, Yong Hu, Zheng-yuan Xue, and Ying Wu (2016d), “Detecting topological phases of microwave photons in a circuit quantum electrodynamics lattice,” *NPJ Quantum Information* **2**, 16015.
- Wang, Zheng, Y. D. Chong, John D. Joannopoulos, and Marin Soljačić (2008), “Reflection-free one-way edge modes in a gyromagnetic photonic crystal,” *Phys. Rev. Lett.* **100**, 013905.
- Wang, Zheng, Yidong Chong, JD Joannopoulos, and Marin Soljačić (2009), “Observation of unidirectional backscattering-immune topological electromagnetic states,” *Nature* **461** (7265), 772–775.
- Wang, Zhuoyuan, Linfang Shen, Zaihe Yu, Xianmin Zhang, and Xiaodong Zheng (2013), “Highly efficient photonic-crystal splitters based on one-way waveguiding,” *JOSA B* **30** (1), 173–176.
- Wang, Zhuoyuan, Linfang Shen, Xianmin Zhang, Yigang Wang, Zaihe Yu, and Xiaodong Zheng (2011), “Photonic crystal cavity with one-way rotating state and its coupling with photonic crystal waveguide,” *Journal of Applied Physics* **110** (4), 043106.
- Waterstraat, Nils (2016), “Fredholm operators and spectral flow,” [arXiv:1603.02009](https://arxiv.org/abs/1603.02009).
- Weimann, S, M Kremer, Y Plotnik, Y Lumer, S Nolte, KG Makris, M Segev, MC Rechtsman, and A Szameit (2017), “Topologically protected bound states in photonic parity-time-symmetric crystals,” *Nature Materials* **16** (4), 433–438.
- Weinstein, C L Fefferman, J P Lee-Thorp, and M I (2016), “Bifurcations of edge states topologically protected and non-protected in continuous 2D honeycomb structures,” *2D Materials* **3** (1), 14008.
- Wen, XG, and A Zee (1992), “Shift and spin vector: New topological quantum numbers for the hall fluids,” *Physical review letters* **69** (6), 953.
- Wertz, E, L. Ferrier, D. D. Solnyshkov, R. Johne, D. Sanvitto, A. Lemaitre, I. Sagnes, R. Grousson, A. V. Kavokin, P. Senellart, G. Malpuech, and J. Bloch (2010), “Spontaneous formation and optical manipulation of extended polariton condensates,” *Nat. Phys.* **6** (11), 860–864.
- Wimmer, Martin, Hannah M Price, Iacopo Carusotto, and Ulf Peschel (2017), “Experimental measurement of the berry curvature from anomalous transport,” *Nature Physics* **13** (6), 545–550.
- Wouters, Michiel (2007), “Resonant polariton-polariton scattering in semiconductor microcavities,” *Phys. Rev. B* **76** (4), 10.1103/PhysRevB.76.045319.
- Wouters, Michiel, Iacopo Carusotto, and Cristiano Ciuti (2008), “Spatial and spectral shape of inhomogeneous nonequilibrium exciton-polariton condensates,” *Phys. Rev. B* **77**, 115340.
- Wu, Congjun, Doron Bergman, Leon Balents, and S. Das Sarma (2007), “Flat Bands and Wigner Crystallization in the Honeycomb Optical Lattice,” *Physical Review Letters* **99** (7), 070401.
- Wu, Long-Hua, and Xiao Hu (2015), “Scheme for achieving a topological photonic crystal by using dielectric material,” *Phys. Rev. Lett.* **114**, 223901.

- Wu, Xiaoxiao, Yan Meng, Jingxuan Tian, Yingzhou Huang, Hong Xiang, Dezhan Han, and Weijia Wen (2017a), “Direct observation of valley-polarized topological edge states in designer surface plasmon crystals,” *Nature communications* **8** (1), 1304.
- Wu, You, Chong Li, Xiaoyong Hu, Yutian Ao, Yifan Zhao, and Qihuang Gong (2017b), “Applications of topological photonics in integrated photonic devices,” *Advanced Optical Materials* **5**, 1700357.
- Xia, Bai-Zhan, Ting-Ting Liu, Guo-Liang Huang, Hong-Qing Dai, Jun-Rui Jiao, Xian-Guo Zang, De-Jie Yu, Sheng-Jie Zheng, and Jian Liu (2017), “Topological phononic insulator with robust pseudospin-dependent transport,” *Physical Review B* **96** (9), 094106.
- Xiao, Bo, Kueifu Lai, Yang Yu, Tzuhsuan Ma, Gennady Shvets, and Steven M. Anlage (2016a), “Exciting reflectionless unidirectional edge modes in a reciprocal photonic topological insulator medium,” *Phys. Rev. B* **94**, 195427.
- Xiao, Di, Ming-Che Chang, and Qian Niu (2010), “Berry phase effects on electronic properties,” *Rev. Mod. Phys.* **82**, 1959–2007.
- Xiao, L, X Zhan, ZH Bian, KK Wang, X Zhang, XP Wang, J Li, K Mochizuki, D Kim, N Kawakami, *et al.* (2017), “Observation of topological edge states in parity–time-symmetric quantum walks,” *Nature Physics* **13** (11), 1117.
- Xiao, Meng, and Shanhui Fan (2017a), “Photonic chern insulator through homogenization of an array of particles,” arXiv preprint arXiv:1707.03365.
- Xiao, Meng, and Shanhui Fan (2017b), “Topologically charged nodal surface,” arXiv preprint arXiv:1709.02363.
- Xiao, Meng, Qian Lin, and Shanhui Fan (2016b), “Hyperbolic weyl point in reciprocal chiral metamaterials,” *Physical review letters* **117** (5), 057401.
- Xiao, Meng, Z. Q. Zhang, and C. T. Chan (2014), “Surface impedance and bulk band geometric phases in one-dimensional systems,” *Phys. Rev. X* **4**, 021017.
- Xu, Gang, Hongming Weng, Zhijun Wang, Xi Dai, and Zhong Fang (2011), “Chern semimetal and the quantized anomalous hall effect in hgr 2 se 4,” *Physical review letters* **107** (18), 186806.
- Xu, Lin, Hai-Xiao Wang, Ya-Dong Xu, Huan-Yang Chen, and Jian-Hua Jiang (2016), “Accidental degeneracy in photonic bands and topological phase transitions in two-dimensional core-shell dielectric photonic crystals,” *Optics Express* **24** (16), 18059.
- Xu, Yong, Sheng-Tao Wang, and L-M Duan (2017), “Weyl exceptional rings in a three-dimensional dissipative cold atomic gas,” *Physical Review Letters* **118** (4), 045701.
- Yablonovitch, Eli (1987), “Inhibited spontaneous emission in solid-state physics and electronics,” *Phys. Rev. Lett.* **58**, 2059–2062.
- Yan, Qinghui, Rongjuan Liu, Zhongbo Yan, Boyuan Liu, Hongsheng Chen, Zhong Wang, and Ling Lu (2018), “Experimental discovery of nodal chains,” *Nature Physics* **14** (5), 461.
- Yan, Zhongbo, Ren Bi, Huitao Shen, Ling Lu, Shou-Cheng Zhang, and Zhong Wang (2017), “Nodal-link semimetals,” *Phys. Rev. B* **96**, 041103.
- Yang, Biao, Qinghua Guo, Ben Tremain, Lauren E Barr, Wenlong Gao, Hongchao Liu, Benjamin Béri, Yuanjiang Xiang, Dianyuan Fan, Alastair P Hibbins, *et al.* (2017a), “Direct observation of topological surface-state arcs in photonic metamaterials,” *Nature Communications* **8** (1), 97.
- Yang, Biao, Qinghua Guo, Ben Tremain, Rongjuan Liu, Lauren E Barr, Qinghui Yan, Wenlong Gao, Hongchao Liu, Yuanjiang Xiang, Jing Chen, *et al.* (2018), “Ideal weyl points and helicoid surface states in artificial photonic crystal structures,” *Science*, eaaq1221.
- Yang, Bing, Tong Wu, and Xiangdong Zhang (2017b), “Engineering topological edge states in two dimensional magnetic photonic crystal,” *Applied Physics Letters* **110** (2), 021109.
- Yang, Bing, Tong Wu, and Xiangdong Zhang (2017c), “Topological properties of nearly flat bands in two-dimensional photonic crystals,” *JOSA B* **34** (4), 831–836.
- Yang, W L, Zhang-qi Yin, Z. X. Chen, Su-Peng Kou, M. Feng, and C. H. Oh (2012), “Quantum simulation of an artificial abelian gauge field using nitrogen-vacancy-center ensembles coupled to superconducting resonators,” *Phys. Rev. A* **86**, 012307.
- Yang, Y, Z. Gao, H. Xue, L. Zhang, M. He, Z. Yang, R. Singh, Y. Chong, B. Zhang, and H. Chen (2018), “Realization of a three-dimensional photonic topological insulator,” ArXiv e-prints arXiv:1804.03595 [physics.class-ph].
- Yang, Yan, Yin Poo, Rui-xin Wu, Yan Gu, and Ping Chen (2013), “Experimental demonstration of one-way slow wave in waveguide involving gyromagnetic photonic crystals,” *Applied Physics Letters* **102**, 231113.
- Yang, Yuting, Yun Fei Xu, Tao Xu, Hai-Xiao Wang, Jian-Hua Jiang, Xiao Hu, and ZhiHong Hang (2018), “Visualization of a Unidirectional Electromagnetic Waveguide Using Topological Photonic Crystals Made of Dielectric Materials,” *Physical Review Letters* **120** (21), 217401.
- Yang, Zhaoju, Meng Xiao, Fei Gao, Ling Lu, Yidong Chong, and Baile Zhang (2017d), “Weyl points in a magnetic tetrahedral photonic crystal,” *Optics Express* **25** (14), 15772–15777.
- Yannopapas, Vassilios (2011), “Gapless surface states in a lattice of coupled cavities: a photonic analog of topological crystalline insulators,” *Physical Review B* **84** (19), 195126.
- Yannopapas, Vassilios (2014), “Dirac points, topological edge modes and nonreciprocal transmission in one-dimensional metamaterial-based coupled-cavity arrays,” *International Journal of Modern Physics B* **28** (02), 1441006.
- Yao, Shunyu, Fei Song, and Zhong Wang (2018), “Non-hermitian chern bands and chern numbers,” arXiv preprint arXiv:1804.04672.
- Yao, Shunyu, and Zhong Wang (2018), “Edge states and topological invariants of non-hermitian systems,” arXiv preprint arXiv:1803.01876.
- Yao, Shunyu, Zhongbo Yan, and Zhong Wang (2017), “Topological invariants of floquet systems: General formulation, special properties, and floquet topological defects,” *Phys. Rev. B* **96**, 195303.
- Yariv, Amnon (1976), *Introduction to optical electronics* (Holt, Rinehart and Winston, Inc., New York, NY).
- Yariv, Amnon, Yong Xu, Reginald K Lee, and Axel Scherer (1999), “Coupled-resonator optical waveguide: a proposal and analysis,” *Optics letters* **24** (11), 711–713.
- Yi, Kexin, and Torsten Karzig (2016), “Topological polaritons from photonic Dirac cones coupled to excitons in a magnetic field,” *Physical Review B* **93** (10), 104303.
- Yin, Chuanhao, Hui Jiang, Linhu Li, Rong Lü, and Shu Chen (2018), “Geometrical meaning of winding number and its characterization of topological phases in one-dimensional chiral non-hermitian systems,” *Physical Review A* **97** (5), 052115.
- Yoshida, Tsuneya, Robert Peters, and Norio Kawakami

- (2018), “Non-hermitian perspective of the band structure in heavy-fermion systems,” arXiv preprint arXiv:1805.01172.
- Yoshioka, Daijiro (2002), *The quantum Hall effect* (Springer).
- You, JQ, and Franco Nori (2011), “Atomic physics and quantum optics using superconducting circuits,” *Nature* **474** (7353), 589–597.
- Yu, Zaihe, Zhuoyuan Wang, Linfang Shen, and Xiaohua Deng (2014), “One-way electromagnetic mode at the surface of a magnetized gyromagnetic medium,” *Electronic Materials Letters* **10** (5), 969–973.
- Yu, Zongfu, and Shanhui Fan (2009), “Complete optical isolation created by indirect interband photonic transitions,” *Nature photonics* **3** (2), 91–94.
- Yu, Zongfu, Georgios Veronis, Zheng Wang, and Shanhui Fan (2008), “One-way electromagnetic waveguide formed at the interface between a plasmonic metal under a static magnetic field and a photonic crystal,” *Physical review letters* **100** (2), 23902.
- Yuan, Luqi, and Shanhui Fan (2015a), “Three-dimensional dynamic localization of light from a time-dependent effective gauge field for photons,” *Phys. Rev. Lett.* **114**, 243901.
- Yuan, Luqi, and Shanhui Fan (2015b), “Topologically non-trivial floquet band structure in a system undergoing photonic transitions in the ultrastrong-coupling regime,” *Phys. Rev. A* **92**, 053822.
- Yuan, Luqi, and Shanhui Fan (2016), “Bloch oscillation and unidirectional translation of frequency in a dynamically modulated ring resonator,” *Optica* **3** (9), 1014–1018.
- Yuan, Luqi, Yu Shi, and Shanhui Fan (2016a), “Photonic gauge potential in a system with a synthetic frequency dimension,” *Optics letters* **41** (4), 741–744.
- Yuan, Luqi, Da-wei Wang, and Shanhui Fan (2017a), “Synthetic gauge potential and effective magnetic field in a raman medium undergoing molecular modulation,” *Phys. Rev. A* **95**, 033801.
- Yuan, Luqi, Meng Xiao, and Shanhui Fan (2016b), “Time reversal of a wave packet with temporal modulation of gauge potential,” *Phys. Rev. B* **94**, 140303.
- Yuan, Luqi, Meng Xiao, Qian Lin, and Shanhui Fan (2017b), “Synthetic space with arbitrary dimensions in a few rings undergoing dynamic modulation,” arXiv preprint arXiv:1710.01373 .
- Yves, Simon, Romain Fleury, Thomas Berthelot, Mathias Fink, Fabrice Lemoult, and Geoffroy Lerosey (2017a), “Crystalline metamaterials for topological properties at subwavelength scales,” *Nature Communications* **8**, 16023.
- Yves, Simon, Romain Fleury, Fabrice Lemoult, Mathias Fink, and Geoffroy Lerosey (2017b), “Topological acoustic polaritons: robust sound manipulation at the subwavelength scale,” *New Journal of Physics* **19** (7), 075003.
- Zak, J (1964), “Magnetic translation group,” *Phys. Rev.* **134**, A1602–A1606.
- Zak, J (1989), “Berry’s phase for energy bands in solids,” *Phys. Rev. Lett.* **62**, 2747–2750.
- Zambon, Nicola Carlon, Philippe St-Jean, Marijana Milićević, Aristide Lemaitre, Abdelmounaim Harouri, Luc LeGratiet, Olivier Bleu, Dmitry Solnyshkov, Guillaume Malpuech, Isabelle Sagnes, *et al.* (2018), “Optically controlling the emission chirality of microlasers,” arXiv preprint arXiv:1806.04590 .
- Zang, Xiaofei, and Chun Jiang (2011), “Edge mode in non-reciprocal photonic crystal waveguide: manipulating the unidirectional electromagnetic pulse dynamically,” *JOSA B* **28** (3), 554–557.
- Zeng, Tian-Sheng, Ce Wang, and Hui Zhai (2015), “Charge pumping of interacting fermion atoms in the synthetic dimension,” *Phys. Rev. Lett.* **115**, 095302.
- Zeuner, Julia M, Mikael C. Rechtsman, Yonatan Plotnik, Yaakov Lumer, Stefan Nolte, Mark S. Rudner, Mordechai Segev, and Alexander Szameit (2015), “Observation of a topological transition in the bulk of a non-hermitian system,” *Phys. Rev. Lett.* **115**, 040402.
- Zhang, Fan, Jeil Jung, Gregory A Fiete, Qian Niu, and Allan H MacDonald (2011), “Spontaneous Quantum Hall States in Chirally Stacked Few-Layer Graphene Systems,” *Phys. Rev. Lett.* **106** (15), 156801.
- Zhang, Le, Dongxiao Yang, Kan Chen, Tao Li, and Song Xia (2013), “Design of nonreciprocal waveguide devices based on two-dimensional magneto-optical photonic crystals,” *Optics & Laser Technology* **50**, 195–201.
- Zhang, Shou-Cheng, and Jiangping Hu (2001), “A four-dimensional generalization of the quantum hall effect,” *Science* **294** (5543), 823–828.
- Zhang, Tiantian, Zhida Song, A. Alexandradinata, Hongming Weng, Chen Fang, Ling Lu, and Zhong Fang (2018), “Double-weyl phonons in transition-metal monosilicides,” *Phys. Rev. Lett.* **120**, 016401.
- Zhang, Xiaogang, Wei Li, and Xunya Jiang (2012), “Confined one-way mode at magnetic domain wall for broadband high-efficiency one-way waveguide, splitter and bend,” *Applied Physics Letters* **100** (4), 041108.
- Zhang, Z, F. Li, G. Malpuech, Y. Zhang, O. Bleu, S. Koniakhin, C. Li, Y. Zhang, M. Xiao, and D. Solnyshkov (2018), “Particle-like behavior of topological defects in linear wave packets in photonic graphene,” ArXiv e-prints arXiv:1806.05540 [physics.optics].
- Zhao, Han, Pei Miao, Mohammad H Teimourpour, Simon Malzard, Ramy El-Ganainy, Henning Schomerus, and Liang Feng (2018), “Topological hybrid silicon microlasers,” *Nature Communications* **9** (1), 981.
- Zhen, Bo, Chia Wei Hsu, Yuichi Igarashi, Ling Lu, Ido Kaminer, Adi Pick, Song-Liang Chua, John D Joannopoulos, and Marin Soljačić (2015), “Spawning rings of exceptional points out of dirac cones,” *Nature* **525** (7569), 354–358.
- Zheng, Wei, and Hui Zhai (2014), “Floquet topological states in shaking optical lattices,” *Phys. Rev. A* **89** (6), 061603.
- Zhou, F, B. Spivak, and B. Altshuler (1999), “Mesoscopic mechanism of adiabatic charge transport,” *Phys. Rev. Lett.* **82**, 608–611.
- Zhou, Hengyun, Chao Peng, Yoseob Yoon, Chia Wei Hsu, Keith A Nelson, Liang Fu, John D Joannopoulos, Marin Soljačić, and Bo Zhen (2018), “Observation of bulk fermi arc and polarization half charge from paired exceptional points,” *Science* , eaap9859.
- Zhou, Ming, Lei Ying, Ling Lu, Lei Shi, Jian Zi, and Zongfu Yu (2017a), “Electromagnetic scattering laws in weyl systems,” *Nature communications* **8** (1), 1388.
- Zhou, Xiang-Fa, Xi-Wang Luo, Su Wang, Guang-Can Guo, Xingxiang Zhou, Han Pu, and Zheng-Wei Zhou (2017b), “Dynamically manipulating topological physics and edge modes in a single degenerate optical cavity,” *Phys. Rev. Lett.* **118**, 083603.
- Zhou, Xin, You Wang, Daniel Leykam, and Y D Chong (2017c), “Optical isolation with nonlinear topological photonics,” *New Journal of Physics* **19** (9), 095002.
- Zhu, Bo, Honghua Zhong, Yongguan Ke, Xizhou Qin, An-

- drey A. Sukhorukov, Yuri S. Kivshar, and Chaohong Lee (2018a), “Topological floquet edge states in periodically curved waveguides,” *Phys. Rev. A* **98**, 013855.
- Zhu, Haibin, and Chun Jiang (2011), “Extraordinary coupling into one-way magneto-optical photonic crystal waveguide,” *Journal of Lightwave Technology* **29** (5), 708–713.
- Zhu, Xuan, Hai-Xiao Wang, Changqing Xu, Yun Lai, Jian-Hua Jiang, and Sajeev John (2018b), “Topological transitions in continuously deformed photonic crystals,” *Phys. Rev. B* **97** (8), 85148.
- Zilberberg, Oded, Sheng Huang, Jonathan Guglielmon, Mohan Wang, Kevin P Chen, Yaacov E Kraus, and Mikael C Rechtsman (2018), “Photonic topological boundary pumping as a probe of 4d quantum hall physics,” *Nature* **553** (7686), 59.